

Brief review related to the foundations of time-dependent density functional theory

Thomas A. Niehaus · Norman H. March

dedicated to Professor Sándor Suhai on the occasion of his 65th birthday

Received: date / Accepted: date

Abstract The electron density $n(\mathbf{r}, t)$, which is the central tool of time-dependent density functional theory, is presently considered to be derivable from a one-body time-dependent potential $V(\mathbf{r}, t)$, via one-electron wave functions satisfying a time-dependent Schrödinger equation. This is here related via a generalized equation of motion to a Dirac density matrix now involving t . Linear response theory is then surveyed, with a special emphasis on the question of causality with respect to the density dependence of the potential. Extraction of $V(\mathbf{r}, t)$ for solvable models is also proposed.

Keywords One-body time-dependent potential · Solvable Moshinsky atom · Time-dependent response function · Current-density functional

PACS 31.15.Ew · 31.25.-v · 31.70.Hq

1 Background

Early work on time-dependent density functional theory can be traced back at least to 1972 [1, 2, 3] followed by various studies [4, 5, 6, 7] in the 1970s / early 1980s, and culminating in the important proposal of Runge and Gross (RG) [8]. In essence, the RG argument generalizes the Hohenberg-Kohn theorem [9] to time-dependent external potentials. Though now known to be far from rigorous [10], the RG work is widely accepted as the basis for the assertion that, for a specified initial state, there is a unique correspondence between the time dependent density $n(\mathbf{r}, t)$ and the external potential $V_{\text{ext}}(\mathbf{r}, t)$. This points the way to construct a time-dependent extension $V(\mathbf{r}, t)$ of a static Slater-Kohn-Sham (SKS) like potential $V(\mathbf{r})$ [11, 12] which will then generate a Slater determinant of non-interacting electron wave functions, $\phi_i(\mathbf{r}, t)$ say, satisfying the time-dependent Schrödinger equation

$$\left(-\frac{\hbar}{2m} \nabla^2 + V(\mathbf{r}, t) \right) \phi_i(\mathbf{r}, t) = i\hbar \frac{\partial}{\partial t} \phi_i(\mathbf{r}, t), \quad (1)$$

T.A. Niehaus
Bremen Center for Computational Materials Science, D-28359 Bremen, Germany
E-mail: t.niehaus@bccms.uni-bremen.de

N.H. March
Dept. of Physics, University of Antwerp, Antwerp, Belgium
Oxford University, Oxford, England

with a specified determinant at time $t = 0$. Then the electron density $n(\mathbf{r}, t)$ introduced above is constructed, formally exactly for N electrons, as

$$n(\mathbf{r}, t) = \sum_i^N \phi_i^*(\mathbf{r}, t) \phi_i(\mathbf{r}, t). \quad (2)$$

Of course, formal exactitude requires precise knowledge of the one-body potential $V(\mathbf{r}, t)$ in eqn(1). At the time of writing, such knowledge is limited for the key exchange and correlation contributions entering V (see also eqn(9) below). These can be formally constructed [13, 14, 15], and perturbative approaches that converge on their exact form are also known [16, 17, 18]. In practical terms, however, accurate exchange-correlation functionals beyond the local density approximation in space and time are still elusive and remain a topic of general interest.

2 The challenge of Schirmer and Dreuw [10] to the RG arguments, and some responses

This is the point to return to the work of [10]. This study contains serious criticism levelled against the very foundations of TDDFT (articles [19, 20, 21] are also concerned with the basis of the theory). In [10], the variational basis of TDDFT proposed by Runge and Gross [8] was not only challenged but seemingly refuted. To be more specific, Schirmer and Dreuw claimed that the variational derivation of the time-dependent SKS equations in [8] is not valid due to an ill-defined action functional proposed there. A non-variational treatment would also encounter difficulties, since in this case the SKS system would permit one to reproduce, but not predict, the exact electron density.

Two contributions involving the present authors [22, 23] have been motivated by the criticism in [10] of the RG work. Both of the contributions accept the challenges of the RG proof, but do not require one to abandon the RG conclusion nonetheless. Let us start by summarizing the content of [22], because this is very specifically focused on time-dependent theory, whereas [23], though also motivated by the challenges in [10], is basically dealing with time-independent DFT.

2.1 Solvable example of a family of two-electron model atoms with general inter-fermion interaction: dynamical generalization

As brief background to the above example, Holas, Howard and March (HHM) [24], obtained analytical solutions for ground-state properties of a whole family of two-electron spin-compensated harmonically confined members characterized by a given interfermionic potential energy $u(r_{12})$ (See also [25, 26]). In [22] a start is made on the dynamic generalization of the harmonic external potential. In the above context, a simplified expression is obtained for the time-dependent electron density for arbitrary inter-particle correlation, which is completely determined by a one-dimensional non-interacting Hamiltonian. That such a construction is generally possible has been shown by Qian and Sahni [13, 14], but follows also from the harmonic potential theorem [27, 28] for this specific example. Furthermore for the simplest case: the Moshinsky atom [29], where the interaction $u(r_{12})$ is also harmonic, a closed solution for the Fourier transform of the density, namely the time-dependent atomic scattering factor, is found.

To summarize the essence of the time-dependent density $n(\mathbf{r}, t)$ calculation in [22], from the above model, we take the special but nevertheless important case of a system which is in its ground-state at $t = 0$. After generalizing the static separation of center of mass (CM) and relative motion (RM) to the dynamic example under consideration, the above assumption at $t = 0$ leads to the square of the CM wave function as the simple Gaussian form

$$|\psi_{000}^{\text{CM},3\text{D}}(\mathbf{c}, t)|^2 = \frac{1}{a_{\text{CM}}^3(t)\pi^{3/2}} \exp\left(-\frac{c^2}{a_{\text{CM}}^2(t)}\right), \quad (3)$$

where the time dependence is determined by the length scale $a_{\text{CM}}(t)$ of the oscillator.

For the Moshinsky example [29] the time dependent atomic scattering factor $f(\mathbf{k}, t)$, defined by

$$f(\mathbf{k}, t) = \int n(\mathbf{r}, t) e^{i\mathbf{k}\mathbf{r}} d\mathbf{r}, \quad (4)$$

is the convenient tool. The total scattering factor turns out, for $u(r_{12}) = -\frac{1}{2}\mathcal{K}r_{12}^2$, to have the form

$$f_{\text{tot}}^{\mathcal{K}}(k, t) = 2f_{\text{CM}}(k, m_{\text{cm}}\dot{\phi}(t)) f_{\text{CM}}(k/2, \tilde{m}_{\text{cm}}\dot{\phi}(t)). \quad (5)$$

where $a_{\text{CM}}(t)$ entering eqn(3) is related to $m_{\text{cm}}\dot{\phi}(t)$ in eqn(5) by

$$a_{\text{CM}}(t) = \frac{1}{m_{\text{cm}}\dot{\phi}(t)}. \quad (6)$$

Or more generally, the dynamic generalization of the static HHM density is obtained in [22] in terms of the time-dependent relative motion wave function as

$$\begin{aligned} n(\mathbf{r}, t) &= \frac{8}{\sqrt{\pi}} \exp\left(-\frac{r^2}{a_{\text{CM}}^2(t)}\right) \\ &\times \int_0^\infty dy y^2 \exp\left(-\frac{y^2}{4}\right) \left| \psi_{000}^{\text{RM},3\text{D}}(a_{\text{CM}}(t)y, t) \right|^2 \\ &\times \frac{\sinh(ry/a_{\text{CM}}(t))}{(ry/a_{\text{CM}}(t))}. \end{aligned} \quad (7)$$

Though this is an admittedly simplistic two-electron correlated time-dependent problem, the time-dependent density $n(\mathbf{r}, t)$ can be got via a one-body time-dependent potential $V(\mathbf{r}, t)$, thereby supporting the original RG assertion. There is no conflict either, we hasten to add, with the Schirmer-Dreuw conclusions. These authors, in spite of questioning the RG derivation [8] of the SKS equations, nowhere claim to have disproved this important assertion! On the contrary, an alternative proof is provided in [10] that the mapping is indeed valid.

To conclude this sub-section, we stress that a correlated two-electron example proposed in the static limit in [24] has been solved exactly in the dynamic generalization in which the system is in its ground state at time $t = 0$. In particular, eqn(7) allows the time-dependent density $n(\mathbf{r}, t)$ of the correlated dynamical problem to be reduced to the single-particle problem of calculating, probably numerically, from a one-body time-dependent Schrödinger equation, the relative motion wave function $\psi_{000}^{\text{RM},3\text{D}}$. This wave function, though calculated from a one-body equation, involves the sum of the harmonic confinement potential and the interparticle potential energy u . Having obtained the exact density, also the generating SKS potential may be obtained by a straightforward, but numerically non-trivial, inversion of the SKS equations in this two-electron case [30,31]. The one-to-one correspondence of densities and potentials in the non-interacting case can hence be verified quite explicitly for this specific example.

2.2 Linear response theory and its inversion

Since the comments [23] are basically time-independent, we shall defer these to follow a brief discussion of the important linear response function in time-dependent theory, already to the forefront in the discussion in [1] and [2]. For the non-interacting SKS system generated by potential $V(\mathbf{r}, t)$ with first order self-consistent change $\Delta V(\mathbf{r}, t)$, with corresponding response function $\chi_s(\mathbf{r}, t, \mathbf{r}', t')$, we have for the density change $\Delta n(\mathbf{r}, t)$ the formal result

$$\Delta n(\mathbf{r}, t) = \int dt' \int d\mathbf{r}' \chi_s(\mathbf{r}, t, \mathbf{r}', t') \Delta V(\mathbf{r}', t'). \quad (8)$$

Here the first order change ΔV in the (now time-dependent) SKS potential is given by

$$\Delta V(\mathbf{r}, t) = V_{\text{ext}}(\mathbf{r}, t) + \int d\mathbf{r}' \frac{\Delta n(\mathbf{r}', t)}{|\mathbf{r} - \mathbf{r}'|} + \int dt' \int d\mathbf{r}' f_{xc}(\mathbf{r}, t, \mathbf{r}', t') \Delta n(\mathbf{r}', t'), \quad (9)$$

where f_{xc} is the as yet unknown, exchange-correlation kernel [32, 33].

In this last cited references by Gross and coworkers, the issue of causality was first raised, which was also taken up later by Amusia and Shaginyan [34, 35] and Harbola [36, 37, 38]. The question posed is quite general: namely whether the potential depends on the density in a causal manner. This, we believe, is another question important to the foundations of TDDFT under discussion here.

The causality issue is important in its own right, but has also implications for the variational formulation of TDDFT: The action principle proposed by Runge and Gross [8] leads to a symmetric and hence unexpected non-causal form of the inverse of the response function χ , giving rise to what has been termed the *symmetry-causality paradox* (see also [39]). Different reformulations of the action principle have appeared in the literature, e.g., [40, 41], up to a recent contribution by Vignale in which the principle of least action in its conventional form is abandoned [42].

Returning to the question of causality, we follow the treatment of Amusia and Shaginyan [35] who write the external potential in terms of the density, by invoking the many-body linear response function χ (in contrast to the one-body χ_s used in eqn(8) above). With the external potential $V_{\text{ext}}(\mathbf{r}, t)$ one can write

$$\Delta n(\mathbf{r}, t) = \int dt' \int d\mathbf{r}' \chi(\mathbf{r}, t, \mathbf{r}', t') V_{\text{ext}}(\mathbf{r}', t'). \quad (10)$$

In [35], it is assumed that eqn(10) can be changed to the Volterra integral equation (see also [38]):

$$\frac{1}{K(t)} \frac{\partial^2 \Delta n(t)}{\partial t^2} = V_{\text{ext}}(t) + \int dt' \frac{1}{K(t)} \frac{\partial^2 \chi(t, t')}{\partial t^2} V_{\text{ext}}(t'), \quad (11)$$

where the spatial variables are omitted for clarity of notation. The function $K(t)$ entering eqn(11) is defined by [35]

$$K(t) = \left. \frac{\partial \chi(t, t')}{\partial t'} \right|_{t=t'}. \quad (12)$$

The solution of eqn(11) has the form [35, 38]

$$V_{\text{ext}}(t) = \frac{1}{K(t)} \frac{\partial^2 \Delta n(t)}{\partial t^2} + \int dt' R(t, t') \frac{1}{K(t')} \frac{\partial^2 \Delta n(t')}{\partial t'^2}, \quad (13)$$

where $R(t, t')$ vanishes for $t < t'$. Harbola [38] argues from the above treatment that while the density depends on the potential in a causal manner, the reverse is not true, leaving

no room for a symmetry-causality paradox. On the other side, Amusia and Shaginyan [34, 35] as well as Cohen and Wasserman [39] provide arguments that causal inverse response functions indeed exist. Our present contention is that all-important for a decisive answer to the causality question resides in the mathematical properties of eqn(13). This is an important future area therefore for rigorous mathematical physics.

2.3 Two further comments pertaining to the Schirmer-Dreuw study

At this point, we return to the comments of Holas et al. [23] on the Schirmer-Dreuw study [10]. As already mentioned above, in [23] the two particular subjects tackled are concerned with the original, time-independent DFT but have implications also for the time-dependent generalization. In the present section, we merely summarize the relevant points in [23] for the present context.

The first focus of [23] was to answer in a positive fashion a question posed in [10]. Whether a local operator can be reconstructed from knowledge of its particle-hole (p-h) matrix elements when the number of particle states exceeds one is the essence of the question. It arises in the context of a linear response treatment within TDDFT, in which only p-h matrix elements of the perturbing operator appear in the relevant equations, although an exact many-body approach requires also the knowledge of p-p and h-h elements. It turns out that there is no conflict with the exactness of TDDFT due to this apparent loss of information [10]. Schirmer and Dreuw formulate and prove the theorem that a local (multiplicative) operator $u = u(\mathbf{r})$ is uniquely determined to within a constant by its p-h and h-p matrix elements with respect to a complete one-particle basis and *any* partitioning of that basis into occupied and unoccupied one-particle orbitals. But in answer to the question to whether it is possible to reconstruct a local operator if only its p-h matrix elements are given, in [10] it is remarked that it seems not possible except for a special case when the number of occupied particle states (n) equals one. It is stressed by Holas et al. [23] that there is a positive answer for any n given in Sec. III of the work of Holas and Cinal [24].

As a second focus arising from the work in [10], Holas et al. [23] refer to the differential equation satisfied by the density amplitude $n(\mathbf{r})^{1/2}$, involving the concept of the Pauli potential. While conceptually satisfying, in [23] some reasons are set out why the implementation of such a *radical Kohn-Sham scheme* [10] is presently hardly computationally competitive with that based on Slater-Kohn-Sham (SKS) orbitals. Being numerically demanding already for the static case, we expect the computational scheme discussed by Schirmer and Dreuw to be even more involved in the full time-dependent case.

3 The differential virial theorem in time-dependent theory

We turn next to an important result for DFT: namely the differential virial theorem (DVT). Following the earlier study of March and Young [43] on the idempotent Dirac density matrix via its equation of motion, which led then in one-dimension to a result which they termed the differential form of the virial theorem, Holas and March [44] some four decades later established the DVT in three dimensions with full account also taken of electron-electron interactions. Ref. [44] has a fairly direct generalization to time-dependent theory [13, 14, 15,

38] and leads to the result (see also [1]):

$$\frac{\partial^2 n(\mathbf{r}, t)}{\partial t^2} = -\frac{1}{4}\nabla^4 n(\mathbf{r}, t) + \nabla \mathbf{z}(\mathbf{r}, t) + 2\nabla \int d\mathbf{r}' n_2(\mathbf{r}, \mathbf{r}', t) \nabla u(\mathbf{r}, \mathbf{r}') + \nabla [n(\mathbf{r}, t) \nabla V_{\text{ext}}(\mathbf{r}, t)]. \quad (14)$$

Here $n_2(\mathbf{r}, \mathbf{r}', t)$ is the pair density, $u(\mathbf{r}, \mathbf{r}')$ is the electron-electron repulsion potential energy, while $\mathbf{z}(\mathbf{r}, t)$ is a vector field defined from the kinetic energy density tensor $t_{\alpha\beta}(\mathbf{r}, t)$ following Holas and March [44]. To be a little more specific, eqn(14) is, in essence a combination of the DVT and the continuity equation relating density $n(\mathbf{r}, t)$ and current density $\mathbf{j}(\mathbf{r}, t)$, namely

$$\frac{\partial n(\mathbf{r}, t)}{\partial t} + \nabla \mathbf{j}(\mathbf{r}, t) = 0, \quad (15)$$

the latter being already invoked in the early work of March and Tosi [1] on TDDFT. What we want to stress is that eqn(14) can be employed at least in principle to construct the applied (external) potential from the density $n(\mathbf{r}, t)$ plus the initial conditions, a matter already touched on in Sec. 1. Prerequisites for such a construction are accurate approximations for the kinetic energy density tensor and the correlated pair density. Efforts to obtain the latter in terms of first-order density matrices [45], should be very useful in this respect.

Rewriting eqn(14) for the non-interacting SKS system, we have

$$\frac{\partial^2 n(\mathbf{r}, t)}{\partial t^2} = -\frac{1}{4}\nabla^4 n(\mathbf{r}, t) + \nabla \mathbf{z}_s(\mathbf{r}, t) + \nabla [n(\mathbf{r}, t) \nabla V(\mathbf{r}, t)], \quad (16)$$

which requires only the knowledge of non-interacting vector field $\mathbf{z}_s(\mathbf{r}, t)$ [14] to extract the Kohn-Sham potential $V(\mathbf{r}, t)$ from a given density $n(\mathbf{r}, t)$. This should be helpful in cases where direct inversion of the Kohn-Sham equations is impossible (cf. sub-section 2.1).

In order to obtain expressions for the exchange-correlation potential as *functional* of the density, admittedly, the pair density $n_2(\mathbf{r}, \mathbf{r}', t)$ still enters in the subtraction of $V_{\text{ext}}(\mathbf{r}, t)$ from $V(\mathbf{r}, t)$. In Ref.[44] dealing with the time-independent problem, the correlated pair density and kinetic energy density tensor were replaced by their non-interacting counterparts, which lead after a combination of the analogues of eqn(14) and eqn(16) to an exchange-only approximation of the exchange-correlation potential beyond the Slater form. A similar proceeding is expected to succeed also in the present time-dependent case. As discussed by Qian and Sahni [13, 14, 15] in their derivation of eqn(14) and (16), the combination of the above equations is also important from a more formal point of view, since it allows to disentangle several contributions to the exchange-correlation functional and interpret these in physical terms.

4 Shortcomings of present calculations by TDDFT on charge-transfer excitations

It is highly relevant to the foundations of TDDFT that current usage leads to substantial errors for charge-transfer excited states [46, 47, 48]. Usually, the excitation energies are severely underestimated. Furthermore, the potential energy curves of such charge transfer states do not display the known $1/R$ dependence along a charge-separation coordinate R [49, 50, 51].

A long-term solution to this problem may well lie in the use of time-dependent current density functional theory, which has recently been implemented [52] (see also [1, 3, 53]). It turns out, as discussed in [52] that a correct description of charge transfer excited states requires non-locality, and current density functionals have this property. However, the high

computational cost of such a current density approach raises doubts as to whether this route will be applicable to large molecules in the foreseeable future.

5 Summary

Serious criticisms of the foundations of TDDFT have recently been made [10], the focus being on the Runge-Gross work [8]. While accepting that there are serious flaws in [8], as correctly demonstrated in [10], the referee has asked us to stress further that the mapping theorem in [8] has never been flawed. The example set out in sub-section 2.1, and culminating in eqn(7), shows in an admittedly simple time-dependent problem with an exact solution, that the time-dependent density can be correctly calculated from a one-body time-dependent potential in support of the mapping theorem [8]. We stress here, again prompted by the referee's comments, that a major point of the Schirmer-Dreuw study [10] is that the time-dependent Kohn-Sham approach has no predictive power due to the lack of a valid variational principle. However, once the time-dependent density, discussed in some detail in sub-sections 2.1 and 2.2 above, is obtained by some other means, and the exact time-dependent exchange-correlation potential-functional would be known, the exact time-evolution of the electron density of the interacting system can be reproduced by the time-dependent Kohn-Sham equations. Since this manuscript was completed, a comment [54] and reply [55] on the Schirmer-Dreuw study [10] have been published.

Further matters discussed involve questions of the causality of the potential in TDDFT, first raised, it would seem, by Gross and co-workers [32,33] and subsequently discussed by Amusia and Shaginyan [34,35] and Harbola [38], and of the need to face the additional complications of current density theory for a specific class of excitations, namely charge-transfer excited states. This is because of the fundamental need for non-locality, which is correctly embedded in current density theory. Questions then arise as to the feasibility of application of such an approach to large molecules, because of the high cost.

Acknowledgements We wish to acknowledge that the present article was brought to fruition during a visit of both authors to the Division of Molecular Biophysics at the German Cancer Research Center (DKFZ). It is a pleasure to thank Professor S. Suhai for generous hospitality and for arranging a Scholarship to support the visit. Finally NHM thanks Prof. A. Rubio for valuable discussions pertaining to charge-transfer excitations.

References

1. March NH, Tosi MP (1972) Plasmons propagating in periodic lattices and their dispersion. Proc Roy Soc A (London) A330:373
2. March NH, Tosi MP (1973) Lattice effects in fast electron-energy losses by plasmon excitation in metals. Phil Mag 28:91
3. See also the review by Gross EKV, Kohn W (1990) Adv Quantum Chem 21:255
4. Ando T (1977) Inter-subband optical-absorption in space-charge layers on semiconductor surfaces. Z f Phys B 26:263
5. Ando T (1977) Inter-subband optical-transitions in a surface space-charge layer. Solid State Commun 21:133
6. Zangwill A, Soven P (1980) Density-functional approach to local-field effects in finite systems - photoabsorption in the rare-gases. Phys Rev A 21:1561
7. Zangwill A, Soven P (1981) Resonant 2-electron excitation in copper. Phys Rev B 24:4121
8. Runge E, Gross EKV (1984) Density-Functional Theory for Time-Dependent Systems. Phys Rev Lett 52:997
9. Hohenberg P, Kohn W (1964) Inhomogeneous Electron Gas. Phys Rev Lett 136:B864

10. Schirmer J, Dreuw A (2007) Critique of the foundations of time-dependent density-functional theory. *Phys Rev A* 75:022513
11. Slater JC (1951) A Simplification of the Hartree-Fock Method. *Phys Rev* 81:385
12. Kohn W, Sham LJ (1965) Self-Consistent Equations Including Exchange and Correlation Effects. *Phys Rev* 140:A1133
13. Qian Z, Sahni V (1998) Quantum-mechanical interpretation of time-dependent density-functional theory. *Phys Lett A* 247:303
14. Qian Z, Sahni V (2000) Time-dependent differential virial theorems. *Int J Quant Chem* 78:341
15. Qian Z, Sahni V (2001) Sum rules and properties in time-dependent density-functional theory. *Phys Rev A* 63:042508
16. Sham LJ, Schlüter M (1983) Density-Functional Theory of the Energy Gap. *Phys Rev Lett* 51:1888
17. van Leeuwen R (1996) The Sham-Schlüter Equation in Time-Dependent Density-Functional Theory. *Phys Rev Lett* 76:3610
18. Görling, A, Levy M (1994) Exact Kohn-Sham scheme based on perturbation theory. *Phys Rev A* 50:196
19. Xu BX, Rajagopal AK (1985) Current-density-functional theory for time-dependent systems. *Phys Rev A* 31:2682
20. Dhara AK, Ghosh SK (1987) Density-functional theory for time-dependent systems. *Phys Rev A* 35:442
21. Pan XY, Sahni V (2003) Corollary to the Hohenberg-Kohn theorem. *Int J Quant Chem* 95:381
22. Niehaus TA, Suhai S, March NH, (2008) Dynamical generalization of a solvable family of two-electron model atoms with general interparticle repulsion. *J Phys A: Math Theor* 41:085304
23. Holas A, Cinal M, March NH (2008) Comment on "Critique of the foundations of time-dependent density-functional theory". *Phys Rev A* 78:016501
24. Holas A, Howard IA, March NH (2003) Wave functions and low-order density matrices for a class of two-electron artificial atoms embracing Hookean and Moshinsky models. *Phys Lett A* 310:451
25. Samanta A, Ghosh SK (1990) Correlation in an exactly solvable two-particle quantum system. *Phys Rev A* 42:1178
26. Ghosh SK, Samanta A (1991) Study of correlation effects in an exactly solvable model two-electron system. *J Chem Phys* 94:517
27. Dobson JF (1994) Harmonic-Potential Theorem: Implications for Approximate Many-Body Theories. *Phys Rev Lett* 73:2244
28. Sahni V (2004) *Quantal Density Functional Theory*, Springer Verlag, Berlin
29. Moshinsky M (1968) How Good is Hartree-Fock Approximation. *Am J Phys* 36:52
30. Lein M, Kümmel S (2005) Exact Time-Dependent Exchange-Correlation Potentials for Strong-Field Electron Dynamics. *Phys Rev Lett* 94:143003
31. D'Amico I, Vignale G (1999) Exact exchange-correlation potential for a time-dependent two-electron system. *Phys Rev B* 59:7876 (1999).
32. Gross EKV, Ulrich CA, Gossmann UJ (1995) in: *Density Functional Theory*, edited by Gross EKV and Dreizler RM, NATO Advanced Studies Institute, Ser. B, Vol. 337, Plenum Press, New York
33. Gross EKV, Dobson JF, Petersilka M (1996) in: *Density Functional Theory*, edited by Nalewajski RF, Springer, New York
34. Amusia MY, Shaginyan VR (1998) Relations between action integral, response functions, and causality in density functional theory. *Phys Lett A* 250:157
35. Amusia MY, Shaginyan VR (2001) Comment on 'Analysis of causality in time-dependent density-functional theory'. *Phys Rev A* 60:056501
36. Harbola MK, Banerjee A (1999) Analysis of causality in time-dependent density-functional theory. *Phys Rev A* 60:5101
37. Harbola MK (2001) Reply to 'Comment on 'Analysis of causality in time-dependent density-functional theory''. *Phys Rev A* 63:056502
38. Harbola MK (2002) *Reviews of Modern Quantum Chemistry*, Vol. II, 1226, World Scientific, Singapore
39. Cohen MH, Wasserman A (2005) N-representability and stationarity in time-dependent density-functional theory. *Phys Rev A* 71:032515
40. Rajagopal AK (1996) Time-dependent variational principle and the effective action in density-functional theory and Berry's phase. *Phys Rev A* 54:3916
41. van Leeuwen R (1998) Causality and Symmetry in Time-Dependent Density-Functional Theory. *Phys Rev Lett* 80:1280
42. Vignale G (2008) Real-time resolution of the causality paradox of time-dependent density-functional theory. *Phys Rev A* 77:062511
43. March NH, Young WH (1959) Approximate Solutions of the Density Matrix Equation for a Local Average Field. *Nucl Phys* 12:237
44. Holas A, March NH (1995) Exact Exchange-Correlation Potential and Approximate Exchange Potential in terms of Density-matrices. *Phys Rev A* 51:2040

-
45. Valdemoro C (1992) Approximating the second-order reduced density matrix in terms of the first-order one. *Phys Rev A* 45:4462
 46. Dreuw A, Fleming GR, Head-Gordon M (2003) Chlorophyll fluorescence quenching by xanthophylls. *Phys Chem Chem Phys* 5:3247
 47. Tozer DJ, Amos RD, Handy NC, Roos BJ, Serrano-Andres L (1999) Does density functional theory contribute to the understanding of excited states of unsaturated organic compounds? *Mol Phys* 97:859
 48. Dreuw A, Head-Gordon M (2005) Single-reference ab initio methods for the calculation of excited states of large molecules. *Chem Rev* 105:4009
 49. Dreuw A, Weismann JL, Head-Gordon M (2003) Long-range charge-transfer excited states in time-dependent. *J Chem Phys* 119:2943
 50. Dreuw A, Head-Gordon M (2004) Failure of time-dependent density functional theory for long-range charge-transfer excited states: The zincbacteriochlorin-bacteriochlorin and bacteriochlorophyll-spheroidene complexes. *J Am Chem Soc* 126:4007
 51. Tozer DJ (2003) Relationship between long-range charge-transfer excitation energy error and integer discontinuity in Kohn-Sham theory. *J Chem Phys* 119:12697
 52. van Faassen M, de Boeij PL, van Leeuwen R, Berger JA, Snijders JG (2003) Application of time-dependent current-density-functional theory to nonlocal exchange-correlation effects in polymers. *J Chem Phys* 118:1044
 53. March NH, Niehaus TA, Suhai S (2006) Influence of the noninteracting density response function on the exchange-only kernel in time-dependent density-functional theory. *Phys Rev A* 74:044502
 54. Maitra NT, van Leeuwen R, Burke K (2008) Comment on "Critique of the foundations of time-dependent density functional theory". *Phys Rev A* 78:056501
 55. Schirmer J, Dreuw A (2008) Reply to "Comment on 'Critique of the foundations of time-dependent density functional theory'". *Phys Rev A* 78:056502