# THE SIGNATURE OF THE CHERN COEFFICIENTS OF LOCAL RINGS

LAURA GHEZZI, JOOYOUN HONG, AND WOLMER V. VASCONCELOS

ABSTRACT. This paper considers the following conjecture: If R is an unmixed, equidimensional local ring that is a homomorphic image of a Cohen-Macaulay local ring, then for any ideal J generated by a system of parameters, the Chern coefficient  $e_1(J) < 0$  is equivalent to R being non Cohen-Macaulay. The conjecture is established if R is a homomorphic image of a Gorenstein ring, and for all universally catenary integral domains containing fields. Criteria for the detection of Cohen-Macaulayness in equi-generated graded modules are derived.

### 1. Introduction

Let  $(R, \mathfrak{m})$  be a Noetherian local ring of dimension d > 0, and let I be an  $\mathfrak{m}$ -primary ideal. We will consider the set of multiplicative, decreasing filtrations of R ideals,  $\mathbf{B} = \{I_n, I_0 = R, I_{n+1} = II_n, n \gg 0\}$ , integral over the I-adic filtration. They are conveniently coded in the corresponding Rees algebra and its associated graded ring

$$\mathcal{R}(\mathbf{B}) = \sum_{n \ge 0} I_n t^n, \quad \operatorname{gr}_{\mathbf{B}}(R) = \sum_{n \ge 0} I_n / I_{n+1}.$$

One of our goals is to study cohomological properties of these filtrations. For that we will focus on the role of the Hilbert polynomial of the Hilbert–Samuel function  $\lambda(R/I_{n+1})$ ,

$$H_{\mathbf{B}}^{1}(n) = P_{\mathbf{B}}^{1}(n) \equiv \sum_{i=0}^{d} (-1)^{i} e_{i}(\mathbf{B}) \binom{n+d-i}{d-i},$$

particularly of the multiplicity,  $e_0(\mathbf{B})$ , and the *Chern coefficient*,  $e_1(\mathbf{B})$ . For Cohen-Macaulay rings, many penetrating relationships between these coefficients have been proved, beginning with Northcott's [7]. More recently, similar questions have been examined in general Noetherian local rings and among those pertinent to our concerns are [3], [10] and [12].

Here we extend several of the results of [12] on the meaning of the sign of  $e_1(\mathbf{B})$ , particularly in the case of *I*-adic filtrations. Our main results are centered around the following question:

AMS 2000 Mathematics Subject Classification. Primary 13A30; Secondary 13B22, 13H10, 13H15. The first author is partially supported by a grant from the City University of New York PSC-CUNY Research Award Program-39. The third author is partially supported by the NSF.

Let J be an ideal generated by a system of parameters. Under which conditions is  $e_1(J) < 0$  equivalent to R being not Cohen-Macaulay? We conjecture that this is so whenever R is an unmixed, equidimensional local ring that is a homomorphic image of a Cohen-Macaulay local ring.

There are reasons for the interest in these numbers. To make the discussion more direct, we assume that the residue field of R is infinite.

- (1) Clarifying the role of the sign of  $e_1(J)$  in the Cohen-Macaulay property of R could be used as a *scale* to classify non-CM rings.
  - (2) In the study of the normalization **B** of R[Jt], the expression (see [8], [12])

$$e_1(\mathbf{B}) - e_1(J)$$

bounds the length of certain computations in the construction of normalizations. It highlights the need to look for upper bounds of  $e_1(\mathbf{B})$  and for lower bounds of  $e_1(J)$ .

(3) The value of  $e_1(J)$  occurs as a correction term in the extensions of several well-known formulas in the theory of the Hilbert polynomials. We highlight two of them. A classical result of Northcott ([7]) asserts that if R is Cohen-Macaulay, then

$$e_1(I) \ge e_0(I) - \lambda(R/I)$$
.

For arbitrary Noetherian rings, if J is a minimal reduction of I, Goto and Nishida ([3, Theorem 3.1]) proved that

$$e_1(I) - e_1(J) \ge e_0(I) - \lambda(R/I),$$

which gives

$$e_1(\mathbf{B}) - e_1(J) \ge e_0(I) - \lambda(R/\overline{I}),$$

since  $e_1(\mathbf{B}) \geq e_1(I)$  for all such ideals I. It is a formula which is relevant to a conjecture of [12], on whether  $e_1(\mathbf{B}) \geq 0$ .

A different kind of relationship given by Huckaba and Marley ([4, Theorem 4.7]) for Cohen-Macaulay rings,

$$e_1(\mathbf{B}) \le \sum_{n>1} \lambda(I_n/JI_{n-1}),$$

is extended by Rossi and Valla ([10, Theorem 2.11]) to general filtrations to an expression that replaces  $e_1(\mathbf{B})$  in the inequality above by  $e_1(\mathbf{B}) - e_1(J)$ .

In our main result (Theorem 3.3) we show that the above Conjecture holds if R is a homomorphic image of a Gorenstein ring, or milder extensions that allow an embedding of R into a (small) Cohen-Macaulay module over a possibly larger ring. The proof is a variation of an argument in [12], but turned more abstract. Another result establishes the Conjecture for universally catenary local domains containing a field (Theorem 4.4).

The same question can be asked about filtrations of modules regarding the negativity of the coefficient  $e_1$  of the corresponding associated graded module. In case R is a

polynomial ring over a field and M is a graded, torsion–free R-module, this extension has a surprising application to the Cohen-Macaulayness of M. In the special case of modules generated in the same degree, the Hilbert coefficient  $e_1(M)$ , which may be different from  $e_1(\operatorname{gr}_{\mathbf{x}}(M))$ , can alone decide whether M is Cohen-Macaulay or not (Corollary 5.4).

We thank Rodney Sharp and Santiago Zarzuela for sharing with us their expertises on balanced big Cohen-Macaulay modules. We are also grateful to Shiro Goto for sharing with us a sketch of his proof of a more extended version of Theorem 3.3. The Yokohama Conference on Commutative Algebra, on March 2008, was the setting where some of the questions arose.

#### 2. Preliminaries

We will assemble quickly here some facts about associated graded modules and their Hilbert functions. As a general reference for unexplained terminology and basic results, we shall use [1].

Let  $(R,\mathfrak{m})$  be a Noetherian local ring, I an  $\mathfrak{m}$ -primary ideal, and M a nonzero finite R-module of dimension d. The associated graded ring  $\operatorname{gr}_I(R) = \bigoplus_{i=0}^\infty I^i/I^{i+1}$  is a standard graded ring with  $[\operatorname{gr}_I(R)]_0 = R/I$  Artinian. The associated graded module  $\operatorname{gr}_I(M) = \bigoplus_{i=0}^\infty I^i M/I^{i+1} M$  of I with respect to M is a finitely generated graded  $\operatorname{gr}_I(R)$ -module. The Hilbert-Samuel function  $\chi_M^I(n)$  of M with respect to I is

$$\chi_M^I(n) = \lambda(M/I^{n+1}M) = \sum_{i=0}^n \lambda(I^i M/I^{i+1}M).$$

For sufficiently large n, the Hilbert-Samuel function  $\chi_M^I(n)$  is of polynomial type:

$$\chi_{M}^{I}(n) = \sum_{i=0}^{d} (-1)^{i} e_{i}(I, M) \binom{n+d-i}{d-i}.$$

For an R-module M of finite length, we denote the length of M by  $\lambda(M)$ .

**Lemma 2.1.** Let  $(R, \mathfrak{m})$  be a Noetherian local ring and let I be an  $\mathfrak{m}$ -primary ideal. Let  $0 \to T \to M \to N \to 0$  be an exact sequence of finitely generated R-modules. Assume that M has dimension  $d \geq 2$  and that T has finite length. Then  $e_1(I, M) = e_1(I, N)$ .

**Proof.** From the following commutative diagram with exact rows

$$0 \longrightarrow T \cap I^{n+1}M \longrightarrow I^{n+1}M \longrightarrow I^{n+1}N \longrightarrow 0$$

$$\downarrow \qquad \qquad \downarrow \qquad \qquad \downarrow$$

$$0 \longrightarrow T \longrightarrow M \longrightarrow N \longrightarrow 0$$

we get an exact sequence

$$0 \longrightarrow T/(T \cap I^{n+1}M) \longrightarrow M/I^{n+1}M \longrightarrow N/I^{n+1}N \longrightarrow 0.$$

By the Artin–Rees Lemma,  $T \cap I^{n+1}M = 0$  for all sufficiently large n. Hence we get for all  $n \gg 0$ 

$$\lambda(T) - \lambda(M/I^{n+1}M) + \lambda(N/I^{n+1}N) = 0.$$

Let d' be the dimension of N. There are Hilbert polynomials such that

$$\lambda(M/I^{n+1}M) = \sum_{i=0}^{d} (-1)^{i} e_{i}(I, M) \binom{d+n-i}{d-i}$$

and

$$\lambda(N/I^{n+1}N) = \sum_{i=0}^{d'} (-1)^i e_i(I, N) \binom{d' + n - i}{d' - i}$$

for all sufficiently large n. Therefore d = d',

$$e_i(I, M) = e_i(I, N)$$
 for all  $i = 0, ..., d-1$  and  $e_d(I, N) = e_d(I, M) + (-1)^{d+1}\lambda(T)$ .  
In particular, since  $d \ge 2$ , we have  $e_1(I, M) = e_1(I, N)$ .

Let R be a ring, I an R-ideal and M an R-module. An element  $h \in I$  is called a superficial element of I with respect to M if there exists a positive integer c such that

$$(I^{n+1}M:_Mh)\cap I^cM=I^nM$$

for all  $n \geq c$ . A detailed discussion on superficial elements can be found in several sources, but we especially benefited from the treatment in [9, Theorem 1.5] of the construction of superficial elements. It depends simply on showing that certain finitely generated modules cannot be written as a union of a finite set of proper submodules. The existence of such elements is guaranteed if the residue field of R is infinite. Its usefulness for our purposes is expressed in the following result.

**Proposition 2.2.** ([6, (22.6)]) Let  $(R, \mathfrak{m})$  be a Noetherian local ring, I an  $\mathfrak{m}$ -primary ideal, and M a nonzero finitely generated R-module of dimension d. Let h be a superficial element of I with respect to M. Then the Hilbert coefficients of M and M/hM satisfy

$$e_i(I, M) = \begin{cases} e_i(I/(h), M/hM) & \text{for } i < d - 1. \\ e_{d-1}(I/(h), M/hM) + (-1)^{d-1}\lambda(0:_M h). & \text{for } i = d - 1. \end{cases}$$

# 3. Cohen-Macaulayness versus the vanishing of the Euler number

In this section we develop an abstract approach to the relationship between the signature of  $e_1$  and the Cohen-Macaulayness of a local ring (see [12, Theorem 3.1]). We also give a more general but still self-contained proof of the main result of [12] that avoids the use of big Cohen-Macaulay modules.

**Lemma 3.1.** Let  $(S, \mathfrak{n})$  be a Cohen–Macaulay local ring of dimension d with infinite residue field and let  $R = S/\mathfrak{p}$ , where  $\mathfrak{p}$  is a minimal prime ideal of S. Let  $x_1, \ldots, x_d$  be a system of parameters of R. Then there exists a system of parameters  $a_1, \ldots, a_d$  of S such that  $x_i = a_i + \mathfrak{p}$  for each i.

**Proof.** Let  $\mathfrak{m}$  denote the maximal ideal of R and let  $\mathfrak{p} = (c_1, \ldots, c_s)$ . Let  $x_1 = b_1 + \mathfrak{p}$  for some  $b_1 \in S$  and let  $\mathfrak{p}_1, \ldots, \mathfrak{p}_n$  be the minimal primes of S different from  $\mathfrak{p}$ . We claim that there exists  $\lambda \in S \setminus \mathfrak{n}$  such that  $b_1 + \lambda c_1 + \cdots + \lambda^s c_s \notin \mathfrak{p}_i$  for all  $i = 1, \ldots, n$ . Suppose not: since  $S/\mathfrak{n}$  is infinite, there exist  $\lambda_1, \ldots, \lambda_{s+1} \in S \setminus \mathfrak{n}$  such that  $\lambda_i + \mathfrak{n} \neq \lambda_j + \mathfrak{n}$  whenever  $i \neq j$  and such that  $b_1 + \lambda_i c_1 + \cdots + \lambda_i^s c_s \in \mathfrak{p}_k$  for some fixed k. Let A be the Vandermonde matrix determined by  $\lambda_i$ ,  $1 \leq i \leq s+1$ . We have

$$A \begin{bmatrix} b_1 \\ c_1 \\ \vdots \\ c_s \end{bmatrix} = \begin{bmatrix} 1 & \lambda_1 & \lambda_1^2 & \cdots & \lambda_1^s \\ 1 & \lambda_2 & \lambda_2^2 & \cdots & \lambda_2^s \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & \lambda_{s+1} & \lambda_{s+1}^2 & \cdots & \lambda_{s+1}^s \end{bmatrix} \begin{bmatrix} b_1 \\ c_1 \\ \vdots \\ c_s \end{bmatrix} = \begin{bmatrix} g_1 \\ g_2 \\ \vdots \\ g_{s+1} \end{bmatrix},$$

where  $g_1, \ldots, g_{s+1} \in \mathfrak{p}_k$ . Since A is invertible,  $c_1, \ldots, c_s \in \mathfrak{p}_k$  so that  $\mathfrak{p} \subseteq \mathfrak{p}_k$ , a contradiction. Let  $a_1 = b_1 + \lambda c_1 + \cdots + \lambda^s c_s$ , with  $\lambda \in S \setminus \mathfrak{n}$ , be such that  $a_1 \notin \mathfrak{p}_i$  for all  $i = 1, \ldots, n$ . Hence  $a_1$  is not contained in any minimal prime ideal of S and  $a_1 + \mathfrak{p} = x_1$ .

Let  $\mathfrak{q}_1, \ldots, \mathfrak{q}_m$  be the minimal primes of  $a_1S$  that do not contain  $\mathfrak{p}$ . Let  $x_2 = b_2 + \mathfrak{p}$  for some  $b_2 \in S$ . Then similarly as shown above, there exists  $\tau \in S \setminus \mathfrak{n}$  such that  $b_2 + \tau c_1 + \cdots + \tau^s c_s \notin \mathfrak{q}_i$  for all i. Let  $a_2 = b_2 + \tau c_1 + \cdots + \tau^s c_s$ . Then  $a_2$  is not contained in any minimal prime ideal of  $a_1S$  and  $a_2 + \mathfrak{p} = x_2$ . Now inductively we show that there exists a system of parameters  $a_1, \ldots, a_d$  of S such that  $a_i + \mathfrak{p} = x_i$  for all i.

**Lemma 3.2.** Let  $(S, \mathfrak{n})$  be a Cohen–Macaulay complete local ring of dimension  $d \geq 2$  and let M be a finitely generated S-module of dimension d satisfying Serre's condition  $(S_1)$ . Then  $H^1_{\mathfrak{n}}(M)$  is a finitely generated S-module.

**Proof.** Let k denote the residue field of S and  $\omega$  the canonical module of S. Since M satisfies Serre's condition  $(S_1)$ ,  $\operatorname{Ext}_S^{d-1}(M,\omega)$  is zero at each localization at  $\mathfrak{p}$  such that  $\mathfrak{p} \neq \mathfrak{n}$ . Thus it is a module of finite length. By Grothendieck duality ([1, 3.5.8]), we have

$$H^1_{\mathfrak{n}}(M) \simeq \operatorname{Hom}_S(\operatorname{Ext}_S^{d-1}(M,\omega), E(k)).$$

By Matlis duality([1, 3.2.13]),  $H_n^1(M)$  is finitely generated.

Let  $(R, \mathfrak{m})$  be a Noetherian local ring of dimension  $d \geq 2$ . The enabling idea is the embedding of R into a Cohen-Macaulay (possibly big Cohen-Macaulay) module E,

$$0 \to R \longrightarrow E \longrightarrow C \to 0.$$

Unfortunately, it may not be always possible to find the appropriate R-module E. Instead we will seek embed R into a Cohen-Macaulay module E over a ring S closely related to R for the purpose of computing associated graded rings of adic-filtrations. The following is our main result.

**Theorem 3.3.** Let  $(R, \mathfrak{m})$  be a Noetherian local domain of dimension  $d \geq 2$ , which is a homomorphic image of a Cohen-Macaulay local ring  $(S, \mathfrak{n})$ . If R is not Cohen-Macaulay, then  $e_1(J) < 0$  for any R-ideal J generated by a system of parameters.

**Proof.** Let  $R = S/\mathfrak{p}$ . We may assume that S has infinite residue field. If height( $\mathfrak{p}$ )  $\geq$  1, we replace S by S/L, where L is the S-ideal generated by a maximal regular sequence in  $\mathfrak{p}$ . This means that we may assume that dim  $R = \dim S$ , and that  $\mathfrak{p}$  is a minimal prime of S. In particular, we have an exact sequence of S-modules

$$0 \longrightarrow R \longrightarrow S \longrightarrow C \longrightarrow 0.$$

Let  $J=(x_1,\ldots,x_d)$  be an R-ideal generated by a system of parameters. Then by Lemma 3.1, there exists a system of parameters  $a_1,\ldots,a_d$  of S such that  $x_i=a_i+\mathfrak{p}$  for each i. Let  $I=(a_1,\ldots,a_d)S$ . Since IR=J, the associated graded ring  $\operatorname{gr}_I(R)$  of I with respect to the S-module R is equal to the associated graded ring  $\operatorname{gr}_I(R)$  of J. In particular,  $e_1(I,R)=e_1(J)$ . Therefore, for the purpose of constructing  $e_1(J)$ , we treat R as an S-module and use the I-adic filtration.

Now proceed as in the proof of [12, Theorem 3.1]. Let a be a superficial element for I with respect to the S-module R which is not contained in any associated prime of C distinct from  $\mathfrak{n}$ . We may assume that  $a=a_1$ . Reduction modulo aS gives rise to the exact sequence

$$0 \to T = \operatorname{Tor}_1^S(S/aS, C) \longrightarrow R/aR \longrightarrow S/aS \longrightarrow C/aC \to 0,$$

where  $T = (0 :_C a) \subset C$ . This shows that the associated primes of T contain a. Therefore T is either zero, or T is a non-zero module of finite support.

Let R' = R/aR, I' = I/(a), and denote the image of R/aR in S/aS by S'.

Now we use induction on  $d \geq 2$  to show that if R is not Cohen–Macaulay, then  $e_1(I,R) < 0$ . Let d=2. Notice that S' is a Cohen-Macaulay ring of dimension 1. We have that depth C=0, and so  $T \neq 0$ . As in the proof of [12, Theorem 3.1], we obtain  $e_1(I',R')=-\lambda(T)$ . Hence by Proposition 2.2,  $e_1(I,R)=-\lambda(T)<0$ .

Suppose d > 2. Consider the exact sequence of S/aS-modules :  $0 \to T \to R' \to S' \to 0$ . By Lemma 2.1, we have  $e_1(I', R') = e_1(I', S')$  since  $\dim(S/aS) = d - 1 \ge 2$ . Now it is enough to show that S' is not a Cohen-Macaulay S/aS-module. Then

since  $\dim(S') = d - 1$ , by induction we get  $e_1(I', S') < 0$ , and we conclude using Proposition 2.2.

Suppose that S' is a Cohen–Macaulay S/aS–module. Let  $\mathfrak{n}$  denote the maximal ideal of S/aS as well and let  $H^i_{\mathfrak{n}}(\cdot)$  denote the *i*th local cohomology. We are going to use the argument of [5, Proposition 2.1]. From the exact sequence  $0 \to T \to R' \to S' \to 0$ , we obtain a long exact sequence:

$$0 \to H^0_{\mathfrak{n}}(T) \to H^0_{\mathfrak{n}}(R^{\,\prime}) \to H^0_{\mathfrak{n}}(S^{\,\prime}) \to H^1_{\mathfrak{n}}(T) \to H^1_{\mathfrak{n}}(R^{\,\prime}) \to H^1_{\mathfrak{n}}(S^{\,\prime}).$$

Since S' is Cohen–Macaulay of dimension  $d-1 \geq 2$ , we have  $H^0_{\mathfrak{n}}(S') = 0 = H^1_{\mathfrak{n}}(S')$ . Since T is a torsion module, we have  $H^0_{\mathfrak{n}}(T) = T$  and  $H^1_{\mathfrak{n}}(T) = 0$ . Therefore  $T \simeq H^0_{\mathfrak{n}}(R')$  and  $H^1_{\mathfrak{n}}(R') = 0$ . Now from the exact sequence of S-modules

$$0 \longrightarrow R \stackrel{\cdot a}{\longrightarrow} R \longrightarrow R/aR \longrightarrow 0$$

we obtain the following exact sequence:

$$0 \longrightarrow T \simeq H_{\mathbf{n}}^{0}(R') \longrightarrow H_{\mathbf{n}}^{1}(R) \stackrel{\cdot a}{\longrightarrow} H_{\mathbf{n}}^{1}(R) \longrightarrow H_{\mathbf{n}}^{1}(R') = 0.$$

Therefore  $H^1_{\mathfrak{n}}(R) = aH^1_{\mathfrak{n}}(R)$ . Moreover, once R is embedded in S, we may assume that S is a complete local ring. By Lemma 3.2,  $H^1_{\mathfrak{n}}(R)$  is finitely generated. By Nakayama Lemma, we have that  $H^1_{\mathfrak{n}}(R) = 0$  so that T = 0. It follows that  $R/aR = R' \simeq S'$ , where S' is Cohen–Macaulay. This means that R is Cohen–Macaulay, which is a contradiction.

**Remark 3.4.** The proof of Theorem 3.3 can be extended from integral domains to more general local rings, R = S/L, where S is Cohen-Macaulay and dim  $R = \dim S$ , if R can be embedded into a maximal Cohen-Macaulay S-module. Notice that in order to embed R into a maximal Cohen-Macaulay S-module at a minimum we need to require that R be unmixed and equidimensional.

There is room for the following problem:

**Problem 3.5.** Let R = S/L (unmixed and equidimensional as above), where S is a Cohen-Macaulay local ring of dimension dim R. Characterize those R that can be embedded into a maximal Cohen-Macaulay S-module. Note that L may be assumed to be a primary ideal: If  $L = \bigcap Q_i$  is a primary decomposition, we have the embedding

$$S/L \hookrightarrow S/Q_1 \oplus \cdots \oplus S/Q_n$$
.

We make an elementary observation of what it takes to embed R into a free S-module (see [11, Theorem A.1]).

**Proposition 3.6.** Let S be a Noetherian ring and L an ideal of codimension zero without embedded components. If R = S/L, there is an embedding  $R \to F$  into a free S-module F if and only if L = 0: (0:L). In particular, this condition always holds if the total ring of fractions of S is a Gorenstein ring.

**Proof.** Let  $\{a_1, \ldots, a_n\}$  be a generating set of 0: L, and consider the mapping  $\varphi: S \to S^n$ ,  $\varphi(1) = (a_1, \ldots, a_n)$ ; its kernel is isomorphic to 0: (0: L). This shows that the equality L = 0: (0: L) is required for the asserted embedding.

Conversely, given an embedding  $\varphi: S/L \to S^n$ , let  $(a_1, \ldots, a_n) \in S^n$  be the image of a generator of S/L. The ideal  $\mathfrak{a}$  these entries generate is annihilated by L, and so  $\mathfrak{a} \subset 0: L$ . Since  $0: \mathfrak{a} = L$ , we have  $0: (0: L) \subset 0: \mathfrak{a} = L$ .

If the total ring of fractions of S is Gorenstein, to prove that  $0:(0:L)\subset L$ , it suffices to localize at the associated primes of L, all of which have codimension zero and a localization which is Gorenstein. But the double annihilator property is characteristic of such rings.

**Corollary 3.7.** Let  $(R, \mathfrak{m})$  be an unmixed and equidimensional Noetherian local ring of dimension  $d \geq 2$ , which is a homomorphic image of a Gorenstein local ring. If R is not Cohen-Macaulay, then  $e_1(J) < 0$  for any R-ideal J generated by a system of parameters.

**Proof.** Let R = S/L. If height $(L) \ge 1$ , we replace S by S/L', where L' is the S-ideal generated by a maximal regular sequence in L. So we may assume that  $\dim R = \dim S$ , and the conclusion follows by Proposition 3.6 and Remark 3.4.  $\square$ 

We give now a family of examples based on a method of [3].

**Example 3.8.** Let  $(S, \mathfrak{m})$  be a regular local ring of dimension four, with an infinite residue field. Let  $P_1, \ldots, P_r$  be a family of codimension two Cohen-Macaulay ideals such that for  $i \neq j$ ,  $P_i + P_j$  is an  $\mathfrak{m}$ -primary ideal. Define  $R = S / \cap_i P_i$ .

Consider the exact sequence of S-modules

$$0 \to R \longrightarrow \bigoplus_{i} S/P_{i} \longrightarrow L \to 0.$$

Note that L is a module of finite support; it may be identified with  $H^1_{\mathfrak{m}}(R)$ . Let J = (a, b) be an ideal of R forming a system of parameters, contained in the annihilator of L. We can assume that  $a, b \in S$  form a regular sequence in each  $S/P_i$ . We are going to determine  $e_1(J)$ .

For each integer n, tensoring by  $S/(a,b)^n$  we get the exact sequence

$$0 \to \operatorname{Tor}_1^S(L, S/(a, b)^n) \to R/(a, b)^n \to \bigoplus_i S/(P_i, (a, b)^n) \to L \otimes_S S/(a, b)^n \to 0.$$

For  $n \gg 0$ ,  $(a,b)^n L = 0$ , so we have that  $L \otimes_S S/(a,b)^n = L$  and  $\operatorname{Tor}_1^S(L,S/(a,b)^n) = L^{n+1}$ , from the Burch-Hilbert  $(n+1) \times n$  resolution of the ideal  $(a,b)^n$ . Since the  $R_i = S/P_i$  are Cohen-Macaulay, we obtain the following Hilbert-Samuel polynomial:

$$e_0(J)\binom{n+2}{2} - e_1(J)\binom{n+1}{1} + e_2(J) = (\sum_{i=1}^r e_0(JR_i))\binom{n+2}{2} + (n+2)\lambda(L) - \lambda(L).$$

<sup>&</sup>lt;sup>1</sup>We thank Jugal Verma for this observation.

It gives

$$e_0(JR) = \sum_{i=1}^r e_0(JR_i),$$
  

$$e_1(JR) = -\lambda(L),$$
  

$$e_2(JR) = 0.$$

# 4. Embedding into balanced big Cohen-Macaulay modules

Let  $(R, \mathfrak{m})$  be a Noetherian local domain. If R has a big Cohen-Macaulay module E, any nonzero element of E allows for an embedding  $R \hookrightarrow E$ . In fact, one may assume that E is a balanced big Cohen-Macaulay module (see [1, Section 8.5] for a discussion). According to the results of Hochster, if R contains a field, then there is a balanced big Cohen-Macaulay R-module E ([1, 8.4.2]).

To use the argument in [12, Theorem 3.2], in the exact sequence

$$0 \to R \longrightarrow E \longrightarrow C \to 0$$
,

we should, given any parameter ideal J of R, pick an element superficial for the purpose of building  $\operatorname{gr}_J(R)$  (if  $\dim R > 2$ ) and not contained in any associated prime of C different from  $\mathfrak{m}$ . This is possible if the cardinality of the residue field is larger than the cardinality of  $\operatorname{Ass}(C)$ .

**Theorem 4.1.** Let  $(R, \mathfrak{m})$  be a Noetherian local integral domain that is not Cohen-Macaulay and let E be a balanced big Cohen-Macaulay module. If the residue field of R has cardinality larger than the cardinality of a generating set for E, then  $e_1(J) < 0$  for any parameter ideal J.

Let **X** be a set of indeterminates of larger cardinality than  $\operatorname{Ass}(C)$ , and consider  $R(\mathbf{X}) = R[\mathbf{X}]_{\mathfrak{m}R[\mathbf{X}]}$ . This is a Noetherian ring ([2]), and we are going to argue that if E is a balanced big Cohen-Macaulay R-module, then  $R(\mathbf{X}) \otimes_R E$  is a balanced big Cohen-Macaulay module over  $R(\mathbf{X})$ . S. Zarzuela has kindly pointed out to us the following result:

**Theorem 4.2** ([13, Theorem 2.3]). Let  $A \to B$  be a flat morphism of local rings  $(A, \mathfrak{m}), (B, \mathfrak{n})$  and M a balanced big Cohen-Macaulay A-module. Then,  $M \otimes_A B$  is a balanced big Cohen-Macaulay B-module if and only if the following two conditions hold:

- (i)  $\mathfrak{n}(M \otimes_A B) \neq M \otimes_A B$  and
- (ii) For any prime ideal  $\mathfrak{q} \in \operatorname{supp}_{B}(M \otimes_{A} B)$ , (1)  $\operatorname{height}(\mathfrak{q}/\mathfrak{p}B) = \operatorname{depth}(C_{\overline{\mathfrak{q}}})$  and (2)  $\operatorname{height}(\mathfrak{q}) + \dim(B/\mathfrak{q}) = \dim(B)$ .

Here, we denote by supp  $_A(M)$  (small support) the set of prime ideals in A with at least one non-zero Bass number in the A-minimal injective resolution of M,  $\mathfrak{p} = \mathfrak{q} \cap A$ ,  $C = B/\mathfrak{p}B$  and  $\overline{\mathfrak{q}} = \mathfrak{q}C$ .

Moreover, if  $\mathfrak{q} \in \operatorname{supp}_B(M \otimes_A B)$  then  $\mathfrak{p} \in \operatorname{supp}_A(M)$ , and  $\operatorname{height}(A/\mathfrak{p}) + \dim(A/\mathfrak{p}) = \dim(A)$ .

Corollary 4.3. Let  $(R, \mathfrak{m})$  be a universally catenary integral domain and let E be a balanced big Cohen-Macaulay R-module. For any set  $\mathbf{X}$  of indeterminates and  $B = R(\mathbf{X}) = R[\mathbf{X}]_{\mathfrak{m}[\mathbf{X}]}, \ B \otimes_R E$  is a balanced big Cohen-Macaulay B-module.

**Theorem 4.4.** Let  $(R, \mathfrak{m})$  be a universally catenary integral domain containing a field. If R is not Cohen-Macaulay, then  $e_1(J) < 0$  for any parameter ideal J.

**Proof.** Let E be a balanced big Cohen–Macaulay R–module ([1, 8.4.2]) and consider the exact sequence  $0 \to R \to E \to C \to 0$ . Let  $\mathbf{X}$  be a set of indeterminates of larger cardinality than  $\mathrm{Ass}(C)$ , and let  $R(\mathbf{X}) = R[\mathbf{X}]_{\mathfrak{m}R[\mathbf{X}]}$ . By applying Theorem 4.1 to the exact sequence  $0 \to R(\mathbf{X}) \to E \otimes R(\mathbf{X}) \to C \otimes R(\mathbf{X}) \to 0$ , the assertion is proved.  $\square$ 

# 5. FILTERED MODULES

The same relationship discussed above between the signature of  $e_1(J)$  and the Cohen-Macaulayness of R holds true when modules are examined. Recall that if a Noetherian local ring R is embedded into either a maximal Cohen-Macaulay module ([12, Theorem 3.1]) or a balanced big Cohen-Macaulay module ([12, proof of Theorem 3.2]), then whenever R is not Cohen-Macaulay, we have  $e_1(J) < 0$  for any parameter ideal J. Now we use the same arguments as in [12, Theorems 3.1, 3.2] in order to extend the validity of Theorem 3.3 in the following manner.

**Theorem 5.1.** Let  $(R, \mathfrak{m})$  be a Noetherian local ring of dimension  $d \geq 1$  and let M be a finitely generated module embedded in a maximal Cohen–Macaulay module E. Then M is Cohen-Macaulay if and only if  $e_1(J, M) \geq 0$  for any ideal J generated by a system of parameters of M.

A variation that uses Theorem 4.4 is the following.

**Theorem 5.2.** Let  $(R, \mathfrak{m})$  be a universally catenary integral domain containing a field and let M be a finitely generated torsion–free R-module. If M is not Cohen-Macaulay, then  $e_1(J, M) < 0$  for any ideal J generated by a system of parameters of M.

**Proof.** By assumption M is a submodule of a finitely generated free R-module, which can be embedded into a finite direct sum of balanced big Cohen-Macaulay modules. The argument of [12, Theorem 3.2] applies again.

Let now  $R = k[x_1, ..., x_d]$  be a ring of polynomials over the field k, and let M be a finitely generated graded R-module. Suppose dim M = d. For  $J = (x_1, ..., x_d)$  we can apply Theorem 5.1 to M in a manner that uses the Hilbert–Samuel function information of the native grading of M.

**Theorem 5.3.** Let  $R = k[\mathbf{x}] = k[x_1, \dots, x_d]$ ,  $d \geq 2$ , be a ring of polynomials over the field k, and let M be a finitely generated graded R-module generated in degree 0. If M is torsion–free, then M is a free R-module if and only if  $e_1(M) = 0$ .

**Proof.** Since M is generated in degree 0,  $M \simeq \operatorname{gr}_{\mathbf{x}}(M)$ . By assumption, M can be embedded in a free R-module E (not necessarily by a homogeneous homomorphism). Now we apply Theorem 5.1.

If M is generated in degree a > 0, we have the equality

$$\lambda(M/(\mathbf{x})^{n+1}M) = \sum_{k=0}^{n} \lambda(M_{a+k}),$$

so the Hilbert coefficients satisfy

$$e_0(\operatorname{gr}_{\mathbf{x}}(M)) = e_0(M[a]) = e_0(M),$$
  
 $e_1(\operatorname{gr}_{\mathbf{x}}(M)) = e_1(M[a]) = e_1(M) - ae_0(M).$ 

**Corollary 5.4.** Let  $R = k[\mathbf{x}] = k[x_1, \dots, x_d]$ ,  $d \geq 2$ , be a ring of polynomials over the field k, and let M be a finitely generated graded R-module generated in degree  $a \geq 0$ . If M is torsion–free, then  $e_1(M) \leq ae_0(M)$ , with equality if and only if M is a free R-module.

## References

- [1] W. Bruns and J. Herzog, Cohen-Macaulay Rings, Cambridge University Press, 1993.
- [2] R. Gilmer and W. Heinzer, The Noetherian property for quotient rings of infinite polynomial rings, *Proc. Amer. Math. Soc.* **76** (1979), 1–7.
- [3] S. Goto and K. Nishida, Hilbert coefficients and Buchsbaumness of associated graded rings, *J. Pure & Applied Algebra* **181** (2003), 61–74.
- [4] S. Huckaba and T. Marley, Hilbert coefficients and the depths of associated graded rings, J. London Math. Soc. 56 (1997), 64–76.
- [5] C. Huneke and B. Ulrich, General hyperplane sections of algebraic varieties, J. Algebraic Geom. 2 (1993), 487–505.
- [6] M. Nagata, Local Rings, Interscience, New York, 1962.
- [7] D. G. Northcott, A note on the coefficients of the abstract Hilbert function, J. London Math. Soc. 35 (1960), 209–214.
- [8] C. Polini, B. Ulrich and W. V. Vasconcelos, Normalization of ideals and Briançon-Skoda numbers, *Math. Research Letters* **12** (2005), 827–842.
- [9] M. E. Rossi and G. Valla, Hilbert Function of Filtered Modules, arXiv:math.AC/07102346.
- [10] M. E. Rossi and G. Valla, On the Chern number of a filtration, *Rendiconti Seminario Matematico Padova*, to appear.
- [11] W. V. Vasconcelos, Reflexive modules over Gorenstein rings, Proc. Amer. Math. Soc. 19 (1968), 1349–1355.
- [12] W. V. Vasconcelos, The Chern coefficients of local rings, Michigan Math. J. 57 (2008), 725–744.
- [13] S. Zarzuela, Balanced big Cohen-Macaulay modules and flat extensions of rings, *Math. Proc. Camb. Phil. Soc.* **102** (1987), 203–209.

DEPARTMENT OF MATHEMATICS, NEW YORK CITY COLLEGE OF TECHNOLOGY-CUNY

300 Jay Street, Brooklyn, NY 11201, U.S.A.

E-mail address: lghezzi@citytech.cuny.edu

DEPARTMENT OF MATHEMATICS, SOUTHERN CONNECTICUT STATE UNIVERSITY

501 Crescent Street, New Haven, CT 06515-1533, U.S.A.

 $E ext{-}mail\ address: hongj2@southernct.edu}$ 

DEPARTMENT OF MATHEMATICS, RUTGERS UNIVERSITY

110 Frelinghuysen Rd, Piscataway, NJ 08854-8019, U.S.A.

E-mail address: vasconce@math.rutgers.edu