# **Explicit tough Ramsey graphs**

Le Anh Vinh
Mathematics Department
Harvard University
Cambridge, MA 02138
vinh@math.harvard.edu

Dang Phuong Dung School of Banking and Finance University of New South Wales Sydney 2052 NSW, Australia phuogdug@yahoo.com

Dedicated to our parents on the occasion of their 30th anniversary.

#### **Abstract**

A graph G is t-tough if any induced subgraph of it with x>1 connected components is obtained from G by deleting at least tx vertices. Chvátal conjectured that there exists an absolute constant  $t_0$  so that every  $t_0$ -tough graph is pancyclic. This conjecture was disproved by Bauer, van den Heuvel and Schmeichel by constructing a  $t_0$ -tough triangle-free graph for every real  $t_0$ . For each finite field  $\mathbb{F}_q$  with q odd, we consider graphs associated to the finite Euclidean plane and the finite upper half plane over  $\mathbb{F}_q$ . These graphs have received serious attention as they have been shown to be Ramanujan (or asymptotically Ramanujan) for large q. We will show that for infinitely many q, these graphs provide further counterexamples to Chvátal's conjecture. They also provide a good constructive lower bound for the Ramsey number R(3,k).

Mathematics Subject Classifications: 05C35, 05C38, 05C55, 05C25. Keywords: finite Euclidean graph, finite non-Euclidean graph, Ramsey graphs, tough graphs, girth, triangle-free, diameter.

# 1. Introduction

The toughness t(G) of a graph G is the largest real t so that for every positive integer  $x \ge 2$  one must delete at least tx vertices from G in order to get an induced subgraph of it with at least x connected components. G is t-tough if  $t(G) \ge t$ . This parameter was introduced by Chvátal in [10]. Chvátal proposed the following conjecture.

**Conjecture 1** ([10]) A simple unlabeled graph on n vertices is called pancyclic if it contains cycles of all lengths  $3, 4, \ldots, n$ . Then there exists an absolute constant  $t_0$  such that every  $t_0$ -tough graph is pancyclic.

This conjecture was disproved by Bauer, van den Heuvel and Schmeichel [7] who constructed, for every real  $t_0$ , a  $t_0$ -tough triangle-free graph. They defined a sequence of triangle-free graphs  $H_1, H_2, H_3, \ldots$  with  $|V(H_j)| = 2^{2j-1}(j+1)!$  and  $t(H_j) \ge \sqrt{2j+4}/2$ . We present here two new explicit constructions based on finite analogues of Euclidean and non-Euclidean planes.

The first construction is based on a remarkable new approach of Wildberger to trigonometry and Euclidean geometry. In [29], Wildberger replaces distance by quadrance and angle by spread, thus allowing the development of Euclidean geometry over any field. The following definition follows from [29].

**Definition 1** Let  $\mathbb{F}_q$  be a finite field of order q where q is an odd prime power. The quadrance Q(X,Y) between the points  $X=(x_1,x_2)$ , and  $Y=(y_1,y_2)$  in  $\mathbb{F}_q^2$  is the number

$$Q(X,Y) = (y_1 - x_1)^2 + (y_2 - x_2)^2.$$
(1)

Let q be an odd prime power and  $\mathbb{F}_q$  be the finite field with q elements. We can associate graphs to the symmetric plane  $\mathbb{F}_q^2$  using the quadrance.

**Definition 2** For a fixed  $a \in \mathbb{F}_q$ , the quadrance graph  $D_q(a)$  has the vertex set  $\mathbb{F}_q^2$ , and  $X, Y \in \mathbb{F}_q^2$  are adjacent if and only if Q(X,Y) = a.

This graph or the so-called finite Euclidean graph was first studied by Moorhouse [18] and then by Medrano et al [17]. Note that for any  $a, b \neq 0 \in \mathbb{F}_q$  then  $D_q(a)$  and  $D_q(b)$  are isomorphic. In usual 2-dimensional Euclidean space  $\mathbb{R}^2$ , then the quadrance between X, Y is unity if and only if the distance between X, Y is unity, so the unit-quadrance graph  $D_q(1)$  becomes the unit-distance graph.

The second construction is based on the well-known finite upper half planes constructed in a similar way using an analogue of Poincaré's non-Euclidean distance. We follow the construction in [22]. Let  $\mathbb{F}_q$  be the finite field with  $q=p^r$  elements, where p is an odd prime. Suppose  $\sigma$  is a generator of the multiplicative group  $\mathbb{F}_q^*$  of nonzero elements in  $\mathbb{F}_q$ . The extension  $\mathbb{F}_q(\sigma)$  is analogous to  $\mathbb{C}=\mathbb{R}[i]$ . We define the *finite Poincaré upper half-plane* as

$$H_q = \{ z = x + y\sqrt{\sigma} : x, y \in \mathbb{F}_q \text{ and } y \neq 0 \}.$$
 (2)

Note that "half-plane" is something of a misnomer since  $y \neq 0$  may not be a good finite analogue of the condition y>0 that defines the usual Poincaré upper half-plane in  $\mathbb C$ . In fact,  $H_q$  is more like a double covering of a finite upper half-plane. We use the familiar notation from complex analysis for  $z=x+y\sqrt{\sigma}\in H_q$ : x=Re(z), y=Im(z),  $\bar z=x-y\sqrt{\sigma}=z^q, N(z)=$  Norm of  $z=z\bar z=z^{1+q}$ . The *Poincaré distance* between  $z,w\in H_q$  is

$$d(z,w) = \frac{N(z-w)}{Im(z)Im(w)}. (3)$$

This distance is not a metric in the sense of analysis, but it is  $GL(2, \mathbb{F}_q)$ -invariant: d(gz, gw) = d(z, w) for all  $g \in GL(2, \mathbb{F}_q)$  and all  $z, w \in H_q$ . We can attach graphs to  $H_q$  by a method analogous to that which led to the unit-quadrance graphs  $D_q$ .

**Definition 3** For a fixed  $a \in \mathbb{F}_q$ , the finite non-Euclidean graph  $V_q(\sigma, a)$  has vertices as the points in  $H_q$  and edges between vertices z, w if and only if d(z, w) = a.

Except when a=0 or  $a=4\sigma$ ,  $P_q(\sigma,a)$  is a connected (q+1)-regular graph. When  $a=0, 4\sigma$  then  $P_q(\sigma,a)$  is disconnected, with one or two nodes, respectively, per connected component. As a varies, we have q-2 (q+1)-regular graphs  $P_q(\sigma,a)$ . The question of whether these graphs are always nonisomorphic or not is still open. These graphs are studied extensively in the literature, see [1-2, 8, 9, 11, 12, 20-22].

The rest of this paper is organized as follows. In Section 2 we determine the girth, diameter and triangle-freeness of these graphs. In Section 3 we establish some useful facts about toughness and cutsets of graphs that we will need in the study of quadrance graphs and finite non-Euclidean graphs. We then show in Sections 4 and 5 that for infinitely many values of q, these graphs constitute counterexamples to Chvátal's conjecture and also provide a good constructive lower bound for the Ramsey number R(3,k). Finally, we make some further remarks on the chromatic numbers of these graphs.

#### 2. Girth and diameter

We define the girth to be the length of a shortest circuit in a graph. The diameter is the maximum length of shortest paths between two vertices of a graph. The main aim of this section is to determine the girth and diameter of finite Euclidean and non-Euclidean graphs.

We first consider the finite Euclidean graph  $D_q$ .

**Definition 4** The circle  $C_k(A_0)$  in  $\mathbb{F}_q^2$  with center  $A_0 \in \mathbb{F}_q \times \mathbb{F}_q$  and quadrance k is the set of all points  $X \in \mathbb{F}_q \times \mathbb{F}_q$  such that

$$Q(A_0, X) = k.$$

The following lemma ([27, 29]) gives us the number of intersections between any two circles in  $\mathbb{F}_q^2$ .

**Lemma 1** Let  $i, j \neq 0$  in  $\mathbb{F}_q$  and let X, Y be two distinct points in  $\mathbb{F}_q^2$  such that  $Q(X, Y) = k \neq 0$ . Then  $|C_i(X) \cap C_j(Y)|$  only depends on i, j and k. Precisely, let

$$f(i, j, k) = ij - (k - i - j)^2/4.$$

Then the number of intersection points is  $p_{ij}^k$ , where

$$p_{ij}^{k} = \begin{cases} 0 & \text{if } f(i,j,k) \text{ is a non-square,} \\ 1 & \text{if } f(i,j,k) = 0, \\ 2 & \text{if } f(i,j,k) \text{ is a square.} \end{cases}$$

$$(4)$$

If -1 is a square in  $\mathbb{F}_q$  then there exist distinct points X, Y in  $\mathbb{F}_q^2$  with Q(X, Y) = 0. The following lemma gives us the intersection number of two circles  $C_i(X)$  and  $C_i(Y)$  in this case.

**Lemma 2** ([27]) For any  $i, j \neq 0$  in  $\mathbb{F}_q$ . Suppose that X, Y are two distinct points in  $\mathbb{F}_q^2$  with Q(X, Y) = 0. Then the circle  $C_i(X)$  intersects  $C_j(Y)$  if and only if  $i \neq j$ . Furthermore, if  $i \neq j$  then two circles intersect at only one point.

The proof of Lemma 2 is similar to the proof of Lemma 1.

**Theorem 1** For a fixed  $a \neq 0 \in \mathbb{F}_q$ , the girth of quadrance graph  $Q_q(a)$  is 3 if 3 is a square in  $\mathbb{F}_q$  and 4 otherwise.

**Proof** We have  $\{(0,0),(0,1),(1,1),(1,0)\}$  is a cycle of length 4 in  $Q_q(1)$  so the girth of unit-quadrance graph is at most 4. Besides,  $Q_q(a), Q_q(b)$  are isomorphic for any  $a,b \neq 0 \in \mathbb{F}_q$ , so all quadrance graphs  $Q_q(a)$  have girth at most 4. From Lemma 1, the girth of quadrance graph  $Q_q(a)$  is 3 if and only if  $f(a,a,a) = 3a^2/4$  is a square in  $\mathbb{F}_q$  or 3 is a square in  $\mathbb{F}_q$ . This concludes the proof of the theorem.

**Theorem 2** If  $q \equiv 3 \mod 4$  then  $D_q$  has diameter 3. Otherwise,  $D_q(a)$  has diameter 3 or 4.

**Proof** Since all quadrance graphs are isomorphic, we only need to prove the theorem for unit-quadrance graph  $D_q = D_q(1)$ . Define g(x) = 4f(x,1,1) = (4-x)x. Then there exist  $u,v \neq 0 \in \mathbb{F}_q$  such that g(u) is a square in  $\mathbb{F}_q$  while g(v) is not. Choose any  $X \in C_v((0,0))$ . By Lemma 1, X is not reachable from (0,0) by two unit-steps. Thus, the diameter of  $D_q$  is at least 3.

Now suppose that  $q \equiv 3 \mod 4$ . Then  $D_q$  is a (q+1)-regular graph. For any vertex  $X \in D_q$ , we define the set V(X) to be all vertices that are adjacent to X and T(X) to be all the vertices that are connected by a path of at most two edges from X. Then

$$T(X) = \bigcup_{Y \in V(X)} V(Y). \tag{5}$$

By Lemma 1, a fixed vertex W can be in V(Y) for at most 2 vertices  $Y \in V(X)$ . Thus,  $|T(X)| \ge (q+1)q/2$ . This implies that T(X) contains elements from no fewer than  $\lceil q/2 \rceil$  circles  $C_i(X)$ . Thus, there are no more than q - (q+1)/2 = (q-1)/2 circles outside of T(X). Let Z be any point then Z is adjacent to at most q-1 points that are not in T(X). Thus Z is connected to X by a path of length at most three. This implies that  $D_q$  has diameter 3.

Suppose that  $q \equiv 1 \mod 4$ . For any two points X,Y with  $Q(X,Y) \neq 1$ , by Lemma 2 we can choose a point Z such that Q(X,Z) = 1, Q(Y,Z) = 0. By Lemma 2 again, we can choose a point W such that Q(Z,W) = 1 and Q(W,Y) = u. Since g(u) is a square in  $\mathbb{F}_q$ , by Lemma 1, we can choose a point T such that Q(W,T) = Q(T,Y) = 1. Thus, Y is connected to X by a path of length at most four. If Q(X,Y) = 1 then Y is adjacent to X in the graph. This concludes the proof of the theorem.

The girth and diameter of finite non-Euclidean graphs have been studied before. The following theorem is due to Celniker [8].

**Theorem 3** ([8]) Let q be odd and  $a \notin \{0, 4\sigma\}$ . The girth of of  $V_q(\sigma, a)$  is either 3 or 4. Furthermore, the girth is 3 if  $a = 2\sigma$  and  $q \equiv 3 \mod 4$  or if a and  $a - 3\sigma$  are squares in  $\mathbb{F}_q$ . The girth is 4 if  $a = 2\sigma$  and  $q \equiv 1 \mod 4$ .

In [2], Angel and Evans obtained the following result for the diameter of finite non-Euclidean graphs  $V_q(\sigma, a)$ .

**Theorem 4** ([2]) Let q be odd and  $a \notin \{0, 2\sigma, 4\sigma\}$ . Then  $V_q(\sigma, a)$  has diameter 3 or 4 according to whether  $\sigma - a$  is a square or a non-square in  $\mathbb{F}_q$ . Besides, the diameter of  $X_{2\sigma}$  is 3, unless q = 3 or 5, in which case the diameter is 2.

### 3. Toughness and cutsets

Let G be a d-regular graph with n eigenvalues  $\lambda_1 \geq \lambda_2 \geq \ldots \geq \lambda_n$ . It is well-known that the largest eigenvalue  $\lambda_1 = d$ . We know that for  $a \neq 0$  the quadrance graph  $D_q(a)$  is a regular graph with degree  $\Delta(D_q(a)) = q - (-1)^{(q-1)/2}$  (see [17]). Medrano et al. [17] give a general bound for eigenvalue of  $D_q(a)$ .

**Lemma 3** [17] Let  $\lambda \neq \Delta(D_q(a))$  be any eigenvalue of graph  $D_q(a)$ . Then

$$|\lambda| \le 2q^{1/2}. (6)$$

The finite non-Euclidean graphs  $V_q(\sigma, a)$  are also known as (q+1)-regular graphs for  $a \neq 0, 4\sigma$ . By combining works of Weil [28], Evans [11, 12], Katz [14, 15], Li [16] and many others, it was proved that the finite non-Euclidean graphs are all Ramanujan graphs, that is, we have the following bound for eigenvalue of  $V_q(\sigma, a)$ .

**Lemma 4** [22] Let  $\lambda \neq q+1$  be any eigenvalue of graph  $V_q(\sigma,a)$ . Then

$$|\lambda| \le 2q^{1/2}.\tag{7}$$

An  $(n, d, \mu)$ -graph is a d-regular graph on n vertices, in which every eigenvalue  $\mu < d$  satisfies  $|\mu| \le \lambda$ . The following result is due to Alon in [4].

**Theorem 5** [4] Let G = (V, E) be an  $(n, d, \lambda)$ -graph. Then the toughness t = t(G) of G satisfies

$$t > \frac{1}{3} \left( \frac{d^2}{\lambda d + \lambda^2} - 1 \right). \tag{8}$$

Let  $\sigma$  be an arbitrary orientation of graph G, and let D be the incidence matrix of  $G^{\sigma}$ . Then the Laplacian of G is the matrix  $Q(G) = DD^T$ . It is easy to show that the Laplacian does not depend on the orientation  $\sigma$ , and hence is well-defined. Let G be a d-regular graph. If the adjacency matrix A of G has eigenvalues  $\lambda_1 \geq \ldots \geq \lambda_n$ , then the Laplacian G has eigenvalues  $\theta_1 = d - \lambda_1 \leq \ldots \leq \theta_n = d - \lambda_n$ . It is well-known (see [13], pages 287-288) that, if G is a graph on G vertices then the minimum value of

$$\frac{\sum_{uv \in E(G)} (x_u - x_v)^2}{\sum_{uv} x_u^2},\tag{9}$$

as x ranges over all nonzero vectors orthogonal to  $(1, \ldots, 1)$ , is  $\theta_2(G)$ . The maximum value is  $\theta_n(G)$ .

If  $S \subset V(G)$ , let  $\delta S$  denote the set of edges with one end in S and the other in V(X) - S. Suppose that |S| = s. Let z be the vector whose value is n - s on the vertices in S and -s on the vertices not in S. Then z is orthogonal to  $(1, \ldots, 1)$ , so we have

$$\theta_2(G) \le \frac{\sum_{uv \in E(G)} (x_u - x_v)^2}{\sum_u x_u^2} = \frac{|\delta S| n^2}{s(n-s)^2 + (n-s)s^2} = \frac{n|\delta S|}{|S|(n-|S|)} \le \theta_n(G). \tag{10}$$

The bisection width of graph G on n vertices is the minimum value of  $|\delta S|$ , for any subset S of size  $\lfloor n/2 \rfloor$ . Let bip(G) denote the maximum number of edges in a spanning bipartite subgraph of G. This equals the maximum value of  $|\delta S|$ , where S ranges over all subsets of V(X) with size at most |V(X)/2|. We have the following lemma which is an immediate consequence of (10)

**Lemma 5** ([13]) If G is a graph with n vertices, then

- 1. the bisection width of G is at least  $n\theta_2(G)/4(1+o(1))$ , and
- 2.  $bip(X) \leq n\theta_n(G)/4$ .

## 4. Properties of quadrance graphs

The following theorem summarizes some of the properties of quadrance graphs.

**Theorem 6** Let q be any prime of the form q = 12k + 7 with  $k \ge 0$  and  $a \ne 0$ . The quadrance graph  $D_q(a)$  is a  $d_q = (q+1)$ -regular graph on  $n_q = q^2$  vertices with the following properties.

- 1.  $D_q(a)$  is triangle-free and has diameter 3.
- 2. The toughness of  $D_q(a)$  is at least  $q^{1/2}/6 = n_q^{1/4}/6$ .
- 3. The bisection width of  $D_q(a)$  is at least  $q^2(q-2q^{1/2})/(4+o(1))$ .
- 4. The maximum number of edges in a spanning bipartite subgraph of  $D_q(a)$  is at most  $bip(D_q(a)) \leq q^2(q + 2a^{1/2})/4$ .
- 5. The independence number of  $D_q(a)$  is at most  $2q^{3/2}=2n_q^{3/4}$ , and hence its chromatic number is at least  $n_q^{1/4}/2$

**Proof** From Lemma 1,  $D_q(a)$  has no triangle as  $f(a, a, a) = 3a^2/4$  is a non-square in  $\mathbb{F}_q$  (If q = 12k + 7 then 3 is a non-square in  $\mathbb{F}_q$ ). Part 1 now follows from Theorem 2.

Part 2 follows directly from Lemma 3 and Theorem 5. Parts 3 and 4 are immediate from Lemma 3 and Lemma 5. Part 5 follows easily from (10) as follows. Let S be a maximum independent set of  $D_q(a)$ , then  $|\delta S| = |S|(q+1)$ . From (10), we have

$$\frac{q^2|S|(q+1)}{|S|(q^2-|S|)} \le \theta_n(D_q(a)) \le q+1+2q^{1/2}.$$

This implies that the independence number  $\alpha(D_a(a))$  is at most

$$\alpha(D_q(a)) = |S| \le q^2 - \frac{q^2(q+1)}{q+1+2q^{1/2}} \le 2q^{3/2} = 2n_q^{3/4}.$$
(11)

Hence, the chromatic number  $\chi(D_q(a))$  is at least

$$\chi(D_q(a)) \ge \frac{|V(D_q(a))|}{\alpha(D_q(a))} \ge \frac{n_q}{2n_q^{3/4}} = n_q^{1/4}/2.$$
(12)

This concludes the proof of the theorem.

Theorem 6 shows that the quadrance graph  $D_q(a)$ , where q is a prime of the form q=12k+7 and  $a\neq 0\in \mathbb{F}_q$ , is an explicit triangle-free graph on  $n_q=q^2$  vertices whose chromatic number exceeds  $0.5n_q^{1/4}$ . Note that the lower bound was already observed in [19]. In addition, the quadrance graph  $D_q(a)$  is an explicit construction showing that  $R(3,k)\geq \Omega(k^{4/3})$ . Moreover,  $D_q(a)$  has  $q^2(q+1)/2$  edges so the graph  $D_q(a)$  is also an explicit construction of a triangle-free graph G with e edges and

$$bip(G) \le \frac{e}{2} + \frac{1}{2}e^{5/6}.$$
 (13)

In general, the quadrance can be defined in any m-dimensional space  $\mathbb{F}_q^m$  for  $m \geq 2$  as follows.

**Definition 5** The quadrance Q(X,Y) between the points  $X=(x_1,\ldots,x_m)$  and  $Y=(y_1,\ldots,y_m)$  in  $\mathbb{F}_q^m$  is the number

$$Q(X,Y) := \sum_{i=1}^{m} (x_i - y_i)^2.$$

For any fixed  $a \neq 0 \in \mathbb{F}_q$ , the quadrance graph  $D_q^m(a)$  has the vertex set  $\mathbb{F}_q^m$ , and  $X, Y \in \mathbb{F}_q^m$  are adjacent if and only if Q(X,Y)=a. Similarly to the above, we have the following theorem.

**Theorem 7** Let q be any odd prime power and  $a \neq 0 \in \mathbb{F}_q$ . The unit-quadrance graph  $D_q^m(a)$  is a  $d_{q,m}$ -regular graph on  $n_{q,m} = q^m$  vertices with the following properties.

1. ([17]) Let  $\chi$  be the quadratic character. Equivalently,  $\chi$  is 1 on squares, 0 at 0 and -1 otherwise. Then

$$d_{q,m} = \begin{cases} q^{m-1} + \chi((-1)^{(m-1)/2})q^{(m-1)/2} & \text{if } m \text{ is odd,} \\ q^{m-1} - \chi((-1)^{n/2})q^{(n-2)/2} & \text{otherwise.} \end{cases}$$

2. ([17]) Let  $\lambda \neq \Delta(D_q^m)$  be any eigenvalue of the graph  $D_q^m$ . Then

$$|\lambda| \le 2q^{(m-1)/2}.$$

3. The toughness of  $D_q$  is at least

$$q^{(m-1)/2}/(6+o(1)) = n_{q,m}^{(m-1)/2m}/(6+o(1)).$$

- 4. The bisection width of  $D_q$  is at least  $q^m(d_{q,m}-2q^{(m-1)/2})/(4+o(1))$ .
- 5. The maximum number of edges in a spanning bipartite subgraph of  $D_q$  is at most  $bip(D_q) \leq q^m(d_{q,m} + 2q^{(m-1)/2})/4$ .
- 6. The independence number of  $D_q$  is at most  $(2+o(1))q^{(m+1)/2}=(2+o(1))n_{q,m}^{(m+1)/2m}$ , and hence its chromatic number is at least  $n_q^{(m-1)/2m}/(2+o(1))$

The proof of this theorem is omitted since it is the same as the proof of Theorem 6. Note that  $D_q^m(a)$  is triangle-free if and only if m=2 and q is a prime of the form  $q=12k\pm 5$ . Thus Theorem 7 is not more useful than Theorem 6 in attacking triangle-free graphs.

### 5. Properties of finite non-Euclidean graphs

The following theorem summarizes some of the properties of finite non-Euclidean graphs.

**Theorem 8** Let q be any prime of the form q = 12k + 5 with  $k \ge 1$ . The finite non-Euclidean graph  $V_q(3,6)$  is a  $d_q = (q+1)$ -regular graph on  $n_q = q^2 - q$  vertices with the following properties.

- 1.  $V_q(3,6)$  is triangle-free and has diameter 3.
- 2. The toughness of  $V_q(3,6)$  is at least  $q^{1/2}/6 > n_q^{1/4}/6$ .
- 3. The bisection width of  $V_q(3,6)$  is at least  $(q^2 q)(q 2q^{1/2})/(4 + o(1))$ .
- 4. The maximum number of edges in a spanning bipartite subgraph of  $V_q(3,6)$  is at most  $bip(V_q(3,6)) \le (q^2 q)(q + 2q^{1/2})/4$ .
- 5. The independence number of  $V_q(3,6)$  is at most  $(2+o(1))n_q^{3/4}$ , and hence its chromatic number is at least  $n_q^{1/4}/(2+o(1))$

**Proof** Part 1 follows from Theorems 3 and 4 combined with the fact that 3 is not a square in  $\mathbb{F}_q$  if q = 12k + 5. Part 2 follows directly from Lemma 4 and Theorem 5.

Parts 3 and 4 are immediate from Lemma 4 and Lemma 5.

The proof of Part 5 is similar to the proof of Theorem 6, Part 5.

Theorem 8 shows that the finite non-Euclidean graph  $V_q(3,6)$ , where q is a prime of the form q=12k+5, is an explicit triangle-free graph on  $n_q=q^2-q$  vertices whose chromatic number exceeds  $(0.5+o(1))n_q^{1/4}$ . In addition, the finite non-Euclidean graph  $V_q(a)$  is an explicit construction showing that  $R(3,k) \geq \Omega(k^{4/3})$ .

#### 6. Further remarks

It is known that the chromatic number of any graph with maximum degree d in which the number of edges in the induced subgraph on the set of all neighbors of any vertex does not exceed  $d^2/f$  is at most  $O(d/\log f)$  (see [6]). Let q be a prime of the form  $q=12k\pm 5$ . Then the induced subgraph on the set of all neighbors of any vertex is empty. So we can set  $f=q^2$ . For any q, the induced subgraph on the set of all neighbors of any vertex of  $D_q(a)$  has at most q+1 edges. This implies that we can set f=d-1. This implies the following upper bound for the chromatic number of quadrance graphs  $D_q(a)$ 

$$\chi(D_q(a)) \le O(q/\log_2 q). \tag{14}$$

This method also gives us a similar bound for higher dimensional cases.

$$\chi(D_q^m(a)) \le O(q^{m-1}/\log_2 q).$$
 (15)

Note that the best known constructive upper bound for the chromatic number of general quadrance graphs is much weaker:

$$\chi(D_q^m) \le q^{m-1}(1/2 + o(1)). \tag{16}$$

This bound is obtained by an explicit coloring in [25].

Finally, the bounds in Theorems 6 and 8 match the bounds obtained by code graphs in Theorem 3.1 in [4]. These latter graphs are Cayley graphs and their construction is based on some of the properties of certain dual BCH errorcorrecting codes. For a positive integer k, let  $F_k = GF(2^k)$  denote the finite field with  $2^k$  elements. The elements of  $F_k$  are represented by binary vectors of length k. If a and b are two such vectors, let (a, b) denote their concatenation. Let  $G_k$  be the graph whose vertices are all  $n=2^{2k}$  binary vectors of length 2k, where two vectors u and v are adjacent if and only if there exists a non-zero  $z \in F_k$  such that  $u+v=(z,z^3) \mod 2$  where  $z^3$  is computed in the field  $F_k$ . Then  $G_k$  is a  $d_k=2^k-1$ -regular graph on  $n_k=2^{2k}$  vertices. Moreover,  $G_k$  is triangle-free with independence number at most  $2n^{3/4}$ . Alon gives the better bound  $R(m,3) \ge \Omega(m^{3/2})$  in [3] by considering a graph whose vertex set is the set of all  $n = 2^{3k}$  binary vectors of length 3k (instead of all binary vectors of length 2k). Suppose that k is not divisible by 3. Let  $W_0$  be the set of all nonzero elements  $\alpha \in F_k$  such that the leftmost bit in the binary representation of  $\alpha^{7}$  is 0, and let  $W_{1}$  be the set of all nonzero elements  $\alpha \in F_{k}$  for which the leftmost bit of  $\alpha^{7}$  is 1. Then  $|W_{0}| = 2^{k-1} - 1$  and  $|W_{1}| = 2^{k-1}$ . Let  $G_{n}$  be the graph whose vertices are all  $n = 2^{3k}$ binary vectors of length 3k, where two vectors u and v are adjacent if and only if there exist  $w_0 \in W_0$  and  $w_1 \in W_1$ such that  $u+v=(w_0,w_0^3,w_0^5)+(w_1,w_1^3,w_1^5)$  where the powers are computed in the field  $F_k$  and the addition is addition modulo 2. Then  $G_n$  is a  $d_n=2^{k-1}(2^{k-1}-1)$ -regular graph on  $n=2^{3k}$  vertices. Moreover,  $G_n$  is a triangle-free graph with independence number at most  $(36 + o(1))n^{2/3}$ . Note that, going to higher dimensional unitquadrance graphs does not give us a better bound for the Ramsey number R(3,k) as for any  $m \ge 3$  the graph  $D_a^m(a)$ is not triangle-free. The problem of finding better bounds for the chromatic number of the graphs  $D_a(a)$  and  $V_a(\sigma,a)$ touches on an important question in graph theory: what is the greatest possible chromatic number for a triangle-free regular graph of order n? A possible approach is to consider the existence of sum-free varieties in high dimensional vector spaces over finite fields. We see that the varieties of degree two only give us triangle-free graphs on vector spaces of dimension two. We hope to address this problem for varieties of higher degrees in higher dimensions in a subsequent paper.

# References

- [1] Angel, J., Finite upper half planes over finite fields, Finite Fields Appl. 2 (1996), 62-68.
- [2] Angel, J. and Evans, R., Diameters of finite upper half plane graphs, Journal of Graph Theory 23 (2) (1996), 129-137.
- [3] Alon, N., Explicit Ramsey graphs and orthonormal labellings, Electronic Journal of Combinatorics 1 (1994), R12, 8pp.
- [4] Alon, N., Tough Ramsey graphs without short cycles, Journal of Algebraic Combinatorics 4 (1995), 189-195.
- [5] Alon, N., Krivelevich, M. and Sudakow, B., List coloring of random and pseudo-random graphs, *Combinatorica* **19** (4) (1999) 453-472.

- [6] Alon, N., Krivelevich, M. and Sudakov, B., Coloring graphs with sparse neighborhoods, *J. Combinatorial Theory, Ser. B* 77 (1999), 73-82.
- [7] Bauer, D., Vandenheuvel, J. and Schmeichel, E., Toughness and triangle-free graphs, *Journal of Combinatorial Theory, Series B*, **65** (2) (1995) 208-221.
- [8] Celniker, N., Eigenvalue bounds and girths of graphs of finite, upper half-planes, *Pacific Journal of Mathematics*, **166** (1), (1994) 1-21.
- [9] Celniker, N., Poulos, S., Terras, A., Trimble, C. and Velasquez, E., Is there life on finite upper half planes?, *Contemp. Math.* **143** (1993), 65-88.
- [10] Chvátal, V., Tough graphs and hamiltonian circuits, Discrete Mathematics 5 (1973), 215-218.
- [11] Evans, R., Character sums as orthogonal eigenfunctions of adjacency operators for Cayley graphs, *Contemporary Mathematics*, **168**, American Mathematical Society, Providence, RI, 1994, 33-50.
- [12] Evans, R., Spherical functions for finite upper half planes with characteristic 2, Finite Fields Appl. 1 (1995), 376-394.
- [13] Godsil, D. and Royle, G., Algebraic Graph Theory, Graduate Texts in Mathematics 207, Springer-Verlag New York, 2001.
- [14] Katz, N. M., Estimates for Soto-Andrade sums, J. Reine Angew. Math. 438 (1993), 143-161.
- [15] Katz, N. M., A note on exponential sums, Finite Fields Appl. 1 (1995), 395-398.
- [16] Li, W.-C.W., Number Theory with Applications, World Scientific, River Edge, NJ, 1996.
- [17] Medrano, A., Myers, P., Stark, H. M., Terras, A., Finite analogues of Euclidean space, *Journal of Computational and Applied Mathematics*, **68** (1996), 221-238.
- [18] Moorhouse, E., On the chromatic number of planes, manuscript, http://www.uwyo.edu/moorhouse/pub/chromatic.pdf
- [19] Problem 6, Problems collected at the AGG2004 Workshop, http://www.ipam.ucla.edu/programs/agg2004/
- [20] Terras, A., Eigenvalue problems related to finite analogues of upper half planes, Discourses in Math. 1 (1991), 237-263.
- [21] Terras, A., Are finite upper half planes ramanujan?, DIMACS Series in Discrete Math and Theoritical Computer Science 10 (1993), 125-142.
- [22] Terras, A., Survey of spectra of laplacians on finite symmetric spaces, Experimental Mathematics, (1996).
- [23] Vinh, L. A., Random walk on hypergroup of circles in a finite field, *Proceedings of Australasian Workshop on Combinatorial Algorithms* 2005, University of Ballarat, Australia, 341-351.
- [24] Vinh, L. A., Random walk on hypergroup of conics over finite fields, to appear on *The Global Journal in Pure and Applied Mathematics*.
- [25] Vinh, L. A., Some coloring problems for unit-quadrance graphs, *Proceedings of Australian Workshop on Combinatorial Algorithms* 2006, Charles Darwin University, Australia, 361-367.
- [26] Vinh, L. A., Quadrance graphs, Australian Mathematical Society Gazette, 33 (5), November 2006, 330-332.
- [27] Vinh, L. A., Explicit Ramsey graphs and Erdos distance problem over finite Euclidean and non-Euclidean spaces, *Electronics Journal of Combinatorics*, **15** (1), 2008, R5, 18 pages.
- [28] Weil, A., On the Riemann hypothesis in function fields, Proc. Natl. Acad. Sci. USA, 27 (1941), 345-347.
- [29] Wildberger, N. J., Divine Proportions: Rational Trigonometry to Universal Geometry, WildEgg, Australia, 2005.