

Classification of 64-element finite semifields

I.F. Rúa*

Elías F. Combarro†

J. Ranilla†

Abstract

A finite semifield D is a finite nonassociative ring with identity such that the set $D^* = D \setminus \{0\}$ is closed under the product. In this paper we obtain a computer-assisted description of all 64-element finite semifields, which completes the classification of finite semifields of order 125 or less.

1 Introduction

A **finite semifield** (or finite division ring) D is a finite nonassociative ring with identity such that the set $D^* = D \setminus \{0\}$ is closed under the product, i.e., it is a loop [1, 2]. Finite semifields have been traditionally considered in the context of finite geometries since they coordinatize projective semifield planes [3]. Recent applications to coding theory [4, 5, 6], combinatorics and graph theory [7], have broaden the potential interest in these rings.

Because of their diversity, the obtention of general theoretical algebraic results seems to be a rather difficult (and challenging) task. On the other hand, because of their finiteness, computational methods can be naturally considered in the study of these objects. So, the classification of finite semifields of a given order is a rather natural problem to use computations. For instance, computers were used in the classification of finite semifields of order 32 [8, 1] and 81 [9].

In this paper we present a classification up to isotopy of finite semifields with **64 elements**. Because of the complexity of the problem, the algorithms used in the papers mentioned above can not be directly used to solve the 64-element case. Our techniques combine several old methods and some crucial observations which allow us to reduce dramatically the computational effort. The resulting classification is very rich. Only one tenth of the existing planes was previously known.

The structure of the paper is as follows. In §2, basic properties of finite semifields are reviewed. Section 3 is devoted to known constructions of semifields of order 64. In §4, we describe the method we used to classify all 64-element finite semifields. Finally, in §5, a complete description of the semifields is given, together with several of their properties.

2 Preliminaries

In this section we collect definitions and facts on finite semifields. Proofs of these results can be found, for instance, in [1, 2].

A finite nonassociative ring D is called **presemifield**, if the set of nonzero elements D^* is closed under the product. If D has an identity element, then it is called **finite semifield**. If D is a finite

*Departamento de Matemáticas, Universidad de Oviedo, rua@uniovi.es . Partially supported by MEC - MTM - 2007 - 67884 C04 - 01

†Artificial Intelligence Center, University of Oviedo, {elias,ranilla}@aic.uniovi.es . Partially supported by MEC -TIN - 2007 - 61273 and MEC- TIN -2007 - 29664 - E

semifield, then D^* is a multiplicative loop. That is, there exists an element $e \in D^*$ (the identity of D) such that $ex = xe = x$, for all $x \in D$ and, for all $a, b \in D^*$, the equation $ax = b$ (resp. $xa = b$) has a unique solution.

Apart from finite fields (which are obviously finite semifields), *proper* finite semifields were first considered by L.E. Dickson [11] and were deeply studied by A.A. Albert [12, 13, 10, 14]. The term *finite semifield* was introduced in 1965 by D.E. Knuth [1]. These rings play an important role in the study of certain projective planes, called *semifield planes* [3, 1].

The characteristic of a finite presemifield D is a prime number p , and D is a finite-dimensional algebra over $GF(q)$ ($q = p^c$) of dimension d , for some $c, d \in \mathbb{N}$, so that $|D| = q^d$. If D is a finite semifield, then $GF(q)$ can be chosen to be its associative-commutative center $Z(D)$. Other relevant subsets of a finite semifield are the left, right, and middle nuclei (N_l, N_r, N_m) , and the nucleus N .

The definition of isomorphism of presemifields is the usual one for algebras, and the classification of finite semifields up to isomorphism can be naturally considered. Because of the connections to finite geometries, we must also consider the following notion. If D_1, D_2 are two presemifields over the same prime field $GF(p)$, then an **isotopy** between D_1 and D_2 is a triple (F, G, H) of bijective linear maps $D_1 \rightarrow D_2$ over $GF(p)$ such that

$$H(ab) = F(a)G(b) \quad \forall a, b \in D_1.$$

It is clear that any isomorphism between two presemifields is an isotopy, but the converse is not necessarily true. Any presemifield is isotopic to a finite semifield [1, Theorem 4.5.4]. From any presemifield D a projective plane $\mathcal{P}(D)$ can be constructed (see [3, 1] for the details of this construction). Theorem 6 in [10] shows that isotopy of finite semifields is the algebraic translation of the isomorphism between the corresponding projective planes. So, two finite semifields D_1, D_2 are isotopic if, and only if, the projective planes $\mathcal{P}(D_1), \mathcal{P}(D_2)$ are isomorphic.

The set of isotopies between a finite semifield D and itself is a group under composition, called the **autotopy** group, and denoted $\text{At}(D)$. This group acts on the *fundamental triangle* of the plane $\mathcal{P}(D)$, that is, it leaves invariant each of the three lines $L_x = \{(1, x, 0) \mid x \in D\} \cup \{(0, 1, 0)\}$, $L_y = \{(1, 0, y) \mid y \in D\} \cup \{(0, 0, 1)\}$, $L_\infty = \{(0, 1, z) \mid z \in D\} \cup \{(0, 0, 1)\}$. If D is a finite semifield, and \mathcal{D} is the set of all nonisomorphic semifields isotopic to D , then

$$(|D| - 1)^2 = |\text{At}(D)| \sum_{E \in \mathcal{D}} \frac{1}{|\text{Aut}(E)|}$$

where $\text{Aut}(E)$ denotes the automorphism group of a finite semifield E . The sum of the right term will be called *the Semifield/Automorphism (S/A) sum* [1, Theorem 3.3.4].

If $\mathcal{B} = [x_1, \dots, x_d]$ is a $GF(q)$ -basis of a presemifield D , then there exists a unique set of constants $\mathbf{A}_{D, \mathcal{B}} = \{A_{i_1 i_2 i_3}\}_{i_1, i_2, i_3=1}^d \subseteq GF(q)$ such that

$$x_{i_1} x_{i_2} = \sum_{i_3=1}^d A_{i_1 i_2 i_3} x_{i_3} \quad \forall i_1, i_2 \in \{1, \dots, d\}$$

This set is called **3-cube** corresponding to D with respect to the basis \mathcal{B} , and it completely determines the multiplication in D .


A remarkable fact is that permutation of the indexes of a 3-cube preserves the absence of nonzero divisors. Namely, if D is a presemifield, and $\sigma \in S_3$ (the symmetric group on the set $\{1, 2, 3\}$), then the set

$$\mathbf{A}_{D, \mathcal{B}}^\sigma = \{A_{i_{\sigma(1)} i_{\sigma(2)} i_{\sigma(3)}}\}_{i_1, i_2, i_3=1}^d \subseteq GF(q)$$

is the 3-cube of a $GF(q)$ -algebra $D_{\mathcal{B}}^{\sigma}$ which has not zero divisors [1, Theorem 4.3.1]. Notice that, in general, different choice of bases $\mathcal{B}, \mathcal{B}'$ lead to nonisomorphic presemifields $D_{\mathcal{B}}^{\sigma}, D_{\mathcal{B}'}^{\sigma}$. However, these presemifields are always isotopic.

The number of projective planes that can be constructed from a given finite semifield D using the transformation of the group S_3 is at most six [1, Theorem 5.2.1]. Actually, S_3 acts on the set of semifield planes of a given order. So, the classification of finite semifields can be reduced to the classification of the corresponding projective planes up to the action of the group S_3 . In this setting, we will consider a plane as *new* when no *known*¹ finite semifield coordinatizes a plane in its S_3 -class.

We shall use a graphical representation to distinguish between the different cases. The vertices of an hexagon will depict the six different planes obtained from a given finite semifield (cf. [1, Theorem 5.2.1]).



A dotted line between two planes shows that the corresponding finite semifields are isotopic. A continuous line represents that a commutative or symplectic [5] coordinatizing finite semifield exists.

The construction of finite semifields of a given order can be rephrased as a matrix problem [17, Proposition 3]. We state this proposition in the particular case of semifields of order 64.

Proposition 1. *There exists a finite semifield D of 64 elements if, and only if, there exists a set of 6 matrices $\{A_1, \dots, A_6\} \subseteq GL(6, 2)$ (a standard basis [9]) such that:*

1. A_1 is the identity matrix;
2. $\sum_{i=1}^6 \lambda_i A_i \in GL(6, 2)$, for all **non-zero tuples** $(\lambda_1, \dots, \lambda_6) \in \mathbb{Z}_2^d$.
3. The first column of the matrix A_i is the column vector with a 1 in the i -th position, and 0 everywhere else.

This result will be used through this paper to represent 64-element finite semifields. Namely, a finite semifield will be given as a tuple of matrices $(A_2, A_3, A_4, A_5, A_6)$. If the last five columns of the matrix A_i which has a one in the i -th position of the first column, and zeroes everywhere else are

$$\begin{pmatrix}
 a_{29} & a_{23} & a_{17} & a_{11} & a_5 \\
 a_{28} & a_{22} & a_{16} & a_{10} & a_4 \\
 a_{27} & a_{21} & a_{15} & a_9 & a_3 \\
 a_{26} & a_{20} & a_{14} & a_8 & a_2 \\
 a_{25} & a_{19} & a_{13} & a_7 & a_1 \\
 a_{24} & a_{18} & a_{12} & a_6 & a_0
 \end{pmatrix}$$

then it is encoded as the number $\sum_{j=0}^{29} a_j 2^j$.

¹We have considered as *known* semifields those appearing in the, up to our knowledge, last survey on the topic, [19]

3 Known semifield planes of order 64

In this section we give a classification of the known semifield planes of order 64. We considered all the constructions in [19] and explored which planes can be coordinatized by those constructions. This yielded to 31 semifield planes, divided into 11 S_3 -classes, that we list below. A semifield representative is given for each plane, together with the order of its automorphism group. Of these planes only two are commutative and can be coordinatized by 14 commutative semifields.

I (Desarguesian plane)

Finite field GF(64) (6 automorphisms)

$$(A_2, A_3, A_4, A_5, A_6) = (135274593, 67639409, 33954937, 25632381, 566730623)$$

II (Twisted field plane)

Twisted Field (1 automorphism) with parameters:

- $j \in \text{GF}(64)$ such that $j^6 + j + 1 = 0$;
- $\alpha \in \text{Aut}(\text{GF}(64))$ such that $\alpha(x) = x^4$, for all $x \in \text{GF}(64)$;
- $\beta \in \text{Aut}(\text{GF}(64))$ such that $\beta(x) = x^{4^2}$, for all $x \in \text{GF}(64)$.

$$(A_2, A_3, A_4, A_5, A_6) = (135274593, 225354480, 673682562, 25632381, 199628676)$$

III

Knuth's semifield of type 2 (1 automorphism) with parameters:

- $f \in \text{GF}(8)$ such that $f^3 + f + 1 = 0$;
- $g \in \text{GF}(8)$ such that $g + 1 = 0$;
- $\sigma \in \text{Aut}(\text{GF}(8))$ such that $\sigma(x) = x^2$, for all $x \in \text{GF}(8)$.

$$(A_2, A_3, A_4, A_5, A_6) = (135274596, 27112887, 35119969, 253266042, 1070246993)$$

IV

Knuth's semifield of type 5 (1 automorphism) with parameters:

- $f \in \text{GF}(8)$ such that $f^3 + f + 1 = 0$;
- $g \in \text{GF}(8)$ such that $g + 1 = 0$;
- $\sigma \in \text{Aut}(\text{GF}(8))$ such that $\sigma(x) = x^2$, for all $x \in \text{GF}(8)$.

$$(A_2, A_3, A_4, A_5, A_6) = (135274593, 189853287, 236639294, 212321269, 624416899)$$

V (II Huang&Johnson plane [20])

Huang&Johnson sporadic semifield of type II (3 automorphisms).

$$(A_2, A_3, A_4, A_5, A_6) = (135274623, 1022013944, 102205750, 429859362, 652592216)$$

VI (III Huang&Johnson plane)

Huang&Johnson sporadic semifield of type III (3 automorphisms).

$$(A_2, A_3, A_4, A_5, A_6) = (135274605, 1022014833, 374827988, 557069354, 336124018)$$

VII (IV Huang&Johnson plane)

Huang&Johnson sporadic semifield of type IV (1 automorphism).

$$(A_2, A_3, A_4, A_5, A_6) = (135274605, 427572234, 1072787891, 401402255, 192290736)$$

VIII (VI Huang&Johnson plane)

Huang&Johnson sporadic semifield of type VI (2 automorphisms).

$$(A_2, A_3, A_4, A_5, A_6) = (135274593, 189853287, 580915984, 793113293, 782199145)$$

IX (VII Huang&Johnson plane)

Huang&Johnson sporadic semifield of type VII (1 automorphism).

$$(A_2, A_3, A_4, A_5, A_6) = (135274605, 67640187, 851743451, 194887306, 617256025)$$

X (VIII Huang&Johnson plane)

Huang&Johnson sporadic semifield of type VIII (1 automorphism).

$$(A_2, A_3, A_4, A_5, A_6) = (135274593, 189853287, 1000703633, 930902659, 782199145)$$

XI

(Commutative plane associated to a Kantor-Williams symplectic presemifield)

Commutative semifield (6 automorphisms) with tuple of matrices

$$(A_2, A_3, A_4, A_5, A_6) = (135274594, 70580276, 37685996, 25345988, 584237329)$$

Apart from the constructions in [19], the 36 nonprimitive [18] finite semifields of [17] were also considered and two new classes (4 planes) were found.

XII (Two-sided nonprimitive plane)

H (Semifield # 1 in [17][page 1423]), the unique nonprimitive semifield of order 64 [17] (6 automorphisms).

$$(A_2, A_3, A_4, A_5, A_6) = (146808934, 811798971, 308657185, 563815286, 374228233)$$

XIII (One-sided nonprimitive plane)

Semifield # 2 in [17][page 1423], one-sided nonprimitive semifield (1 automorphism) with tuple of matrices

$$(A_2, A_3, A_4, A_5, A_6) = (135274600, 518296613, 253216863, 778190320, 47879003)$$

For each of these 13 semifield representatives, we computed the order of the center and nuclei $ZN = (Z, N, N_i, N_m, N_r)$, the list of all principal isotopes, and the order of their isomorphism groups. Some information on the autotopy group was computed as well as the length of the orbits in the fundamental triangle (L_x, L_∞, L_y) , given in the form $\sum_{i=1}^r a_i[b_i]$, if a_i cycles of length b_i ($i = 1, \dots, r$) exist. All these data are collected in Table 1.














Plane	S_3 - class	At	(L_x, L_∞, L_y)	S/A sum	ZN
I		23814	2[1] + 1[80] 2[1] + 1[80] 2[1] + 1[80]	$\frac{1}{6}$	(64, 64, 64, 64, 64)
II		567 Solvable	2[1] + 1[63] 2[1] + 1[63] 2[1] + 1[63]	$\frac{7}{1}$	(4, 4, 4, 4, 4)
III		49 \mathbb{Z}_7^2	2[1] + 2[7] + 1[49] 2[1] + 2[7] + 1[49] 2[1] + 2[7] + 1[49]	$\frac{81}{1}$	(2, 2, 2, 2, 2)
IV		441 Solvable	2[1] + 1[63] 2[1] + 1[63] 2[1] + 2[7] + 1[49]	$\frac{9}{1}$	(2, 2, 8, 4, 8)
V		42 Solvable	3[1] + 1[2] + 2[3] + 9[6] 2[1] + 1[7] + 1[14] + 1[42] 2[1] + 1[7] + 1[14] + 1[42]	$\frac{92}{1} + \frac{2}{2} + \frac{4}{3} + \frac{1}{6}$	(2, 2, 8, 2, 2)
VI		42 Solvable	3[1] + 1[2] + 2[3] + 9[6] 2[1] + 1[7] + 1[14] + 1[42] 2[1] + 1[7] + 1[14] + 1[42]	$\frac{92}{1} + \frac{2}{2} + \frac{4}{3} + \frac{1}{6}$	(2, 2, 8, 2, 2)
VII		14 Cyclic	9[1] + 28[2] 2[1] + 1[7] + 4[14] 2[1] + 1[7] + 4[14]	$\frac{280}{1} + \frac{7}{2}$	(2, 2, 8, 2, 2)
VIII		126 Solvable	2[1] + 4[3] + 1[6] + 3[9] + 1[18] 2[1] + 1[21] + 1[42] 2[1] + 1[21] + 1[42]	$\frac{28}{1} + \frac{7}{2}$	(2, 2, 8, 4, 4)
IX		42 Solvable	3[1] + 1[2] + 6[3] + 7[6] 2[1] + 1[21] + 1[42] 2[1] + 1[21] + 1[42]	$\frac{91}{1} + \frac{7}{2}$	(2, 2, 8, 2, 2)
X		42 Solvable	5[1] + 6[2] + 4[3] + 6[6] 2[1] + 1[21] + 1[42] 2[1] + 1[21] + 1[42]	$\frac{91}{1} + \frac{7}{2}$	(2, 2, 8, 2, 2)
XI		18 Solvable	2[1] + 1[3] + 1[6] + 2[9] + 2[18] 2[1] + 1[3] + 1[6] + 2[9] + 2[18] 3[1] + 1[2] + 2[3] + 9[6]	$\frac{211}{1} + \frac{16}{2} + \frac{4}{3} + \frac{1}{6}$	(2, 2, 2, 4, 2)
XII		6 Symmetric	3[1] + 1[2] + 6[3] + 7[6] 3[1] + 1[2] + 6[3] + 7[6] 3[1] + 1[2] + 6[3] + 7[6]	$\frac{636}{1} + \frac{48}{2} + \frac{4}{3} + \frac{1}{6}$	(2, 2, 2, 2, 2)
XIII		3 Cyclic	17[1] + 16[3] 2[1] + 21[3] 2[1] + 21[3]	$\frac{1323}{1}$	(2, 2, 2, 2, 2)

Table 1: Known finite semifields of 64 elements and their properties

Remark 1. 1. Let us notice that the Huang&Johnson plane of type V [20] is S_3 -equivalent to the plane **VII** above. Namely, it is the plane **VII**⁽²³⁾.

2. All these planes can be coordinatized by a finite semifield containing a primitive element (even the planes **XII** and **XIII**).

Let us introduce presentations of the autotopism group of some semifields in the table above.

$$\text{At}_{\mathbf{II}} = \langle x, y, z, t, u \mid x^3 = z, y^3 = t^2, z^3 = t^3 = u^7 = 1, y^x = yt, u^x = u^2 \rangle$$

$$\text{At}_{\mathbf{IV}} = \langle x, y, z, t \mid x^3 = y, y^3 = z^7 = t^7 = 1, z^x = z^2, t^x = t^2 \rangle$$

$$\text{At}_{\mathbf{V}} = \text{At}_{\mathbf{VI}} = \langle x, y, z \mid x^2 = y^3 = z^7 = 1, z^y = z^2 \rangle$$

$$\text{At}_{\mathbf{VIII}} = \langle x, y, z, t \mid x^2 = y^7 = z^3 = t^3 = 1, z^x = z^2, t^x = t^2 \rangle$$

$$\text{At}_{\mathbf{IX}} = \text{At}_{\mathbf{X}} = \langle x, y, z \mid x^2 = z^3 = y^7 = 1, z^x = z^2 \rangle$$

$$\text{At}_{\mathbf{XI}} = \langle x, y, z \mid x^2 = y^3 = z^3 = 1, z^x = z^2 \rangle$$

Since the same plane can be coordinatized by several constructions, we include Table 2, where every plane is followed by a list of those (pre)semifield constructions that coordinatize it:

- FF: Finite Field
- TF: Twisted Field
- K: Knuth's semifield of types 1 to 5
- JJ: Jha-Johnson semifield constructed over $GF(4)$ or $GF(8)$
- HJ: Huang-Johnson esporadic semifield II to VII
- KW: Kantor-Williams symplectic presemifield or associated commutative presemifield
- NP: Nonprimitive semifield (1 or 2-sided)

4 Search algorithm

In this section we describe an algorithm that generates all finite semifields of order 64. As we have previously observed (Proposition 1), any 64-element semifield can be described as a tuple of 6 matrices satisfying certain conditions. So, the output of the algorithm will be tuples of matrices which correspond to finite semifields. Not all possible tuples satisfying the conditions of Proposition 1 will be listed. It is only necessary to obtain representatives of all S_3 -classes of equivalence.

Our method is based on the algorithm proposed in [17]. This algorithm proved to be extraordinary efficient in the computation of all nonprimitive semifields of order 64, and it can be easily adapted to our case. It first takes matrices A_1 and A_2 , the first equal to the identity matrix, the second one chosen among a small amount of matrices (we will comment on this latter). Then, it computes 15 lists

Plane \ Permutation	1	(1 2)	(1 3)	(1 2 3)	(2 3)	(1 3 2)
I	FF	–	–	–	–	–
II	TF	–	–	–	–	–
III	K2	K2	–	–	–	–
IV	K5	–	K3 / JJ8	–	K4	–
V	HJII	NONE	–	NONE	–	–
VI	HJIII	NONE	–	NONE	–	–
VII	HJIV	–	–	–	HJV	–
VIII	HJVI / JJ4	NONE	–	NONE	–	–
IX	HJVII	NONE	–	NONE	–	–
X	HJVIII	NONE	–	NONE	–	–
XI	KWC	–	KWS	–	NONE	–
XII	NP2	–	–	–	–	–
XIII	NP1	NP1	–	NONE	–	–

Table 2: Known planes and their coordinatizing constructions

$L_{(\lambda_3, \lambda_4, \lambda_5, \lambda_6)}$ (where $(\lambda_3, \lambda_4, \lambda_5, \lambda_6) \in \mathbb{Z}_2^4 \setminus \{(0, 0, 0, 0)\}$) containing matrices B of $GL(6, 2)$ such that the first column of B is the vector $(0, 0, \lambda_3, \lambda_4, \lambda_5, \lambda_6)^t$, and such that the matrices $B + A_2, B + A_1$ and $B + A_2 + A_1$ are elements of $GL(6, 2)$. If $(A_1, A_2, A_3, A_4, A_5, A_6)$ is a tuple corresponding to a finite semifield of 64 elements, then any nonzero linear combination $\sum_{i=3}^6 \lambda_i A_i$ must be contained in the corresponding list $L_{(\lambda_3, \lambda_4, \lambda_5, \lambda_6)}$. This provides a fast test to check if a tuple of matrices corresponds to a finite semifield.

Then, these lists are compared to produce consistent tuples of matrices (A_3, A_4, A_5, A_6) . This procedure, which is described in detail in [17], essentially sieves the lists of matrices A_4, A_5 and A_6 with the help of the other lists (the matrix A_3 is fixed at this point). The resulting tiny lists are used to generate tuples of 6 matrices, and it is checked whether these tuples are related to a finite semifield or not. Once all these tuples are generated, a classification algorithm produces the different isomorphism classes of nonprimitive finite semifields.

This method produced satisfactory results in [17], even though the computational effort was remarkable big (if run in a single machine, 10 months of computing time). However, it can not be directly applied to our case. The main obstruction is the size of the lists $L_{(\lambda_3, \lambda_4, \lambda_5, \lambda_6)}$. The lists created in [17], because of some extra conditions, contained approximately 2 million matrices each. In the new situation, the lists contain more than 7 million matrices each. If we applied directly the method, it would take approximately 466 months to complete the task. This means that, in a certain sense, the classification problem of 64-element finite semifields is almost 50 times more difficult than the primitivity problem. The main drawback of the method is that all tuples of matrices must be computed, and classification is only achieved after search.

The algorithm which was used in [9] to classify 81-element finite semifields combines search and classification. It works as follows. Tuples of four matrices (A_1, A_2, A_3, A_4) associated to finite semifields of order 81 are obtained in 4 steps. In the i -th step a list of *nonequivalent partial standard bases of size i* is computed [9]. This means that a list of tuples (A_1, \dots, A_i) of i matrices is produced, with the following two conditions. First, any tuple in the list can be potentially extended to get a tuple of 4 matrices associated to a finite semifield. Second, none of them can be obtained from any other by means of an isotopy transformation. This ensures that the resulting tuples of 4 matrices coordinatize

nonisotopic finite semifields. This method lead to the classification of 81-element finite semifields *in a few days on a PC* [9].

This algorithm can not be directly used in our setting either. First, because it is necessary to obtain tuples of 6 matrices, which represents 6 steps instead of 4. Moreover, these matrices have order 6×6 , and not 4×4 . This means a significant increase of computation time. Also, some storing problems do appear in the intermediate steps, since the number of nonequivalent partial standard bases of size 4 is huge.

We managed to combine these two methods in an effective way. Namely, we used first the method of [9] to produce nonequivalente partial standard bases of size 3. Since S_3 -classes are enough for the classification of finite semifields, we considered nonequivalent partial standard bases up to S_3 -equivalence: the matrices of a partial standard basis of a semifield D , when transposed, are a partial standard basis of $D^{(1,3)}$. Then, the algorithm of [17] was used to complete these bases to tuples of 6 matrices. This combination of methods produced satisfactory results since there was a 95% reduction in the number of cases to be explored. Specifically, a total amount of 399866 nonequivalent partial standard bases of size 3 were considered. Of these, 377675 had matrix $A_2 = C(x^6 + x + 1)$ (companion matrix), and the rest had matrix $A_2 = C(x^6 + x^5 + x^3 + x^2 + 1)$. Notice that, because of [17, Section 5] and Remark 1 above, the matrix A_2 can be always chosen to be a companion matrix of a primitive polynomial. The implemented algorithm in language C required 30 days on a 12 2.5 GHz CPU linux cluster (1 computer year on a single PC). The output consisted in 95877 tuples. Classification of these matrices up to S_3 -equivalence lead to the results of the next section.

Let us remark that the extraordinary feature of our algorithm is that it allowed us to solve the classification problem in approximately the same time it was needed in [17] to solve the primitivity problem. As noticed above, this problem is, in a certain sense, 50 times simpler.

5 New Semifield Planes of order 64: a classification

The classification of 64-element finite semifields that we present in this section, completes the classification of finite semifields of order 125 or less [23, 8, 1, 9, 21, 22]. Let us compare the number of S_3 -equivalence classes, semifield planes, and coordinatizing finite semifields which were found, with those previously known (Table 3).

Number of classes	S_3 -action	Isotopy	Isomorphism
Previously known	13	35	3058
Actual number	80	332	87714

Table 3: Number of 64-element finite semifields

As we can see approximately one tenth of the semifield planes which exist were previously known. The matrices of the coordinatizing semifields, from S_3 -classes XIV to LXXX are collected in Table 4. A semifield representative with maximal number of automorphisms was chosen for each S_3 -class.

Table 4: Matrices of new 64-element finite semifields

#	A_2	A_3	A_4	A_5	A_6
XIV	135274594	833399958	260289148	1031543734	289062724
XV	135274608	929017139	43661225	236902583	58939658

Table 4 – continued from previous page

#	A_2	A_3	A_4	A_5	A_6
XVI	135274599	780494300	790514579	230544263	113930782
XVII	135274613	129115169	42444851	21901924	898304298
XVIII	135274600	61518615	173580196	594359470	490618435
XIX	135274596	127101463	659059070	333784654	916173704
XX	135274594	71778966	584275037	294214292	47830349
XXI	135274623	210604913	369031692	869572955	513934562
XXII	135274614	298719213	371084575	198518457	1064230576
XXIII	135274593	861665485	782365624	171288969	901459391
XXIV	135274599	329921844	727137562	76464013	227638817
XXV	135274611	502051974	921041233	799525002	1031595865
XXVI	135274608	245456463	960054086	892794578	298620733
XXVII	135274605	685174186	978351539	30452336	770639372
XXVIII	135274605	287116498	229601764	116769706	659794001
XXIX	135274600	391353891	935896110	611263392	219544639
XXX	135274605	253357853	818841952	214536771	376932474
XXXI	135274605	927823728	1043775209	180241271	355149199
XXXII	135274594	830091251	496454571	294214292	937736232
XXXIII	135274594	625566993	820897994	1041833019	527914637
XXXIV	135274605	1015682606	420972778	331973660	602075080
XXXV	135274605	760830577	942032486	331689540	1059713288
XXXVI	135274600	1013759534	967035803	19745382	54171530
XXXVII	135274618	702966560	581719755	551123260	248773288
XXXVIII	135274594	615209500	653723442	597803524	64203530
XXXIX	135274594	766078631	695326175	915348146	1069733783
XL	135274623	151599276	350755192	628485436	451306380
XLI	135274603	977768416	817386359	868142796	1019542329
XLII	135274618	815130072	525371889	729166901	111689247
XLIII	135274606	106019670	834133431	978786631	810064469
XLIV	135274605	127050072	1067943835	187195280	46720452
XLV	135274611	1029317650	520088048	219035419	1059375133
XLVI	135274593	331766335	841368844	1036592040	609803946
XLVII	135274608	153781051	728397374	250468680	37621084
XLVIII	135274613	194046320	574609178	123846514	234580732
XLIX	135274606	769446985	220872112	247312724	317866821
L	135274620	658083501	394974963	168700243	63130518
LI	135274623	834590458	173628128	932232776	129367971
LII	135274623	836224690	207465030	645936619	243736264
LIII	135274613	799452523	265350121	339974318	626807932
LIV	135274617	1021351255	508837061	331738527	774541306
LV	135274603	685722191	420150503	99266464	259409008
LVI	135274594	256099125	620378737	948416852	976951193
LVII	135274596	709090705	611750851	500630203	252344113
LVIII	135274599	296969012	813505114	902870605	253337313
LIX	135274620	1001102903	554659815	887933802	199592399
LX	135274617	1033514144	124586730	857117276	1057874229

Table 4 – continued from previous page

#	A_2	A_3	A_4	A_5	A_6
LXI	135274596	24792404	923093719	241684897	311785609
LXII	135274600	335867443	64403431	539538757	1048273875
LXIII	135274599	379947884	468255421	166163471	737134591
LXIV	135274608	675671366	173741043	1064349970	330692042
LXV	135274599	773727160	109764209	24488523	96297681
LXVI	135274603	116650580	995749484	402467592	539691559
LXVII	135274617	17661426	345529328	1017107050	175067510
LXVIII	135274618	213725930	722775913	632723229	390126804
LXIX	135274614	298218413	614428189	791447010	595936461
LXX	135274617	885643496	663907199	176085274	942663142
LXXI	135274620	508419752	1024350995	67020677	581629534
LXXII	135274606	559395716	262131527	86612471	850591393
LXXIII	135274593	808819530	46309136	231311151	1016044841
LXXIV	135274608	382084651	723363276	222713684	40992056
LXXV	135274605	844321142	295097610	835339006	805165097
LXXVI	135274617	342415979	844418594	360800935	877880127
LXXVII	135274596	61984053	93250846	172201558	898164556
LXXVIII	135274613	844649372	514653314	956814412	890109555
LXXIX	135274608	580417165	384216079	701045724	922685158
LXXX	135274614	599051760	670195531	755889110	1021850782

We processed these semifield representatives to obtain the order of the center and nuclei, the list of all principal isotopes, and the order of their isomorphism groups. Also, the length of the orbits in the fundamental triangle and some information on the autotopy group was computed. All these data are collected in Table 5.

Table 5: Properties of new finite semifields of 64 elements

Plane	S_3 – class	$ \text{At} $	$(\mathbf{L}_x, \mathbf{L}_\infty, \mathbf{L}_y)$	\mathbf{S}/\mathbf{A} sum	\mathbf{ZN}
XIV		63 Solvable	$2[1] + 1[63]$ $3[1] + 2[3] + 2[7] + 2[21]$ $2[1] + 1[63]$	$\frac{63}{1}$	$(2, 2, 2, 2, 4)$
XV		21 Solvable	$2[1] + 3[7] + 2[21]$ $2[1] + 3[7] + 2[21]$ $2[1] + 3[7] + 2[21]$	$\frac{186}{1} + \frac{9}{3}$	$(2, 2, 2, 2, 2)$
XVI		21 Solvable	$2[1] + 3[7] + 2[21]$ $2[1] + 3[7] + 2[21]$ $2[1] + 3[7] + 2[21]$	$\frac{186}{1} + \frac{9}{3}$	$(2, 2, 2, 2, 2)$
XVII		15 Cyclic	$2[1] + 1[3] + 4[15]$ $2[1] + 1[3] + 4[15]$ $5[1] + 12[5]$	$\frac{264}{1} + \frac{3}{5}$	$(2, 2, 2, 4, 2)$
XVIII		15 Cyclic	$2[1] + 1[3] + 3[5] + 3[15]$ $2[1] + 1[3] + 4[15]$ $2[1] + 1[3] + 4[15]$	$\frac{264}{1} + \frac{3}{5}$	$(2, 2, 2, 2, 2)$

Table 5 – continued from previous page















Plane	S_3 – class	$ At $	(L_x, L_∞, L_y)	S/A sum	ZN
XIX		14 Cyclic	$2[1] + 1[7] + 4[14]$ $2[1] + 1[7] + 4[14]$ $2[1] + 1[7] + 4[14]$	$\frac{280}{1} + \frac{7}{2}$	(2, 2, 2, 2, 2)
XX		9 Z_3^2	$2[1] + 6[3] + 5[9]$ $2[1] + 6[3] + 5[9]$ $2[1] + 6[3] + 5[9]$	$\frac{440}{1} + \frac{3}{3}$	(2, 2, 2, 2, 2)
XXI		9 Z_3^2	$2[1] + 6[3] + 5[9]$ $2[1] + 6[3] + 5[9]$ $2[1] + 6[3] + 5[9]$	$\frac{440}{1} + \frac{3}{3}$	(2, 2, 2, 2, 2)
XXII		9 Cyclic	$2[1] + 7[9]$ $2[1] + 7[9]$ $5[1] + 20[3]$	$\frac{441}{1}$	(2, 2, 2, 4, 2)
XXIII		9 Cyclic	$5[1] + 20[3]$ $2[1] + 7[9]$ $2[1] + 7[9]$	$\frac{441}{1}$	(2, 2, 4, 2, 2)
XXIV		9 Z_3^2	$2[1] + 21[3]$ $2[1] + 21[3]$ $2[1] + 6[3] + 5[9]$	$\frac{441}{1}$	(2, 2, 4, 2, 4)
XXV		9 Cyclic	$2[1] + 7[9]$ $5[1] + 20[3]$ $2[1] + 7[9]$	$\frac{441}{1}$	(2, 2, 2, 2, 4)
XXVI		9 Z_3^2	$2[1] + 21[3]$ $2[1] + 21[3]$ $2[1] + 6[3] + 5[9]$	$\frac{441}{1}$	(2, 2, 4, 2, 4)
XXVII		9 Z_3^2	$2[1] + 21[3]$ $2[1] + 21[3]$ $2[1] + 6[3] + 5[9]$	$\frac{441}{1}$	(2, 2, 4, 2, 4)
XXVIII		9 Z_3^2	$2[1] + 6[3] + 5[9]$ $2[1] + 6[3] + 5[9]$ $2[1] + 6[3] + 5[9]$	$\frac{440}{1} + \frac{3}{3}$	(2, 2, 2, 2, 2)
XXIX		9 Z_3^2	$2[1] + 6[3] + 5[9]$ $2[1] + 6[3] + 5[9]$ $2[1] + 6[3] + 5[9]$	$\frac{440}{1} + \frac{3}{3}$	(2, 2, 2, 2, 2)
XXX		9 Z_3^2	$2[1] + 6[3] + 5[9]$ $2[1] + 6[3] + 5[9]$ $2[1] + 6[3] + 5[9]$	$\frac{440}{1} + \frac{3}{3}$	(2, 2, 2, 2, 2)
XXXI		9 Z_3^2	$2[1] + 6[3] + 5[9]$ $2[1] + 6[3] + 5[9]$ $2[1] + 6[3] + 5[9]$	$\frac{440}{1} + \frac{3}{3}$	(2, 2, 2, 2, 2)
XXXII		9 Z_3^2	$2[1] + 6[3] + 5[9]$ $2[1] + 6[3] + 5[9]$ $2[1] + 6[3] + 5[9]$	$\frac{440}{1} + \frac{3}{3}$	(2, 2, 2, 2, 2)

Table 5 – continued from previous page

Plane	S_3 – class	$ At $	(L_x, L_∞, L_y)	S/A sum	ZN
XXXIII		7 Cyclic	$2[1] + 9[7]$ $9[1] + 8[7]$ $2[1] + 9[7]$	$\frac{567}{1}$	(2, 2, 2, 2, 2)
XXXIV		6 Symmetric	$2[1] + 7[3] + 7[6]$ $2[1] + 7[3] + 7[6]$ $5[1] + 6[2] + 4[3] + 6[6]$	$\frac{637}{1} + \frac{49}{2}$	(2, 2, 2, 2, 2)
XXXV		6 Cyclic	$3[1] + 1[2] + 2[3] + 9[6]$ $3[1] + 1[2] + 2[3] + 9[6]$ $3[1] + 1[2] + 2[3] + 9[6]$	$\frac{652}{1} + \frac{16}{2} + \frac{4}{3} + \frac{1}{6}$	(2, 2, 2, 2, 2)
XXXVI		6 Cyclic	$3[1] + 1[2] + 2[3] + 9[6]$ $3[1] + 1[2] + 2[3] + 9[6]$ $3[1] + 1[2] + 2[3] + 9[6]$	$\frac{652}{1} + \frac{16}{2} + \frac{4}{3} + \frac{1}{6}$	(2, 2, 2, 2, 2)
XXXVII		6 Symmetric	$2[1] + 7[3] + 7[6]$ $2[1] + 7[3] + 7[6]$ $5[1] + 6[2] + 4[3] + 6[6]$	$\frac{637}{1} + \frac{49}{2}$	(2, 2, 2, 2, 2)
XXXVIII		6 Symmetric	$2[1] + 7[3] + 7[6]$ $2[1] + 7[3] + 7[6]$ $5[1] + 6[2] + 4[3] + 6[6]$	$\frac{637}{1} + \frac{49}{2}$	(2, 2, 2, 2, 2)
XXXIX		5 Cyclic	$5[1] + 12[5]$ $5[1] + 12[5]$ $5[1] + 12[5]$	$\frac{792}{1} + \frac{9}{5}$	(2, 2, 2, 2, 2)
XL		3 Cyclic	$2[1] + 21[3]$ $2[1] + 21[3]$ $65[1]$	$\frac{1323}{1}$	(2, 2, 2, 4, 2)
XLI		3 Cyclic	$65[1]$ $2[1] + 21[3]$ $2[1] + 21[3]$	$\frac{1323}{1}$	(2, 2, 4, 2, 2)
XLII		3 Cyclic	$2[1] + 21[3]$ $65[1]$ $2[1] + 21[3]$	$\frac{1323}{1}$	(2, 2, 2, 2, 4)
XLIII		3 Cyclic	$65[1]$ $2[1] + 21[3]$ $2[1] + 21[3]$	$\frac{1323}{1}$	(2, 2, 4, 2, 2)
XLIV		3 Cyclic	$2[1] + 21[3]$ $2[1] + 21[3]$ $65[1]$	$\frac{1323}{1}$	(2, 2, 2, 4, 2)
XLV		3 Cyclic	$65[1]$ $2[1] + 21[3]$ $2[1] + 21[3]$	$\frac{1323}{1}$	(2, 2, 4, 2, 2)
XLVI		3 Cyclic	$2[1] + 21[3]$ $17[1] + 16[3]$ $2[1] + 21[3]$	$\frac{1323}{1}$	(2, 2, 2, 2, 2)

Table 5 – continued from previous page















Plane	S_3 – class	$ At $	(L_x, L_∞, L_y)	S/A sum	ZN
XLVII		3 Cyclic	$2[1] + 21[3]$ $2[1] + 21[3]$ $17[1] + 16[3]$	$\frac{1323}{1}$	(2, 2, 2, 2, 2)
XLVIII		3 Cyclic	$2[1] + 21[3]$ $2[1] + 21[3]$ $17[1] + 16[3]$	$\frac{1323}{1}$	(2, 2, 2, 2, 2)
XLIX		3 Cyclic	$2[1] + 21[3]$ $17[1] + 16[3]$ $2[1] + 21[3]$	$\frac{1323}{1}$	(2, 2, 2, 2, 2)
L		3 Cyclic	$2[1] + 21[3]$ $2[1] + 21[3]$ $17[1] + 16[3]$	$\frac{1323}{1}$	(2, 2, 2, 2, 2)
LI		3 Cyclic	$17[1] + 16[3]$ $2[1] + 21[3]$ $2[1] + 21[3]$	$\frac{1323}{1}$	(2, 2, 2, 2, 2)
LII		3 Cyclic	$17[1] + 16[3]$ $2[1] + 21[3]$ $2[1] + 21[3]$	$\frac{1323}{1}$	(2, 2, 2, 2, 2)
LIII		3 Cyclic	$65[1]$ $2[1] + 21[3]$ $2[1] + 21[3]$	$\frac{1323}{1}$	(2, 2, 4, 2, 2)
LIV		3 Cyclic	$65[1]$ $2[1] + 21[3]$ $2[1] + 21[3]$	$\frac{1323}{1}$	(2, 2, 4, 2, 2)
LV		3 Cyclic	$2[1] + 21[3]$ $65[1]$ $2[1] + 21[3]$	$\frac{1323}{1}$	(2, 2, 2, 2, 4)
LVI		3 Cyclic	$17[1] + 16[3]$ $2[1] + 21[3]$ $2[1] + 21[3]$	$\frac{1323}{1}$	(2, 2, 2, 2, 2)
LVII		3 Cyclic	$17[1] + 16[3]$ $2[1] + 21[3]$ $2[1] + 21[3]$	$\frac{1323}{1}$	(2, 2, 2, 2, 2)
LVIII		3 Cyclic	$65[1]$ $2[1] + 21[3]$ $2[1] + 21[3]$	$\frac{1323}{1}$	(2, 2, 4, 2, 2)
LIX		3 Cyclic	$17[1] + 16[3]$ $2[1] + 21[3]$ $2[1] + 21[3]$	$\frac{1323}{1}$	(2, 2, 2, 2, 2)
LX		3 Cyclic	$2[1] + 21[3]$ $17[1] + 16[3]$ $2[1] + 21[3]$	$\frac{1323}{1}$	(2, 2, 2, 2, 2)

Table 5 – continued from previous page




















Plane	S_3 – class	$ At $	(L_x, L_∞, L_y)	S/A sum	ZN
LXI		3 Cyclic	$17[1] + 16[3]$ $2[1] + 21[3]$ $2[1] + 21[3]$	$\frac{1323}{1}$	$(2, 2, 2, 2, 2)$
LXII		3 Cyclic	$2[1] + 21[3]$ $2[1] + 21[3]$ $17[1] + 16[3]$	$\frac{1323}{1}$	$(2, 2, 2, 2, 2)$
LXIII		3 Cyclic	$65[1]$ $2[1] + 21[3]$ $2[1] + 21[3]$	$\frac{1323}{1}$	$(2, 2, 4, 2, 2)$
LXIV		3 Cyclic	$2[1] + 21[3]$ $2[1] + 21[3]$ $17[1] + 16[3]$	$\frac{1323}{1}$	$(2, 2, 2, 2, 2)$
LXV		3 Cyclic	$17[1] + 16[3]$ $2[1] + 21[3]$ $2[1] + 21[3]$	$\frac{1323}{1}$	$(2, 2, 2, 2, 2)$
LXVI		3 Cyclic	$2[1] + 21[3]$ $17[1] + 16[3]$ $2[1] + 21[3]$	$\frac{1323}{1}$	$(2, 2, 2, 2, 2)$
LXVII		3 Cyclic	$65[1]$ $2[1] + 21[3]$ $2[1] + 21[3]$	$\frac{1323}{1}$	$(2, 2, 4, 2, 2)$
LXVIII		3 Cyclic	$65[1]$ $2[1] + 21[3]$ $2[1] + 21[3]$	$\frac{1323}{1}$	$(2, 2, 4, 2, 2)$
LXIX		3 Cyclic	$65[1]$ $2[1] + 21[3]$ $2[1] + 21[3]$	$\frac{1323}{1}$	$(2, 2, 4, 2, 2)$
LXX		3 Cyclic	$2[1] + 21[3]$ $2[1] + 21[3]$ $65[1]$	$\frac{1323}{1}$	$(2, 2, 2, 4, 2)$
LXXI		2 Cyclic	$9[1] + 28[2]$ $9[1] + 28[2]$ $9[1] + 28[2]$	$\frac{1960}{1} + \frac{49}{2}$	$(2, 2, 2, 2, 2)$
LXXII		2 Cyclic	$9[1] + 28[2]$ $9[1] + 28[2]$ $9[1] + 28[2]$	$\frac{1960}{1} + \frac{49}{2}$	$(2, 2, 2, 2, 2)$
LXXIII		2 Cyclic	$9[1] + 28[2]$ $9[1] + 28[2]$ $9[1] + 28[2]$	$\frac{1960}{1} + \frac{49}{2}$	$(2, 2, 2, 2, 2)$
LXXIV		2 Cyclic	$9[1] + 28[2]$ $9[1] + 28[2]$ $9[1] + 28[2]$	$\frac{1960}{1} + \frac{49}{2}$	$(2, 2, 2, 2, 2)$

Table 5 – continued from previous page

Plane	S_3 – class	$ \text{At} $	(L_x, L_∞, L_y)	S/A sum	ZN
LXXV		1 Cyclic	65[1] 65[1] 65[1]	$\frac{3969}{1}$	(2, 2, 2, 2, 2)
LXXVI		1 Cyclic	65[1] 65[1] 65[1]	$\frac{3969}{1}$	(2, 2, 2, 2, 2)
LXXVII		1 Cyclic	65[1] 65[1] 65[1]	$\frac{3969}{1}$	(2, 2, 2, 2, 2)
LXXVIII		1 Cyclic	65[1] 65[1] 65[1]	$\frac{3969}{1}$	(2, 2, 2, 2, 2)
LXXIX		1 Cyclic	65[1] 65[1] 65[1]	$\frac{3969}{1}$	(2, 2, 2, 2, 2)
LXXX		1 Cyclic	65[1] 65[1] 65[1]	$\frac{3969}{1}$	(2, 2, 2, 2, 2)

The autotopism group of the semifield XIV is $\langle x, y, z \mid x^3 = y, y^3 = z^7 = 1, z^x = z^2 \rangle$. The autotopism group of the semifields XV and XVI is $\langle x, y \mid x^3 = y^7 = 1, y^x = y^2 \rangle$.

6 Concluding remarks

In this paper we present a classification of finite semifields of order 64, with the help of computational tools. The resulting finite semifield *zoo* is very rich, and it completes the classification of finite semifields up to order 125. The algorithm used is a combination of known methods and some crucial observations. It proved to be quite suitable to the problem considered. We hope this classification will be helpful in the discovery of new semifield constructions, a problem which has not been addressed in this paper.

Appendix

Using the Oyama representation of matrices [24] we managed to obtain different algebraic descriptions for the semifield representatives. As a complement we include these presentations here, in the hope that they can be helpful in the discovery of new constructions. There are three different types of algebraic representations:

1. $D = GF(8)^2$, with $j^3 + j + 1 = 0$, and multiplication rule $(x_1, x_2) * (a_1, a_2)$.
2. $D = GF(4)^3$, with $j^2 + j + 1 = 0$, and multiplication rule $(x_1, x_2, x_3) * (a_1, a_2, a_3)$.
3. $D = GF(64)$, with $j^6 + j + 1 = 0$, and multiplication rule $x * a$.

Next we list the multiplication rules of the semifield representatives.

- **XIV: (1)** $(x_1a_1 + x_2(j^4a_1 + j^4a_1^4 + a_2^4) + x_2^2(j^3a_1 + j^3a_1^4 + j^3a_2^4) + x_2^4(j^6a_1 + j^6a_1^4 + j^6a_2^4) , x_1a_2 + x_2(a_1^4 + j^4a_2) + x_2^2(j^3a_2 + a_2^4) + x_2^4(j^6a_2 + j^3a_2^4))$
- **XV: (1)** $(x_1a_1 + x_2(ja_2 + j^6a_2^4) + x_2^2j^2a_2^2 + x_2^4(j^2a_1 + j^2a_1^2 + j^6a_2 + j^3a_2^2) , x_1a_2 + x_2(a_1 + j^2a_2^4) + x_2^2j^5a_2^2 + x_2^4j^2a_2)$
- **XVI: (1)** $(x_1a_1 + x_2(j^2a_2 + a_2^2) + x_2^2(j^6a_1 + j^6a_1^4 + j^3a_2^4) + x_2^4(j^5a_1 + j^5a_1^2 + a_2^2 + j^3a_2^4) , x_1a_2 + x_2(a_1 + j^6a_2^2) + x_2^2j^6a_2 + x_2^4j^5a_2)$
- **XVII: (2)** $(x_1a_1 + x_2(j^2a_2 + j^2a_3) + x_2^2j^2a_2^2 + x_3(ja_2 + j^2a_3) + x_3^2(j^2a_2^2 + j^2a_3^2) , x_1a_2 + x_2a_1 + x_2^2j^2a_3^2 + x_3^2(j^2a_2^2 + j^2a_3^2) , x_1a_3 + x_2j^2a_3 + x_2^2(a_2^2 + j^2a_3^2) + x_3(a_1 + j^2a_2) + x_3^2(j^2a_2^2 + a_3^2))$
- **XVIII: (2)** $(x_1a_1 + x_2(j^2a_1 + j^2a_1^2 + a_2^2 + j^2a_3 + ja_3^2) + x_2^2(a_2^2 + j^2a_3) + x_3(ja_1 + ja_1^2 + j^2a_2 + j^2a_2^2 + a_3^2) + x_3^2(a_2 + a_3 + j^2a_3^2) , x_1a_2 + x_2(a_1^2 + j^2a_2 + ja_2^2 + j^2a_3^2) + x_2^2(j^2a_2 + ja_2^2 + j^2a_3 + ja_3^2) + x_3ja_2 + x_3^2(ja_2 + ja_2^2 + j^2a_3) , x_1a_3 + x_2j^2a_3 + x_2^2(a_2 + j^2a_3 + j^2a_3^2) + x_3(a_1^2 + ja_2^2 + ja_3 + j^2a_3^2) + x_3^2(j^2a_2^2 + ja_3 + ja_3^2))$
- **XIX: (3)** $(x_1(j^{59}a_1 + j^7a_1^2 + j^{30}a_1^4 + j^8a_1^8 + j^{11}a_1^{16} + j^{17}a_1^{32}) + x_1^2(j^5a_1 + j^{39}a_1^2 + j^6a_1^4 + j^{13}a_1^8 + j^{57}a_1^{16} + j^{62}a_1^{32}) + x_1^4(j^4a_1 + j^7a_1^2 + j^9a_1^4 + j^{53}a_1^8 + j^{49}a_1^{16} + j^{57}a_1^{32}) + x_1^8(j^{29}a_1 + j^{37}a_1^2 + j^{22}a_1^4 + j^{23}a_1^8 + j^{51}a_1^{16} + j^{11}a_1^{32}) + x_1^{16}(j^{16}a_1 + j^{12}a_1^2 + j^{43}a_1^4 + j^{16}a_1^8 + j^{51}a_1^{16} + j^{45}a_1^{32}) + x_1^{32}(j^{47}a_1 + j^{56}a_1^2 + j^{53}a_1^4 + j^{36}a_1^8 + j^{26}a_1^{16} + j^6a_1^{32}))$
- **XX: (2)** $(x_1a_1 + x_2(a_2^2 + a_3^2) + x_2^2(a_2^2 + ja_3^2) + x_3(a_1 + a_1^2 + a_2^2 + a_3^2) + x_3^2(ja_2^2 + a_3^2) , x_1a_2 + x_2(a_1^2 + a_3^2) + x_3a_2 + x_3^2ja_3^2 , x_1a_3 + x_2^2(a_2^2 + ja_3^2) + x_3(a_1^2 + a_3 + a_3^2) + x_3^2(ja_2^2 + a_3^2))$
- **XXI: (2)** $(x_1a_1 + x_2(a_1 + a_1^2 + ja_3^2) + x_3ja_3^2 + x_3^2(j^2a_2^2 + j^2a_3^2) , x_1a_2 + x_2(a_1^2 + a_2 + a_2^2) + x_3^2(j^2a_2^2 + j^2a_3^2) , x_1a_3 + x_2a_3 + x_2^2a_2^2 + x_3(a_1^2 + a_2^2) + x_3^2(ja_2^2 + a_3^2))$
- **XXII: (2)** $(x_1a_1 + x_2(j^2a_1 + j^2a_1^2 + a_2^2 + j^2a_3) + x_2^2(j^2a_2 + ja_2^2 + ja_3 + a_3^2) + x_3j^2a_3 + x_3^2(ja_2^2 + a_3^2) , x_1a_2 + x_2(a_1^2 + j^2a_3^2) + x_2^2(ja_2 + ja_2^2 + ja_3 + j^2a_3^2) + x_3j^2a_2 + x_3^2(ja_2^2 + j^2a_3^2) , x_1a_3 + x_2(a_1 + a_1^2 + ja_2 + ja_2^2 + a_3) + x_2^2(ja_2 + j^2a_2^2 + ja_3 + ja_3^2) + x_3(a_1 + ja_2 + ja_3) + x_3^2(j^2a_2^2 + ja_3^2))$
- **XXIII: (2)** $(x_1a_1 + x_2(ja_1 + ja_1^2 + ja_2 + ja_2^2 + ja_3^2) + x_3(a_2 + ja_3 + ja_3^2) , x_1a_2 + x_2(ja_1 + j^2a_1^2 + j^2a_2 + j^2a_2^2 + ja_3 + ja_3^2) + x_3(ja_1 + ja_1^2 + a_2 + j^2a_2^2 + a_3) , x_1a_3 + x_2(j^2a_1 + j^2a_1^2 + j^2a_2 + ja_2^2 + j^2a_3 + ja_3^2) + x_3(a_1^2 + a_2 + j^2a_3^2))$
- **XXIV: (2)** $(x_1a_1 + x_2(ja_1 + ja_1^2 + j^2a_2 + j^2a_2^2 + ja_3 + j^2a_3^2) + x_3(a_1 + a_1^2 + j^2a_2 + a_2^2 + j^2a_3^2) , x_1a_2 + x_2(j^2a_1 + ja_1^2 + a_2 + a_2^2 + a_3 + j^2a_3^2) + x_3(a_1 + a_1^2 + a_2 + a_2^2 + j^2a_3 + a_3^2) , x_1a_3 + x_2(ja_1 + ja_1^2 + j^2a_2 + ja_2^2 + a_3 + j^2a_3^2) + x_3(a_1 + ja_2 + ja_3))$
- **XXV: (1)** $(x_1a_1 + x_2(j^5a_1 + j^3a_1^2 + j^2a_1^4 + a_2 + ja_2^2 + j^5a_2^4) + x_2^2(j^2a_1 + a_1^2 + j^6a_1^4 + j^5a_2 + ja_2^2) + x_2^4(j^4a_1 + a_1^2 + j^5a_1^4 + j^3a_2 + j^5a_2^4) , x_1a_2 + x_2(ja_1 + j^6a_1^2 + j^4a_1^4 + a_2 + ja_2^2 + ja_2^4) + x_2^2(j^6a_1 + j^6a_1^2 + j^2a_2^2 + ja_2^4) + x_2^4(j^4a_1 + j^4a_1^4 + j^5a_2 + j^3a_2^2 + j^6a_2^4))$
- **XXVI: (2)** $(x_1a_1 + x_2(a_1 + a_1^2 + a_2 + a_2^2 + ja_3) + x_3(a_1 + a_1^2 + ja_2 + a_3 + j^2a_3^2) , x_1a_2 + x_2(a_1 + j^2a_3) + x_3(a_1 + a_1^2 + j^2a_2 + ja_2^2 + j^2a_3 + a_3^2) , x_1a_3 + x_2(a_2^2 + ja_3 + ja_3^2) + x_3(a_1 + ja_2 + j^2a_2^2 + j^2a_3^2))$
- **XXVII: (2)** $(x_1a_1 + x_2(ja_1 + ja_1^2 + ja_2 + ja_2^2 + a_3 + a_3^2) + x_3(ja_1 + ja_1^2 + j^2a_2 + a_2^2 + j^2a_3) , x_1a_2 + x_2(a_1 + a_2 + a_2^2 + a_3) + x_3(ja_1 + ja_1^2 + j^2a_2 + j^2a_2^2 + ja_3^2) , x_1a_3 + x_2(j^2a_1 + j^2a_1^2 + a_2 + ja_2^2) + x_3(a_1 + ja_2 + ja_3))$

- **XXVIII: (2)** $(x_1a_1 + x_2ja_2^2 + x_2^2(ja_2^2 + ja_3^2) + x_3(ja_1 + ja_1^2 + ja_2^2 + ja_3^2), x_1a_2 + x_2(a_1^2 + j^2a_3^2) + x_2^2ja_3^2 + x_3ja_2 + x_3^2(j^2a_2^2 + j^2a_3^2), x_1a_3 + x_2^2(a_2^2 + a_3^2) + x_3(a_1^2 + ja_3 + j^2a_3^2))$
- **XXIX: (2)** $(x_1a_1 + x_2(a_1 + a_1^2 + a_2^2 + j^2a_3^2) + x_2^2(a_2^2 + a_3^2) + x_3(a_1 + a_1^2 + j^2a_2^2 + a_3^2) + x_3^2a_2^2, x_1a_2 + x_2(a_1^2 + a_2 + a_2^2 + a_3^2) + x_2^2j^2a_3^2 + x_3a_2 + x_3^2(a_2^2 + a_3^2), x_1a_3 + x_2a_3 + x_2^2(a_2^2 + ja_3^2) + x_3(a_1^2 + a_2^2 + a_3 + a_3^2) + x_3^2a_3^2)$
- **XXX: (2)** $(x_1a_1 + x_2(a_1 + a_1^2) + x_2^2ja_3^2 + x_3(a_2^2 + j^2a_3^2) + x_3^2a_3^2, x_1a_2 + x_2(a_1^2 + a_2 + a_2^2) + x_2^2ja_3^2 + x_3^2a_3^2, x_1a_3 + x_2a_3 + x_2^2(a_2^2 + j^2a_3^2) + x_3(a_1^2 + a_2^2) + x_3^2(j^2a_2^2 + j^2a_3^2))$
- **XXXI: (2)** $(x_1a_1 + x_2(a_1 + a_1^2 + j^2a_2^2 + j^2a_3^2) + x_2^2(j^2a_2^2 + a_3^2) + x_3(j^2a_1 + j^2a_1^2 + ja_2^2) + x_3^2a_3^2, x_1a_2 + x_2(a_1^2 + a_2 + a_2^2 + ja_3^2) + x_3j^2a_2 + x_3^2(ja_2^2 + j^2a_3^2), x_1a_3 + x_2a_3 + x_2^2(a_2^2 + ja_3^2) + x_3(a_1^2 + a_2^2 + j^2a_3 + ja_3^2) + x_3^2(j^2a_2^2 + j^2a_3^2))$
- **XXXII: (2)** $(x_1a_1 + x_2(a_1 + a_1^2) + x_2^2ja_3^2 + x_3(a_2^2 + j^2a_3^2) + x_3^2a_3^2, x_1a_2 + x_2(a_1^2 + a_2 + a_2^2) + x_2^2ja_3^2 + x_3^2a_3^2, x_1a_3 + x_2a_3 + x_2^2(a_2^2 + a_3^2) + x_3(a_1^2 + a_2^2) + x_3^2(j^2a_2^2 + ja_3^2))$
- **XXXIII: (1)** $(x_1a_1 + x_2ja_2^2 + x_2^2(j^6a_1 + j^6a_1^2 + j^6a_2^2) + x_2^4(j^6a_1 + j^6a_1^2 + j^3a_2^2), x_1a_2 + x_2(a_1^2 + j^5a_2^2) + x_2^2(j^6a_2 + ja_2^2) + x_2^4(j^6a_2 + j^5a_2^2))$
- **XXXIV: (2)** $(x_1a_1 + x_2(a_1 + a_1^2 + a_2 + ja_3 + a_3^2) + x_2^2(a_2^2 + j^2a_3^2) + x_3(j^2a_2 + a_2^2 + a_3) + x_3^2ja_2^2, x_1a_2 + x_2(a_1^2 + a_2^2 + ja_3) + x_2^2(a_2^2 + j^2a_3^2) + x_3(j^2a_2 + a_3) + x_3^2ja_2^2, x_1a_3 + x_2(a_2 + j^2a_3) + x_3(a_1^2 + a_2 + a_2^2 + ja_3) + x_3^2(j^2a_2^2 + j^2a_3^2))$
- **XXXV: (2)** $(x_1a_1 + x_2(j^2a_1 + j^2a_1^2 + ja_2^2) + x_2^2(a_2 + j^2a_2^2 + a_3^2) + x_3(ja_2^2 + ja_3 + ja_3^2) + x_3^2(j^2a_1 + j^2a_1^2 + ja_2 + j^2a_2^2 + a_3), x_1a_2 + x_2(a_1^2 + j^2a_2 + ja_2^2) + x_2^2a_2^2 + x_3(j^2a_2^2 + j^2a_3) + x_3^2(a_1 + a_1^2 + j^2a_2 + ja_2^2), x_1a_3 + x_2(j^2a_2 + j^2a_3 + a_3^2) + x_2^2(ja_1 + ja_1^2 + a_2 + j^2a_2^2) + x_3(a_1^2 + ja_2^2) + x_3^2a_3^2)$
- **XXXVI: (2)** $(x_1a_1 + x_2(ja_1 + ja_1^2 + a_2 + ja_2^2) + x_2^2(j^2a_2 + a_3) + x_3(j^2a_1 + j^2a_1^2 + j^2a_3 + ja_3^2) + x_3^2ja_2, x_1a_2 + x_2(a_1^2 + a_2 + a_2^2 + a_3 + j^2a_3^2) + x_2^2j^2a_3 + x_3(ja_2 + j^2a_2^2) + x_3^2(a_2 + ja_3), x_1a_3 + x_2(a_3 + j^2a_3^2) + x_2^2a_2 + x_3(a_1^2 + j^2a_2 + a_2^2 + ja_3 + j^2a_3^2) + x_3^2a_3)$
- **XXXVII: (2)** $(x_1a_1 + x_2(j^2a_1 + j^2a_1^2 + a_2^2 + j^2a_3^2) + x_2^2(ja_1 + ja_1^2 + ja_2 + j^2a_2^2 + j^2a_3 + j^2a_3^2) + x_3(a_1 + a_1^2 + ja_2^2 + ja_3^2) + x_3^2(ja_1 + ja_1^2 + ja_2^2 + a_3 + ja_3^2), x_1a_2 + x_2(a_1^2 + j^2a_2 + ja_2^2 + ja_3^2) + x_2^2(j^2a_1 + j^2a_1^2 + j^2a_2 + ja_2^2 + ja_3 + j^2a_3^2) + x_3(a_2 + j^2a_2^2) + x_3^2(ja_1 + ja_1^2 + ja_2^2 + a_3 + ja_3^2), x_1a_3 + x_2(a_2^2 + j^2a_3 + j^2a_3^2) + x_2^2(j^2a_1 + j^2a_1^2 + j^2a_2 + j^2a_2^2 + a_3 + ja_3^2) + x_3(a_1^2 + a_2^2 + a_3 + ja_3^2) + x_3^2(j^2a_1 + j^2a_1^2 + j^2a_2^2 + ja_3 + j^2a_3^2))$
- **XXXVIII: (2)** $(x_1a_1 + x_2(a_1 + a_1^2 + ja_2^2 + j^2a_3 + j^2a_3^2) + x_2^2(ja_1 + ja_1^2 + ja_2^2 + j^2a_3^2) + x_3(a_1 + a_1^2 + a_3) + x_3^2(a_1 + a_1^2 + j^2a_2^2 + j^2a_3 + a_3^2), x_1a_2 + x_2(a_1^2 + a_2 + j^2a_2^2) + x_2^2(ja_2 + j^2a_3^2) + x_3(a_2 + j^2a_2^2) + x_3^2(a_2 + ja_3), x_1a_3 + x_2(a_2^2 + a_3 + j^2a_3^2) + x_2^2(ja_3 + ja_3^2) + x_3(a_1^2 + j^2a_2^2 + a_3 + a_3^2) + x_3^2a_3^2)$
- **XXXIX: (2)** $(x_1a_1 + x_2(ja_1 + ja_1^2 + ja_2^2 + j^2a_3 + ja_3^2) + x_2^2(a_2 + a_2^2 + a_3 + a_3^2) + x_3(ja_2^2 + j^2a_3 + ja_3^2) + x_3^2(ja_2 + ja_2^2 + a_3), x_1a_2 + x_2(ja_1 + j^2a_1^2 + a_2^2 + j^2a_3 + j^2a_3^2) + x_2^2(j^2a_2 + j^2a_2^2) + x_3(j^2a_1 + j^2a_1^2 + ja_2 + a_3) + x_3^2(j^2a_2 + a_3 + a_3^2), x_1a_3 + x_2(a_1 + a_1^2 + a_2^2 + ja_3 + a_3^2) + x_2^2(j^2a_2 + ja_2^2 + j^2a_3 + j^2a_3^2) + x_3(a_1 + a_2^2 + a_3^2) + x_3^2(a_2 + j^2a_2^2 + j^2a_3 + ja_3^2))$
- **XL: (2)** $(x_1a_1 + x_2(j^2a_2 + j^2a_3) + x_2^2(ja_1 + ja_1^2 + j^2a_2^2 + ja_3^2) + x_3(j^2a_1 + j^2a_1^2 + ja_2 + a_2^2 + a_3 + j^2a_3^2) + x_3^2(j^2a_1 + j^2a_1^2 + ja_2 + j^2a_2^2 + ja_3 + j^2a_3^2), x_1a_2 + x_2(a_1 + ja_2) + x_2^2(ja_2 + j^2a_3^2) + x_3(a_3 + a_3^2) + x_3^2(j^2a_1 + j^2a_1^2 + ja_2 + j^2a_2^2 + a_3^2), x_1a_3 + x_2ja_3 + x_2^2(a_2^2 + ja_3) + x_3(a_1 + j^2a_2 + ja_2^2 + j^2a_3) + x_3^2(ja_3 + j^2a_3^2))$
- **XLI: (2)** $(x_1a_1 + x_2(j^2a_2 + j^2a_2^2 + a_3 + a_3^2) + x_3(a_1 + a_1^2 + ja_2 + a_3 + ja_3^2), x_1a_2 + x_2(a_1^2 + a_2 + a_2^2 + ja_3) + x_3(ja_2^2 + a_3^2), x_1a_3 + x_2(ja_1 + ja_1^2 + a_2 + j^2a_3 + ja_3^2) + x_3(a_1 + a_3 + j^2a_3^2))$

- **XLII: (2)** $(x_1a_1 + x_2(j^2a_1 + j^2a_1^2 + ja_2 + a_2^2 + j^2a_3 + a_3^2) + x_2^2(j^2a_1 + j^2a_1^2 + j^2a_2^2 + j^2a_3^2) + x_3(ja_1 + ja_1^2 + ja_3 + ja_3^2) + x_3^2(ja_1 + ja_1^2 + a_2 + j^2a_2^2 + a_3^2) , x_1a_2 + x_2(ja_1 + j^2a_1^2 + j^2a_2 + a_2^2 + a_3^2) + x_2^2(ja_1 + ja_1^2 + a_2 + j^2a_2^2) + x_3(ja_2 + j^2a_3 + j^2a_3^2) + x_3^2(a_1 + a_1^2 + ja_2 + j^2a_2^2 + j^2a_3^2) , x_1a_3 + x_2(a_1 + a_1^2 + a_2 + j^2a_3 + j^2a_3^2) + x_2^2j^2a_3 + x_3(a_1^2 + j^2a_3 + j^2a_3^2) + x_3^2(j^2a_1 + j^2a_1^2 + j^2a_2 + j^2a_3 + ja_3^2))$
- **XLIII: (2)** $(x_1a_1 + x_2(j^2a_1 + j^2a_1^2 + a_2 + a_2^2 + ja_3 + ja_3^2) + x_3(a_1 + a_1^2 + ja_2^2 + ja_3 + a_3^2) , x_1a_2 + x_2(a_1 + a_2 + a_2^2 + ja_3 + a_3^2) + x_3(ja_1 + ja_1^2 + a_2^2 + a_3) , x_1a_3 + x_2(j^2a_1 + j^2a_1^2 + ja_2 + j^2a_2^2 + j^2a_3^2) + x_3(j^2a_1 + ja_1^2 + a_2^2 + a_3 + j^2a_3^2))$
- **XLIV: (2)** $(x_1a_1 + x_2(j^2a_1 + j^2a_1^2 + j^2a_2 + a_2^2 + ja_3 + j^2a_3^2) + x_2^2(a_2 + j^2a_3) + x_3(ja_2 + a_3) + x_3^2(a_1 + a_1^2 + a_2 + ja_2^2 + j^2a_3 + a_3^2) , x_1a_2 + x_2(a_1^2 + a_2 + ja_3) + x_2^2(j^2a_1 + j^2a_1^2 + a_2 + j^2a_2^2 + a_3 + j^2a_3^2) + x_3(ja_1 + ja_1^2 + ja_2^2 + ja_3^2) + x_3^2(ja_2 + ja_2^2 + j^2a_3 + ja_3^2) , x_1a_3 + x_2(j^2a_1 + j^2a_1^2 + a_2 + ja_2^2 + a_3 + j^2a_3^2) + x_2^2j^2a_2 + x_3(a_1 + j^2a_3) + x_3^2(a_1 + a_1^2 + ja_2 + j^2a_2^2 + ja_3 + a_3^2))$
- **XLV: (2)** $(x_1a_1 + x_2(ja_1 + ja_1^2 + j^2a_2 + j^2a_2^2 + j^2a_3 + ja_3^2) + x_3(j^2a_2 + a_2^2 + a_3 + a_3^2) , x_1a_2 + x_2(a_1^2 + a_2 + a_2^2 + ja_3) + x_3(a_2^2 + j^2a_3) , x_1a_3 + x_2(a_2 + a_3 + a_3^2) + x_3(a_1^2 + a_2 + a_3 + ja_3^2))$
- **XLVI: (2)** $(x_1a_1 + x_2(ja_1 + ja_1^2 + j^2a_2^2) + x_2^2(j^2a_1 + j^2a_1^2 + j^2a_2^2 + ja_3^2) + x_3(ja_1 + ja_1^2 + j^2a_2^2) + x_3^2(a_2^2 + j^2a_3^2) , x_1a_2 + x_2(a_1^2 + ja_2 + j^2a_3^2) + x_2^2(j^2a_2 + a_2^2 + a_3^2) + x_3(ja_2 + ja_2^2 + ja_3^2) + x_3^2a_2^2 , x_1a_3 + x_2(ja_3 + a_3^2) + x_2^2(a_2^2 + j^2a_3) + x_3(a_1^2 + a_2^2 + ja_3))$
- **XLVII: (2)** $(x_1a_1 + x_2(a_2 + j^2a_3) + x_2^2(a_2^2 + a_3 + ja_3^2) + x_3(a_2^2 + j^2a_3 + ja_3^2) + x_3^2(ja_1 + ja_1^2 + j^2a_2 + ja_3 + ja_3^2) , x_1a_2 + x_2(a_1^2 + ja_2 + ja_2^2 + j^2a_3 + j^2a_3^2) + x_2^2(a_1 + a_1^2 + a_2 + a_2^2 + a_3 + j^2a_3^2) + x_3(a_1 + a_1^2 + j^2a_2 + ja_3 + j^2a_3^2) + x_3^2(a_1 + a_1^2 + a_2^2 + ja_3 + a_3^2) , x_1a_3 + x_2(j^2a_2 + a_2^2 + ja_3 + j^2a_3^2) + x_2^2(a_1 + a_1^2 + ja_2 + a_2^2 + ja_3 + a_3^2) + x_3(a_1 + a_2 + j^2a_2^2 + a_3^2) + x_3^2(a_1 + a_1^2 + a_2 + a_2^2 + j^2a_3 + j^2a_3^2))$
- **XLVIII: (2)** $(x_1a_1 + x_2(a_2^2 + ja_3^2) + x_2^2(a_1 + a_1^2 + ja_2 + j^2a_2^2 + a_3 + ja_3^2) + x_3(ja_2 + a_2^2 + a_3) + x_3^2(ja_1 + ja_1^2 + j^2a_2 + j^2a_2^2 + ja_3) , x_1a_2 + x_2(j^2a_1 + ja_1^2 + ja_2 + ja_2^2 + ja_3 + a_3^2) + x_2^2(j^2a_2 + j^2a_2^2 + a_3 + ja_3^2) + x_3(ja_1 + ja_1^2 + j^2a_2 + j^2a_2^2 + ja_3^2) + x_3^2(ja_2 + a_3 + j^2a_3^2) , x_1a_3 + x_2(a_1 + a_1^2 + ja_2 + a_2^2 + j^2a_3^2) + x_2^2(j^2a_2 + a_2^2 + j^2a_3 + j^2a_3^2) + x_3(j^2a_1 + ja_1^2 + ja_2 + ja_3) + x_3^2(ja_2 + ja_2^2 + a_3))$
- **XLIX: (2)** $(x_1a_1 + x_2(j^2a_1 + j^2a_1^2 + j^2a_2 + a_2^2 + a_3 + ja_3^2) + x_2^2(a_2 + j^2a_2^2 + ja_3^2) + x_3(j^2a_1 + j^2a_1^2 + a_2 + j^2a_3 + ja_3^2) + x_3^2(ja_1 + ja_1^2 + ja_3^2) , x_1a_2 + x_2(j^2a_1 + ja_1^2 + a_2 + j^2a_2^2 + j^2a_3 + ja_3^2) + x_2^2(j^2a_1 + j^2a_1^2 + ja_2 + a_3 + a_3^2) + x_3(a_1 + a_1^2) + x_3^2(ja_1 + ja_1^2 + a_2 + ja_2^2 + j^2a_3) , x_1a_3 + x_2(a_1 + a_1^2 + j^2a_2 + ja_2^2 + j^2a_3^2) + x_2^2(ja_1 + ja_1^2 + j^2a_2 + j^2a_2^2 + ja_3 + ja_3^2) + x_3(a_1^2 + a_2 + a_2^2 + j^2a_3 + a_3^2) + x_3^2(ja_1 + ja_1^2 + ja_2 + ja_3 + ja_3^2))$
- **L: (2)** $(x_1a_1 + x_2(a_2 + j^2a_2^2 + a_3^2) + x_2^2(j^2a_1 + j^2a_1^2 + ja_2) + x_3(j^2a_1 + j^2a_1^2 + a_2 + j^2a_2^2 + j^2a_3^2) + x_3^2(a_1 + a_1^2 + j^2a_3 + j^2a_3^2) , x_1a_2 + x_2(a_1^2 + a_2) + x_2^2(ja_1 + ja_1^2 + a_2 + j^2a_3 + a_3^2) + x_3(j^2a_1 + j^2a_1^2 + ja_2 + ja_2^2 + a_3 + j^2a_3^2) + x_3^2(a_2^2 + a_3 + ja_3^2) , x_1a_3 + x_2(a_1 + a_1^2 + a_2 + a_3 + ja_3^2) + x_2^2(a_3 + j^2a_3^2) + x_3(ja_1 + j^2a_1^2 + ja_2 + a_2^2 + ja_3 + a_3^2) + x_3^2(j^2a_1 + j^2a_1^2 + j^2a_2 + j^2a_3 + j^2a_3^2))$
- **LI: (2)** $(x_1a_1 + x_2(ja_1 + ja_1^2 + a_3 + ja_3^2) + x_2^2(a_2 + a_2^2 + a_3) + x_3(ja_1 + ja_1^2 + ja_2 + a_2^2 + j^2a_3^2) + x_3^2(ja_2 + ja_2^2 + j^2a_3^2) , x_1a_2 + x_2(a_1^2 + ja_2 + j^2a_2^2 + j^2a_3^2) + x_2^2(j^2a_2 + ja_2^2 + j^2a_3^2) + x_3ja_2 + x_3^2(j^2a_2 + ja_2^2 + ja_3) , x_1a_3 + x_2ja_3 + x_2^2(a_2^2 + j^2a_3 + j^2a_3^2) + x_3(a_1^2 + j^2a_2^2 + ja_3 + j^2a_3^2) + x_3^2(ja_2 + j^2a_2^2 + ja_3 + ja_3^2))$
- **LII: (2)** $(x_1a_1 + x_2(a_1 + a_1^2 + j^2a_2 + ja_2^2 + ja_3) + x_2^2(ja_1 + ja_1^2 + j^2a_2 + ja_2^2) + x_3(j^2a_1 + j^2a_1^2 + ja_3 + a_3^2) + x_3^2(ja_1 + ja_1^2 + ja_2^2) , x_1a_2 + x_2(j^2a_1 + ja_1^2 + j^2a_2 + ja_3) + x_2^2(a_1 + a_1^2 + a_2 + ja_2^2 + a_3) + x_3(a_1 + a_1^2 + a_2 + ja_2^2 + ja_3 + a_3^2) + x_3^2(ja_1 + ja_1^2 + ja_2^2 + j^2a_3 + a_3^2) , x_1a_3 + x_2(j^2a_1 + j^2a_1^2 + j^2a_2 + ja_2^2 + ja_3 + a_3^2) + x_2^2(ja_1 + ja_1^2 + ja_2 + ja_2^2 + a_3^2) + x_3(a_1 + ja_2 + ja_2^2 + a_3 + a_3^2) + x_3^2(ja_2 + ja_2^2 + ja_3^2))$

- **LIII: (2)** $(x_1a_1 + x_2(j^2a_1 + j^2a_1^2 + j^2a_2 + j^2a_2^2 + a_3) + x_3(ja_1 + ja_1^2 + ja_2 + ja_3 + ja_3^2)) , x_1a_2 + x_2(ja_1 + j^2a_1^2 + j^2a_2 + j^2a_2^2 + ja_3) + x_3(j^2a_1 + j^2a_1^2 + a_2^2 + j^2a_3^2) , x_1a_3 + x_2(a_1 + a_1^2 + a_2 + ja_3 + a_3^2) + x_3(a_1 + j^2a_2 + j^2a_2^2 + a_3^2))$
- **LIV: (2)** $(x_1a_1 + x_2(j^2a_1 + j^2a_1^2 + a_3) + x_3(a_1 + a_1^2 + ja_2^2 + ja_3 + a_3^2)) , x_1a_2 + x_2(ja_1 + j^2a_1^2 + j^2a_2 + j^2a_2^2 + j^2a_3) + x_3(ja_1 + ja_1^2 + a_2^2 + j^2a_3 + j^2a_3^2) , x_1a_3 + x_2(a_1 + a_1^2 + a_2 + j^2a_3 + a_3^2) + x_3(a_1^2 + j^2a_2 + j^2a_2^2 + a_3 + ja_3^2))$
- **LV: (2)** $(x_1a_1 + x_2(ja_1 + ja_1^2 + ja_2 + j^2a_2^2 + ja_3^2) + x_2^2(a_1 + a_1^2 + j^2a_2 + ja_2^2 + a_3^2) + x_3(a_1 + a_1^2 + ja_2^2 + a_3^2) + x_3^2ja_2 , x_1a_2 + x_2(ja_1 + j^2a_1^2 + ja_2 + a_2^2 + j^2a_3^2) + x_2^2j^2a_2 + x_3(ja_1 + ja_1^2 + j^2a_2^2 + ja_3^2) + x_3^2ja_2 , x_1a_3 + x_2(ja_1 + ja_1^2 + a_2^2 + a_3) + x_2^2(ja_2 + ja_2^2 + a_3 + a_3^2) + x_3(j^2a_1 + ja_1^2 + ja_2 + a_2^2 + j^2a_3) + x_3^2(j^2a_2 + j^2a_2^2))$
- **LVI: (2)** $(x_1a_1 + x_2(ja_2 + j^2a_3 + j^2a_3^2) + x_2^2(ja_2^2 + a_3) + x_3(a_1 + a_1^2 + a_2 + j^2a_3 + j^2a_3^2) + x_3^2(ja_1 + ja_1^2 + j^2a_2 + j^2a_2^2 + a_3 + j^2a_3^2)) , x_1a_2 + x_2(j^2a_1 + ja_1^2 + ja_2 + ja_3 + j^2a_3^2) + x_2^2(ja_1 + ja_1^2 + j^2a_2 + a_2^2 + j^2a_3^2) + x_3(ja_1 + ja_1^2 + a_2^2 + j^2a_3 + j^2a_3^2) + x_3^2(ja_2 + j^2a_2^2 + j^2a_3 + ja_3^2) , x_1a_3 + x_2(j^2a_2 + ja_2^2 + a_3) + x_2^2(a_1 + a_1^2 + ja_2 + ja_2^2 + j^2a_3 + a_3^2) + x_3(j^2a_1 + ja_1^2 + j^2a_2 + a_2^2 + j^2a_3 + a_3^2) + x_3^2(j^2a_2 + ja_2^2 + ja_3))$
- **LVII: (2)** $(x_1a_1 + x_2(j^2a_1 + j^2a_1^2 + j^2a_2 + ja_2^2 + ja_3 + j^2a_3^2) + x_2^2(ja_1 + ja_1^2 + a_2^2 + a_3^2) + x_3(j^2a_2 + ja_2^2 + ja_3 + j^2a_3^2) + x_3^2(ja_1 + ja_1^2 + j^2a_2^2 + a_3 + j^2a_3^2)) , x_1a_2 + x_2(a_1 + j^2a_2 + a_2^2 + j^2a_3 + j^2a_3^2) + x_2^2(a_1 + a_1^2 + j^2a_2 + a_2^2 + a_3^2) + x_3(a_2 + a_2^2 + j^2a_3 + j^2a_3^2) + x_3^2(j^2a_1 + j^2a_1^2 + j^2a_2 + ja_2^2 + ja_3^2) , x_1a_3 + x_2(j^2a_1 + j^2a_1^2 + ja_2 + j^2a_2^2) + x_2^2(ja_2 + ja_2^2 + j^2a_3) + x_3(j^2a_1 + ja_1^2 + ja_3) + x_3^2(j^2a_2 + ja_3 + j^2a_3^2))$
- **LVIII: (2)** $(x_1a_1 + x_2(ja_2 + ja_2^2 + j^2a_3 + ja_3^2) + x_3(a_1 + a_1^2 + a_2 + j^2a_3 + a_3^2)) , x_1a_2 + x_2(a_1 + ja_2 + ja_2^2 + ja_3^2) + x_3(a_1 + a_1^2 + ja_2 + ja_2^2 + ja_3 + ja_3^2) , x_1a_3 + x_2(j^2a_1 + j^2a_1^2 + a_2 + j^2a_3^2) + x_3(ja_1 + j^2a_1^2 + a_2 + j^2a_2^2 + ja_3 + ja_3^2))$
- **LIX: (2)** $(x_1a_1 + x_2(j^2a_1 + j^2a_1^2 + j^2a_2 + a_2^2 + a_3 + ja_3^2) + x_2^2(j^2a_1 + j^2a_1^2 + ja_2^2 + j^2a_3^2) + x_3(j^2a_1 + j^2a_1^2 + ja_2 + j^2a_2^2 + j^2a_3 + ja_3^2) + x_3^2(ja_2 + j^2a_2^2 + j^2a_3) , x_1a_2 + x_2(j^2a_1 + ja_1^2 + j^2a_2^2 + ja_3 + j^2a_3^2) + x_2^2(ja_1 + ja_1^2 + j^2a_2^2 + j^2a_3) + x_3(ja_1 + ja_1^2 + j^2a_2) + x_3^2(j^2a_1 + j^2a_1^2 + j^2a_2^2 + a_3) , x_1a_3 + x_2(j^2a_1 + j^2a_1^2 + a_2 + ja_2^2 + j^2a_3 + a_3^2) + x_2^2(j^2a_1 + j^2a_1^2 + a_2 + j^2a_2^2) + x_3(a_1^2 + a_2 + ja_2^2 + a_3 + ja_3^2) + x_3^2(ja_1 + ja_1^2 + j^2a_2 + ja_2^2 + ja_3 + ja_3^2))$
- **LX: (2)** $(x_1a_1 + x_2(a_1 + a_1^2 + ja_2 + j^2a_2^2 + j^2a_3 + a_3^2) + x_2^2(ja_1 + ja_1^2 + ja_2 + j^2a_2^2 + a_3 + a_3^2) + x_3(a_1 + a_1^2 + j^2a_2 + a_2^2 + j^2a_3) + x_3^2(j^2a_1 + j^2a_1^2 + ja_2^2 + ja_3 + a_3^2)) , x_1a_2 + x_2(ja_1 + j^2a_1^2 + a_2^2 + ja_3) + x_2^2(j^2a_1 + j^2a_1^2 + a_2 + j^2a_3 + j^2a_3^2) + x_3(j^2a_1 + j^2a_1^2 + j^2a_2^2 + ja_3 + a_3^2) + x_3^2(j^2a_1 + j^2a_1^2 + ja_2 + ja_2^2 + a_3^2) , x_1a_3 + x_2(j^2a_1 + j^2a_1^2 + j^2a_2 + a_2^2 + j^2a_3 + ja_3^2) + x_2^2(j^2a_1 + j^2a_1^2 + ja_2 + a_2^2 + ja_3 + a_3^2) + x_3(ja_1 + j^2a_1^2 + ja_2 + j^2a_2^2 + ja_3 + j^2a_3^2) + x_3^2(j^2a_1 + j^2a_1^2 + j^2a_2 + a_3))$
- **LXI: (2)** $(x_1a_1 + x_2(a_2 + ja_2^2 + j^2a_3 + ja_3^2) + x_2^2(ja_1 + ja_1^2 + j^2a_2 + a_3 + ja_3^2) + x_3(j^2a_1 + j^2a_1^2 + j^2a_2 + j^2a_2^2 + j^2a_3) + x_3^2(j^2a_1 + j^2a_1^2 + ja_2 + a_2^2 + ja_3^2) , x_1a_2 + x_2(a_1^2 + ja_2 + ja_2^2 + ja_3^2) + x_2^2(a_1 + a_1^2 + ja_2 + ja_2^2 + j^2a_3) + x_3(a_2 + ja_3 + j^2a_3^2) + x_3^2(a_1 + a_1^2 + j^2a_2 + j^2a_3 + j^2a_3^2) , x_1a_3 + x_2(ja_2 + ja_2^2 + a_3^2) + x_2^2(j^2a_1 + j^2a_1^2 + j^2a_2 + ja_2^2 + a_3^2) + x_3(j^2a_1 + ja_1^2 + a_2 + a_3 + j^2a_3^2) + x_3^2(a_3 + ja_3^2))$
- **LXII: (2)** $(x_1a_1 + x_2(ja_1 + ja_1^2 + a_2^2 + a_3^2) + x_2^2(ja_1 + ja_1^2 + ja_2 + j^2a_2^2 + ja_3 + a_3^2) + x_3(a_1 + a_1^2 + a_2 + a_3 + a_3^2) + x_3^2(ja_1 + ja_1^2 + j^2a_2 + j^2a_3 + a_3^2)) , x_1a_2 + x_2(a_1^2 + ja_2 + ja_2^2) + x_2^2(a_1 + a_1^2 + a_2 + a_2^2 + a_3) + x_3(j^2a_1 + j^2a_1^2 + j^2a_2 + a_2^2 + j^2a_3 + j^2a_3^2) + x_3^2(ja_1 + ja_1^2 + j^2a_2 + ja_2^2 + ja_3 + a_3^2) , x_1a_3 + x_2(a_2^2 + ja_3 + j^2a_3^2) + x_2^2(ja_1 + ja_1^2 + ja_2 + ja_2^2 + ja_3 + ja_3^2) + x_3(a_1 + a_2 + j^2a_2^2 + a_3 + ja_3^2) + x_3^2(j^2a_1 + j^2a_1^2 + j^2a_2 + j^2a_2^2 + ja_3 + a_3^2))$
- **LXIII: (2)** $(x_1a_1 + x_2(a_1 + a_1^2 + a_2 + a_2^2 + ja_3 + ja_3^2) + x_3(ja_2 + a_2^2 + ja_3 + j^2a_3^2) , x_1a_2 + x_2(a_1 + a_2 + a_2^2 + ja_3) + x_3(ja_1 + ja_1^2 + j^2a_2 + a_2^2) , x_1a_3 + x_2(a_1 + a_1^2 + a_2^2 + ja_3 + j^2a_3^2) + x_3(j^2a_1 + ja_1^2 + j^2a_2 + j^2a_3))$

- **LXIV: (2)** $(x_1a_1 + x_2(ja_1 + ja_1^2 + j^2a_2 + j^2a_3 + ja_3^2) + x_2^2(a_1 + a_1^2 + j^2a_2^2 + j^2a_3^2) + x_3(a_1 + a_1^2 + a_2 + a_3 + j^2a_3^2) + x_3^2(a_1 + a_1^2 + j^2a_2 + ja_2^2 + j^2a_3) , x_1a_2 + x_2(a_1^2 + ja_2^2) + x_2^2(j^2a_1 + j^2a_1^2 + ja_2 + ja_3 + a_3^2) + x_3(a_1 + a_1^2 + ja_2^2) + x_3^2(j^2a_2 + j^2a_2^2 + ja_3) , x_1a_3 + x_2(a_2 + j^2a_2^2 + a_3 + a_3^2) + x_2^2(j^2a_1 + j^2a_1^2 + j^2a_2 + j^2a_3 + a_3^2) + x_3(j^2a_1 + ja_1^2 + ja_2 + ja_3 + j^2a_3^2) + x_3^2(a_2 + a_2^2 + ja_3^2))$
- **LXV: (2)** $(x_1a_1 + x_2(a_2 + j^2a_2^2 + ja_3 + ja_3^2) + x_2^2(j^2a_2 + a_2^2 + ja_3^2) + x_3(a_1 + a_1^2 + a_2 + j^2a_3^2) + x_3^2(j^2a_1 + j^2a_1^2 + ja_2 + j^2a_2^2 + a_3) , x_1a_2 + x_2(a_1 + a_2 + ja_2^2 + a_3 + a_3^2) + x_2^2(ja_2 + a_2^2) + x_3(j^2a_1 + j^2a_1^2 + j^2a_2 + j^2a_2^2) + x_3^2(j^2a_1 + j^2a_1^2 + j^2a_2 + ja_2^2 + j^2a_3) , x_1a_3 + x_2(j^2a_1 + j^2a_1^2 + ja_2^2 + ja_3) + x_2^2(j^2a_1 + j^2a_1^2 + a_2 + ja_2^2 + ja_3 + a_3^2) + x_3(ja_1 + j^2a_1^2 + ja_2^2 + ja_3^2) + x_3^2(j^2a_1 + j^2a_1^2 + ja_2 + j^2a_2^2 + a_3 + a_3^2))$
- **LXVI: (2)** $(x_1a_1 + x_2(ja_2^2 + ja_3 + a_3^2) + x_2^2(ja_1 + ja_1^2 + j^2a_2 + a_2^2 + j^2a_3) + x_3(a_1 + a_1^2 + a_2 + j^2a_2^2 + a_3) + x_3^2(ja_1 + ja_1^2 + ja_2 + ja_2^2 + a_3 + ja_3^2) , x_1a_2 + x_2(a_1^2 + ja_2^2 + a_3) + x_2^2(ja_2 + ja_3 + a_3^2) + x_3(ja_1 + ja_1^2 + j^2a_3 + j^2a_3^2) + x_3^2(ja_1 + ja_1^2 + a_2^2 + j^2a_3 + a_3^2) , x_1a_3 + x_2(j^2a_1 + j^2a_1^2 + ja_2^2 + j^2a_3 + a_3^2) + x_2^2(j^2a_1 + j^2a_1^2 + a_2 + ja_2^2 + a_3 + a_3^2) + x_3(ja_1 + j^2a_1^2 + ja_2 + j^2a_2^2 + a_3 + j^2a_3^2) + x_3^2(j^2a_1 + j^2a_1^2 + j^2a_2 + j^2a_3 + j^2a_3^2))$
- **LXVII: (2)** $(x_1a_1 + x_2(ja_1 + ja_1^2 + ja_2 + ja_2^2 + a_3^2) + x_3(a_1 + a_1^2 + j^2a_2 + ja_2^2 + j^2a_3 + j^2a_3^2) , x_1a_2 + x_2(a_1 + j^2a_2 + j^2a_2^2 + ja_3 + j^2a_3^2) + x_3(j^2a_1 + j^2a_1^2 + ja_2 + j^2a_3^2) , x_1a_3 + x_2(a_2 + ja_3) + x_3(a_1^2 + a_2 + ja_2^2 + ja_3 + j^2a_3^2))$
- **LXVIII: (2)** $(x_1a_1 + x_2(j^2a_1 + j^2a_1^2 + ja_2 + ja_2^2 + a_3 + a_3^2) + x_3(a_1 + a_1^2 + a_2 + j^2a_2^2 + ja_3 + a_3^2) , x_1a_2 + x_2(ja_1 + j^2a_1^2 + a_2 + a_2^2 + ja_3^2) + x_3(j^2a_1 + j^2a_1^2 + a_2^2 + ja_3^2) , x_1a_3 + x_2(a_1 + a_1^2 + a_2^2 + ja_3^2) + x_3(a_1 + a_2 + a_2^2))$
- **LXIX: (2)** $(x_1a_1 + x_2(j^2a_1 + j^2a_1^2 + a_2 + a_2^2 + ja_3 + ja_3^2) + x_3(a_1 + a_1^2 + j^2a_2 + a_2^2 + ja_3 + a_3^2) , x_1a_2 + x_2(a_1^2 + a_2 + a_2^2 + j^2a_3) + x_3(a_1 + a_1^2 + j^2a_2 + j^2a_2^2 + a_3 + ja_3^2) , x_1a_3 + x_2(ja_1 + ja_1^2 + ja_2 + j^2a_2^2 + j^2a_3^2) + x_3(ja_1 + j^2a_1^2 + a_2^2 + j^2a_3 + ja_3^2))$
- **LXX: (2)** $(x_1a_1 + x_2(ja_1 + ja_1^2 + j^2a_2 + j^2a_3^2) + x_2^2(ja_1 + ja_1^2 + j^2a_2 + a_3^2) + x_3(a_1 + a_1^2 + j^2a_2) + x_3^2(ja_1 + ja_1^2 + a_2 + ja_3^2) , x_1a_2 + x_2(j^2a_1 + ja_1^2 + j^2a_2 + ja_2^2 + j^2a_3 + j^2a_3^2) + x_2^2(a_1 + a_1^2 + a_2 + a_3 + j^2a_3^2) + x_3(ja_1 + ja_1^2 + a_2 + j^2a_2^2 + ja_3 + ja_3^2) + x_3^2(a_2 + j^2a_2^2 + j^2a_3^2) , x_1a_3 + x_2(ja_2^2 + ja_3 + j^2a_3^2) + x_2^2(j^2a_1 + j^2a_1^2 + a_2 + ja_2^2 + ja_3 + j^2a_3^2) + x_3(ja_1 + j^2a_1^2 + j^2a_2 + a_2^2 + a_3 + ja_3^2) + x_3^2(ja_1 + ja_1^2 + j^2a_2 + ja_2^2 + ja_3 + j^2a_3^2))$
- **LXXI: (3)** $(x_1(j^5a_1 + j^{24}a_1^2 + j^{43}a_1^4 + j^{18}a_1^8 + j^{58}a_1^{16} + j^{43}a_1^{32}) + x_1^2(j^2a_1 + j^{62}a_1^2 + j^{45}a_1^4 + j^{53}a_1^8 + j^{14}a_1^{16} + ja_1^{32}) + x_1^4(j^{16}a_1 + j^{16}a_1^2 + j^{45}a_1^4 + j^{49}a_1^8 + j^{50}a_1^{16} + j^{43}a_1^{32}) + x_1^8(j^{50}a_1 + j^{42}a_1^4 + j^4a_1^8 + j^{57}a_1^{16} + j^5a_1^{32}) + x_1^{16}(j^{40}a_1 + j^{16}a_1^2 + j^{38}a_1^4 + j^{18}a_1^8 + j^{17}a_1^{16} + j^{27}a_1^{32}) + x_1^{32}(j^{44}a_1 + j^{10}a_1^2 + j^{36}a_1^4 + j^5a_1^8 + j^{26}a_1^{16} + j^{37}a_1^{32}))$
- **LXXII: (2)** $(x_1a_1 + x_2(ja_2 + a_2^2 + ja_3 + j^2a_3^2) + x_2^2(a_2 + ja_2^2 + ja_3 + a_3^2) + x_3(a_1 + a_1^2 + j^2a_2 + j^2a_2^2 + ja_3 + a_3^2) + x_3^2(ja_2 + a_2^2 + ja_3 + ja_3^2) , x_1a_2 + x_2(a_1 + ja_2 + j^2a_2^2 + a_3 + ja_3^2) + x_2^2(a_2^2 + a_3) + x_3(ja_2 + a_3) + x_3^2(a_1 + a_1^2 + j^2a_2^2 + ja_3 + j^2a_3^2) , x_1a_3 + x_2(a_2 + ja_3) + x_3(a_1^2 + ja_2 + j^2a_2^2 + a_3 + a_3^2) + x_3^2(a_2 + ja_3 + a_3^2))$
- **LXXIII: (3)** $(x_1(j^{57}a_1 + j^{16}a_1^2 + j^{50}a_1^4 + j^{37}a_1^8 + j^{39}a_1^{16} + ja_1^{32}) + x_1^2(j^{29}a_1 + j^{38}a_1^2 + j^8a_1^4 + j^{30}a_1^8 + j^{35}a_1^{16} + j^{54}a_1^{32}) + x_1^4(j^{15}a_1 + j^4a_1^2 + j^{26}a_1^4 + j^{32}a_1^8 + j^{39}a_1^{16}) + x_1^8(j^{56}a_1 + j^{37}a_1^2 + j^{23}a_1^4 + j^{35}a_1^8 + j^6a_1^{16} + j^{33}a_1^{32}) + x_1^{16}(j^4a_1^2 + j^{55}a_1^4 + j^4a_1^8 + j^4a_1^{16} + j^{57}a_1^{32}) + x_1^{32}(j^{10}a_1 + j^{34}a_1^2 + j^{29}a_1^4 + j^{48}a_1^8 + j^{55}a_1^{16} + j^{18}a_1^{32}))$
- **LXXIV: (3)** $(x_1(j^{21}a_1 + j^{46}a_1^2 + j^{25}a_1^4 + j^{53}a_1^8 + j^{44}a_1^{16} + j^{44}a_1^{32}) + x_1^2(j^{28}a_1 + j^{26}a_1^2 + j^{10}a_1^4 + j^{61}a_1^8 + j^{34}a_1^{16} + j^{57}a_1^{32}) + x_1^4(j^{10}a_1 + j^{18}a_1^2 + j^{57}a_1^4 + j^{48}a_1^8 + j^{33}a_1^{16} + j^{19}a_1^{32}) + x_1^8(j^{23}a_1 + j^{44}a_1^2 + j^{37}a_1^4 + j^7a_1^8 + j^{40}a_1^{16} + a_1^{32}) + x_1^{16}(j^6a_1 + j^{14}a_1^2 + j^{48}a_1^4 + j^{17}a_1^8 + j^{57}a_1^{16} + ja_1^{32}) + x_1^{32}(j^{20}a_1 + j^6a_1^2 + j^{48}a_1^4 + j^{39}a_1^8 + j^{13}a_1^{16} + j^{27}a_1^{32}))$

- **LXXV: (3)** $(x_1(j^{53}a_1 + j^{33}a_1^2 + j^{14}a_1^4 + j^{18}a_1^8 + j^{29}a_1^{16} + j^{49}a_1^{32}) + x_1^2(j^{22}a_1 + j^{30}a_1^2 + j^{50}a_1^4 + j^{53}a_1^8 + j^{52}a_1^{16} + j^{48}a_1^{32}) + x_1^4(j^{31}a_1 + j^2a_1^2 + j^{47}a_1^4 + j^{48}a_1^8 + j^{10}a_1^{16} + j^{21}a_1^{32}) + x_1^8(j^{18}a_1 + j^{25}a_1^2 + j^{26}a_1^4 + a_1^8 + j^{30}a_1^{16} + j^9a_1^{32}) + x_1^{16}(j^{16}a_1 + j^{52}a_1^2 + j^{29}a_1^4 + j^{10}a_1^8 + j^{38}a_1^{16} + j^{52}a_1^{32}) + x_1^{32}(j^{47}a_1 + j^{33}a_1^2 + j^{29}a_1^4 + j^{45}a_1^8 + j^{22}a_1^{16} + j^{62}a_1^{32}))$
- **LXXVI: (2)** $(x_1a_1 + x_2(j^2a_1 + j^2a_1^2 + ja_2 + a_2^2 + a_3 + j^2a_3^2) + x_2^2(a_1 + a_1^2 + a_2 + ja_2^2 + j^2a_3) + x_3(a_1 + a_1^2 + j^2a_2^2) + x_3^2(ja_1 + ja_1^2 + ja_3^2) , x_1a_2 + x_2(a_1^2 + j^2a_2 + j^2a_3 + ja_3^2) + x_2^2(a_2 + ja_2^2 + ja_3 + a_3^2) + x_3(a_2 + j^2a_2^2) + x_3^2(ja_2 + j^2a_2^2 + a_3^2) , x_1a_3 + x_2(ja_1 + ja_1^2 + j^2a_2 + a_2^2 + j^2a_3 + a_3^2) + x_2^2(a_1 + a_1^2 + ja_2 + a_2^2 + a_3 + j^2a_3^2) + x_3(a_1^2 + a_3) + x_3^2(ja_2^2 + ja_3 + a_3^2))$
- **LXXVII: (3)** $(x_1(j^{53}a_1 + j^{38}a_1^2 + j^5a_1^4 + j^{39}a_1^8 + j^7a_1^{16} + j^5a_1^{32}) + x_1^2(j^{14}a_1 + j^{26}a_1^2 + j^{17}a_1^4 + j^{27}a_1^8 + j^6a_1^{16} + j^{29}a_1^{32}) + x_1^4(j^{38}a_1 + j^{44}a_1^2 + j^{42}a_1^4 + j^{48}a_1^8 + j^{19}a_1^{16} + j^{56}a_1^{32}) + x_1^8(j^{30}a_1 + j^7a_1^2 + j^{10}a_1^4 + j^{26}a_1^8 + j^7a_1^{16} + j^2a_1^{32}) + x_1^{16}(j^{15}a_1 + j^{28}a_1^2 + j^{34}a_1^4 + j^{61}a_1^8 + j^{55}a_1^{16} + j^{18}a_1^{32}) + x_1^{32}(j^{58}a_1 + j^{51}a_1^2 + j^{15}a_1^4 + j^6a_1^8 + j^{30}a_1^{16} + j^8a_1^{32}))$
- **LXXVIII: (3)** $(x_1(j^{27}a_1 + j^{10}a_1^2 + j^{39}a_1^4 + j^{13}a_1^8 + j^{22}a_1^{16} + j^{39}a_1^{32}) + x_1^2(j^4a_1 + j^{48}a_1^2 + j^{14}a_1^4 + j^{23}a_1^8 + j^{47}a_1^{16} + j^{26}a_1^{32}) + x_1^4(j^{11}a_1 + j^{45}a_1^2 + j^{35}a_1^4 + j^{38}a_1^8 + j^{30}a_1^{16} + j^{45}a_1^{32}) + x_1^8(j^{49}a_1 + ja_1^2 + j^{11}a_1^4 + j^4a_1^8 + j^{58}a_1^{16} + j^{14}a_1^{32}) + x_1^{16}(j^{49}a_1 + j^{48}a_1^2 + j^{34}a_1^4 + j^{18}a_1^8 + j^{46}a_1^{16} + j^{27}a_1^{32}) + x_1^{32}(j^{20}a_1 + j^{51}a_1^2 + j^{55}a_1^4 + j^{47}a_1^8 + j^{37}a_1^{16} + j^{46}a_1^{32}))$
- **LXXIX: (3)** $(x_1(j^{31}a_1 + j^{32}a_1^2 + j^{29}a_1^4 + j^{57}a_1^8 + j^{26}a_1^{16} + j^{37}a_1^{32}) + x_1^2(j^{62}a_1 + j^{40}a_1^2 + j^{55}a_1^4 + j^3a_1^8 + j^{15}a_1^{16}) + x_1^4(j^{31}a_1 + j^3a_1^2 + j^{60}a_1^4 + j^{29}a_1^8 + j^{16}a_1^{16} + j^{58}a_1^{32}) + x_1^8(j^9a_1 + j^{52}a_1^2 + j^{54}a_1^4 + j^5a_1^{16} + j^{42}a_1^{32}) + x_1^{16}(j^{51}a_1 + ja_1^2 + j^{43}a_1^4 + j^{32}a_1^8 + j^{34}a_1^{16} + j^{30}a_1^{32}) + x_1^{32}(j^{16}a_1 + j^{60}a_1^2 + j^{18}a_1^4 + j^{26}a_1^8 + j^{29}a_1^{16} + j^{49}a_1^{32}))$
- **LXXX: (3)** $(x_1(j^{41}a_1 + j^{24}a_1^2 + j^{59}a_1^4 + j^8a_1^8 + j^{43}a_1^{16} + j^{35}a_1^{32}) + x_1^2(j^{34}a_1 + j^{39}a_1^2 + j^{13}a_1^4 + j^{35}a_1^8 + j^{27}a_1^{16} + j^{41}a_1^{32}) + x_1^4(j^{18}a_1 + j^{26}a_1^2 + j^2a_1^4 + a_1^8 + j^2a_1^{16} + j^{32}a_1^{32}) + x_1^8(j^{50}a_1 + j^{34}a_1^2 + j^{29}a_1^4 + j^6a_1^8 + j^{40}a_1^{16} + j^2a_1^{32}) + x_1^{16}(j^{52}a_1 + j^{28}a_1^2 + j^{21}a_1^4 + j^{48}a_1^8 + j^4a_1^{16} + j^{40}a_1^{32}) + x_1^{32}(j^{36}a_1 + j^9a_1^2 + j^{10}a_1^4 + j^3a_1^8 + ja_1^{16} + j^{50}a_1^{32}))$

References

- [1] D.E. Knuth, *Finite semifields and projective planes*, Journal of Algebra **2** (1965), 182-217.
- [2] M. Cordero, G. P. Wene, *A survey of finite semifields*, Discrete Mathematics **208/209** (1999), 125-137.
- [3] M. Hall(Jr.), *The theory of groups*, Macmillan, (1959).
- [4] A. R. Calderbank, P. J. Cameron, W. M. Kantor, J. J. Seidel, \mathbb{Z}_4 -Kerdock codes, orthogonal spreads, and extremal Euclidean line-sets, Proc. London Math. Soc **75** (1997), 436–480.
- [5] W. M. Kantor, M. E. Williams, *Symplectic semifield planes and \mathbb{Z}_4 -linear codes*, Transactions of the American Mathematical Society **356** (2004), 895–938.
- [6] S. González, C. Martínez, I.F. Rúa, *Symplectic Spread based Generalized Kerdock Codes*, Designs, Codes and Cryptography **42 (2)** (2007), 213–226.
- [7] J. P. May, D. Saunders, Z. Wan, *Efficient Matrix Rank Computation with Applications to the Study of Strongly Regular Graphs*, Proceedings of ISSAC 2007, 277-284, ACM, New-York, 2007

- [8] R. J. Walker, *Determination of division algebras with 32 elements*, Proceedings in Symposia of Applied Mathematics **75** (1962), 83-85.
- [9] U. Dempwolff, *Semifield Planes of Order 81*, J. of Geometry (to appear).
- [10] A. A. Albert, *Finite division algebras and finite planes*, Proceedings of Symposia in Applied Mathematics **10** (1960), 53-70.
- [11] L. E. Dickson, *Linear algebras in which division is always uniquely possible*, Transactions of the American Mathematical Society **7** (1906), 370-390.
- [12] A. A. Albert, *On nonassociative division algebras*, Transactions of the American Mathematical Society **72** (1952), 296-309.
- [13] A. A. Albert, *Finite noncommutative division algebras*, Proceedings of the American Mathematical Society **9** (1958), 928-932.
- [14] A. A. Albert, *Generalized twisted fields*, Pacific Journal of Mathematics **11** (1961), 1-8.
- [15] J. H. M. Wedderburn, *A theorem on finite algebras*, Transactions of the American Mathematical Society **6** (1905), 349-352.
- [16] W. M. Kantor, *Commutative semifields and symplectic spreads*, Journal of Algebra **270** (2003), 96-114.
- [17] I.R. Hentzel, I. F. Rúa, *Primitivity of Finite Semifields with 64 and 81 elements*, International Journal of Algebra and Computation **17** (7) (2007), 1411-1429.
- [18] G. P. Wene, *On the multiplicative structure of finite division rings*, Aequationes Mathematicae **41** (1991), 222-233.
- [19] W. M. Kantor, *Finite semifields*, Finite Geometries, Groups, and Computation (Proc. of Conf. at Pingree Park, CO Sept. 2005), de Gruyter, Berlin-New York (2006).
- [20] H. Huang, N.L. Johnson, *8 semifield planes of order 8^2* , Discrete Mathematics **80** (1990), 69-79.
- [21] Menichetti, G. *Algebre Tridimensionali su un campo di Galois*. Ann. Mat. Pura Appl. **1973**, 97, 293-302.
- [22] Menichetti, G. *On a Kaplansky conjecture concerning three-dimensional division algebras over a finite field*. Journal of Algebra **1977**, 47, 400-410.
- [23] Kleinfeld, E. *Techniques for enumerating Veblen-Wedderburn systems*. J. Assoc. Comp. Mach. **1960**, 7, 330-337.
- [24] Oyama, T. *On Quasifields*. Osaka J. Math. **1985**, 22, 35-54.