

A new look at Bell's inequalities and Nelson's theorem

Benjamin Schulz *

January 31, 2021

Abstract

In 1985, Edward Nelson, who is known for the foundation of a theory called „stochastic mechanics“, made an interesting remark about Bell's theorem. Nelson analysed the theorem in the light of classical fields, which behave randomly. He found that if a stochastic hidden variable theory fulfils certain conditions, the inequalities of John Bell could be violated. Moreover, Nelson was able to prove that this could happen without any instantaneous communication between the two spatially separated measurement stations. Since Nelson's paper got almost overlooked by physicists, we try to review his comments on the theorem. We argue, that a modification of Nelson's stochastic mechanics, published recently in „Annalen der Physik“, can be extended to a theory which fulfils the requirements from Nelson's analysis. The article proceeds to derive the quantum mechanical formalism of spinning particles and the Pauli equation from the said formalism. Then, we investigate Bohm's version of the EPR experiment. Additionally, other setups like entanglement swapping or time and position correlations are shortly explained from the viewpoint of our local model. Finally, we mention, that the theory could perhaps be relativistically extended and useful for the formulation of quantum mechanics in curved space-times.

Contents

1	Introduction	2
2	Review of mathematical probability theory	3
2.1	Probability spaces	3
2.2	Random variables and stochastic processes	4
2.3	Enlarged probability spaces	5
3	Nelson's analysis of Bell's theorem	6
3.1	The setup of the EPR experiment in theory	6
3.2	Active locality	8
3.3	Passive locality	9
3.4	Nelson's theorem	9
4	What the failure of passive locality implies - a deeper analysis	11
5	The locality problem in the hidden variable theory of Fritsche and Haugk	12
5.1	Derivation of the single particle Schrödinger equation from Brownian motion	12
5.2	Quantum states in superpositions	15
5.3	Locality in the theory of Fritsche and Haugk with single particle systems	17
5.4	Derivation of the many particle Schrödinger equation	18
5.5	Locality in the theory of Fritsche and Haugk with many particle systems	19
5.6	Entangled states in the model of Fritsche and Haugk	20

*Department für Physik, Ludwig-Maximilians-Universität München, Theresienstrasse 37, 80333 München, Germany, Benjamin.Schulz@physik.uni-muenchen.de

6 Spin in a Fritsche-Haugk like hidden variable theory	21
6.1 Simple estimations	21
6.2 The Stern-Gerlach effect	22
6.3 The Pauli equation	23
6.4 The spin operator	25
7 Entangled systems with spin	26
7.1 Transformation of the entangled state into equations of motion	26
7.2 How the correlations of entangled states emerge in the local hidden variable theory	28
8 Objections against Nelson’s theorem	31
8.1 The argument of Ballentine and Zeilinger	31
8.2 The argument of Dürr	32
8.3 The argument of Redhead, stability against disturbances	33
9 Conclusions and outlook	34
9.1 What has been achieved	34
9.2 Possibility of relativistic extension	34
9.3 Problems of standard quantum mechanics in curved space-times, which might be absent in the model of Fritsche and Haugk	35

1 Introduction

In 1935, Einstein, Podolsky and Rosen (EPR) wrote an article, in which they denied that quantum theory would be a complete theory of nature [1]. Around 1951, Bohm gave a more testable outline of the so called “EPR paradox” [2,3]. He described a thought experiment with one source that ejects particles having opposite spin to two separate Stern-Gerlach magnets of variable orientations. Then, in 1964 John S. Bell published a theorem about this paradox in the form of an inequality. It made clear, that hidden variable theories which fulfil certain conditions would be in difference with quantum mechanics. Bell called these conditions “locally causal” and the most detailed explanation of this term was made by him in [7].

In his first publication about that topic, Bell wrote: “If hidden parameters would be added to quantum mechanics, there must be a mechanism, whereby the setting of one measuring device can influence another spatially separated device”, and the signal involved has to “propagate instantaneously” [4]. However, Bell’s first contribution underwent several modifications. Over the years, more and more instructive proofs of his inequality were constructed by him and others. A collection with all of Bell’s fundamental articles can be found in [5]. Finally in 1969, the inequality was brought by Clauser, Holt, Shimony and Horne (CHSH) into a form, suitable for experimental investigation [6].

In 1985, Princeton mathematician Edward Nelson tried to analyse Bell’s theorem with full mathematical rigour. The result of his analysis was, that Bell’s definition of “locally causal” which is the starting point to derive Bell’s inequalities, could be divided into two separate conditions. Both were necessary to derive the inequalities, but only one of them has to hold, if a hidden parameter theory should not include instantaneous signals. Nelson published his result twice [8,9] (with a small correction in [10] to make the theorem compatible with Mermin’s presentation [11]) and there exist works from mathematicians e.g. [12], giving further insightful analysis of it. Unfortunately, Nelson’s articles got almost overlooked by physicists. The reason might be, that they were written in the language of advanced mathematical probability theory. Another reason was, that Edward Nelson found his own interpretation of quantum mechanics [17, 18] to be in difference with the requirements of his theorem for a model without instantaneous signalling effects.

Since the author is aware, that physicists are sometimes not trained in mathematical probability theory, this article begins with an overview of the necessary mathematical concepts in section 2. After that, we review Nelson’s contribution towards a mathematically rigorous understanding of Bell’s inequalities in section 3. The physical implications of that analysis are discussed in section 4. A recent modification of Nelson’s stochastic mechanics, published in “Annalen der Physik” [19], is the topic of section 5. There it is shown, that this theory fulfils Nelson’s requirements for a local hidden variable model which is compatible with quantum mechanics. In section 6, we extend this theory to include spinning particles described by the Pauli equation. Entangled spin states are discussed in section 7, where it is shown in detail how their correlations emerge without any instantaneous signalling effects. In section 8, we review some arguments

against Nelson's thoughts about Bell's theorem which have occurred in the literature. We come to the conclusion that these arguments are unfounded. Finally, we mention the possible use of our theory for a reasonable foundation of quantum mechanics in curved space-times at section 9.

2 Review of mathematical probability theory

2.1 Probability spaces

Probabilistic experiments are described with a so called probability space (Ω, \mathcal{F}, P) . The set Ω consists of the theoretically possible outcomes $\omega \in \Omega$. Sets of outcomes are called events. The latter can be combined, to form new events by intersection, union and complement. If A is an event, then the complementary event $\complement A$ denotes the set of all outcomes not in A . If A and B are events, the event $A \cap B$ defines the outcomes, which are in both A and B . Similarly, the event $A \cup B$ consists of all outcomes, which are in A or in B .

Often, one wants to consider a collection of events that may be freely combined by these operations. Therefore, we define the terminology of a sigma algebra \mathcal{F} on Ω . Whenever new events are formed by intersection, union or complement, the resulting event is still in \mathcal{F} . A sigma algebra \mathcal{F} is determined by the following properties:

- The set $\Omega \in \mathcal{F}$
- For all sets $A \in \mathcal{F} \Rightarrow \complement A \in \mathcal{F}$
- For every sequence $(A_n)_{n \in \mathbb{N}}$ of sets in \mathcal{F} , the union $\bigcup_n A_n$ is in \mathcal{F} .

Every system γ of subsets from Ω determines a so called smallest sigma algebra $\sigma(\gamma)$ on Ω . This sigma algebra is defined by:

- $\gamma \subseteq \sigma(\gamma)$
- For all sigma algebras \mathcal{F} on Ω , where $\gamma \subseteq \mathcal{F}$, we have $\sigma(\gamma) \subseteq \mathcal{F}$

We say, that $\sigma(\gamma)$ is generated by γ .

For example, if γ is just one set $\gamma \subseteq \Omega$, we have $\sigma(\gamma) = \{\Omega, \emptyset, \gamma, \complement \gamma\}$.

A second example may be given with the experiment of tossing two coins that come up heads $\{H\}$ or tails $\{T\}$. The corresponding set of theoretically possible outcomes is $\Omega = \{HH, HT, TH, TT\}$. We can define γ as a partition of Ω into 4 events $\{HH\}$, $\{HT\}$, $\{TH\}$ and $\{TT\}$. Accordingly, the sigma algebra $\sigma(\gamma)$ consists of the sure event Ω , the impossible event \emptyset , the events in the partition and the 10 possible different unions of them. Another system of subsets γ' would be a partition of Ω into three events $\{HH\}$, $\{HT \cup TH\}$ and $\{TT\}$. Similarly, the sigma algebra $\sigma(\gamma')$ generated by γ' consists of Ω , \emptyset , the three events in the partition and their three possible different unions. We observe, that $\sigma(\gamma') \subset \sigma(\gamma)$. For this reason, $\sigma(\gamma')$ is called a sub-sigma algebra of $\sigma(\gamma)$.

Finally, one needs a probability measure P to define a probability space. It is a function $P : \Omega \mapsto [0, 1]$ that fulfils the properties below:

- For the sure event Ω , we have $P(\Omega) \equiv 1$.
- The probability of the impossible event \emptyset is $P(\emptyset) \equiv 0$.
- $\forall A \in \mathcal{F}$, we get $0 \leq P(A) \leq 1$.
- $\forall A, B \in \mathcal{F}$, we have $P(A \cap B) + P(A \cup B) = P(A) + P(B)$

In later sections, we will often investigate events which are called equivalent. Those are two events $A, B \in \mathcal{F}$, where the event that A happens is determined by B happening and vice-versa. Equivalence of events can be expressed as

$$P(A) = P(A \cap B) = P(B). \quad (1)$$

2.2 Random variables and stochastic processes

Let Ω be a set and \mathcal{F} be a sigma algebra on Ω . Then we say, that (Ω, \mathcal{F}) is a measurable space and all sets $A \in \mathcal{F}$ are called measurable sets. With two measurable spaces (Ω, \mathcal{F}) and (Ω', \mathcal{F}') , a mapping $T : \Omega \mapsto \Omega'$ is called \mathcal{F} -measurable, if $T^{-1}(A') \in \mathcal{F}$ for all $A' \in \mathcal{F}'$.

Often, one is not interested in a single event, but on more complex data related to a probabilistic experiment. This maybe the sum of the numbers on a dice, after it was rolled for two times. Using a probability space (Ω, \mathcal{F}, P) and a measurable space (Ω', \mathcal{F}') , these values can be described with an \mathcal{F} -measurable mapping:

$$X : \Omega \mapsto \Omega' \quad (2)$$

Such a mapping is called an (Ω', \mathcal{F}') -random variable. In case X is integrable on Ω with respect to P , it has an expectation value

$$EX[X] \equiv \int_{\Omega} X dP. \quad (3)$$

A simple random variable for events $A \in \mathcal{F}$ is given by the indicator function:

$$1_A(\omega) \equiv \begin{cases} 1 & \text{if } \omega \in A \\ 0 & \text{if } \omega \notin A \end{cases} \quad (4)$$

Obviously, $1_A(\omega)$ is \mathcal{F} -measurable. Furthermore, it can be used, to compute the probability of an event A :

$$EX(1_A) = \int_{\Omega} 1_A dP = P(A) \quad (5)$$

We define for any set $A' \in \Omega'$ the event $(X^{-1}(A')) \equiv \{X \in A'\}$ and write $\{A'\}$ for short. We call $\{A'\}$ the event, that X lies in A' and say $\{A'\}$ is generated by X . The probability of $\{A'\}$ to happen is $P(X^{-1}(A'))$ and we will abbreviate it as $P_X \{A'\}$.

With this notation, we say that two random variables X and Y are equivalent under P if, for any set $A' \in \mathcal{F}'$, the following relation holds:

$$P_X \{A'\} = P_Y \{A'\} \quad (6)$$

Sometimes, we have partial knowledge of an experiment and that knowledge is contained in a sub sigma algebra $\mathcal{F}_{sub} \subset \mathcal{F}$ on a probability space (Ω, \mathcal{F}, P) . It may be, that we want to compute the probability of an event $A \in \mathcal{F}$ given the information in \mathcal{F}_{sub} . Therefore, one defines the so called conditional probability with respect to the hypothesis \mathcal{F}_{sub} .

We assume, that \mathcal{F}_{sub} is generated by a partition of Ω into $I = \{1, \dots, n\}$ finite or $I = \mathbb{N}$ countable infinite disjoint sets $(B_i)_{i \in I} \in \mathcal{F}$, where $\Omega = \bigcup_{i \in I} B_i$, and $P(B_i) > 0$. The conditional probability of an event $A \in \mathcal{F}$ with respect to an event $B_i \in \mathcal{F}_{sub}$, where $\mathcal{F}_{sub} = \sigma(B_i; i \in I)$, is given by

$$P_{B_i}(A) \equiv \frac{P(A \cap B_i)}{P(B_i)} = \frac{1}{P(B_i)} \int 1_A 1_B dP = \frac{1}{P_{B_i}}(1_{B_i} P)(A). \quad (7)$$

We observe, that this expression basically defines a new probability measure. We can compute the expectation value of a random variable X with respect to P_{B_i} and obtain

$$EX_{B_i}[X] \equiv \int X dP_{B_i} = \frac{1}{P(B_i)} \int_{B_i} X dP. \quad (8)$$

Equation (8) can be used, to define a new random variable:

$$EX[X | \sigma(B_i; i \in I)] \equiv \sum_{i \in I} EX_{B_i}[X] 1_{B_i} \quad (9)$$

The latter is called the conditional expectation value of X with respect to the hypothesis $\sigma(B_i; i \in I)$. Obviously, $EX[X | \sigma(B_i; i \in I)]$ is measurable with respect to $\sigma(B_i; i \in I)$. Moreover, we can integrate $EX[X | \sigma(B_i; i \in I)]$ on any set $B_i \in \sigma(B_i; i \in I)$ and observe, that

$$\int_{B_i} EX[X | \sigma(B_i; i \in I)] dP = EX_{B_i}[X] P(B_i) = \int_{B_i} X dP. \quad (10)$$

With the conditional expectation value, we are in the position to define the conditional probability of an event $A \in \mathcal{F}$ with respect to the hypothesis $\sigma(B_i; i \in I)$ by

$$P(A | \sigma(B_i; i \in I)) \equiv EX[1_A | \sigma(B_i; i \in I)] = \sum_{i \in I} P_{B_i}(A) 1_{B_i} = \sum_{i \in I} \frac{P(A \cap B_i)}{P(B_i)} 1_{B_i}. \quad (11)$$

It is a random variable, which can be interpreted as the revised probability corresponding to a given outcome $\omega \in \Omega$, that an event $A \in \mathcal{F}$ will occur, given the extra information about which events in $\sigma(B_i; i \in I)$ occur for ω . Obviously, $P(A | \sigma(B_i; i \in I))$ fulfils the following properties:

- $0 \leq P(A | \sigma(B_i; i \in I)) \leq 1$ for all events $A \in \mathcal{F}$ and outcomes in Ω .
- $P(A | \sigma(B_i; i \in I))$ is measurable with respect to $\sigma(B_i; i \in I)$.
- Using eq. (10), we get $EX[P(A | \sigma(B_i; i \in I))] = P(A)$ for all $A \in \mathcal{F}$.
- For an event $B_i \in \sigma(B_i; i \in I)$, one has $P(A \cap B_i | \sigma(B_i; i \in I)) = P(A)$ for outcomes in B_i and $P(A \cap B_i | \sigma(B_i; i \in I)) = 0$ for outcomes not in B_i .

In physical situations, one often has randomness which varies over a parameter, for example time. This gives rise to the definition of a stochastic process. With a probability space (Ω, \mathcal{F}, P) and a measurable space (Ω', \mathcal{F}') , a stochastic process is defined as a family of (Ω', \mathcal{F}') -random variables

$$(X_t)_{t \in I} : \Omega \mapsto \Omega'. \quad (12)$$

The parameter t is called parameter-set and lies in an interval I . The measurable space (Ω', \mathcal{F}') is called state space. We say, that a stochastic process is markovian, if the likely-hood of a given future state depends only on the present state of the process.

2.3 Enlarged probability spaces

Often, an experiment happens in more than one probability space. For example, we might toss a coin and a dice. The coin has two theoretically possible outcomes whereas the dice has six. Hence, their events take place in two different probability spaces. Nevertheless, there exists a joint probability for the events of the two experiments. In such situations, it is necessary to define an enlarged probability space $(\Omega, \mathcal{F}, P) \equiv \left(\prod_{i=1}^N \Omega_i, \bigotimes_{i=1}^N \mathcal{F}_i, P \right)$, where $N \in \mathbb{N}$.

On the enlarged probability space, the set

$$\Omega \equiv \prod_{i=1}^N \Omega_i \equiv \Omega_1 \times \dots \times \Omega_N \quad (13)$$

is a Cartesian product. It consists of theoretically possible outcomes $\omega \in \Omega$, where $\omega \equiv \prod_{i=1}^N \omega_i$ and $\omega_i \in \Omega_i$.

Assume, we have N measurable spaces $(\Omega_i, \mathcal{F}_i)$, where $1 \leq i \leq N$, a set Ω and N mappings $p_i : \Omega \mapsto \Omega_i$. Now, we want to consider the system of subsets on Ω which is defined through

$$p_i^{-1}(\mathcal{F}_i) \equiv \{p_i^{-1}(A_i) : A_i \in \mathcal{F}_i\}. \quad (14)$$

Obviously, the sigma algebra generated by the system $\bigcup_{i=1}^N p_i^{-1}(\mathcal{F}_i)$ is the smallest one, which all the N functions p_i are measurable with respect to. We denote this sigma algebra by

$$\mathcal{F} \equiv \bigotimes_{i=1}^N \mathcal{F}_i \equiv \sigma \left(\bigcup_{i=1}^N p_i^{-1}(\mathcal{F}_i) \right) \quad (15)$$

and we will use it on the enlarged probability space.

Again, sets of outcomes on the enlarged probability space are called events and the operations union, intersection and complement can be applied on events to form new ones. For two events $A = \prod_{i=1}^N A_i$

and $B \equiv \prod_{i=1}^N B_i$, where $A, B \in \otimes_{i=1}^N \mathcal{F}_i$, the union event is defined as $A \cup B = \prod_{i=1}^N (A_i \cup B_i)$. The intersection event is given by $A \cap B = \prod_{i=1}^N (A_i \cap B_i)$ and the complement of A is $\mathcal{C}A = \prod_{i=1}^N \mathcal{C}A_i$.

Finally, the probability measure on the enlarged probability space is a function $P : \prod_{i=1}^N \Omega_i \mapsto [0, 1]$. For an event $A = \prod_{i=1}^N A_i$ with $A_i \in \mathcal{F}_i$, we write

$$P(A) \equiv P \left(\bigcap_{i=1}^N A_i \right). \quad (16)$$

Let $X_i : \Omega \mapsto \Omega'_i$ be a family of N $(\Omega'_i, \mathcal{F}'_i)$ -random variables, where $1 \leq i \leq N$, for events on an arbitrary probability space (Ω, \mathcal{F}, P) . These random variables can be used, to define an $(\prod_{n=1}^N \Omega'_i, \otimes_{i=1}^N \mathcal{F}'_i)$ -product random variable

$$Y \equiv \bigotimes_{i=1}^N X_i : \Omega \mapsto \prod_{i=1}^N \Omega'_i \quad (17)$$

for events $A \in \mathcal{F}$ by

$$Y(A) \equiv (X_1(A), \dots, X_N(A)). \quad (18)$$

In the above definitions, (Ω, \mathcal{F}, P) can well be an enlarged probability space. In this case, it may be, that there is a family of N random variables $X_i : \Omega_i \mapsto \Omega'_i$, where $1 \leq i \leq N$, which only map the i -th component of events $A = \prod_{i=1}^N A_i$ on the enlarged space $(\prod_{i=1}^N \Omega_i, \otimes_{i=1}^N \mathcal{F}_i, P)$ to the measurable space $(\Omega'_i, \mathcal{F}'_i)$. Then, we can define another $(\prod_{n=1}^N \Omega'_i, \otimes_{i=1}^N \mathcal{F}'_i)$ -product random variable

$$Y \equiv \bigotimes_{i=1}^N X_i : \prod_{i=1}^N \Omega_i \mapsto \prod_{i=1}^N \Omega'_i \quad (19)$$

for events $A \in \otimes_{i=1}^N \mathcal{F}_i$ by

$$Y(A) \equiv (X_1(A_1), \dots, X_N(A_N)). \quad (20)$$

For all sets $A' \equiv \prod_{i=1}^N A'_i$ on the measurable space $(\prod_{i=1}^N \Omega'_i, \otimes_{i=1}^N \mathcal{F}'_i)$, the event that a product random variable $Y = \bigotimes_{i=1}^N X_i$ lies in A' is given by

$$\left\{ Y \in \prod_{i=1}^N A'_i \right\} \equiv \left\{ \bigcap_{i=1}^N X_i \in A'_i \right\} \equiv \left(\bigcap_{i=1}^N X_i^{-1}(A'_i) \right). \quad (21)$$

We abbreviate this event by $\{A'\}$ and say that it is generated by Y . The probability of its occurrence is

$$P \left\{ Y \in \prod_{i=1}^N A'_i \right\} = P \left(\bigcap_{i=1}^N X_i^{-1}(A'_i) \right). \quad (22)$$

It will be written as $P_Y \{A'\}$ or $P_{\otimes_{i=1}^N X_i} \left\{ \bigcap_{i=1}^N A'_i \right\}$.

3 Nelson's analysis of Bell's theorem

3.1 The setup of the EPR experiment in theory

An EPR experiment consists of two measurement devices 1 and 2 which are space-like separated, as well as a particle source in the intersection of their past cones. The source ejects j pairs of particles, where $1 \leq j \leq S \in \mathbb{N}$, to the two detectors. The particles in each pair have opposite spin and opposite momentum. At each device, the direction of the spins is measured with a Stern-Gerlach magnet. These magnets can be




Figure 1: Drawing of an EPR experiment with Stern-Gerlach magnets 1, 2 and axes $\vec{\mu}$, $\vec{\nu}$. The black dots illustrate the particles moving to the detectors. Their spin direction with respect to the \vec{z} axis is indicated by the small arrows around them.




Figure 2: Space-time diagram of an EPR experiment. The dashed arrows display particle trajectories

rotated around an arbitrary direction, perpendicular to the axis of the particle trajectory \vec{z} . The direction of the magnets is called $\vec{\mu}$ for detector 1 and $\vec{\nu}$ for detector 2 (see figs. 1, 2).

We set up two families of stochastic processes $\phi_{j1t_z} : \Omega_{j1} \mapsto \Omega'_{j1}$ and $\phi_{j2t_z} : \Omega_{j2} \mapsto \Omega'_{j2}$. The processes ϕ_{j1t_z} and ϕ_{j2t_z} define the time dependent spin values of the j -th particles, which are sent to detector 1 and 2. They are called “spin trajectories” and their outcomes in the measurable spaces are confined to $\Omega'_{j1} = \Omega'_{j2} \equiv \Omega'_j \equiv \{\uparrow, \downarrow\}$.

As parameter of the spin trajectories, the vector \vec{z} corresponds to the axis around which the described particle rotates. If it rotates in the left direction with respect to \vec{z} , we have $\phi_{j1(2)t_z} = \uparrow$ and a rotation in the right direction is defined by $\phi_{j1(2)t_z} = \downarrow$. The value t is the time, during which the particle flies. Both particles start at time $t = 0$ from the source and arrive at time $t = T$, where $T > 0$, in a Stern-Gerlach magnet.

The measurement devices could change the particle’s properties, including the spin. In an EPR experiment, the parts of the detectors that might influence the particles are the axes $\vec{\mu}$ and $\vec{\nu}$. They can be chosen by the experimenter at will. With the Stern-Gerlach magnets, we have to describe our measurement results by device dependent random variables. For the two detectors 1 and 2, we define two axis dependent families of random variables $D_{j1\vec{\mu}} : \Omega_{j1D} \mapsto \Omega'_{j1D}$ and $D_{j2\vec{\nu}} : \Omega_{j2D} \mapsto \Omega'_{j2D}$, where $\Omega'_{j1D} = \Omega'_{j2D} = \Omega'_j = \{\uparrow, \downarrow\}$. Since the random variables depend on the setting of the Stern-Gerlach magnets, we write their outcomes on the measurable space in an axis dependent notation as $\sigma_{j1\vec{\mu}}$ and $\sigma_{j2\vec{\nu}}$. However, the way, how the events generated by $D_{j1\vec{\mu}}$ and $D_{j2\vec{\nu}}$ are connected to those generated by the processes ϕ_{j1t_z} and ϕ_{j2t_z} will be postponed at the moment.

The results at the detectors could depend on the preparation of the particles by the source. We say, that the corresponding events happen at preparation stage. They are generated by a product random variable

$$S_{j12} \equiv \phi_{j1,t=0_z} \otimes \phi_{j2,t=0_z} : (\Omega_{j1} \times \Omega_{j2}) \mapsto (\Omega'_{j1} \times \Omega'_{j2}). \quad (23)$$

The axis dependent family of product random variables

$$X_{j1\vec{\mu}2\vec{\nu}} \equiv D_{j1\vec{\mu}} \otimes S_{j12} \otimes D_{j2\vec{\nu}} : (\Omega_{j1D} \times \Omega_{j1} \times \Omega_{j2} \times \Omega_{j2D}) \mapsto (\Omega'_{j1D} \times \Omega'_{j1} \times \Omega'_{j2} \times \Omega'_{j2D}) \quad (24)$$

generates events on an enlarged probability space. We use that probability space to define two new families of random variables

$$\phi_{j1\vec{\mu}} : (\Omega_{j1D} \times \Omega_{j1} \times \Omega_{j2} \times \Omega_{j2D}) \mapsto \Omega'_{j1D}, \quad (25)$$

$$\phi_{j2\vec{\nu}} : (\Omega_{j1D} \times \Omega_{j1} \times \Omega_{j2} \times \Omega_{j2D}) \mapsto \Omega'_{j2D}. \quad (26)$$

They provide information about the measurement results at the detectors. Since $\phi_{j1\vec{\mu}}$ and $\phi_{j2\vec{\nu}}$ are functions of events on the enlarged probability space, their construction will make it finally possible to investigate the relationship between the events happening at the detectors and the ones at preparation stage. The definition of $\phi_{j1\vec{\mu}}$ is

$$\{X_{j1\vec{\mu}2\vec{\nu}} = (\sigma_{j1\vec{\mu}} \cap \Omega'_{j1} \cap \Omega'_{j2} \cap \Omega'_{j2D})\} \equiv \{\phi_{j1\vec{\mu}} = \sigma_{j1\vec{\mu}}\} \quad (27)$$

and similarly, $\phi_{j2\bar{v}}$ is defined through

$$\left\{ X_{j1\bar{\mu}2\bar{v}} = \left(\Omega'_{j1D} \cap \Omega'_{j1} \cap \Omega'_{j2} \cap \sigma_{j2\bar{v}} \right) \right\} \equiv \{ \phi_{j2\bar{v}} = \sigma_{j2\bar{v}} \}. \quad (28)$$

The events generated by $\phi_{j1\bar{\mu}}$ will be denoted by $\{ \sigma_{j1\bar{\mu}} \}$. These events generate a sigma algebra, which will be called \mathcal{F}_{j1} . Similarly, we say that the random variable $\phi_{j2\bar{v}}$ generates events $\{ \sigma_{j2\bar{v}} \}$ which themselves generate the sigma algebra \mathcal{F}_{j2} .

Due to the definitions of $\{ \sigma_{j1\bar{\mu}} \}$ and $\{ \sigma_{j2\bar{v}} \}$, the two sigma algebras \mathcal{F}_{j1} and \mathcal{F}_{j2} only contain information about the events in the left and right detector.

Additionally, we define a third sigma algebra \mathcal{F}_{jS} which only contains the knowledge about the events at preparation stage. It is generated by $\{ X_{j1\bar{\mu}2\bar{v}} = (\Omega'_{j1D} \cap \uparrow \cap \downarrow \cap \Omega'_{j2D}) \}$ and $\{ X_{j1\bar{\mu}2\bar{v}} = (\Omega'_{j1D} \cap \downarrow \cup \uparrow \cap \Omega'_{j2D}) \}$.

Particularly, we will be interested in the joint probability of the events that happen at the detectors. They are generated by a family of product random variables

$$\phi_{j1\bar{\mu}} \otimes \phi_{j2\bar{v}} : (\Omega_{j1D} \times \Omega_{j1} \times \Omega_{j2} \times \Omega_{j2D}) \mapsto (\Omega'_{j1D} \times \Omega'_{j2D}), \quad (29)$$

whose definition is

$$\left\{ X_{j1\bar{\mu}2\bar{v}} = \left(\sigma_{j1\bar{\mu}} \cap \Omega'_{j1} \cap \Omega'_{j2} \cap \sigma_{j2\bar{v}} \right) \right\} \equiv \left\{ \phi_{j1\bar{\mu}} \otimes \phi_{j2\bar{v}} = \left(\sigma_{j1\bar{\mu}} \cap \sigma_{j2\bar{v}} \right) \right\}. \quad (30)$$

Since this family of random variables depends on the axes of both detectors, we must use an axis dependent family of probability measures $P_{\phi_{j1\bar{\mu}} \otimes \phi_{j2\bar{v}}}$ for it.

There may still be situations, where we are interested in only one event $\{ \sigma_{j1\bar{\mu}} \}$ or $\{ \sigma_{j2\bar{v}} \}$ at a single detector. Hence, we define the notations

$$P_{\phi_{j1\bar{\mu}} \otimes \phi_{j2\bar{v}}} \{ \sigma_{j1\bar{\mu}} \} \equiv P_{\phi_{j1\bar{\mu}} \otimes \phi_{j2\bar{v}}} \left\{ \sigma_{j1\bar{\mu}} \cap \Omega'_{j2D} \right\} \quad (31)$$

and

$$P_{\phi_{j1\bar{\mu}} \otimes \phi_{j2\bar{v}}} \{ \sigma_{j2\bar{v}} \} \equiv P_{\phi_{j1\bar{\mu}} \otimes \phi_{j2\bar{v}}} \left\{ \Omega'_{j1D} \cap \sigma_{j2\bar{v}} \right\}. \quad (32)$$

Now, the mathematical structure needed to analyse the EPR experiment in full has been completely defined.

3.2 Active locality

At first, Nelson defined two different forms of locality: Active locality and passive locality. The meaning of active locality is:

Whatever axes the experimenter selects at one measurement device, e.g. at 2, it does not change the probabilities of an event happening at 1, as long as 1 does not lie in the future cone of 2. Hence, active locality corresponds to a “non-signalling” condition. If it is violated, one can send instantaneous signals between two spatially separated locations. We can mathematically define active locality as follows:

The area of measurement station 1 has to be disjoint from the future cone of station 2. At 2, different axes $\bar{v} \neq \bar{v}'$ of the Stern-Gerlach magnet are chosen. These axes influence the measurement results through the random variables $\phi_{j2\bar{v}}$ and $\phi_{j2\bar{v}'}$. We define a theory to be actively local, if $\phi_{j2\bar{v}}$ and $\phi_{j2\bar{v}'}$ are equivalent under $P_{\phi_{j1\bar{\mu}} \otimes \phi_{j2\bar{v}}} \{ \sigma_{j1\bar{\mu}} \}$, or

$$P_{\phi_{j1\bar{\mu}} \otimes \phi_{j2\bar{v}}} \{ \sigma_{j1\bar{\mu}} \} = P_{\phi_{j1\bar{\mu}} \otimes \phi_{j2\bar{v}'}} \{ \sigma_{j1\bar{\mu}} \}. \quad (33)$$

A similar condition should be true at 2:

$$P_{\phi_{j1\bar{\mu}} \otimes \phi_{j2\bar{v}}} \{ \sigma_{j2\bar{v}} \} = P_{\phi_{j1\bar{\mu}} \otimes \phi_{j2\bar{v}'}} \{ \sigma_{j2\bar{v}} \} \quad (34)$$

In eq. (34), the spatial region of 2 must be disjoint from the future cone of 1 and we should have $\bar{\mu}' \neq \bar{\mu}$ for the axes of detector 1.

The events in \mathcal{F}_{jS} contain no information about the results at the detectors. Instead, they only define, what has happened at the source. The latter is located disjoint from the future cones of the two measurement

devices. Therefore, in an actively local theory, the probabilities of the events in \mathcal{F}_{jS} cannot be changed by the detector's axes.

We can condition the probabilities of $\{\sigma_{j1\vec{\mu}}\} \in \mathcal{F}_{j1}$ and $\{\sigma_{j2\vec{\mu}}\} \in \mathcal{F}_{j2}$ with respect to the hypothesis \mathcal{F}_{jS} . Since the events in \mathcal{F}_{jS} do not depend on any detector setting, $\mathbb{P}_{\phi_{j1\vec{\mu}} \otimes \phi_{j2\vec{\nu}}} \{\sigma_{j1\vec{\mu}} | \mathcal{F}_{jS}\}$ and $\mathbb{P}_{\phi_{j1\vec{\mu}} \otimes \phi_{j2\vec{\nu}}} \{\sigma_{j2\vec{\nu}} | \mathcal{F}_{jS}\}$ will have similar properties as in the eqs. (33) and (34):

$$\begin{aligned} \mathbb{P}_{\phi_{j1\vec{\mu}} \otimes \phi_{j2\vec{\nu}}} \{\sigma_{j1\vec{\mu}} | \mathcal{F}_{jS}\} &= \mathbb{P}_{\phi_{j1\vec{\mu}} \otimes \phi_{j2\vec{\nu}'}} \{\sigma_{j1\vec{\mu}} | \mathcal{F}_{jS}\} \\ \mathbb{P}_{\phi_{j1\vec{\mu}} \otimes \phi_{j2\vec{\nu}}} \{\sigma_{j2\vec{\nu}} | \mathcal{F}_{jS}\} &= \mathbb{P}_{\phi_{j1\vec{\mu}'} \otimes \phi_{j2\vec{\nu}}} \{\sigma_{j2\vec{\nu}} | \mathcal{F}_{jS}\} \end{aligned} \quad (35)$$

3.3 Passive locality

In this subsection, we consider the joint probability of $\{\sigma_{j1\vec{\mu}}\}$ and $\{\sigma_{j2\vec{\nu}}\}$. It gives information about events which happen simultaneously at spatially separated locations 1 and 2. We say, that passive locality holds, if

$$\mathbb{P}_{\phi_{j1\vec{\mu}} \otimes \phi_{j2\vec{\nu}}} \left\{ \sigma_{j1\vec{\mu}} \cap \sigma_{j2\vec{\nu}} \middle| \mathcal{F}_{jS} \right\} = \mathbb{P}_{\phi_{j1\vec{\mu}} \otimes \phi_{j2\vec{\nu}}} \{\sigma_{j1\vec{\mu}} | \mathcal{F}_{jS}\} \mathbb{P}_{\phi_{j1\vec{\mu}} \otimes \phi_{j2\vec{\nu}}} \{\sigma_{j2\vec{\nu}} | \mathcal{F}_{jS}\} \quad (36)$$

for every pair of axes $\vec{\mu}$ and $\vec{\nu}$.

It is possible, to have active locality without passive locality. Furthermore, a theory which violates passive locality does not have to incorporate any signalling mechanism between the spatially separated measurement stations 1 and 2. However, the true physical implications of the passive locality condition will be postponed at the moment and analysed more closely in section 4.

3.4 Nelson's theorem

With the definitions above, quantum mechanics fulfils the following properties:

$$\mathbb{P}_{\phi_{j1\vec{\mu}} \otimes \phi_{j2\vec{\mu}}} \left\{ \sigma_{j1\vec{\mu}} = \uparrow \cap \sigma_{j2\vec{\mu}} = \downarrow \right\} = \mathbb{P}_{\phi_{j1\vec{\mu}} \otimes \phi_{j2\vec{\mu}}} \left\{ \sigma_{j1\vec{\mu}} = \downarrow \cap \sigma_{j2\vec{\mu}} = \uparrow \right\} = 1 \quad (37)$$

$$|\mathbb{E}(\vec{\mu}, \vec{\nu}) + \mathbb{E}(\vec{\mu}, \vec{\nu}') + \mathbb{E}(\vec{\mu}', \vec{\nu}) - \mathbb{E}(\vec{\mu}', \vec{\nu}')| > 2 \quad (38)$$

In eq. (38), $\mathbb{E}(\vec{\mu}, \vec{\nu})$ is the so called ‘‘correlation coefficient’’, a function defined through:

$$\begin{aligned} \mathbb{E}(\vec{\mu}, \vec{\nu}) &\equiv \mathbb{P}_{\phi_{j1\vec{\mu}} \otimes \phi_{j2\vec{\nu}}} \{\sigma_{j1\vec{\mu}} = \uparrow \cap \sigma_{j2\vec{\nu}} = \uparrow\} + \mathbb{P}_{\phi_{j1\vec{\mu}} \otimes \phi_{j2\vec{\nu}}} \{\sigma_{j1\vec{\mu}} = \downarrow \cap \sigma_{j2\vec{\nu}} = \downarrow\} \\ &\quad - \mathbb{P}_{\phi_{j1\vec{\mu}} \otimes \phi_{j2\vec{\nu}}} \{\sigma_{j1\vec{\mu}} = \uparrow \cap \sigma_{j2\vec{\nu}} = \downarrow\} - \mathbb{P}_{\phi_{j1\vec{\mu}} \otimes \phi_{j2\vec{\nu}}} \{\sigma_{j1\vec{\mu}} = \downarrow \cap \sigma_{j2\vec{\nu}} = \uparrow\}. \end{aligned} \quad (39)$$

Equation (37) implies, that if the axes of the Stern-Gerlach detectors are the same, the spin values measured at 1 and 2 are always opposite. Eq. (38) is, as shall be shown below, in conflict with theories where active and passive locality hold. In Nelson's form, Bell's theorem states: If active and passive locality hold eq. (37) and eq. (38) cannot be fulfilled simultaneously. Nelson's proof from [8, 9] (with corrections in [10]) goes as follows:

Passive locality demands for events $\{\sigma_{j1\vec{\mu}} = \uparrow\}$ at 1 and $\{\sigma_{j2\vec{\nu}} = \downarrow\}$ at 2, that:

$$\begin{aligned} \mathbb{P}_{\phi_{j1\vec{\mu}} \otimes \phi_{j2\vec{\nu}}} \left\{ \sigma_{j1\vec{\mu}} = \uparrow \cap \sigma_{j2\vec{\nu}} = \downarrow \middle| \mathcal{F}_{jS} \right\} &= \mathbb{P}_{\phi_{j1\vec{\mu}} \otimes \phi_{j2\vec{\nu}}} \{\sigma_{j1\vec{\mu}} = \uparrow | \mathcal{F}_{jS}\} \\ &\quad \cdot \mathbb{P}_{\phi_{j1\vec{\mu}} \otimes \phi_{j2\vec{\nu}}} \{\sigma_{j2\vec{\nu}} = \downarrow | \mathcal{F}_{jS}\} \end{aligned} \quad (40)$$

Active locality implies, that the actions which happen in 2 can only affect anything in its future cone. Since 1 does not lie in that cone, nothing what happens in 2 can affect the events measured at 1. Therefore, the random variables $\phi_{j2\vec{\nu}}$ and $\phi_{j2\vec{\mu}}$ are equivalent for any result $\{\sigma_{j1\vec{\mu}}\}$:

$$\mathbb{P}_{\phi_{j1\vec{\mu}} \otimes \phi_{j2\vec{\nu}}} \{\sigma_{j1\vec{\mu}} = \uparrow | \mathcal{F}_{jS}\} = \mathbb{P}_{\phi_{j1\vec{\mu}} \otimes \phi_{j2\vec{\mu}}} \{\sigma_{j1\vec{\mu}} = \uparrow | \mathcal{F}_{jS}\} \equiv P_{\vec{\mu}} \quad (41)$$

An analogous expression is true for the events at 2:

$$\mathbb{P}_{\phi_{j1\vec{\mu}} \otimes \phi_{j2\vec{\nu}}} \{\sigma_{j2\vec{\nu}} = \downarrow | \mathcal{F}_{jS}\} = \mathbb{P}_{\phi_{j1\vec{\nu}} \otimes \phi_{j2\vec{\nu}}} \{\sigma_{j2\vec{\nu}} = \downarrow | \mathcal{F}_{jS}\} \quad (42)$$

On the other hand, eq. (37) yields $\{\sigma_{j1\vec{\nu}} = \uparrow\}$ in the right side of eq. (42). Hence, we can write for the coordinates ν :

$$\mathbb{P}_{\phi_{j1\vec{\mu}} \otimes \phi_{j2\vec{\nu}}} \{\sigma_{j2\vec{\nu}} = \downarrow | \mathcal{F}_{jS}\} = \mathbb{P}_{\phi_{j1\vec{\mu}} \otimes \phi_{j2\vec{\nu}}} \{\sigma_{j1\vec{\nu}} = \uparrow | \mathcal{F}_{jS}\} \equiv P_{\vec{\nu}} \quad (43)$$

Plugging eq. (41) and eq. (43) back to eq. (40), we arrive at

$$P_{\phi_{j1\bar{\mu}} \otimes \phi_{j2\bar{\nu}}} \left\{ \sigma_{j1\bar{\mu}} = \uparrow \cap \sigma_{j2\bar{\nu}} = \downarrow \middle| \mathcal{F}_{jS} \right\} = P_{\bar{\mu}} P_{\bar{\nu}}. \quad (44)$$

The sum of the probabilities for two complementary measurement results $\{\sigma_{j2\bar{\nu}} = \uparrow\}$ and $\{\sigma_{j2\bar{\nu}} = \downarrow\}$ at 2 must be unity. Therefore, with eq. (43), we get

$$P_{\phi_{j1\bar{\mu}} \otimes \phi_{j2\bar{\nu}}} \left\{ \sigma_{j2\bar{\nu}} = \uparrow \middle| \mathcal{F}_{jS} \right\} = 1 - P_{\bar{\nu}}. \quad (45)$$

Since active locality holds, a similar result can be written for the measurement results $\{\sigma_{j1\bar{\mu}} = \downarrow\}$ at 1:

$$\begin{aligned} P_{\phi_{j1\bar{\mu}} \otimes \phi_{j2\bar{\nu}}} \left\{ \sigma_{j1\bar{\mu}} = \downarrow \middle| \mathcal{F}_{jS} \right\} &= P_{\phi_{j1\bar{\mu}} \otimes \phi_{j2\bar{\mu}}} \left\{ \sigma_{j1\bar{\mu}} = \downarrow \middle| \mathcal{F}_{jS} \right\} \\ &= 1 - P_{\phi_{j1\bar{\mu}} \otimes \phi_{j2\bar{\mu}}} \left\{ \sigma_{j1\bar{\mu}} = \uparrow \middle| \mathcal{F}_{jS} \right\} = 1 - P_{\bar{\mu}} \end{aligned} \quad (46)$$

Due to passive locality, we have for the events $\{\sigma_{j1\bar{\mu}} = \downarrow\}$ and $\{\sigma_{j2\bar{\nu}} = \uparrow\}$, by using eq. (46) and eq. (45):

$$P_{\phi_{j1\bar{\mu}} \otimes \phi_{j2\bar{\nu}}} \left\{ \sigma_{j1\bar{\mu}} = \downarrow \cap \sigma_{j2\bar{\nu}} = \uparrow \middle| \mathcal{F}_{jS} \right\} = (1 - P_{\bar{\mu}}) (1 - P_{\bar{\nu}}) \quad (47)$$

In the same way, we can derive the relations

$$P_{\phi_{j1\bar{\mu}} \otimes \phi_{j2\bar{\nu}}} \left\{ \sigma_{j1\bar{\mu}} = \uparrow \cap \sigma_{j2\bar{\nu}} = \uparrow \middle| \mathcal{F}_{jS} \right\} = P_{\bar{\mu}} (1 - P_{\bar{\nu}}) \quad (48)$$

and

$$P_{\phi_{j1\bar{\mu}} \otimes \phi_{j2\bar{\nu}}} \left\{ \sigma_{j1\bar{\mu}} = \downarrow \cap \sigma_{j2\bar{\nu}} = \downarrow \middle| \mathcal{F}_{jS} \right\} = (1 - P_{\bar{\mu}}) P_{\bar{\nu}}. \quad (49)$$

Because of eq. (48), eq. (49), eq. (44) and eq. (47), the correlation coefficient defined in eq. (39), but with conditional probabilities, becomes:

$$E(\bar{\mu}, \bar{\nu} | \mathcal{F}_{jS}) = P_{\bar{\mu}} (1 - P_{\bar{\nu}}) + (1 - P_{\bar{\mu}}) P_{\bar{\nu}} - P_{\bar{\mu}} P_{\bar{\nu}} - (1 - P_{\bar{\mu}}) (1 - P_{\bar{\nu}}) \quad (50)$$

Now, using four arbitrary axes $\bar{\mu}, \bar{\mu}'$ and $\bar{\nu}, \bar{\nu}'$ at the two stations 1 and 2, the following inequality can be computed:

$$|E(\bar{\mu}, \bar{\nu} | \mathcal{F}_{jS}) + E(\bar{\mu}, \bar{\nu}' | \mathcal{F}_{jS}) + E(\bar{\mu}', \bar{\nu} | \mathcal{F}_{jS}) - E(\bar{\mu}', \bar{\nu}' | \mathcal{F}_{jS})| \leq 2 \quad (51)$$

We got this result, because the conditional probabilities in $E(\bar{\mu}, \bar{\nu} | \mathcal{F}_{jS})$ are all in the range $0 \leq P_{\phi_{j1\bar{\mu}} \otimes \phi_{j2\bar{\nu}}} \left\{ \sigma_{j1\bar{\mu}} \cap \sigma_{j2\bar{\nu}} \middle| \mathcal{F}_{jS} \right\} \leq 1$. The unconditional probabilities can be calculated with the expectation value of the conditional probabilities. Since the resulting probability measures are also in the range $0 \leq P_{\phi_{j1\bar{\mu}} \otimes \phi_{j2\bar{\nu}}} \left\{ \sigma_{j1\bar{\mu}} \cap \sigma_{j2\bar{\nu}} \right\} \leq 1$, an analogous inequality must be true for them:

$$|E(\bar{\mu}, \bar{\nu}) + E(\bar{\mu}, \bar{\nu}') + E(\bar{\mu}', \bar{\nu}) - E(\bar{\mu}', \bar{\nu}')| \leq 2 \quad (52)$$

Eq. (52) is the so called Clauser-Holt-Shimony-Horne inequality [6]. It is a version of Bells statement in [4]

$$|E(\bar{\mu}, \bar{\nu}) - E(\bar{\mu}, \bar{\nu}')| - E(\bar{\nu}, \bar{\nu}') \leq 1, \quad (53)$$

which can be similarly derived. Both eq. (52) and eq. (53) are violated in quantum mechanics and this violation was confirmed experimentally [20].

The analysis of Nelson leads to two possibilities: Since Bell's inequalities can only be derived assuming that **active and passive locality** holds, in a theory which reproduces quantum mechanics, either **active locality or passive locality** has to be violated. A violation of active locality would mean that action at a distance is possible. The physical implications of a violation of passive locality will be explored in the next section.

It should be remarked, that Jarret arrived at a similar conclusion in [13], just one year before Nelson published his analysis. Despite Jarret was not able to identify the correct probability space given by quantum mechanics for the EPR experiment, one can find a preliminary definition of active and passive locality in his publication.

4 What the failure of passive locality implies - a deeper analysis

Stochastic processes can be constructed such, that they act on each particle actively local but still are violating passive locality. This was shown at first by Nelson with two examples in [8,9]. One was a rather ad-hoc construction of a stochastic process fulfilling the Klein-Gordon equation and the other was a variant of Nelson's stochastic mechanics applied to relativistic field theory. Unfortunately the latter had no tempting particle interpretation as Nelson's original theory [17, 18] had, where particles move on stochastic paths. Hence, this „stochastic field theory” was never developed extensively. The present article aims to show, that a more physically motivated hidden variable theory published in [19] violates Bell's inequalities without any communication between the two measurement stations in the EPR setup. To reach our goal, we need to analyse the implications of Nelson's theorem further.

We begin by evaluating some deeper consequences of passive locality. Faris showed in [12], that passive locality, combined with relation (37) from quantum mechanics, leads immediately to another condition, which he calls „deterministic passive locality”. Suppose, as in the EPR experiment, two spatially separated events happen or not happen always together with probability 1. Then, deterministic passive locality states that this must be, because they are determined by some random event at the prior preparation stage. Therefore, all randomness lies in the preparation. Deterministic passive locality is defined mathematically as follows:

Let the events $\{\sigma_{j1\bar{\mu}}\} \in \mathcal{F}_{j1}$ at 1 and $\{\sigma_{j2\bar{\mu}}\} \in \mathcal{F}_{j2}$ at 2 be equivalent with respect to $P_{\phi_{j1\bar{\mu}} \otimes \phi_{j2\bar{\mu}}}$ and passive locality hold. Then, there is an event $\{x_{j\bar{z}}\} \in \mathcal{F}_{jS}$ at preparation stage, which is equivalent to both $\{\sigma_{j1\bar{\mu}}\}$ and $\{\sigma_{j2\bar{\mu}}\}$.

In his contribution, Faris states that a similar result is presented by Redhead in [52] at pp. 101-102. Redhead claims, it was discovered at first by Suppes and Zanotti in [14]. Recently, another derivation was given by Conway and Kochen (who called it “free will theorem“ to emphasise its implication of the particle's random behaviour) [15]. They relate the statement given below to the famous Kochen-Specker theorem [16]. The works of Conway, Kochen and Specker are using geometric arguments. In contrast, the derivation of this article relies on probability theory and will explicitly use the property of passive locality. The two different methods to prove the same lemma demonstrate an interesting connection between Bell's theorem and the one of Kochen-Specker. Our derivation, will follow closely the lines of Faris in [12]:

Passive locality demands, that

$$P_{\phi_{j1\bar{\mu}} \otimes \phi_{j2\bar{\mu}}} \left\{ \sigma_{j1\bar{\mu}} \cap \sigma_{j2\bar{\mu}} \mid \mathcal{F}_{jS} \right\} = P_{\phi_{j1\bar{\mu}} \otimes \phi_{j2\bar{\mu}}} \left\{ \sigma_{j1\bar{\mu}} \mid \mathcal{F}_{jS} \right\} P_{\phi_{j1\bar{\mu}} \otimes \phi_{j2\bar{\mu}}} \left\{ \sigma_{j2\bar{\mu}} \mid \mathcal{F}_{jS} \right\} \quad (54)$$

and because of $0 \leq P_{\phi_{j1\bar{\mu}} \otimes \phi_{j2\bar{\mu}}} \left\{ \sigma_{j1\bar{\mu}} \cap \sigma_{j2\bar{\mu}} \mid \mathcal{F}_{jS} \right\} \leq 1$, we have

$$P_{\phi_{j1\bar{\mu}} \otimes \phi_{j2\bar{\mu}}} \left\{ \sigma_{j1\bar{\mu}} \cap \sigma_{j2\bar{\mu}} \mid \mathcal{F}_{jS} \right\} \leq P_{\phi_{j1\bar{\mu}} \otimes \phi_{j2\bar{\mu}}} \left\{ \sigma_{j1\bar{\mu}} \mid \mathcal{F}_{jS} \right\}. \quad (55)$$

With section 2, we can compute the unconditional probabilities of $\{\sigma_{j1\bar{\mu}}\}$ and $\{\sigma_{j1\bar{\mu}} \cap \sigma_{j2\bar{\nu}}\}$ by:

$$P_{\phi_{j1\bar{\mu}} \otimes \phi_{j2\bar{\mu}}} \left\{ \sigma_{j1\bar{\mu}} \right\} = \text{EX} \left[P_{\phi_{j1\bar{\mu}} \otimes \phi_{j2\bar{\mu}}} \left\{ \sigma_{j1\bar{\mu}} \mid \mathcal{F}_{jS} \right\} \right] \quad (56)$$

and

$$P_{\phi_{j1\bar{\mu}} \otimes \phi_{j2\bar{\mu}}} \left\{ \sigma_{j1\bar{\mu}} \cap \sigma_{j2\bar{\mu}} \right\} = \text{EX} \left[P_{\phi_{j1\bar{\mu}} \otimes \phi_{j2\bar{\mu}}} \left\{ \sigma_{j1\bar{\mu}} \cap \sigma_{j2\bar{\mu}} \mid \mathcal{F}_{jS} \right\} \right]. \quad (57)$$

The events $\{\sigma_{j1\bar{\mu}}\}$ and $\{\sigma_{j2\bar{\mu}}\}$ are defined such, that they are equivalent under $P_{\phi_{j1\bar{\mu}} \otimes \phi_{j2\bar{\nu}}}$, or

$$P_{\phi_{j1\bar{\mu}} \otimes \phi_{j2\bar{\mu}}} \left\{ \sigma_{j1\bar{\mu}} \right\} = P_{\phi_{j1\bar{\mu}} \otimes \phi_{j2\bar{\mu}}} \left\{ \sigma_{j1\bar{\mu}} \cap \sigma_{j2\bar{\mu}} \right\} = P_{\phi_{j1\bar{\mu}} \otimes \phi_{j2\bar{\mu}}} \left\{ \sigma_{j2\bar{\mu}} \right\}. \quad (58)$$

Hence, it follows from eq. (56) and eq. (57), that their expectation values are equal. Accordingly, we can conclude from eq. (55), that

$$P_{\phi_{j1\bar{\mu}} \otimes \phi_{j2\bar{\mu}}} \left\{ \sigma_{j1\bar{\mu}} \cap \sigma_{j2\bar{\mu}} \mid \mathcal{F}_{jS} \right\} = P_{\phi_{j1\bar{\mu}} \otimes \phi_{j2\bar{\mu}}} \left\{ \sigma_{j1\bar{\mu}} \mid \mathcal{F}_{jS} \right\} \quad (59)$$

and similarly

$$P_{\phi_{j1\bar{\mu}} \otimes \phi_{j2\bar{\mu}}} \left\{ \sigma_{j1\bar{\mu}} \cap \sigma_{j2\bar{\mu}} \mid \mathcal{F}_{jS} \right\} = P_{\phi_{j1\bar{\mu}} \otimes \phi_{j2\bar{\mu}}} \left\{ \sigma_{j2\bar{\mu}} \mid \mathcal{F}_{jS} \right\}. \quad (60)$$

Using the assumption of passive locality, we get with eq. (59) and eq. (60):

$$\begin{aligned}
P_{\phi_{j1\bar{\mu}} \otimes \phi_{j2\bar{\mu}}} \{ \sigma_{j1\bar{\mu}} | \mathcal{F}_{jS} \} &= P_{\phi_{j1\bar{\mu}} \otimes \phi_{j2\bar{\mu}}} \{ \sigma_{j1\bar{\mu}} \cap \sigma_{j2\bar{\mu}} | \mathcal{F}_{jS} \} \\
&= P_{\phi_{j1\bar{\mu}} \otimes \phi_{j2\bar{\mu}}} \{ \sigma_{j1\bar{\mu}} | \mathcal{F}_{jS} \} P_{\phi_{j1\bar{\mu}} \otimes \phi_{j2\bar{\mu}}} \{ \sigma_{j2\bar{\mu}} | \mathcal{F}_{jS} \} \\
&= (P_{\phi_{j1\bar{\mu}} \otimes \phi_{j2\bar{\mu}}} \{ \sigma_{j1\bar{\mu}} | \mathcal{F}_{jS} \})^2.
\end{aligned} \tag{61}$$

This implies, that the conditional probability $P_{\phi_{j1\bar{\mu}} \otimes \phi_{j2\bar{\mu}}} \{ \sigma_{j1\bar{\mu}} | \mathcal{F}_{jS} \}$ can only have the two values 0 or 1. We denote the event, when this random variable has the value one by $\{x_{j1\bar{z}}\}$. Since $P_{\phi_{j1\bar{\mu}} \otimes \phi_{j2\bar{\mu}}} \{ \sigma_{j1\bar{\mu}} | \mathcal{F}_{jS} \}$ is measurable with respect to \mathcal{F}_{jS} , it follows that $\{x_{j1\bar{z}}\} \in \mathcal{F}_{jS}$.

If we compute the unconditional probability of $\{x_{j1\bar{z}}\}$ with the expectation value, we observe, that

$$P_{\phi_{j1\bar{\mu}} \otimes \phi_{j2\bar{\mu}}} \{ \sigma_{j1\bar{\mu}} \} = P_{\phi_{j1\bar{\mu}} \otimes \phi_{j2\bar{\mu}}} \{ x_{j1\bar{z}} \} \tag{62}$$

and

$$P_{\phi_{j1\bar{\mu}} \otimes \phi_{j2\bar{\mu}}} \{ \sigma_{j1\bar{\mu}} \cap x_{j1\bar{z}} \} = P_{\phi_{j1\bar{\mu}} \otimes \phi_{j2\bar{\mu}}} \{ x_{j1\bar{z}} \}. \tag{63}$$

The eqs. (62) and (63) lead to the conclusion that $\{x_{j1\bar{z}}\} \in \mathcal{F}_{jS}$ is equivalent to the event $\{ \sigma_{j1\bar{\mu}} \} \in \mathcal{F}_{j1}$. From

$$P_{\phi_{j1\bar{\mu}} \otimes \phi_{j2\bar{\mu}}} \{ \sigma_{j1\bar{\mu}} | \mathcal{F}_{jS} \} = (P_{\phi_{j1\bar{\mu}} \otimes \phi_{j2\bar{\mu}}} \{ \sigma_{j2\bar{\mu}} | \mathcal{F}_{jS} \})^2 \tag{64}$$

it follows, that the conditional probability $P_{\phi_{j1\bar{\mu}} \otimes \phi_{j2\bar{\mu}}} \{ \sigma_{j2\bar{\mu}} | \mathcal{F}_{jS} \}$ is equal to $P_{\phi_{j1\bar{\mu}} \otimes \phi_{j2\bar{\mu}}} \{ \sigma_{j1\bar{\mu}} | \mathcal{F}_{jS} \}$. The same calculation as in the eq. (63) for $\{ \sigma_{j2\bar{\mu}} \}$ then shows, that $\{x_{j1\bar{z}}\}$ is equivalent to $\{ \sigma_{j2\bar{\mu}} \}$ also.

Deterministic passive locality, which is a consequence of passive locality plus the observed equivalent events of quantum mechanics states: Every event at the detectors must be determined by another one at preparation stage.

The failure of deterministic passive locality implies for the spin trajectories in a hidden variable description of the EPR experiment: During the time of its flight from the source to the measurement apparatus, each particle has to undergo a stochastic process, constantly changing its spin orientation from up to down and vice-versa, with all events not determined by an earlier one at the source. Additionally, because of eq. (37), the spin directions of the two particles in every pair have to be opposite at any time. For a theory compatible with relativity theory, this all must happen such, that active locality is preserved.

If these conditions can be fulfilled, Bell's inequalities will be violated without instantaneous signalling between the two measurement stations 1 and 2. As shall be shown in the next sections of this article, there is no reason to believe, that hidden variable theories are unable to do so.

5 The locality problem in the hidden variable theory of Fritsche and Haugk

5.1 Derivation of the single particle Schrödinger equation from Brownian motion

In 2003, L. Fritsche and M. Haugk [19] published a modification of Nelson's stochastic mechanics. It is, in contrast to Nelson's original contribution [17, 18], an actively local theory with a non-markovian stochastic process. Moreover, it can, if slightly modified, be made to violate passive locality only. Here, we will not discuss in detail how the (many-particle) Schrödinger equation (and the corresponding quantum mechanical operators) can be derived from this process, because it is fully described in the original article. Instead, the focus of this section is on superposed states and we show the theory to be actively local, even if many particle systems are considered. In turn, we will have enough material to explain how the correlations in entangled states emerge.

The starting point to derive the modified stochastic mechanics of [19] is, to interpret the energy time uncertainty relation $\Delta E \Delta t \approx \hbar/2$ as an indication, that in quantum mechanics, energy is only conserved strictly in physical interactions, but not otherwise. Particles are assumed to be under statistical energy fluctuations. In the theory of Fritsche and Haugk, there occurs a deviation ΔE of the particles energy E during the time interval Δt , after which the initial energy E has to be restored on the average. Therefore, kinetic energy is thought to be conserved on the average only. The second assumption is in spirit of Ballentine's

famous statistical ensemble interpretation [51]. In this interpretation, the quantum state vector $|\psi\rangle$ does not represent an individual particle, but a statistical ensemble of infinitely many of them.

For this reason, the probability to find the electron of a hydrogen atom in a volume Ω at time t , given by

$$p(\Omega, t) \equiv \int_{\Omega} \psi^*(\vec{r}, t)\psi(\vec{r}, t)d^3r, \quad (65)$$

where $\psi(\vec{r}, t)$ is the electron's wave-function, describes the following situation: To get all the information of $p(\Omega, t)$ experimentally, one has to prepare infinitely many identical hydrogen atoms with the same state (including the same orientation in space). Then, one has to measure on each atom, if one finds the electron in Ω at time t after the preparation of the state. Finally, one must compute the electron's probability of being there in the infinitely large ensemble.

Accordingly, to describe the probabilistic time evolution of the statistical ensemble of electrons in a quantum mechanical system, one has to deal with an infinitely large number of sample trajectories. To model energy conservation on the average for this ensemble, Fritsche and Haugk divide these trajectories into two sub-ensembles with equal average velocities, and therefore the same average kinetic energy. In one sub-ensemble, every particle then undergoes a Brownian motion with a negative friction coefficient during a time-step Δt , and in the other one, each particle underlies the same process, but with a positive friction coefficient. After each time step, some of the particles are interchanged between the two sub-ensembles, until the kinetic energy of the latter agrees again. From this point, the entire procedure is repeated.

For a single particle of rest mass m_0 , we start our discussion with two Langevin equations for j -th trajectories on a coarse grained timescale τ , at which the Brownian motion process takes place:

$$\begin{aligned} m_0 \ddot{\vec{x}}_j^B(t) + \frac{m_0}{\tau} \dot{\vec{x}}_j^B(t) &= \vec{F}_j^{\text{ext}}(t) + \vec{F}_j^{\text{Brown}}(t) \\ m_0 \ddot{\vec{x}}_j^A(t) - \frac{m_0}{\tau} \dot{\vec{x}}_j^A(t) &= \vec{F}_j^{\text{ext}}(t) + \vec{F}_j^{\text{Brown}}(t) \end{aligned} \quad (66)$$

The three dimensional vector $\vec{x}_j^{\text{A(B)}}(t)$ defines the hidden position variable for the j -th sample trajectory of a particle in the $\{A\}$ ($\{B\}$) ensemble at time t , where $1 \leq j \leq S \in \mathbb{N}$. The vector $\vec{F}_j^{\text{ext}}(t)$ is an arbitrary external force field and $\vec{F}_j^{\text{Brown}}(t)$ denotes the random force the j -th particle is subjected to. This random force is assumed to have a Gaussian distribution

$$P\{\vec{F}_{jk}^{\text{Brown}}(t)\} = \frac{1}{\sqrt{\pi C}} e^{-\left(\vec{F}_{jk}^{\text{Brown}}(t)/C\right)^2} \quad (67)$$

in each coordinate, where C is a constant. The Gaussian distribution should be independent for every coordinate $1 \leq k \leq 3$ and trajectory j of the random force field. One can separate the sample trajectories $\vec{x}_j^{\text{A(B)}}$ into a sum, caused by the random and the external force:

$$\vec{x}_j^{\text{A(B)}}(t) = \vec{x}_{rj}^{\text{A(B)}}(t) + \vec{x}_{cj}^{\text{A(B)}}(t) \quad (68)$$

Because $\vec{F}_j^{\text{Brown}}(t)$ is independent in each coordinate, there does not exist any dependence between different components of the velocities $\dot{x}_{rj1}^{\text{A(B)}}(t)$, $\dot{x}_{rj2}^{\text{A(B)}}(t)$, and $\dot{x}_{rj3}^{\text{A(B)}}(t)$, as well as there is no dependence between $\vec{x}_{cj}^{\text{A(B)}}(t)$ and $\vec{x}_{rj}^{\text{A(B)}}(t)$. With this property in mind, the authors of [19] average in each of the two sub-ensembles $\{A\}$ and $\{B\}$ about all j -th sample trajectories $\vec{x}_j^{\text{A(B)}}(t)$, after the ensemble was made infinitely large ($S \rightarrow \infty$). Finally, they arrive at two differential equations:

$$\begin{aligned} \frac{\partial}{\partial t} (\vec{v}^B + \vec{u}^B) + (\vec{v}^B - \vec{u}^B) \cdot \vec{\nabla} (\vec{v}^B + \vec{u}^B) + \nu \Delta (\vec{v}^B + \vec{u}^B) &= \frac{1}{m_0} \vec{F}^{\text{ext}} \\ \frac{\partial}{\partial t} (\vec{v}^A - \vec{u}^A) + (\vec{v}^A + \vec{u}^A) \cdot \vec{\nabla} (\vec{v}^A - \vec{u}^A) - \nu \Delta (\vec{v}^A - \vec{u}^A) &= \frac{1}{m_0} \vec{F}^{\text{ext}} \end{aligned} \quad (69)$$

In eq. (69), $\vec{F}^{\text{ext}} \equiv (\vec{F}_1^{\text{ext}}(\vec{r}, t), \vec{F}_2^{\text{ext}}(\vec{r}, t), \vec{F}_3^{\text{ext}}(\vec{r}, t))$ denotes the average external force at point $\vec{r} \equiv (r_1, r_2, r_3)$ and time t . The vector valued entity

$$\vec{v}^{\text{A(B)}} \equiv \frac{\sum_{j=1}^{\infty} \epsilon_j(\vec{r}, t) \dot{\vec{x}}_j^{\text{A(B)}}(t)}{\sum_{j=1}^{\infty} \epsilon_j(\vec{r}, t)} = \left(v_1^{\text{A(B)}}(\vec{r}, t), v_2^{\text{A(B)}}(\vec{r}, t), v_3^{\text{A(B)}}(\vec{r}, t) \right) \quad (70)$$

is the average velocity of the ensemble, where:

$$\epsilon_j(\vec{r}, t) \equiv \begin{cases} 1 & \text{if } \vec{x}_j^{A(B)}(t) \text{ at time } t \text{ is in a small volume } d^3r \text{ around } \vec{r} \\ 0 & \text{otherwise} \end{cases}$$

Additionally, we have the so called osmotic average velocity $\vec{u}^{A(B)}$. It is connected to the probability density $\rho^{A(B)}(\vec{r}, t)d\vec{r}$ of finding a particle in the ensemble at point (\vec{r}, t) via

$$\vec{u}^{A(B)} \equiv \mp \nu \vec{\nabla} \ln \left(\rho^{A(B)}(\vec{r}, t) / \rho_0 \right) = \left(u_1^{A(B)}(\vec{r}, t), u_2^{A(B)}(\vec{r}, t), u_3^{A(B)}(\vec{r}, t) \right). \quad (71)$$

In eq. (71), ρ_0 is a constant reference density without physical importance and $\nu \equiv \frac{k_B T \tau}{m_0}$ is the friction coefficient of the Brownian motion, depending on the temperature T of the heat-bath and Boltzmann's constant k_B . Without loss of generalisation, ν can be written as

$$\nu \equiv \frac{\hbar}{2m_0}. \quad (72)$$

Since we derive a non-relativistic system, it is possible that the random and the external forces are for some times near infinity. Under this circumstances, a particle may be driven to $\vec{r} = \infty$ at a certain time-step t with an infinitely large velocity. However, in physical situations important for the non-locality issues discussed later in this article, relativistic effects certainly will occur, in the sense that particles are not allowed to move faster than the speed of light. Then, if an ensemble of quantum particles is emitted at a point (\vec{r}_0, t_0) , for example by a decaying atomic nucleus, it can only reach points (\vec{r}, t) within the future cone of (\vec{r}_0, t_0) . Accordingly, we restrict the functions $\vec{u}, \vec{v}, \rho(\vec{r}, t)$ such, that they are defined only for these (\vec{r}, t) in the future cone.

Before the two ensembles underwent a Brownian motion, it was ensured, that their average velocity was equal:

$$\vec{v}^B = \vec{v}^A \quad (73)$$

It is shown in [19], that this constraint implies $\vec{u}^B = \vec{u}^A$ also. After the first time step, eq. (73) does not hold anymore, since the {A} system gains energy compared to the {B} system. To prevent this, we have to interchange j_1, j_2, j_3, \dots trajectories of fast particles from the {A} ensemble, with as many trajectories j'_1, j'_2, j'_3, \dots of slow particles from the {B} system, until

$$\vec{v}^B(\vec{r}, t + \Delta t) = \vec{v}^A(\vec{r}, t + \Delta t) \equiv \vec{v}(\vec{r}, t + \Delta t)$$

holds. After this was done, we restart the diffusion process again for the next time-step, but with the new division into two sub-ensembles, where a condition similar to eq. (73) holds.

At every time step, we can compute the arithmetic mean of the two diffusion equations for the {A} and {B} systems. As is shown in [19], the time evolution of the overall system is described by the following diffusion equation:

$$\frac{\partial}{\partial t} \vec{v} + \left(\vec{v} \cdot \vec{\nabla} \right) \vec{v} - \left(\vec{u} \cdot \vec{\nabla} \right) \vec{u} + \nu \Delta \vec{u} = \frac{d}{dt} \vec{v} - \left(\vec{u} \cdot \vec{\nabla} \right) \vec{u} + \nu \Delta \vec{u} = \frac{1}{m_0} \vec{F}^{\text{ext}} \quad (74)$$

If the Brownian forces were absent, we would have $\vec{u} = 0$ with eq. (74) becoming the equation of motion of a classical particle.

For the description of quantum systems, will assume the random forces to be present. We try to simplify eq. (74) with the definition of a function

$$\psi(\vec{r}, t) \equiv \pm \sqrt{\rho(\vec{r}, t)} e^{i\varphi(\vec{r}, t)}.$$

This function is connected to the ensemble average velocity \vec{v} through $\vec{v} \equiv \frac{\hbar}{m_0} \vec{\nabla} \varphi(\vec{r}, t)$. (We note, that the latter equation can only be written, since one can show that \vec{v} is curl free.) After some computations, the authors of [19] are able to cast eq. (74) directly in form of the time dependent Schrödinger equation:

$$i\hbar \frac{\partial \psi(\vec{r}, t)}{\partial t} = \hat{H}(\vec{r}, t) \psi(\vec{r}, t) \quad (75)$$

In eq. (75), $\hat{H}(\vec{r}, t)$ denotes the Hamiltonian operator

$$\hat{H}(\vec{r}, t) \equiv -\frac{\hbar^2 \Delta}{2m_0} + \hat{V}^{\text{ext}}(\vec{r}, t), \quad (76)$$

with $\hat{V}^{\text{ext}}(\vec{r}, t)$ as the potential of the external forces $\vec{F}^{\text{ext}} \equiv -\vec{\nabla} \hat{V}^{\text{ext}}(\vec{r}, t)$. The stochastic process for the infinitely large ensemble of single particles with trajectories $\vec{x}_j(t)$ leads to a differential equation, encoding the ensemble average velocities \vec{v} and \vec{u} in a wave-function. For the interpretation of a quantum mechanical state $|\psi\rangle$, this implies, that all the information $|\psi\rangle$ carries, is about the evolution of average velocities from a statistical ensemble. It does reveal nothing about individual particles. One should mention, that the quantum mechanical operators, their eigenvalues and uncertainty relations can be derived from the given formalism, as is fully documented in [19].

5.2 Quantum states in superpositions

Before we can discuss the locality properties of the Fritsche-Haugk theory and the physics behind entanglement in many particle systems, we turn our attention to superpositions of two states in a single particle system. The Schrödinger equation is linear. Therefore, the sum of two solutions $\psi_\uparrow(\vec{r}, t)$ and $\psi_\downarrow(\vec{r}, t)$ i.e.

$$\Psi(\vec{r}, t) \equiv a\psi_\uparrow(\vec{r}, t) + b\psi_\downarrow(\vec{r}, t) \quad (77)$$

is also a solution of eq. (75). With eq. (77), one would get mixing terms in $\rho(\vec{r}, t)$:

$$\rho(\vec{r}, t) = a^2 \psi_\uparrow^2(\vec{r}, t) + b^2 \psi_\downarrow^2(\vec{r}, t) + a^* b \psi_\uparrow^*(\vec{r}, t) \psi_\downarrow(\vec{r}, t) + b^* a \psi_\downarrow^*(\vec{r}, t) \psi_\uparrow(\vec{r}, t) \quad (78)$$

In turn, the equations for eq. (71) would become quite complicated. Accordingly, we need to make a basis change to get simpler equations.

The physical information of the superposed state (77) is, that for a given particle of the ensemble, we have a probability of a^2 that it belongs to the $\psi_\uparrow(\vec{r}, t)$ ensemble and a probability of b^2 to find it in in the $\psi_\downarrow(\vec{r}, t)$ system. Since both states are solutions of the same Schrödinger equation, we can write it without loss of information in the form given below:

$$\begin{aligned} i\hbar \frac{\partial \Psi(\vec{r}, t)}{\partial t} &= \hat{H}(\vec{r}, t) \Psi(\vec{r}, t) \\ i\hbar \frac{\partial (b\psi_\uparrow(\vec{r}, t) + a\psi_\downarrow(\vec{r}, t))}{\partial t} &= \hat{H}(\vec{r}, t) (a\psi_\uparrow(\vec{r}, t) + b\psi_\downarrow(\vec{r}, t)) \\ &= \begin{cases} i\hbar \frac{\partial a\psi_\uparrow(\vec{r}, t)}{\partial t} = \hat{H}(\vec{r}, t) a\psi_\uparrow(\vec{r}, t) \\ + \\ i\hbar \frac{\partial b\psi_\downarrow(\vec{r}, t)}{\partial t} = \hat{H}(\vec{r}, t) b\psi_\downarrow(\vec{r}, t) \end{cases} \end{aligned} \quad (79)$$

$$(80)$$

With basis vectors in Dirac notation defined through $|\psi_\uparrow\rangle \equiv \begin{pmatrix} 1 \\ 0 \end{pmatrix} \equiv |+\rangle$ and $|\psi_\downarrow\rangle \equiv \begin{pmatrix} 0 \\ 1 \end{pmatrix} \equiv |-\rangle$, we get the following coefficients in coordinate representation:

$$c_\uparrow(\vec{r}, t) \equiv \langle \psi_\uparrow | \Psi(\vec{r}, t) \rangle = a\psi_\uparrow(\vec{r}, t) \quad \text{and} \quad c_\downarrow(\vec{r}, t) \equiv \langle \psi_\downarrow | \Psi(\vec{r}, t) \rangle = b\psi_\downarrow(\vec{r}, t)$$

Hence, the overall state vector $|\Psi(\vec{r}, t)\rangle$ becomes

$$|\Psi(\vec{r}, t)\rangle = c_\uparrow |\psi_\uparrow\rangle + c_\downarrow |\psi_\downarrow\rangle \quad (81)$$

and the probability density can be written without mixing terms:

$$\begin{aligned} \rho(\vec{r}, t) &= \langle \Psi(\vec{r}, t) | \Psi(\vec{r}, t) \rangle \\ &= a^2 \psi_\uparrow^*(\vec{r}, t) \psi_\uparrow(\vec{r}, t) + b^2 \psi_\downarrow^*(\vec{r}, t) \psi_\downarrow(\vec{r}, t) \\ &= a^2 \rho_\uparrow(\vec{r}, t) + b^2 \rho_\downarrow(\vec{r}, t) \end{aligned} \quad (82)$$

For a^2 and b^2 , it follows from $\int \rho_{\uparrow(\downarrow)}(\vec{r}_1, t) d^3r = 1$ and $\int \rho(\vec{r}, t) d^3\vec{r} = 1$, that $a^2 + b^2 = 1$.

We can derive the two Schrödinger equations for the states \uparrow and \downarrow in eq. (80) with the same methods as given before. The derivation begins with four Langevin equations

$$\begin{aligned}
m_0 \ddot{\vec{x}}_{j\uparrow}^B(t) + \frac{m_0}{\tau} \dot{\vec{x}}_{j\uparrow}^B(t) &= \vec{F}_j^{\text{ext}}(t) + \vec{F}_{j\uparrow}^{\text{Brown}}(t) \\
m_0 \ddot{\vec{x}}_{j\uparrow}^A(t) - \frac{m_0}{\tau} \dot{\vec{x}}_{j\uparrow}^A(t) &= \vec{F}_j^{\text{ext}}(t) + \vec{F}_{j\uparrow}^{\text{Brown}}(t) \\
m_0 \ddot{\vec{x}}_{j\downarrow}^B(t) + \frac{m_0}{\tau} \dot{\vec{x}}_{j\downarrow}^B(t) &= \vec{F}_j^{\text{ext}}(t) + \vec{F}_{j\downarrow}^{\text{Brown}}(t) \\
m_0 \ddot{\vec{x}}_{j\downarrow}^A(t) - \frac{m_0}{\tau} \dot{\vec{x}}_{j\downarrow}^A(t) &= \vec{F}_j^{\text{ext}}(t) + \vec{F}_{j\downarrow}^{\text{Brown}}(t)
\end{aligned} \tag{83}$$

and appropriate hidden trajectory variables for the $\{A \uparrow\}$, $\{B \uparrow\}$, $\{A \downarrow\}$ and $\{B \downarrow\}$ ensembles. After some steps, we are led to the following differential equations:

$$\begin{aligned}
\frac{\partial}{\partial t} (\vec{v}_{\uparrow}^B + \vec{u}_{\uparrow}^B) + (\vec{v}_{\uparrow}^B - \vec{u}_{\uparrow}^B) \cdot \vec{\nabla} (\vec{v}_{\uparrow}^B + \vec{u}_{\uparrow}^B) + \nu \Delta (\vec{v}_{\uparrow}^B + \vec{u}_{\uparrow}^B) &= \frac{1}{m_0} \vec{F}_{\uparrow}^{\text{ext}} \\
\frac{\partial}{\partial t} (\vec{v}_{\uparrow}^A - \vec{u}_{\uparrow}^A) + (\vec{v}_{\uparrow}^A + \vec{u}_{\uparrow}^A) \cdot \vec{\nabla} (\vec{v}_{\uparrow}^A - \vec{u}_{\uparrow}^A) - \nu \Delta (\vec{v}_{\uparrow}^A - \vec{u}_{\uparrow}^A) &= \frac{1}{m_0} \vec{F}_{\uparrow}^{\text{ext}} \\
\frac{\partial}{\partial t} (\vec{v}_{\downarrow}^B + \vec{u}_{\downarrow}^B) + (\vec{v}_{\downarrow}^B - \vec{u}_{\downarrow}^B) \cdot \vec{\nabla} (\vec{v}_{\downarrow}^B + \vec{u}_{\downarrow}^B) + \nu \Delta (\vec{v}_{\downarrow}^B + \vec{u}_{\downarrow}^B) &= \frac{1}{m_0} \vec{F}_{\downarrow}^{\text{ext}} \\
\frac{\partial}{\partial t} (\vec{v}_{\downarrow}^A - \vec{u}_{\downarrow}^A) + (\vec{v}_{\downarrow}^A + \vec{u}_{\downarrow}^A) \cdot \vec{\nabla} (\vec{v}_{\downarrow}^A - \vec{u}_{\downarrow}^A) - \nu \Delta (\vec{v}_{\downarrow}^A - \vec{u}_{\downarrow}^A) &= \frac{1}{m_0} \vec{F}_{\downarrow}^{\text{ext}}
\end{aligned} \tag{84}$$

The vectors $\vec{v}_{\uparrow(\downarrow)}$, $\vec{u}_{\uparrow(\downarrow)}$ are the ensemble average velocities of the states \uparrow (\downarrow). In this notation, the four equations in eq. (84) would define two separate particle systems, with the particles belonging to one ensemble $\{\uparrow\}$ or $\{\downarrow\}$ for all the time. For a given single particle of a superposed state, there has to be a probability greater than 0 for it, to be found in each one of these states. Therefore, the eqs. (84) cannot describe a state in a superposition and we have to modify the derivation of eq. (80).

The two systems are no more separate, if we demand, that the \vec{v}_{\uparrow}^B (\vec{v}_{\downarrow}^B) ensemble is averaged over with the \vec{v}_{\downarrow}^A (\vec{v}_{\uparrow}^A) system. To ensure energy conservation on the average, we are forced to do an interchange procedure, where the slower particles j_1, j_2, j_3, \dots from the $\{B \uparrow\}$ ($\{B \downarrow\}$) ensemble are interchanged with the same number of faster particles j'_1, j'_2, j'_3, \dots from the $\{A \downarrow\}$ ($\{A \uparrow\}$) system. This has to be done with as many particles until

$$\vec{v}_{\uparrow}^B(\vec{r}, t) = \vec{v}_{\downarrow}^A(\vec{r}, t), \quad \vec{u}_{\uparrow}^B(\vec{r}, t) = \vec{u}_{\downarrow}^A(\vec{r}, t) \quad \text{and} \quad \vec{v}_{\downarrow}^B(\vec{r}, t) = \vec{v}_{\uparrow}^A(\vec{r}, t), \quad \vec{u}_{\downarrow}^B(\vec{r}, t) = \vec{u}_{\uparrow}^A(\vec{r}, t) \tag{85}$$

holds. If we add the $\{A \uparrow\}$ and $\{B \downarrow\}$, as well as the $\{A \downarrow\}$ and the $\{B \uparrow\}$ ensembles, we arrive at two equations:

$$\begin{aligned}
\frac{\partial}{\partial t} (\vec{v}_{\uparrow}^B + \vec{v}_{\downarrow}^A) + [(\vec{v}_{\uparrow}^B \cdot \vec{\nabla}) \vec{v}_{\uparrow}^B - (\vec{u}_{\uparrow}^B \cdot \vec{\nabla}) \vec{u}_{\uparrow}^B] + [(\vec{v}_{\downarrow}^A \cdot \vec{\nabla}) \vec{v}_{\downarrow}^A - (\vec{u}_{\downarrow}^A \cdot \vec{\nabla}) \vec{u}_{\downarrow}^A] + \nu \Delta (\vec{u}_{\uparrow}^B + \vec{u}_{\downarrow}^A) \\
= \frac{1}{m_0} (\vec{F}_{\uparrow}^{\text{ext}} + \vec{F}_{\downarrow}^{\text{ext}}) \\
\frac{\partial}{\partial t} (\vec{v}_{\uparrow}^A + \vec{v}_{\downarrow}^B) + [(\vec{v}_{\uparrow}^A \cdot \vec{\nabla}) \vec{v}_{\uparrow}^A - (\vec{u}_{\uparrow}^A \cdot \vec{\nabla}) \vec{u}_{\uparrow}^A] + [(\vec{v}_{\downarrow}^B \cdot \vec{\nabla}) \vec{v}_{\downarrow}^B - (\vec{u}_{\downarrow}^B \cdot \vec{\nabla}) \vec{u}_{\downarrow}^B] + \nu \Delta (\vec{u}_{\downarrow}^B + \vec{u}_{\uparrow}^A) \\
= \frac{1}{m_0} (\vec{F}_{\uparrow}^{\text{ext}} + \vec{F}_{\downarrow}^{\text{ext}})
\end{aligned} \tag{86}$$

They can be summed together, yielding:

$$\begin{aligned}
\frac{\partial}{\partial t} (\vec{v}_{\uparrow}^B + \vec{v}_{\uparrow}^A + \vec{v}_{\downarrow}^B + \vec{v}_{\downarrow}^A) + [(\vec{v}_{\uparrow}^B \cdot \vec{\nabla}) \vec{v}_{\uparrow}^B - (\vec{u}_{\uparrow}^B \cdot \vec{\nabla}) \vec{u}_{\uparrow}^B] + [(\vec{v}_{\uparrow}^A \cdot \vec{\nabla}) \vec{v}_{\uparrow}^A - (\vec{u}_{\uparrow}^A \cdot \vec{\nabla}) \vec{u}_{\uparrow}^A] \\
+ [(\vec{v}_{\downarrow}^B \cdot \vec{\nabla}) \vec{v}_{\downarrow}^B - (\vec{u}_{\downarrow}^B \cdot \vec{\nabla}) \vec{u}_{\downarrow}^B] + [(\vec{v}_{\downarrow}^A \cdot \vec{\nabla}) \vec{v}_{\downarrow}^A - (\vec{u}_{\downarrow}^A \cdot \vec{\nabla}) \vec{u}_{\downarrow}^A] + \nu \Delta (\vec{u}_{\uparrow}^B + \vec{u}_{\uparrow}^A + \vec{u}_{\downarrow}^B + \vec{u}_{\downarrow}^A) \\
= \frac{2}{m_0} (\vec{F}_{\uparrow}^{\text{ext}} + \vec{F}_{\downarrow}^{\text{ext}})
\end{aligned} \tag{87}$$

Both equations in eq. (86) contain $\{A\}$ and $\{B\}$, as well as $\{\uparrow\}$ and $\{\downarrow\}$ systems. They describe the average behaviour of equations of motion under the same random and external forces. Hence, we have:

$$\vec{v}_{\uparrow}^B = \vec{v}_{\uparrow}^A \equiv \vec{v}_{\uparrow}, \quad \vec{v}_{\downarrow}^B = \vec{v}_{\downarrow}^A \equiv \vec{v}_{\downarrow} \quad \text{and} \quad \vec{u}_{\uparrow}^B = \vec{u}_{\uparrow}^A \equiv \vec{u}_{\uparrow}, \quad \vec{u}_{\downarrow}^B = \vec{u}_{\downarrow}^A \equiv \vec{u}_{\downarrow} \tag{88}$$

With eq. (88), we can write $2\vec{v}_\uparrow = \vec{v}_\uparrow^B + \vec{v}_\uparrow^A$ and $2\vec{v}_\downarrow = \vec{v}_\downarrow^B + \vec{v}_\downarrow^A$ as well as $2\vec{u}_\uparrow = \vec{u}_\uparrow^B + \vec{u}_\uparrow^A$ and $2\vec{u}_\downarrow = \vec{u}_\downarrow^B + \vec{u}_\downarrow^A$. Now, eq. (87) becomes:

$$\begin{aligned} \frac{\partial}{\partial t} (\vec{v}_\uparrow + \vec{v}_\downarrow) + \left[(\vec{v}_\uparrow \cdot \vec{\nabla}) \vec{v}_\uparrow - (\vec{u}_\uparrow \cdot \vec{\nabla}) \vec{u}_\uparrow \right] + \left[(\vec{v}_\downarrow \cdot \vec{\nabla}) \vec{v}_\downarrow - (\vec{u}_\downarrow \cdot \vec{\nabla}) \vec{u}_\downarrow \right] + \nu \Delta (\vec{u}_\uparrow + \vec{u}_\downarrow) \\ = \frac{1}{m_0} (\vec{F}^{\text{ext}} + \vec{F}^{\text{ext}}) \end{aligned} \quad (89)$$

This is a sum of two separate differential equations for the ensemble averages of the \uparrow and the \downarrow state. It may be written as

$$\begin{cases} \frac{d}{dt} \vec{v}_\uparrow - (\vec{u}_\uparrow \cdot \vec{\nabla}) \vec{u}_\uparrow + \nu \Delta \vec{u}_\uparrow = \frac{1}{m_0} \vec{F}^{\text{ext}} \\ + \\ \frac{d}{dt} \vec{v}_\downarrow - (\vec{u}_\downarrow \cdot \vec{\nabla}) \vec{u}_\downarrow + \nu \Delta \vec{u}_\downarrow = \frac{1}{m_0} \vec{F}^{\text{ext}} \end{cases} \quad (90)$$

from which the two Schrödinger equations eq. (80) can be derived.

The two conditions eq. (85) and eq. (88) imply, that the average velocities $\vec{v}_\uparrow = \vec{v}_\downarrow \equiv \vec{v}$ and $\vec{u}_\uparrow = \vec{u}_\downarrow \equiv \vec{u}$ are equal. Hence, the equations in (80) describe two oscillating systems with identical dynamics.

The procedure given above also works with an arbitrary number of states $|\psi_1(\vec{r}, t)\rangle, |\psi_2(\vec{r}, t)\rangle, \dots, |\psi_{\mathcal{N}}(\vec{r}, t)\rangle$ summed up to a general superposition with weighting factors a_i :

$$|\Psi(\vec{r}, t)\rangle \equiv \sum_i^{\mathcal{N}} a_i |\psi_i(\vec{r}, t)\rangle \quad (91)$$

The weighting factors of these states define different probabilities of a given particle to be found in one particular state. Therefore, the ensembles in the corresponding \mathcal{N} diffusion equations must contain different numbers of particles. The states $|\psi_i(\vec{r}, t)\rangle$ have to be computed with an ensemble of Sa_i^2 trajectories $j_1, j_2, j_2, \dots, j_{Sa_i^2}$. This ensures, that there is a chance of a_i^2 to find a particle in the ensemble associated with $|\psi_i(\vec{r}, t)\rangle$.

Furthermore, the exchange procedure must be defined differently. One has to interchange the trajectories j and j' between the ensembles $\{Bi\}, \{Ak\}$, with the state indices $1 \leq i, k \leq \mathcal{N}$. The numbers i, k have to be chosen randomly, but with equal probability for each state. The different probabilities of finding a given particle in the separate states $|\psi_i(\vec{r}, t)\rangle$ and $|\psi_k(\vec{r}, t)\rangle$ only arise due to the unequal size of their corresponding ensembles.

After the necessary interchanges have been taken place, one may add the \mathcal{N} differential equations in one step and split them up into separate equations for the $1, 2, 3 \dots, \mathcal{N}$ systems analogous to eq. (90). We remark, that because of eq. (71), nothing about the weighting factors a_i^2 in the probability density appears in \vec{v}, \vec{u} . Hence, one has to re-introduce this information after one gets from the separated system in eq. (80) to the superposed eq. (79).

5.3 Locality in the theory of Fritsche and Haugk with single particle systems

In this section, we turn to the locality properties of the stochastic process described above. We observe, that in the Langevin equations (66), there is no dynamical coupling between different trajectories $\vec{x}_{j\uparrow(\downarrow)}^{A(B)}(t)$ and $\vec{x}_{j'\uparrow(\downarrow)}^{A(B)}(t)$, where $j \neq j'$. $\vec{F}_j^{\text{ext}}(t)$ is an arbitrary force field, only acting on the j -trajectory and $\vec{F}_j^{\text{Brown}}(t)$ is required to be a Gaussian distribution independent for each j . Nevertheless, the time evolution of different trajectories will be correlated because of the interchange mechanism.

If something happens to a large set of trajectories $\vec{x}_{j'\uparrow(\downarrow)}^{A(B)}(t)$ at a point (\vec{r}', t') , and if this point lies in the past cone of another point (\vec{r}, t) , some of the changed trajectories may arrive at (\vec{r}, t) . The ensemble average $\vec{v}(\vec{r}, t)$ is computed with all trajectories in (\vec{r}, t) . Therefore, what has happened at (\vec{r}', t') can modify $\vec{v}(\vec{r}, t)$. The interchange procedure then compares the velocity of an arbitrary single particle trajectory $\vec{x}_{j\uparrow(\downarrow)}^{A(B)}(t)$, which has arrived at (\vec{r}, t) , with the changed $\vec{v}(\vec{r}, t)$. For the next time step, dependent on the result of the comparison, the particle might be placed in a new differential equation $\{B\} (\{A\})$, where $\vec{x}_{j\uparrow(\downarrow)}^{A(B)}(t)$ will be renamed to $\vec{x}_{j\downarrow(\uparrow)}^{B(A)}(t)$.

As a consequence, the particle trajectory $\vec{x}_{j\uparrow(\downarrow)}^{A(B)}(t)$ will be influenced at (\vec{r}, t) by what happened at (\vec{r}', t') , even if it never was at (\vec{r}', t') . This effect is important in double slit experiments, where a particle's trajectory, after the passage of one slit, depends on whether the second slit is open or not, no matter if the particle went through the second slit.

It is important, that the trajectories $\vec{x}_{j\uparrow(\downarrow)}^{A(B)}(t)$ at (\vec{r}, t) can only be influenced through the exchange mechanism, if the place (\vec{r}', t') where some intervention was made, lies in the past cone of (\vec{r}, t) . In turn, there does not exist any violation of active locality. Quantum effects like interference patterns arise in this theory solely due to the non-markovian property of the process that governs the trajectories.

However, by writing this, one should be aware, that we are still discussing a non relativistic system, where external (non-covariant) forces are able to produce instantaneous effects, even if we restricted the speed of the particles in section 5.1. To remove this last problem, one would need a covariant formulation of the Fritsche-Haugk theory. Fortunately, as we will see in section 5.3, the discussion of entanglement does not depend on this problem. The different particle ensembles of a many particle system will turn out to be dynamically uncoupled, at least if inter-particle forces are not present.

We emphasise, that the exchange procedure does not introduce a direct dynamical coupling between the trajectories. Instead, the comparison of the particle's velocity with the ensemble averages is only a model to account for the idea of kinetic energy conservation on the average in a stochastic theory. That there are in fact no dynamical couplings between the individual trajectories of the ensemble can be seen experimentally in so called self-interferometry experiments. The typical example is neutron interferometry [21], where "usually, at one time, there is only one neutron in the interferometer, if at all because at that time, the next neutron has not yet been born and is still contained in the uranium nuclei of the reactor fuel" [22].

5.4 Derivation of the many particle Schrödinger equation

In this section, we want to explore, whether active locality persists when we consider many particle systems. The many particle Schrödinger equation can be derived, if the three dimensional coordinate space of the single particle problem is expanded into a $3N$ dimensional space for N particles. We start with the Langevin equations of the j -th particle trajectory in the ensemble for an n -th particle state, where $1 \leq n \leq N$:

$$\begin{aligned} m_0 \ddot{\vec{x}}_{jn}^B(t) + \frac{m_0}{\tau} \dot{\vec{x}}_{jn}^B(t) &= \vec{F}_{jn}^{\text{ext}}(t) + \vec{F}_{jn}^{\text{Brown}}(t) + \sum_{\substack{n'=1 \\ n' \neq n}}^N \vec{F}_{jn}^{\text{ext}}(\vec{x}_{jn}^{A(B)}(t), \vec{x}_{j'n'}^{A(B)}(t), t) \\ m_0 \ddot{\vec{x}}_{jn}^A(t) - \frac{m_0}{\tau} \dot{\vec{x}}_{jn}^A(t) &= \vec{F}_{jn}^{\text{ext}}(t) + \vec{F}_{jn}^{\text{Brown}}(t) + \sum_{\substack{n'=1 \\ n' \neq n}}^N \vec{F}_{jn}^{\text{ext}}(\vec{x}_{jn}^{A(B)}(t), \vec{x}_{j'n'}^{A(B)}(t), t) \end{aligned} \quad (92)$$

In eq. (92), a sum of additional force terms $\vec{F}_{jn}^{\text{ext}}(\vec{x}_{jn}^{A(B)}(t), \vec{x}_{j'n'}^{A(B)}(t), t)$, corresponding to inter-particle forces between the particle states n and $n' \neq n$ may be present. From the two Langevin equations (92), the authors of [19] derive differential equations of the $\{A\}$ and $\{B\}$ systems, completely analogous to the single particle case:

$$\begin{aligned} \frac{\partial}{\partial t} (\vec{v}^{\text{NB}} + \vec{u}^{\text{NB}}) + (\vec{v}^{\text{NB}} - \vec{u}^{\text{NB}}) \cdot \vec{\nabla}^N (\vec{v}^{\text{NB}} + \vec{u}^{\text{NB}}) + \nu \Delta^N (\vec{v}^{\text{NB}} + \vec{u}^{\text{NB}}) &= \vec{F}_{\text{ext}}^N \\ \frac{\partial}{\partial t} (\vec{v}^{\text{NA}} - \vec{u}^{\text{NA}}) + (\vec{v}^{\text{NA}} + \vec{u}^{\text{NA}}) \cdot \vec{\nabla}^N (\vec{v}^{\text{NA}} - \vec{u}^{\text{NA}}) - \nu \Delta^N (\vec{v}^{\text{NA}} - \vec{u}^{\text{NA}}) &= \vec{F}_{\text{ext}}^N \end{aligned} \quad (93)$$

In contrast to the single particle case, \vec{F}_{ext}^N is a $3N$ dimensional vector:

$$\vec{F}_{\text{ext}}^N \equiv \left(\left(\frac{\vec{F}_{n=1}^{\text{ext}}(\vec{r}_1, t)}{m_{01}} + \sum_{\substack{n'=1 \\ n' \neq n}}^N \frac{\vec{F}_{n=1}^{\text{ext}}(\vec{r}_1, \vec{r}_{n'}, t)}{m_{01}} \right), \dots, \left(\frac{\vec{F}_{n=N}^{\text{ext}}(\vec{r}_N, t)}{m_{0N}} + \sum_{\substack{n'=1 \\ n' \neq n}}^N \frac{\vec{F}_{n=N}^{\text{ext}}(\vec{r}_N, \vec{r}_{n'}, t)}{m_{0N}} \right) \right)$$

It consists of the average single particle force $\vec{F}_n^{\text{ext}}(\vec{r}_n, t)$ and of an average inter-particle force $\vec{F}_n^{\text{ext}}(\vec{r}_n, \vec{r}_{n'}, t)$. Both forces are acting on the n -th particle state with rest mass m_{0n} . Similarly, the three dimensional ensemble averages \vec{u} and \vec{v} are replaced by $3N$ dimensional entities. They depend on the $3N$ coordinates

$\vec{r}^N \equiv (\vec{r}_1, \vec{r}_2, \dots, \vec{r}_N)$ from the various particles in the following way:

$$\vec{v}^{NA(B)} \equiv \left(\vec{v}_1^{A(B)}(\vec{r}_1, t), \dots, \vec{v}_N^{A(B)}(\vec{r}_N, t) \right) \quad \text{and} \quad \vec{u}^{NA(B)} \equiv \left(\vec{u}_1^{A(B)}(\vec{r}_1, t), \dots, \vec{u}_N^{A(B)}(\vec{r}_N, t) \right)$$

The components of the ensemble averages $\vec{u}_n^N(\vec{r}_n^N, t)$ and $\vec{v}_n^N(\vec{r}_n^N, t)$ are computed with all the j -th trajectories for each single particle state n separately. This is the reason, why \vec{v}_n^N and \vec{u}_n^N only depend on the same n -th component of \vec{r}_n^N .

To derive the many particle Schrödinger equation, an interchange between the trajectories corresponding to slow particles from the $\{B\}$ ensemble and fast particles from the $\{A\}$ ensemble is necessary after each time-step until

$$\vec{v}^{NA}(\vec{r}^N, t) = \vec{v}^{NB}(\vec{r}^N, t) \quad (94)$$

holds.

Here, we have to note an additional subtlety which does only arise in many particle systems. It has to be ensured, that exchanges between two trajectories $\vec{x}_{jn}^{A(B)}(t)$ and $\vec{x}_{j'n'}^{B(A)}(t)$ are confined within the same particle state $n = n'$. An interchange between trajectories of different states n and $n' \neq n$ would couple the corresponding particle ensembles, even if no inter-particle forces are present. This would cause a general impossibility to define single particle operators which act on only one state n of a many particle system. Fortunately, due to the specific coordinate dependence of the ensemble averages \vec{v}^N and \vec{u}^N , eq. (94) is defined for every state n separately. Therefore, the exchange procedure can well be confined within each single particle state only.

After the trajectory exchange, one can compute the arithmetic mean of the two differential equations in eq. (93) and gets, analogous to the single particle case, a simple diffusion equation:

$$\frac{d}{dt} \vec{v}^N - \left(\vec{u}^N \cdot \vec{\nabla}^N \right) \vec{u}^N + \nu \Delta^N \vec{u}^N = \vec{F}_{\text{ext}}^N. \quad (95)$$

From eq. (95), the many particle Schrödinger equation

$$i\hbar \frac{\partial \Psi(\vec{r}^N, t)}{\partial t} = \hat{H}(\vec{r}^N, t) \Psi(\vec{r}^N, t) \quad (96)$$

can be derived, similar to the single particle case before. In eq. (96) the Hamiltonian operator becomes

$$\hat{H}(\vec{r}^N, t) \equiv \sum_{n=1}^N \frac{-\hbar^2 \Delta_n^N}{2m_{0n}} + \hat{V}_{\text{ext}}(\vec{r}^N, t). \quad (97)$$

Its external potential

$$\hat{V}_{\text{ext}}(\vec{r}^N, t) \equiv \sum_{n=1}^N \hat{V}_{\text{ext}}(\vec{r}_n, t) + \frac{1}{2} \sum_{n=1}^N \sum_{\substack{n'=1 \\ n' \neq n}}^N \hat{V}_{\text{ext}}(\vec{r}_n, \vec{r}_{n'}, t) \quad (98)$$

is connected to the n -th component of \vec{F}_{ext}^N via $\vec{F}_{\text{ext},n}^N \equiv -\frac{\vec{\nabla}_n^N}{m_{0n}} V_{\text{ext}}(\vec{r}^N, t)$.

5.5 Locality in the theory of Fritsche and Haugk with many particle systems

In this section, we assume all inter-particle terms to be absent. We observe, that with the absence of inter-particle forces like $\vec{F}_{jn}^{\text{ext}}(\vec{x}_{jn}^{A(B)}(t), \vec{x}_{j'n'}^{B(A)}(t), t)$, there are no terms in eq. (92) which interconnect trajectories $\vec{x}_{jn}^{A(B)}(t)$ and $\vec{x}_{j'n'}^{B(A)}(t)$, from different particle states $n' \neq n$. This is true for the ensemble averages $\vec{v}_n^N(\vec{r}_n^N, t)$ and $\vec{u}_n^N(\vec{r}_n^N, t)$ also, since they are calculated with the $\vec{x}_{jn}^{A(B)}(t)$ trajectories for each single particle state n separately.

The decision, if a trajectory $\vec{x}_{jn}^{A(B)}(t)$ changes to $\vec{x}_{jn}^{B(A)}(t)$ during the interchange procedure only relies on the (j, n) -th particle's velocity $\left| \dot{\vec{x}}_{jn}^{A(B)}(t) \right|$ and the ensemble average $\vec{v}_n^N(\vec{r}_n^N, t)$ of its state n . Therefore,

the trajectories $\vec{x}_{jn}^{A(B)}(t)$ are never coupled in any way to trajectories $\vec{x}_{j'n'}^{A(B)}(t)$ from other particle states $n' \neq n$.

The ensemble average $\vec{v}_n^N(\vec{r}_n^N, t)$ for the n -th particle may depend on j different sample trajectories $\vec{x}_{jn}^{A(B)}(t)$, as long as they belong to the state n and lie in the past cone of (\vec{r}_n^N, t) . However, even such a dependence does not exist between $\vec{v}_n^N(\vec{r}_n^N, t)$ and the trajectories $\vec{x}_{j'n'}^{A(B)}(t)$ of other states $n' \neq n$ in a many particle system. As a result, we conclude, that active locality holds in the hidden variable theory of Fritsche and Haugk.

5.6 Entangled states in the model of Fritsche and Haugk

That there does not exist any dynamical coupling between different particle states which a many particle system is composed of, doesn't imply, that correlations between them are forbidden. In section 8, we will introduce further conditions on the Gaussian distributed Brownian forces $\vec{F}_{jn}^{\text{Brown}}(t)$ in a completely local framework. These conditions will be such, that under certain circumstances, the trajectories $\vec{x}_{jn}^{A(B)}(t)$ change their ensemble from $\{A\}$ ($\{B\}$) to $\{B\}$ ($\{A\}$) simultaneously with the trajectories $\vec{x}_{j'=j, n'}^{A(B)}(t)$ from another state $n' \neq n$. Then, correlations between the particle ensembles, as well as their wave-functions will emerge. The resulting many particle states are of the form

$$|\Psi(\vec{r}_1, \vec{r}_2, t)\rangle \equiv \frac{1}{\sqrt{2}} (|\psi_1(\vec{r}_1, t)\rangle \otimes |\psi_2(\vec{r}_2, t)\rangle - |\psi_2(\vec{r}_1, t)\rangle \otimes |\psi_1(\vec{r}_2, t)\rangle) \quad (99)$$

and commonly called entangled. They were observed in massive fermionic systems for the first time in flavour states of B-Mesons [23]. Those particles have the unwanted property to decay rapidly and the violation of Bell's inequality could only be confirmed after an interpolation. A later experiment used spin states of entangled protons [24]. They are stable particles and therefore, no interpolation was necessary to ascertain a violation of the CHSH inequalities on a confidence level of 99.3%.

The overall state vector $|\Psi(\vec{r}_1, \vec{r}_2, t)\rangle$ is an element in the product Hilbert space $\mathcal{H} \equiv \mathcal{H}_1 \otimes \mathcal{H}_2$. It is a superposition of the two product states

$$|\chi_1(\vec{r}_1, \vec{r}_2, t)\rangle \equiv |\psi_1(\vec{r}_1, t)\rangle \otimes |\psi_2(\vec{r}_2, t)\rangle \quad (100)$$

and

$$|\chi_2(\vec{r}_1, \vec{r}_2, t)\rangle \equiv |\psi_2(\vec{r}_1, t)\rangle \otimes |\psi_1(\vec{r}_2, t)\rangle, \quad (101)$$

where $|\psi_1(\vec{r}_1, t)\rangle, |\psi_2(\vec{r}_1, t)\rangle \in \mathcal{H}_1$ and $|\psi_1(\vec{r}_2, t)\rangle, |\psi_2(\vec{r}_2, t)\rangle \in \mathcal{H}_2$ denote the single particle solutions. The entangled state solves a many-particle Schrödinger equation with an Hamiltonian operator

$$\hat{H}(\vec{r}_1, \vec{r}_2, t) \equiv \sum_{n=1}^2 \frac{-\hbar^2 \Delta_n^N}{2m_{0n}} + \sum_{n=1}^2 \hat{V}_{\text{ext}}(\vec{r}_n, t), \quad (102)$$

which has no inter-particle forces and whose external potentials agree at each time t :

$$\hat{V}_{\text{ext}}(\vec{r}_1, t) = \hat{V}_{\text{ext}}(\vec{r}_2, t) \quad (103)$$

The corresponding Schrödinger equation can be written, analogous to eq. (80), as:

$$\begin{aligned} \frac{i\hbar \partial \Psi(\vec{r}_1, \vec{r}_2, t)}{\partial t} &= \hat{H}(\vec{r}_1, \vec{r}_2, t) \Psi(\vec{r}_1, \vec{r}_2, t) \\ \frac{i\hbar}{\sqrt{2}} \frac{\partial \chi_1(\vec{r}_1, \vec{r}_2, t)}{\partial t} - \frac{i\hbar}{\sqrt{2}} \frac{\partial \chi_2(\vec{r}_1, \vec{r}_2, t)}{\partial t} &= \frac{1}{\sqrt{2}} \hat{H}(\vec{r}_1, \vec{r}_2, t) \chi_1(\vec{r}_1, \vec{r}_2, t) - \frac{1}{\sqrt{2}} \hat{H}(\vec{r}_1, \vec{r}_2, t) \chi_2(\vec{r}_1, \vec{r}_2, t) \\ &= \begin{cases} \frac{i\hbar}{\sqrt{2}} \frac{\partial \chi_1(\vec{r}_1, \vec{r}_2, t)}{\partial t} = \frac{1}{\sqrt{2}} \hat{H}(\vec{r}_1, \vec{r}_2, t) \chi_1(\vec{r}_1, \vec{r}_2, t) \\ - \\ \frac{i\hbar}{\sqrt{2}} \frac{\partial \chi_2(\vec{r}_1, \vec{r}_2, t)}{\partial t} = \frac{1}{\sqrt{2}} \hat{H}(\vec{r}_1, \vec{r}_2, t) \chi_2(\vec{r}_1, \vec{r}_2, t) \end{cases} \quad (104) \end{aligned}$$

The system consists of two ensembles \vec{r}_1 and \vec{r}_2 , the particles of which can be in different states $|\psi_1(\vec{r}_1)\rangle$ or $|\psi_2(\vec{r}_1)\rangle$ for the \vec{r}_1 ensemble and $|\psi_1(\vec{r}_2)\rangle$ or $|\psi_2(\vec{r}_2)\rangle$ for the \vec{r}_2 system. To describe the dynamics of

the \vec{r}_1 particles, the $\{A1\}$ ($\{B1\}$) ensemble of $|\psi_1(\vec{r}_1)\rangle$ must be averaged with the $\{B2\}$ ($\{A2\}$) ensemble of $|\psi_2(\vec{r}_1)\rangle$ and the same has to be done with the \vec{r}_2 particles. During this procedure, one has to ensure that each j -th pair of particle trajectories in the \vec{r}_1 and \vec{r}_2 system changes its state simultaneously. We will show in the context with Bohm's EPR experiment, how this can be achieved, even if the trajectories of the two different ensembles are disconnected from each other. For now, we only remark, that without inter-particle-forces, it is in fact a requirement for the particle trajectories $\vec{x}_{j1}^{A(B)}(t)$ and $\vec{x}_{j'2}^{A(B)}(t)$ not to be interconnected directly in any way. This may also be seen with the following example by Edward Nelson [8]:

If one induces an external measurement process on the n -th particle of a multi-particle system, one may get information about the hidden variable $\vec{x}_{jn}^{A(B)}(t)$. The measurement device may change $\vec{x}_{jn}^{A(B)}(t)$ in a way, that correlations with all the other particle trajectories $\vec{x}_{j'n'}^{A(B)}(t)$ are lost. Since the different particle systems are uncoupled, an interaction happening on the n -th particle ensemble can in no way influence the properties of the particles in the n' system.

Let $\mathcal{H} \equiv \mathcal{H}_1 \otimes \mathcal{H}_2$ be the product Hilbert space above, with a Hamiltonian operator

$$\hat{H} \equiv \hat{H}_1 \otimes 1_2 + 1_1 \otimes \hat{H}_2, \quad (105)$$

where $\hat{H}_1 \in \mathcal{H}_1$ and $\hat{H}_2 \in \mathcal{H}_2$. The time evolution for any observable $\hat{A}_1 \in \mathcal{H}_1$ in the Heisenberg picture

$$\hat{A}_{\hat{H}}(t) \equiv e^{it\hat{H}} \hat{A}_1 e^{-it\hat{H}} = e^{it\hat{H}_1} \hat{A}_1 e^{-it\hat{H}_1} \otimes 1_2 \quad (106)$$

is completely independent of \mathcal{H}_2 , even if $|\Psi(\vec{r}_1, \vec{r}_2, t)\rangle \in \mathcal{H}$ is described by the entangled state of eq. (99). As Nelson put it: „no matter how systems may be correlated, there is no way of telling, what [external] forces the second system is being subjected to, as long as they are not dynamically coupled.”

In this context, we mention that Nelson's original theory could not reproduce the example above [8]. The reason was, that Nelson identified $\vec{v}(\vec{r}, t)$ and $\vec{u}(\vec{r}, t)$ not as average velocities of an ensemble, but as real particle properties. With this assertion, neither the splitting of the many particle Schrödinger equation in eq. (104), nor any sort of interchanging procedure between the particles in the $\{A1\}$ ($\{B1\}$) and $\{B2\}$ ($\{A2\}$) ensemble was possible. Accordingly, active locality was lost in Nelson's original formulation of stochastic mechanics, similarly as in Bohmian mechanics. For the latter (see [48] for a general introduction), it is well known, that the derivation of the many particle Schrödinger equation is only possible, if a highly non-trivial dependency between the Bohmian “trajectories” of different particle states has to be introduced, which is at odds with the theory of relativity and without any physical foundation. Interestingly, Nelson himself has shown recently in [26], that $\vec{v}(\vec{r}, t)$ and $\vec{u}(\vec{r}, t)$ cannot correspond to properties of individual particles, since this would lead to measurable differences between his original theory and ordinary quantum mechanics. A similar observation was made for Bohmian “trajectories” earlier in [25].

6 Spin in a Fritsche-Haug like hidden variable theory

6.1 Simple estimations

Nelson's derivation of the Schrödinger equation was quickly extended to include particle spin, but without proper physical interpretation [27]. Over the years, more physically motivated derivations of the Pauli equation were developed. One by Faris [28, 29] and another one by the authors of the modification of Nelson's stochastic mechanics discussed above [30]. Unfortunately, these frameworks did either violate active locality, or they did not contain any mechanism, which would change a particle's spin state after its emission. Thereby, none of these theories led to a violation of deterministic passive locality only. In this section, we propose a theory of spinning particles providing such a mechanism. Beginning with an example of [30], we try to interpret the spin of a particle as rotation around a chosen axis \vec{z} (see fig. 3). We let the particle rotate around of this axis under energy fluctuations $\Delta E \Delta t \approx \hbar/2$. For the rotational energy, we have from classical mechanics $E = \frac{\Theta}{2} \omega_0^2$. The inertial tensor is given by $\Theta = m_0 r^2$ with a distance r from the chosen axis. The angular velocity is $\omega_0 = \frac{2\pi}{T}$ and T is the time of a complete rotation. If we identify Δt with T and ΔE with E , we may write

$$\Delta E \Delta t = \frac{1}{2} m_0 r \cdot r \frac{2\pi}{T} \frac{2\pi}{T} T = m_0 v r \pi = \pi s \approx \hbar/2. \quad (107)$$




Figure 3: A particle with spin \uparrow , rotating in the left direction with respect to the \vec{z} axis

Here, $v = \frac{2\pi r}{T}$ is the magnitude of the particle's velocity and

$$s = m_0 v r \approx \frac{\hbar}{2\pi} \quad (108)$$

defines the magnitude of its angular momentum. Apparently, s is independent of the particle's mass and the radius of the rotation. The arguments above were only estimations. For later, it will be useful, if we set the magnitude of the spin angular momentum to $s = \hbar/2$ on the average. From the time dependent Schrödinger equation, one can derive the so called Ehrenfest theorem $\frac{d}{dt} \langle \hat{v} \rangle = \langle \vec{F}^{\text{ext}} \rangle$, where $\langle \hat{v} \rangle$ is the expectation value of the velocity operator. This equation implies, that the angular momentum operator $\hat{L} = \hat{r} \times m_0 \hat{v}$ may not vanish in the case of $\hbar \rightarrow 0$. Since the so called spin should go to zero in the classical limit, s must be independent of \hat{L} and we must expand our description.

6.2 The Stern-Gerlach effect

If particles of spin $1/2$ are not specially prepared, they occur in a superposition of \uparrow and \downarrow states. To derive such a superposition, we have to make a similar computation as in section 5.2. Here however, the trajectories $\vec{x}_{j\uparrow(\downarrow)}^{A(B)}(t)$ of eq. (83) describe particles in rotational motion. Their rotation can be in the left- or in the right direction with respect to the chosen axis \vec{z} . Particles rotating in the left direction are denoted by \uparrow , whereas particles rotating in the right direction are identified by \downarrow . Eventually, we arrive after some calculations at the four differential equations of eq. (84).

As explained before, the interpretation of a state in a superposition is, that a given particle from the superposed system must have a non-vanishing probability to be found in each one of the component states. Accordingly, we have to interchange trajectories between the $\{A \uparrow\}$ ($\{A \downarrow\}$) and $\{B \downarrow\}$ ($\{B \uparrow\}$) ensembles, before we get to the differential equations in eq. (86).

For a single particle, the effect of this exchange procedure is, that it does as often belong to the \uparrow state as to the \downarrow state. A nice drawing of this process is given by Penrose for Dirac particles in [34]. It is this procedure, whereby the possibility for a violation of deterministic passive locality does arise.

At the end of the calculation in section 5.2, we were able to describe the behaviour of the overall system by a probability density

$$\rho(\vec{r}, t) = a^2 \rho_{\uparrow}(\vec{r}, t) + b^2 \rho_{\downarrow}(\vec{r}, t) \quad (109)$$

and a quantum state

$$|\Psi(\vec{r}, t)\rangle \equiv \begin{pmatrix} a\psi_{\uparrow}(\vec{r}, t) \\ b\psi_{\downarrow}(\vec{r}, t) \end{pmatrix} = a\psi_{\uparrow}(\vec{r}, t) |+\rangle + b\psi_{\downarrow}(\vec{r}, t) |-\rangle, \quad (110)$$

both in form of a sum composed of the two rotating systems.

From here on, we can closely follow [30] to derive the rest of quantum mechanics for spinning particles. If an external magnetic field $B_{\vec{z}}(\vec{r}, t)$ in the direction of \vec{z} is applied, the energy of $\psi_{\uparrow}(\vec{r}, t)$ decreases with magnitude $\mu_B B_{\vec{z}}(\vec{r}, t)$, whereas the energy of $\psi_{\downarrow}(\vec{r}, t)$ increases by the same magnitude. The constant μ_B is called Bohr magneton

$$\mu_B \equiv \frac{e\hbar}{2m_0}. \quad (111)$$

It emerges, since the relativistic Dirac equation is, in contrast to the Pauli equation to be derived here, linear in all its derivatives. The necessity of this linearity, which is absent in the non-relativistic theory, changes the interaction of angular momentum with magnetic fields. If one does linearise the Schrödinger equation in a similar manner as it is typically done for Dirac's equation, one gets the same value for μ_B [31].

Under $B_{\vec{z}}(\vec{r}, t)$, the change of the particle's energy densities caused by their interaction with the magnetic field is

$$\mu_{\uparrow}(\vec{r}, t) \equiv -\mu_B B_{\vec{z}}(\vec{r}, t) |a|^2 \psi_{\uparrow}^*(\vec{r}, t) \psi_{\uparrow}(\vec{r}, t) \quad (112)$$

and

$$\mu_{\downarrow}(\vec{r}, t) \equiv \mu_B B_{\vec{z}}(\vec{r}, t) |b|^2 \psi_{\downarrow}^*(\vec{r}, t) \psi_{\downarrow}(\vec{r}, t). \quad (113)$$

With their sum

$$\mu(\vec{r}, t) \equiv \mu_{\uparrow}(\vec{r}, t) + \mu_{\downarrow}(\vec{r}, t), \quad (114)$$

the total change $\mu(\vec{r}, t)$ for both spinors can be cast as

$$\mu(\vec{r}, t) = -\Psi^+(\vec{r}, t) \mu_B \mathbf{B}_{\vec{z}} \Psi(\vec{r}, t), \quad (115)$$

where

$$\mathbf{B}_{\vec{z}} \equiv B_{\vec{z}}(\vec{r}, t) \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \quad (116)$$

The different coupling of the systems $\psi_{\uparrow(\downarrow)}(\vec{r}, t)$ to magnetic fields gives rise to the Stern-Gerlach effect [32]. If atoms from an incident beam with equally distributed magnetic energy density enter a strong magnet, the two spin systems are subjected to different forces $\vec{F}_{j\uparrow}^{\text{ext}}(t) \neq \vec{F}_{j\downarrow}^{\text{ext}}(t)$. Accordingly, they get deflected in opposite directions. Due to active locality, the interchanging procedure can only happen between trajectories $\vec{x}_{j\uparrow(\downarrow)}^{\text{A(B)}}(t)$ which are located at the same point (\vec{r}, t) in space-time. However, the Stern-Gerlach magnet separates trajectories, which belong to different spin states spatially. Then, the exchange procedure can take place between trajectories of the same spin ensemble only. This is observed in so called double Stern-Gerlach experiments [33]. If one would place a second Stern-Gerlach magnet behind the first and let it have the same directional orientation, the parts of the beam which were previously separated, would not split up again. In this way, the particle ensembles of a beam can be prepared to contain one spin state only. The repeated Stern-Gerlach experiment demonstrates both the locality and the non-markovian property of the stochastic process involved. The stochastic differential equation that a particle obeys at a given time depends, due to the comparison with the ensemble averages and the interchanging procedure, on what has happened to the ensemble before. Therefore, the stochastic process acting on a single particle has a kind of memory about the past interactions between the magnets and the ensemble the individual particle is placed in. Nevertheless, the stochastic process is a local one. After the two spin systems are spatially separated, they do not influence each other anymore.

6.3 The Pauli equation

In [19], the Schrödinger equation under a magnetic field with its Hamiltonian

$$\hat{\mathbf{H}}(\vec{r}, t) \equiv \frac{\left(-i\hbar\vec{\nabla} - e\hat{\mathbf{A}}(\vec{r}, t)\right)^2}{2m_0} + e\Phi(\vec{r}, t) + \hat{\mathbf{V}}^{\text{ext}}(\vec{r}, t) \quad (117)$$

was derived from a Brownian motion process. If the two rotating systems undergo this process, their overall energy density maybe written as superposition

$$w(\vec{r}, t) = \psi_{\uparrow}^*(\vec{r}, t) \hat{\mathbf{H}}(\vec{r}, t) \psi_{\uparrow}(\vec{r}, t) + \psi_{\downarrow}^*(\vec{r}, t) \hat{\mathbf{H}}(\vec{r}, t) \psi_{\downarrow}(\vec{r}, t) = \Psi^+(\vec{r}, t) \hat{\mathbf{H}} \Psi(\vec{r}, t), \quad (118)$$

where we have used the definition

$$\hat{\mathbf{H}} \equiv \hat{\mathbf{H}}(\vec{r}, t) \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}. \quad (119)$$




Figure 4: Two coordinate systems with unit vectors $(\vec{x}, \vec{y}, \vec{z})$ and $(\vec{x}', \vec{y}', \vec{z}')$ related by Euler angles ψ , ϕ and θ

To include the energy from the interaction of the spin rotation with $B_z(\vec{r}, t)$, we must add $\mu(\vec{r}, t)$ and get the Hamiltonian $\hat{\mathbf{H}} - \mu_B \mathbf{B}_z$. We note, that the effect of $\mu_B \mathbf{B}_z$ on each ensemble is that of a potential. Since the time dependent Schrödinger equation was derived under an arbitrary external potential, we can write a first equation for a spinning particle:

$$i\hbar \frac{\partial \Psi(\vec{r}, t)}{\partial t} = \left(\hat{\mathbf{H}} - \mu_B \mathbf{B}_z \right) \Psi(\vec{r}, t) \quad (120)$$

If we want to measure the particle's spin rotation around an axis \vec{z}' that is different from the axis defined by the z component $B_z(\vec{r}, t)$ of the magnetic field, we must change the coordinate system of $\vec{B}(\vec{r}, t)$. This can be done via linear transformations $\vec{B}'(\vec{r}, t) \equiv \mathbf{Q} \vec{B}(\vec{r}, t)$, where \mathbf{Q} defines a 3×3 rotation matrix which transforms $(B_x(\vec{r}, t), B_y(\vec{r}, t), B_z(\vec{r}, t))$ into $\vec{B}'(\vec{r}, t) = (B_{x'}(\vec{r}, t), B_{y'}(\vec{r}, t), B_{z'}(\vec{r}, t))$. It is known from classical mechanics (see e.g. [35]), that by using Euler angles ψ , ϕ and θ as shown in fig. 4, any 3×3 rotation matrix can be converted in 2×2 form to

$$\mathbf{Q} \equiv \begin{pmatrix} e^{\frac{i}{2}(\psi+\phi)} \cos(\theta/2) & i e^{\frac{i}{2}(\psi-\phi)} \sin(\theta/2) \\ i e^{-\frac{i}{2}(\psi-\phi)} \sin(\theta/2) & e^{-\frac{i}{2}(\psi+\phi)} \cos(\theta/2) \end{pmatrix}. \quad (121)$$

To apply the 2×2 matrix \mathbf{Q} on $\vec{B}(\vec{r}, t)$, we must rewrite the magnetic field:

$$\mathbf{B} \equiv \begin{pmatrix} B_z(\vec{r}, t) & B_x(\vec{r}, t) - i B_y(\vec{r}, t) \\ B_x(\vec{r}, t) + i B_y(\vec{r}, t) & -B_z(\vec{r}, t) \end{pmatrix} \quad (122)$$

Then, one can compute \mathbf{B}' via

$$\mathbf{B}' = \mathbf{Q} \mathbf{B} \mathbf{Q}^+. \quad (123)$$

We note, that a vector $\vec{B}(\vec{r}, t) = (0, 0, B_z(\vec{r}, t))$ has a matrix form equal to \mathbf{B}_z in eq. (116). Furthermore, it can be observed that

$$\mathbf{B} \equiv B_x(\vec{r}, t) \sigma_x + B_y(\vec{r}, t) \sigma_y + B_z(\vec{r}, t) \sigma_z, \quad (124)$$

where $\sigma_x, \sigma_y, \sigma_z$ are the so called Pauli matrices. Writing them as vector $\vec{\sigma} \equiv (\sigma_x, \sigma_y, \sigma_z)$, we have

$$\mathbf{B} = \vec{B}(\vec{r}, t) \vec{\sigma}. \quad (125)$$

It has to be derived, how $\Psi(\vec{r}, t)$ will change in a rotation of the coordinate system. At each point (\vec{r}, t) , the change of the particle's energy density which is caused by the interaction with the magnetic field, should be invariant under the rotation:

$$\mu(\vec{r}, t) = \mu'(\vec{r}, t) \quad (126)$$

Making the assumption that

$$\Psi'(\vec{r}, t) \equiv \mathbf{Q} \Psi(\vec{r}, t), \quad (127)$$

we get with a general magnetic field in a rotated coordinate system the following expression for $\mu'(\vec{r}, t)$:

$$\mu'(\vec{r}, t) = \Psi^\dagger(\vec{r}, t) \mathbf{Q}^\dagger \mathbf{B}' \mathbf{Q} \Psi(\vec{r}, t). \quad (128)$$

This is the same as $\mu(\vec{r}, t)$, if

$$\mathbf{B} = \mathbf{Q}^\dagger \mathbf{B}' \mathbf{Q}. \quad (129)$$

Since \mathbf{Q} in eq. (121) is unitary, the following identity holds with eq. (129):

$$\mathbf{Q} \mathbf{Q}^\dagger \mathbf{B}' \mathbf{Q} \mathbf{Q}^\dagger = \mathbf{Q} \mathbf{B} \mathbf{Q}^\dagger = \mathbf{B}' \quad (130)$$

Eq. (130) exactly defines the transformation law of \mathbf{B} in eq. (123).

With eq. (128), eq. (129) and eq. (125), the overall energy density of the particle ensemble under a general magnetic field is

$$w(\vec{r}, t) = \Psi^\dagger(\vec{r}, t) \left(\hat{\mathbf{H}} - \mu_B \vec{B}(\vec{r}, t) \vec{\sigma} \right) \Psi(\vec{r}, t). \quad (131)$$

Again, the energy from the interaction of spin and magnetic field has the form of a potential. Hence, we can write the time dependent Pauli equation for an arbitrary magnetic field as:

$$i\hbar \frac{\partial \Psi(\vec{r}, t)}{\partial t} = \left(\hat{\mathbf{H}} - \mu_B \vec{B}(\vec{r}, t) \vec{\sigma} \right) \Psi(\vec{r}, t) \quad (132)$$

A common situation is, that the magnetic field $\vec{B}(\vec{r}, t)$ of a Stern-Gerlach apparatus lies in a coordinate system, which is rotated with respect to the orientation of an incoming spinor $\Psi'(\vec{r}, t)$. Especially in EPR experiments, one has to compute $\Psi(\vec{r}, t) = \mathbf{Q}^\dagger \Psi'(\vec{r}, t)$. To describe the direction of the spin rotation around the axis \vec{z}' , only the Euler angles θ and ϕ are necessary and one can select $\psi = -\pi/2$ without loss of generalisation. The projection of the rotated axis \vec{x}' in the \vec{z}/\vec{y} plane of the original, non-rotated coordinate system, encloses an angle φ with the \vec{x} axis. This angle is related to the Euler angle ϕ by $\phi \equiv \varphi + \pi/2$. Applying the adjoint of \mathbf{Q} in eq. (121) with those simplifications on a spinor $\psi'_\uparrow(\vec{r}, t) |+\rangle$ yields

$$\psi(\vec{r}, t) = \psi'_\uparrow(\vec{r}, t) \left(e^{-i\varphi/2} \cos(\theta/2) |+\rangle + e^{i\varphi/2} \sin(\theta/2) |-\rangle \right) \quad (133)$$

and for a $\psi'_\downarrow(\vec{r}, t) |-\rangle$ state, we get

$$\psi(\vec{r}, t) = \psi'_\downarrow(\vec{r}, t) \left(-e^{-i\varphi/2} \sin(\theta/2) |+\rangle + e^{i\varphi/2} \cos(\theta/2) |-\rangle \right). \quad (134)$$

6.4 The spin operator

Finally, we will derive the so called spin-operator. Both particle ensembles have an average spin angular momentum of $\pm\hbar/2$. Therefore, the overall expectation value of spin in the \vec{z} direction can be written as weighed sum

$$\langle S_{\vec{z}} \rangle \equiv \frac{\hbar}{2} \int \left(|a|^2 |\psi_\uparrow(\vec{r}, t)|^2 - |b|^2 |\psi_\downarrow(\vec{r}, t)|^2 \right) d^3r. \quad (135)$$

We can simplify this expression with a matrix

$$\mathbf{S}_{\vec{z}} \equiv \frac{\hbar}{2} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \quad (136)$$

and obtain

$$\langle S_{\vec{z}} \rangle = \int \psi^\dagger(\vec{r}, t) \mathbf{S}_{\vec{z}} \psi(\vec{r}, t) d^3r. \quad (137)$$

For the spin with respect to the direction of an unit vector $\vec{z}' = \vec{\mu}$ from an arbitrary rotated coordinate system, we have, analogous to eq. (125) the relation $\mathbf{S}_{\vec{z}'} = \frac{\hbar}{2} \vec{\mu} \vec{\sigma}$. It is convenient to write the axis $\vec{\mu}$ in

spherical coordinates as $\vec{\mu} \equiv (\cos(\varphi)\sin(\theta), \sin(\varphi)\sin(\theta), \cos(\theta))$ and we observe, that $\vec{\mu}$ depends on the same angles which are used for the spinors of eq. (133) and eq. (134).

Using this notation, a ‘‘spin operator’’ can be defined as

$$\hat{\vec{S}} \equiv \frac{\hbar}{2}\vec{\sigma} \quad (138)$$

and we get $\mathbf{S}_{\vec{z}'} = \vec{\mu}\hat{\vec{S}}$. Finally, the average value $\langle S_{\vec{z}'} \rangle$ becomes

$$\langle S_{\vec{z}'} \rangle = \int \psi^\dagger(\vec{r}, t) \mathbf{S}_{\vec{z}'} \psi(\vec{r}, t) d^3r = \vec{\mu} \langle \vec{S} \rangle, \quad (139)$$

where

$$\langle \vec{S} \rangle = \int \psi^\dagger(\vec{r}, t) \hat{\vec{S}} \psi(\vec{r}, t) d^3r. \quad (140)$$

7 Entangled systems with spin

7.1 Transformation of the entangled state into equations of motion

As in the case of the many particle Schrödinger equation, we can extend the coordinate space to derive the N particle Pauli equation. From now on, we will assume that there are no forces which interconnect the different particle systems and that their external forces coincide at each time.

We start our discussion with a state

$$\begin{aligned} \Psi(\vec{r}_1, \vec{r}_2, t) &\equiv \frac{1}{\sqrt{2}} (\psi(\vec{r}_1, t) |-\rangle \otimes \psi(\vec{r}_2, t) |+\rangle - \psi(\vec{r}_1, t) |+\rangle \otimes \psi(\vec{r}_2, t) |-\rangle) \\ &= \frac{1}{\sqrt{2}} (\chi_1(\vec{r}_1, \vec{r}_2, t) - \chi_2(\vec{r}_1, \vec{r}_2, t)). \end{aligned} \quad (141)$$

It shall describe two ensembles of entangled particles with the same rest mass m_0 and opposite spin. The one particle spin states in the first line of eq. (141) are all eigenstates with respect to a chosen axis \vec{z} and in the second line, we have defined the functions

$$\chi_1(\vec{r}_1, \vec{r}_2, t) \equiv \psi(\vec{r}_1, t) |-\rangle \otimes \psi(\vec{r}_2, t) |+\rangle \quad \text{and} \quad \chi_2(\vec{r}_1, \vec{r}_2, t) \equiv \psi(\vec{r}_1, t) |+\rangle \otimes \psi(\vec{r}_2, t) |-\rangle.$$

The state in eq. (141) is the solution of a two particle Pauli equation without external electromagnetic fields. This leads to a Hamiltonian operator

$$\hat{\mathbf{H}}(\vec{r}_1, \vec{r}_2, t) \equiv \left(\sum_{n=1}^2 \frac{-\hbar^2 \Delta_n^N}{2m_0} + \sum_{n=1}^2 \hat{V}_{\text{ext}}(\vec{r}_n, t) \right) \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad (142)$$

where the external potentials for the two particle systems agree at each time t :

$$\hat{V}_{\text{ext}}(\vec{r}_1, t) = \hat{V}_{\text{ext}}(\vec{r}_2, t) \quad (143)$$

Similar as in eq. (104), the Pauli equation for the state of eq. (141) can be written as:

$$\begin{aligned} i\hbar \frac{\partial}{\partial t} \Psi(\vec{r}_1, \vec{r}_2, t) &= \hat{\mathbf{H}}(\vec{r}_1, \vec{r}_2, t) \Psi(\vec{r}_1, \vec{r}_2, t) \\ &= \begin{cases} \frac{1}{\sqrt{2}} i\hbar \frac{\partial \chi_1(\vec{r}_1, \vec{r}_2)}{\partial t} = \frac{1}{\sqrt{2}} \hat{\mathbf{H}}(\vec{r}_1, \vec{r}_2, t) \chi_1(\vec{r}_1, \vec{r}_2) \\ - \\ \frac{1}{\sqrt{2}} i\hbar \frac{\partial \chi_2(\vec{r}_1, \vec{r}_2)}{\partial t} = \frac{1}{\sqrt{2}} \hat{\mathbf{H}}(\vec{r}_1, \vec{r}_2, t) \chi_2(\vec{r}_1, \vec{r}_2) \end{cases} \end{aligned} \quad (144)$$

For the particle ensembles described by eq. (144), the overall probability density is given by

$$\rho(\vec{r}_1, \vec{r}_2, t) \equiv \frac{1}{2} \rho_1(\vec{r}_1, \vec{r}_2, t) + \frac{1}{2} \rho_2(\vec{r}_1, \vec{r}_2, t), \quad (145)$$

with

$$\begin{aligned}
\rho_1(\vec{r}_1, \vec{r}_2, t) &\equiv \chi_1^+(\vec{r}_1, \vec{r}_2, t)\chi_1(\vec{r}_1, \vec{r}_2, t) \\
&= \psi^*(\vec{r}_1, t)\psi(\vec{r}_1, t)\langle - | - \rangle \otimes \psi^*(\vec{r}_2, t)\psi(\vec{r}_2, t)\langle + | + \rangle \\
&\equiv \rho_{1\downarrow}(\vec{r}_1, t) \otimes \rho_{1\uparrow}(\vec{r}_2, t)
\end{aligned} \tag{146}$$

and

$$\begin{aligned}
\rho_2(\vec{r}_1, \vec{r}_2, t) &\equiv \chi_2^+(\vec{r}_1, \vec{r}_2, t)\chi_2(\vec{r}_1, \vec{r}_2, t) \\
&= \psi^*(\vec{r}_1, t)\psi(\vec{r}_1, t)\langle + | + \rangle \otimes \psi^*(\vec{r}_2, t)\psi(\vec{r}_2, t)\langle - | - \rangle \\
&\equiv \rho_{2\uparrow}(\vec{r}_1, t) \otimes \rho_{2\downarrow}(\vec{r}_2, t).
\end{aligned} \tag{147}$$

Probability densities of many particle states are connected to the corresponding osmotic velocities via

$$\vec{u}^N(\vec{r}^N, t) \equiv -\frac{\hbar}{2m_0}\vec{\nabla}^N \ln(\rho(\vec{r}^N, t)/\rho_0). \tag{148}$$

Hence, after the tensor product in $\rho_1(\vec{r}_1, \vec{r}_2, t)$ is converted by the logarithm laws into a sum, we get for $\vec{u}_{\chi_1}^N(\vec{r}_1, \vec{r}_2, t)$:

$$\begin{aligned}
\vec{u}_{\chi_1}^N(\vec{r}_1, \vec{r}_2, t) &= -\frac{\hbar}{2m_0}\vec{\nabla}^N (\ln((\rho_{1\downarrow}(\vec{r}_1, t))_1/\rho_0) + \ln((\rho_{1\uparrow}(\vec{r}_2, t))_2/\rho_0)) \\
&= \begin{pmatrix} -\frac{\hbar}{2m_0}\vec{\nabla} \ln((\rho_{1\downarrow}(\vec{r}_1, t))_1/\rho_0) \\ -\frac{\hbar}{2m_0}\vec{\nabla} \ln((\rho_{1\uparrow}(\vec{r}_2, t))_2/\rho_0) \end{pmatrix} = \begin{pmatrix} \vec{u}_{\downarrow}(\vec{r}_1, t) \\ \vec{u}_{\uparrow}(\vec{r}_2, t) \end{pmatrix}.
\end{aligned} \tag{149}$$

Similarly, $\vec{u}_{\chi_2}^N$ in the Pauli equation for $\chi_2(\vec{r}_1, \vec{r}_2, t)$ is of the form $\vec{u}_{\chi_2}^N(\vec{r}^N, t) \equiv \begin{pmatrix} \vec{u}_{\uparrow}(\vec{r}_1, t) \\ \vec{u}_{\downarrow}(\vec{r}_2, t) \end{pmatrix}$.

The definition of the average velocity $\vec{v}^N(\vec{r}^N, t)$ for a many particle ensemble is

$$\vec{v}^N(\vec{r}^N, t) \equiv \frac{\hbar}{m_0}\vec{\nabla}^N \varphi(\vec{r}^N, t), \tag{150}$$

where $\varphi(\vec{r}^N, t)$ is the phase of the many particle wave-function

$$\Psi(\vec{r}^N, t) \equiv \pm \sqrt{\rho(\vec{r}^N)}e^{i\varphi(\vec{r}^N)}. \tag{151}$$

Writing $\chi_1(\vec{r}_1, \vec{r}_2, t)$ in the form of eq. (151)

$$\chi_1(\vec{r}_1, \vec{r}_2, t) = \sqrt{\rho_{\downarrow}(\vec{r}_1, t)}e^{i\varphi_{\downarrow}(\vec{r}_1)} \otimes \sqrt{\rho_{\uparrow}(\vec{r}_2, t)}e^{i\varphi_{\uparrow}(\vec{r}_2)}, \tag{152}$$

we can identify the overall phase of $\chi_1(\vec{r}_1, \vec{r}_2, t)$ as

$$\varphi_1(\vec{r}_1, \vec{r}_2, t) = \varphi_{\downarrow}(\vec{r}_1, t) + \varphi_{\uparrow}(\vec{r}_2, t). \tag{153}$$

This yields

$$\vec{v}_{\chi_1}^N(\vec{r}_1, \vec{r}_2, t) = \frac{\hbar}{m_0}\vec{\nabla}^N \varphi_1(\vec{r}_1, \vec{r}_2, t) = \begin{pmatrix} \vec{v}_{\downarrow}(\vec{r}_1, t) \\ \vec{v}_{\uparrow}(\vec{r}_2, t) \end{pmatrix} \tag{154}$$

and we get $\vec{v}_{\chi_2}^N(\vec{r}_1, \vec{r}_2, t) \equiv \begin{pmatrix} \vec{v}_{\uparrow}(\vec{r}_1, t) \\ \vec{v}_{\downarrow}(\vec{r}_2, t) \end{pmatrix}$ in completely the same way.

The expressions for $\vec{u}_{\chi_1}^N, \vec{u}_{\chi_2}^N, \vec{v}_{\chi_1}^N, \vec{v}_{\chi_2}^N$ imply, that the Pauli equation eq. (144) with the entangled state in eq. (141) can be decomposed into 2 two component differential equations, where $\vec{F}_2^{\text{ext}}(\vec{r}_2, t) = \vec{F}_1^{\text{ext}}(\vec{r}_1, t)$:

$$\left\{ \begin{array}{l} \left(\begin{array}{l} \frac{d}{dt}\vec{v}_{\downarrow}(\vec{r}_1, t) - \left(\vec{u}_{\downarrow}(\vec{r}_1, t) \cdot \vec{\nabla} \right) \vec{u}_{\downarrow}(\vec{r}_1, t) + \nu \Delta \vec{u}_{\downarrow}(\vec{r}_1, t) = \frac{1}{m_0}\vec{F}_1^{\text{ext}}(\vec{r}_1, t) \\ \frac{d}{dt}\vec{v}_{\uparrow}(\vec{r}_2, t) - \left(\vec{u}_{\uparrow}(\vec{r}_2, t) \cdot \vec{\nabla} \right) \vec{u}_{\uparrow}(\vec{r}_2, t) + \nu \Delta \vec{u}_{\uparrow}(\vec{r}_2, t) = \frac{1}{m_0}\vec{F}_2^{\text{ext}}(\vec{r}_2, t) \end{array} \right) \\ \left(\begin{array}{l} \frac{d}{dt}\vec{v}_{\uparrow}(\vec{r}_1, t) - \left(\vec{u}_{\uparrow}(\vec{r}_1, t) \cdot \vec{\nabla} \right) \vec{u}_{\uparrow}(\vec{r}_1, t) + \nu \Delta \vec{u}_{\uparrow}(\vec{r}_1, t) = \frac{1}{m_0}\vec{F}_1^{\text{ext}}(\vec{r}_1, t) \\ \frac{d}{dt}\vec{v}_{\downarrow}(\vec{r}_2, t) - \left(\vec{u}_{\downarrow}(\vec{r}_2, t) \cdot \vec{\nabla} \right) \vec{u}_{\downarrow}(\vec{r}_2, t) + \nu \Delta \vec{u}_{\downarrow}(\vec{r}_2, t) = \frac{1}{m_0}\vec{F}_2^{\text{ext}}(\vec{r}_2, t) \end{array} \right) \end{array} \right. \tag{155}$$

Eq. (155) is equivalent to eq. (144), but it contains the ensemble averages \bar{u} and \bar{v} of each particle system directly. All equations in eq. (155) can be similarly derived as the two equations in eq. (90), but the derivation must be done for the \vec{r}_1 and \vec{r}_2 particle ensembles separately.

7.2 How the correlations of entangled states emerge in the local hidden variable theory

Having converted the entangled state into equations of motion that can be derived from the previously discussed hidden variable model, we will investigate in detail how the correlations of entangled systems emerge from the viewpoint of these hidden variables.

The source in the EPR experiment produces particles of a defined spin direction with respect to the axis \vec{z} . However, it is not able to prepare the spin states in separate beams like a Stern-Gerlach magnet, because it does not divide the two spin ensembles spatially. Hence, the spin states of the particles emerging from the source are a superposition of an \uparrow and a \downarrow state. As described in section 7.2 the individual particles in such a system have to undergo an exchange procedure between the $\{A \uparrow\}$ and $\{B \downarrow\}$ or the $\{B \uparrow\}$ and $\{A \downarrow\}$ ensembles. The trajectories $\vec{x}_{j\uparrow(\downarrow)1}^{A(B)}(t)$ of the \vec{r}_1 particles have to switch between their $\vec{v}_{\downarrow}^A(\vec{r}_1, t)$ and $\vec{v}_{\uparrow}^B(\vec{r}_1, t)$ system as well as their $\vec{v}_{\uparrow}^A(\vec{r}_1, t)$ and $\vec{v}_{\downarrow}^B(\vec{r}_1, t)$ ensemble in eq. (155) during their flight. An analogous procedure has to happen with the trajectories $\vec{x}_{j\uparrow(\downarrow)2}^{A(B)}(t)$ of the \vec{r}_2 ensembles.

We observe, that it is possible, to treat the \vec{r}_1 and \vec{r}_2 states completely disconnected, because in eq. (155) they represent uncoupled statistical ensembles. They may even be spatially separated, in a way that there is no contact between them. A measurement on one particle ensemble, say \vec{r}_1 , might change the potential $\hat{V}_{\text{ext}}(\vec{r}_1, t)$ in eq. (142). Along with this goes a change of the external force $\vec{F}_1^{\text{ext}}(\vec{r}_1, t)$. If the action of the device was such, that $\vec{F}_1^{\text{ext}}(\vec{r}_1, t) \neq \vec{F}_2^{\text{ext}}(\vec{r}_2, t)$ at a certain time t , the equations governing the \vec{r}_1 system would be different from the equations for \vec{r}_2 . As a result, correlations in eq. (155) between both ensembles would be lost. Obviously, the dynamics of the \vec{r}_2 system is not be changed by something that happens with the \vec{r}_1 ensemble.

For the discussion of the EPR experiment, it is important to note, that there will still be correlations between the two particle systems, if the experimenters at both measurement stations 1 and 2 choose the settings of their devices in a way, that $\vec{F}_1^{\text{ext}}(\vec{r}_1, t) = \vec{F}_2^{\text{ext}}(\vec{r}_2, t)$ for all times t . This is exactly the situation, when the two separated Stern-Gerlach magnets are rotated in the same axis $\vec{\mu} = \vec{v}$ and the experiment is done with (infinitely) many particle pairs.

To connect the present discussion with the content of section 3, we recall the definition of the spin trajectories as stochastic processes ϕ_{j1t_z} and ϕ_{j2t_z} . We have $\phi_{j1(2)t_z} = \uparrow$, if the trajectory $\vec{x}_{j1(2)}^{A(B)}(t)$ at time t is in an \uparrow ensemble with respect to \vec{z} and $\phi_{j1(2)t_z} = \downarrow$, if the trajectory is in the \downarrow system at this time. One gets exact anti-correlations between the spins of the j -th particle pair during its flight, if one can ensure that

$$\phi_{j1t_z} \neq \phi_{j2t_z} \quad (156)$$

for all times t . This property gives rise to the state in eq. (141). It is a state, which describes the solution of a Pauli equation with two separate particle ensembles whose spin states are oscillating in an opposite way.

During the flight of the particles to the detectors, the trajectory exchange has to be done such, that if a trajectory changes its $\{A\}$ ($\{B\}$) ensemble, its spin state has to change to the opposite value at the same time. Similarly, if a particle changes its spin, its $\{A\}$ ($\{B\}$) ensemble must change also. Transitions $\{B \uparrow\} \rightarrow \{A \uparrow\}$, $\{B \downarrow\} \rightarrow \{A \downarrow\}$, $\{A \uparrow\} \rightarrow \{A \downarrow\}$ or $\{B \uparrow\} \rightarrow \{B \downarrow\}$ are not allowed. Otherwise, the trajectories affected would not be described any more by a superposition of an \uparrow and a \downarrow state, where energy conservation on the average holds.

The particles are interchanged from $\{A \uparrow\}$ ($\{A \downarrow\}$) to $\{B \downarrow\}$ ($\{B \uparrow\}$), when they are moving too fast in the $\{A\}$ system. Similarly, a change from $\{B \uparrow\}$ ($\{B \downarrow\}$) to $\{A \downarrow\}$ ($\{A \uparrow\}$) occurs, when their velocity is too slow in the $\{B\}$ system.

In each pair, the particles start having opposite spin and the same initial speed, due to momentum conservation. Therefore, at time $t = 0$, the trajectories from each pair will be placed in the same $\{A\}$ ($\{B\}$) ensemble, but in the opposite spin system.

With inter-particle forces being absent and equal external forces, any differences in the velocity of the two entangled particles can be caused by the random forces $\vec{F}_{j1(2)}^{\text{Brown}}$ only. Hence, we may demand that

$$\vec{F}_{j1}^{\text{Brown}}(t) = \vec{F}_{j2}^{\text{Brown}}(t) \quad (157)$$

for all j and each time t . Then, the velocities of the two entangled trajectories from every particle pair will coincide at all times, after they started with the same speed in the identical $\{A\}$ ($\{B\}$) ensemble. Accordingly, the particles will switch their $\{A\}$ ($\{B\}$) system in the same way during their flight. Each placement in another $\{A\}$ ($\{B\}$) ensemble changes the particle's spin state to the opposite. Since the spin ensemble of the particles is opposite from the beginning at $t = 0$, the spin trajectories of both particles will take opposite values $\phi_{j1t_z} \neq \phi_{j2t_z}$ for all times $t > 0$. To fulfil the requirement of eq. (157), a theory must not necessarily incorporate a signalling mechanism between (\vec{r}_1, t) and (\vec{r}_2, t) . This can be easily seen by an explicit construction of a local model, where eq. (157) holds.

The only constraint on the random forces was, that they have a Gaussian distribution. Now consider the following mechanism: A real random number generator sets a value $\lambda_j \in \mathbb{R}$ for each j -th particle pair at preparation stage. This value λ_j is similar to the hidden parameter in Bell's original work [4], but independent for each j -th particle pair generated by the source.

In our model, λ_j serves as the starting value for two pseudo random number generators of the same type. Pseudo random number generators are deterministic procedures which, if initialised with the same starting value λ_j , always produce the same series. After they were initialised at $t = 0$, we let the two generators produce Gaussian distributed and time dependent series $\vec{F}_{j1}^{\text{Brown}}(t, \lambda_j)$ and $\vec{F}_{j2}^{\text{Brown}}(t, \lambda_j)$. These series are the random forces of the j -th trajectories for the \vec{r}_1 and \vec{r}_2 particles. Because they only depend on their common starting value $\lambda_j \in \mathbb{R}$, eq. (157) holds at each time. There is no signalling mechanism between the two disconnected forces, since they simply depend on their common past.

At first sight, one may think, that our a procedure would set up a deterministic passively local theory. Indeed, for $\vec{F}_{j1}^{\text{Brown}}(t, \lambda_j)$ and $\vec{F}_{j2}^{\text{Brown}}(t, \lambda_j)$, there is a preparation event λ_j , which determines all later values of these forces. However, one does not have access to the Gaussian distributed random forces on the probability space, but only to the two spin values on the measurable space of the random variables.

The proof in section 4 was with regards the sigma algebras \mathcal{F}_{jS} , \mathcal{F}_{j1} and \mathcal{F}_{j2} . They contain those events, that determine the accessible spin results in the measurable spaces of the EPR experiment for sure. However, the set of theoretically possible events in the probability space for the Gaussian distributed force fields $\vec{F}_{j1(2)}^{\text{Brown}}(t, \lambda_j)$ is considerably larger than the set of the two spin values in the measurable space. Hence, every element in \mathcal{F}_{jS} , \mathcal{F}_{j1} and \mathcal{F}_{j2} must be a union event of many theoretically possible outcomes of the force fields, which all lead to the same spin result in the measurable space.

For an entangled particle that flies to a detector, the condition of deterministic passive locality would imply, that a union event of certain outcomes of the random force happens at the first time step. All events in this union must lead to the same spin state of the particle at that time. By deterministic passive locality, the spin result of a given particle at preparation stage has to determine the particle's spin value at the detector. Accordingly, the union event for the Brownian forces at the first time step has to determine another union of events at later times. The latter consists of outcomes for the random force at a time, where the particle has arrived in the Stern-Gerlach magnet. In particular, all events in this predetermined union must be such, that they lead to the same particular spin state. It can be seen from a simple example, that even if the theoretically possible events of the time dependent random forces are predetermined for each j -th particle pair by a hidden parameter λ_j , deterministic passive locality can be easily violated.

Let the spin trajectory $\phi_{j=1,1,t_z}$ of a particle, which is in a pair $j = 1$ and will fly to detector 1, start in the $\{A \uparrow\}$ ensemble. The velocity of this particle at time $t = 0$ can be denoted by $\left| \dot{\vec{x}}_{j=1,\uparrow 1}^A(t=0) \right|$ and the particle's spin trajectory at that time is $\phi_{j=1,1,t=0_z} = \uparrow$. The particle will have to be placed in the $\{B \downarrow\}$ ensemble, if its velocity $\left| \dot{\vec{x}}_{j=1,\uparrow 1}^A(t) \right|$ is higher than the arithmetic mean

$$v^{\text{AB}}(\vec{r}_1, t) \equiv \frac{|\vec{v}^A(\vec{r}_1, t)| + |\vec{v}^B(\vec{r}_1, t)|}{2} \quad (158)$$

of the ensemble average velocities from the $\{A \uparrow\}$ and the $\{B \downarrow\}$ system.

The velocity $\left| \dot{\vec{x}}_{j=1,\uparrow 1}^A(t) \right|$ is a function of $\vec{F}_{j=1,1}^{\text{Brown}}(t, \lambda_{j=1})$. For a given random starting value $\lambda_{j=1} \in \mathbb{R}$ of the pseudo random number generators, we let the Brownian force at time $t = 1$ take a value

$$\vec{F}_{j=1,1}^{\text{Brown}}(t = 1, \lambda_{j=1}) \equiv \xi_1. \quad (159)$$

It may be, that ξ_1 leads to a particle velocity of $\left| \dot{\vec{x}}_{j=1,\uparrow 1}^A(t = 1) \right| \leq v^{\text{AB}}(\vec{r}_1, t = 1)$. Then, no change of the spin trajectory takes place and we get:

$$\phi_{j=1,1,t=0_z} = \phi_{j=1,1,t=1_z}.$$

Similarly, at time $t = 2$, let the pseudo random number generator produce the value

$$\vec{F}_{j=1,1}^{\text{Brown}}(t_2, \lambda_{j=1}) \equiv \xi_2 \quad (160)$$

such, that $\left| \dot{\vec{x}}_{j=1,\uparrow 1}^{\text{A}}(t = 2) \right| \leq v^{\text{AB}}(\vec{r}_1, t = 2)$. Still, no change of $\phi_{j=1,1,t=2_z}$ happens and we can write

$$\phi_{j=1,1,t=0_z} = \phi_{j=1,1,t=1_z} = \phi_{j=1,1,t=2_z}. \quad (161)$$

There could be another pair $j = 2$, for which the particle 1 that goes through detector 1 starts in the $\{A \uparrow\}$ ensemble. The forces related to this particle may be driven by another random starting value $\lambda_{j=2} \neq \lambda_{j=1}$. Obviously, it is well possible, that we get for $\lambda_{j=2}$ at $t = 1$:

$$\vec{F}_{j=2,1}^{\text{Brown}}(t = 1, \lambda_{j=2}) \equiv \delta_1 \approx \xi_1, \quad (162)$$

with δ_1 such, that $\left| \dot{\vec{x}}_{j=2,\uparrow 1}^{\text{A}}(t = 1) \right| \leq v^{\text{AB}}(\vec{r}_1, t = 1)$, and thereby

$$\phi_{j=2,1,t=0_z} = \phi_{j=2,1,t=1_z} \quad (163)$$

again. Yet, for $t = 2$, we may get with $\lambda_{j=2}$

$$\vec{F}_{j=2,1}^{\text{Brown}}(t = 2, \lambda_{j=2}) \equiv \delta_2 \gg \xi_2, \quad (164)$$

where δ_2 causes a significant change in the velocity, leading to $\left| \dot{\vec{x}}_{j=2,\uparrow 1}^{\text{A}}(t = 2) \right| > v^{\text{AB}}(\vec{r}_1, t = 2)$. As a consequence, the particle in the $\{A \uparrow\}$ ensemble would be placed in the $\{B \downarrow\}$ ensemble at this time, taking the velocity $\left| \dot{\vec{x}}_{j=2,\downarrow 1}^{\text{B}}(t = 2) \right|$. If so, the spin trajectory would become:

$$\phi_{j=2,1,t=0_z} = \phi_{j=2,1,t=1_z} \neq \phi_{j=2,1,t=2_z}. \quad (165)$$

When the particles arrive in the Stern-Gerlach magnets of the EPR experiment, their spin states transform with axis dependent probabilities to an up or to a down state once again, due to the application of \mathbf{Q} from eq. (121). By eq. (133) and eq. (134), the axes of the magnets can be selected in a way, that for a given outcome, these probabilities are different from unity. In this case, the decision, to what spin ensemble an individual trajectory $\vec{x}_{j1(2)}^{\text{A(B)}}(t)$ will change, is random. Since the only probabilistic elements in our theory are the Brownian forces, they must be solely responsible for that random decision. The requirement of eq. (157) is, that for two entangled trajectories $\vec{x}_{j1}^{\text{A(B)}}(t)$ and $\vec{x}_{j2}^{\text{A(B)}}(t)$, the time dependent Brownian forces coincide at each time t . If the two detectors in the EPR experiment have identical settings $\vec{\mu} = \vec{\nu}$, the external forces in eq. (155) also agree at every time: $\vec{F}_1^{\text{ext}}(\vec{r}_1, t) = \vec{F}_2^{\text{ext}}(\vec{r}_2, t)$

In other words: If the settings of the detectors in an EPR experiment coincide, the entangled trajectories $\vec{x}_{j1}^{\text{A(B)}}(t)$ and $\vec{x}_{j2}^{\text{A(B)}}(t)$ are under equal forces at each time. Accordingly, as described above, the two trajectories would change their positions, velocities, as well as their $\{A\}$ and $\{B\}$ ensemble simultaneously in the same, but their spin ensemble in an exactly opposite way. It is this mechanism, which gives rise to the observed correlations at the detectors.

That our theory reproduces the results of quantum mechanics by the violation of passive locality can be easily seen, if we select an angle $\theta = 2\pi$ at the measurement devices in eq. (133) and eq. (134). For the up state in eq. (161) at $t = 2$, one would then get a spin up result in detector 1 for sure. Similarly, for spin state in eq. (165) at $t = 2$, a down state would be measured at station 1 for sure. These are opposite measurement results for two particles $j = 1$ and $j = 2$, that started both with spin up at the source and went to detector 1. Hence, there is no event at preparation stage, which determines a specific results at the detector. This corresponds to the failure of deterministic passive locality.

We have to remark, that even if the model with the pseudo random number generators may look terribly artificial, it would nevertheless be difficult to test. Especially, because the initial value λ_j is assumed to be a real random number, the statistics of $\{\sigma_{1\vec{\mu}}\}$ and $\{\sigma_{2\vec{\nu}}\}$ would equal the properties of real random numbers, if infinitely many particle pairs j are considered.

An objection against such a model could seem to be experiments with entanglement swapping [36]. In those setups, the pairs of entangled particles are generated by two spatially separated sources. To describe these experiments with the model above, two changes have to be made:

The area of the particle source in section 3.1 has to be replaced by a slice in which the two particle sources are. The slice must be disjoint from the future cones of the two Stern-Gerlach magnets and all proofs in section 3 and 4 work as before. In section 7.2, eq. (157) has to hold for the Brownian forces at the two separated particle sources. Nevertheless, it is possible to find a spatial region, where the past cones of the two sources intersect. At this place, a common cause $\lambda_j \in \mathbb{R}$ could well be established, leading to eq. (157). Hence, even with entangled particles which are produced by two independent sources, there is no need for hidden variable theories to include an instantaneous connection between spatially separated regions.

It is well known, that entanglement phenomena are not restricted to spin states. For example, according to quantum mechanics, exact correlations between positions of entangled particles should be observed, when they are emitted by the source at almost the same time. A first outline of an experiment to investigate this behaviour was presented in [37] and the analogue of this setup for photons was realised in [38]. Lastly in [39], a concrete proposal for an experiment with material particles has been made, which still waits for its realisation. Here, we only mention, that the correlations predicted by quantum mechanics for these situations are naturally explained by our local model. Obviously, if two particles are ejected within an interval smaller than the course grained timescale $\tau = \frac{\hbar}{2k_B T}$ of the Brownian motion process, the discussion above leads to exactly correlated behaviour of the two particle's trajectories in position and time.

To violate Bell's inequalities in the EPR experiment with a local model, it does suffice to assume that the Gaussian distributed random forces $\bar{F}_{j1}^{\text{Brown}}(t), \bar{F}_{j2}^{\text{Brown}}(t)$ have a kind of "memory" of their past. Nevertheless, the author has to admit the somewhat incomplete nature of this article because the model developed above is artificial. To give a more physical reason for eq. (157), one would need further studies. In fact, passive locality may be violated with many other hidden variable theories, all sharing the property, that the events at one measurement apparatus do not influence the events at another spatially separated one. Besides from Edward Nelson, who gave two examples of such theories, a different model with similar locality features was published recently in [40]. The authors of this article do not study Bell's theorem in the language of advanced mathematical probability, as it was done by Nelson, but they seem to share his opinion about the meaning of passive locality.

Here, we wish to leave it at that, with a citation by Edward Nelson from a popular talk. It shows, that behaviour which is similar to entanglement is in fact commonly observed in stochastically evolving systems and not at all confined to quantum mechanical objects:

"The snowflake problem is this: The inexhaustible variety of snowflakes makes it evident that chance plays a major role in their development, yet they always preserve hexagonal symmetry – how does a portion of the snowflake growing at random on one side know to grow in precisely the same fashion as its partner all the way over on the other side? This is mysterious but not beyond understanding" [41]. "Similar phenomena are well known to occur in the study of random fields. [...] Perhaps the results of correlated spin experiments are ultimately no more, and no less, mysterious than is the random growth of snowflakes with hexagonal symmetry" [9].

8 Objections against Nelson's theorem

8.1 The argument of Ballentine and Zeilinger

The basic argument in Nelson's (and Jarret's) reasoning is, that two disconnected random variables $\phi_{j1\bar{\mu}}$ and $\phi_{j2\bar{\nu}}$ may generate events $\{\sigma_{j1\bar{\mu}}\}$ and $\{\sigma_{j2\bar{\nu}}\}$ for which passive locality fails, or

$$P_{\phi_{j1\bar{\mu}} \otimes \phi_{j2\bar{\nu}}} \left\{ \sigma_{j1\bar{\mu}} \cap \sigma_{j2\bar{\nu}} \mid \mathcal{F}_{jS} \right\} \neq P_{\phi_{j1\bar{\mu}} \otimes \phi_{j2\bar{\nu}}} \left\{ \sigma_{j1\bar{\mu}} \mid \mathcal{F}_{jS} \right\} P_{\phi_{j1\bar{\mu}} \otimes \phi_{j2\bar{\nu}}} \left\{ \sigma_{j2\bar{\nu}} \mid \mathcal{F}_{jS} \right\}. \quad (166)$$

Moreover, Nelson and Jarret explicitly wrote, that this can happen without any a signalling mechanism in the theory which produces the events in question. The latter statement, although mathematically correct, has raised criticisms from physicists since it was made. For example, in his famous introductory textbook, Ballentine [42] writes:

„There have been some debates as to whether the factorisation of the probability is justified by the locality condition alone, or whether it requires some additional stronger assumptions. If so, the contradiction may be blamed on those additional assumptions. The issue is subtle, but fortunately it is now irrelevant, since the new proof in [43] does not make use of probability”.

That the work in [43] does not make use of probability is true, however, it only does account for deterministic hidden variable theories of a sort, where the events at the particle source are equivalent to the ones at the detectors. The authors are aware of this, noting at page 1138 in their article:

„The salvaging strategy could consist in [...] a stochastic local theory in contrast to a deterministic local theory which has been considered so far in this paper”.

They then go on, writing that these so called “stochastic local theories“ would be impossible. They claim, this would have been thoroughly investigated in [44, 45]. Furthermore, they mention the existence of so called „equivalence theorems” [46, 47]. Those are said to show, that each stochastic local hidden variable theory could be duplicated by a deterministic passively local one. But in all the articles they cite, every author only considers theories, where passive locality is assumed to hold ([44] makes this assumption in his eq. (5), [45] make it at pp. 528 in their eqs. (2), (2'), [46] at page 776 in eq. (7a), and [47] at pp. 1306 and 1308). In fact, it is only for this case, where an equivalence to deterministic passive locality can be proven. We conclude, that the proof in [43] does not call actively local hidden variable theories into question, where only passive locality fails.

8.2 The argument of Dürr

Furthermore, there exist claims, which try to derive Bell’s theorem in its original form of eq. (53), but without the assumption of passive locality. These „derivations” (e.g. [48] at page 182) do not write the inequality with a whole family of axis dependent probability measures $P_{\phi_{j1\vec{\mu}} \otimes \phi_{j2\vec{\nu}}}$ for device dependent events $\{\sigma_{j1\vec{\mu}}\} \in \mathcal{F}_{j1}$ and $\{\sigma_{j2\vec{\nu}}\} \in \mathcal{F}_{j2}$. Instead, they are using one single and detector independent measure $P_{j\vec{z}}$ for events, which are thought to be produced by the source and as, nevertheless, dependent on the axes of the detectors.

This is problematic, because if we compute the correlation coefficient

$$E(\vec{\mu}, \vec{\nu}) \equiv \langle \Psi | (\vec{\mu}\vec{\sigma})_1 \otimes (\vec{\nu}\vec{\sigma})_2 | \Psi \rangle \quad (167)$$

with the state of eq. (141), we are definitely led to eq. (39), and to a family of axis dependent measures $P_{\phi_{j1\vec{\mu}} \otimes \phi_{j2\vec{\nu}}}$.

By using one probability measure for axis dependent events generated by the particle source, we would only be able to investigate theories, where an intrinsic particle property is defined at preparation stage, that determines the results at the two detectors for each conceivable pair of axes.

Yet it may be possible, that the source generates particles whose properties are changed later on by the Stern-Gerlach magnets. In this case, a hidden variable model of the EPR experiment has to use an enlarged probability space. The events at preparation stage take place in a smaller space and their probabilities do not depend on the detector’s axes, whereas the events at the detectors happen in the enlarged probability space. A natural assumption is, that the events at the two separated measurement stations are generated by two families of completely uncoupled but device dependent random variables. Accordingly, we arrive at a family of axis dependent probability measures $P_{\phi_{j1\vec{\mu}} \otimes \phi_{j2\vec{\nu}}}$ for the joint probabilities of events at the two Stern-Gerlach magnets. Because the probability measures from the formalism of quantum mechanics are of the same structure, we believe that Nelson’s mathematical analysis of the EPR experiment is the correct one. At this point, we want to remark, that the impossibility to write the EPR experiment on one single probability space and the need for the inclusion of the detector settings in the corresponding random variables, was also mentioned recently by Khrennikov in [49].

Nevertheless, it is possible to confine oneself to a single and axis independent measure $P_{j\vec{z}}$ on one probability space, if passive locality is explicitly stated to hold. With that proposition and the proof in section 4, one can trace back the experimentally relevant axis dependent events $\{\sigma_{j1\vec{\mu}}\} \in \mathcal{F}_{j1}$ and $\{\sigma_{j2\vec{\nu}}\} \in \mathcal{F}_{j2}$ at the detectors to equivalent events $\{x_{j1\vec{z}}\} \in \mathcal{F}_{jS}$ at the common preparation stage. Due to active locality, all events in \mathcal{F}_{jS} cannot depend on the settings at the detectors. Therefore,

$$P_{\phi_{j1\vec{\mu}} \otimes \phi_{j2\vec{\nu}}} \{x_{j1\vec{z}}\} = P_{\phi_{j1\vec{\mu}'} \otimes \phi_{j2\vec{\nu}'}} \{x_{j1\vec{z}}\} \quad (168)$$

is fulfilled with $\vec{\mu}'$ and $\vec{\nu}'$ as arbitrary axes and we are able to write an axis independent measure $P_{j\vec{z}}$ for the events $\{x_{j1\vec{z}}\}$. We conclude, that in a theory, where passive and active locality holds, one can work with a single measure $P_{j\vec{z}}$ for detector independent events. A proof of Bell’s theorem that goes exactly this way can be found in [12].




Figure 5: Velocity distribution of the particles in the $\{A\}$ ensemble. Only those particles within a very small velocity interval of width $|\vec{\delta}_{j1}(t)|$ around $v^{AB}(\vec{r}, t)$ will be interchanged because of the disturbance.

8.3 The argument of Redhead, stability against disturbances

A different argument against stochastic hidden variable theories was raised by Redhead in [52] (who presents in his book at pp. 98 a proof of Bell’s theorem which is very similar to that of Nelson, although with some differences). He argues, that stochastic hidden parameter theories would be impossible and advances this opinion with the following observation: Redhead asserts, that under natural conditions, one can never exclude small disturbances, which might happen erroneously to a single particle (maybe through a random disturbance by the gases of the air in the laboratory). Redhead then defines, what he calls “the most general perturbation” that “is not a phase shift” as

$$u(\vec{n}, \vartheta) \equiv e^{(\vec{\sigma}\vec{n})_2 \vartheta/2}, \quad (169)$$

where \vec{n} is an arbitrary unit vector and ϑ is an angle about $0 \leq \vartheta < 4\pi$. This perturbation should be confined to one part of the overall Hilbert space, e.g. \mathcal{H}_2 only. Then, the entangled state in eq. (141) will transform to

$$\Psi'(\vec{r}_1, \vec{r}_2, t) \rightarrow \frac{1}{\sqrt{2}} (\psi(\vec{r}_1, t) |-\rangle \otimes u(\vec{n}, \vartheta) \psi(\vec{r}_2, t) |+\rangle - \psi(\vec{r}_1, t) |+\rangle \otimes u(\vec{n}, \vartheta) \psi(\vec{r}_2, t) |-\rangle). \quad (170)$$

Redhead concludes, that with $\Psi'(\vec{r}_1, \vec{r}_2, t)$, the exact correlations observed in EPR experiments will be lost for certain axes of the detectors. Nevertheless, entanglement can be experimentally investigated through long distances over 100 kilometres in the air [50], where small perturbations acting on single particles cannot be avoided.

Yet, in spirit of Ballentine’s ensemble interpretation [51], a quantum mechanical state may not describe individual particles, but rather an infinitely large statistical ensemble of trajectories. In this case, a perturbation like eq. (169) would correspond to a disturbance of an entire statistical ensemble of infinitely many particles for a sufficiently long time. Moreover, if defined as above, $u(\vec{n}, \vartheta)$ acts on every particle of that system in exactly the same way. This has nothing to do with a small erroneous disturbance that may happen accidentally to individual particles in the laboratory.

In this context, we want to show, that our theory is in some sense especially robust against small disturbances, at least if they act on individual particles. For a long distance entanglement experiment, it is a reasonable requirement, to expect large uncorrelated external forces $\vec{F}_2^{\text{ext}}(\vec{r}_2, t)$ and $\vec{F}_1^{\text{ext}}(\vec{r}_1, t)$, which act on the particles during their flight, to be absent. It is well known, that without external forces, the Langevin equations (66) lead to a Maxwell distribution of the particle’s velocity. Accordingly, most \vec{r}_1 particles in the $\{A\}$ or $\{B\}$ ensembles have a velocity around one centre $\approx |\vec{v}^A(\vec{r}_1, t)|$ or $\approx |\vec{v}^B(\vec{r}_1, t)|$.

Without loss of generalisation, we assume that a small disturbance $\vec{\delta}_{j1}(t)$ induces a slightly different velocity of the j -th \vec{r}_1 particle in the $\{A \uparrow\}$ ensemble at a point (\vec{r}_1, t) in space-time:

$$\dot{\vec{x}}_{j\uparrow 1}^A(t) \equiv \dot{\vec{x}}_{j\uparrow 1}^A(t) + \vec{\delta}_{j1}(t) \quad (171)$$

We define $\vec{\delta}_{j1}(t)$ to be small, if the half-width of the particle’s velocity distribution is far greater than $|\vec{\delta}_{j1}(t)|$. Moreover, the disturbance should be small enough, that the perturbation can’t change the statistics

of the entire ensemble through the trajectory exchange mechanism. The latter requirement is equivalent to the condition

$$|v^{AB}(\vec{r}_1, t) - |\vec{v}^A(\vec{r}_1, t)|| \gg |\vec{\delta}_{j1}(t)|. \quad (172)$$

Additionally, we expect the components of $\vec{\delta}_{j1}(t)$ to be fluctuating with zero mean.

A particle is replaced from the $\{A \uparrow\}$ to the $\{B \downarrow\}$ system, if its velocity $|\dot{\vec{x}}_{j\uparrow 1}^A(t)|$ is higher than $v^{AB}(\vec{r}_1, t)$. Since the average of each component in $\vec{\delta}_{j1}(t)$ is assumed to be zero, the disturbance does not contribute to $v^{AB}(\vec{r}_1, t)$. Hence,

$$|v^{AB}(\vec{r}_1, t) - |\dot{\vec{x}}_{j\uparrow 1}^A(t)|| < |\vec{\delta}_{j1}(t)| \quad (173)$$

has to hold, if the perturbation $\vec{\delta}_{j1}(t)$ is the only reason for $\dot{\vec{x}}_{j\uparrow 1}^A(t)$ to be placed in an ensemble. The particles fulfilling eq. (173) are in a small velocity interval of width $|\vec{\delta}_{j1}(t)|$ around $v^{AB}(\vec{r}_1, t)$ (see fig. 5).

Since $|\vec{\delta}_{j1}(t)|$ is assumed to be much smaller than the half-width of the particle's velocity distribution, the number of trajectories for which eq. (173) holds, will be low. Because of

$$|v^B(\vec{r}_1, t)| < v^{AB}(\vec{r}_1, t) < |v^A(\vec{r}_1, t)|, \quad (174)$$

eq. (172) and eq. (173), the interval in which the disturbance is able to change the ensemble of a particle, is far away from the centre $|\vec{v}^A(\vec{r}_1, t)|$, where most trajectories are found. This further reduces the number of particles that can be affected by $\vec{\delta}_{j1}(t)$.

We conclude, that only a very small number of trajectories changes its (spin-) ensemble solely because of $\vec{\delta}_{j1}(t)$. Hence, the disturbance merely results in a reduced efficiency of the entanglement, but not to its breakdown. For this reason, the theory we have constructed is especially robust against small disturbances unavoidable in a laboratory, in contrast to the claims by [52].

9 Conclusions and outlook

9.1 What has been achieved

The most popular argument against hidden variable theories with particle trajectories is, as commonly believed from Bell's words: That [those theories] "require a mechanism, whereby the setting of one measuring device can influence the reading of another device" and "moreover, the signal involved must propagate instantaneously". As shown by Edward Nelson whose analysis we reviewed here, this assertion is physically unfounded. Specially designed stochastic processes can violate Bell's inequalities at two separated locations without any contact at all. We have mathematically analysed, what such a behaviour really does imply for a stochastic theory and we have constructed a corresponding theory by ourselves.

9.2 Possibility of relativistic extension

Unfortunately, our theory has, at least till now, no relativistic extension. Stochastic mechanics of the original form in which Nelson wrote it down, had severe conceptual difficulties in the relativistic domain. These problems arose due to the Markov property of the stochastic process one had to use. In fact it was shown, that the definition of a Markov process on Minkowski space would lead either to determinism or the violation of causality [53–55]. Nevertheless, with some different assumptions for the process involved, the Klein-Gordon equation in four dimensions [56–61] as well as the Dirac equation in 1+1 dimensions [62–64] could be derived in various works.

We emphasise, that unlike Nelson's original formulation, the theory of Fritsche and Haugk describes a non-markovian stochastic process and thereby it avoids the no-go theorems of [53–55]. However, to rewrite the Fritsche-Haugk model in a manifestly covariant form, one would need a fully formulated theory of special relativistic Brownian motion (at least if one would try to derive the theory from Langevin equations). Regrettably, although such a theory certainly must exist (since non-relativistic Brownian motion does), it seems, that besides from some recent attempts [65–67], no one was able to write it down completely.

At this point, we want to mention, that the authors of [19] show at pp. 386, their eqs. (69) can, after an approximation, directly be casted into the Navier-Stokes equations. We have in eq. (69) for the $\{A\}$ system:

$$\frac{\partial}{\partial t} (\vec{v}^A - \vec{u}^A) + (\vec{v}^A + \vec{u}^A) \cdot \vec{\nabla} (\vec{v}^A - \vec{u}^A) - \nu \Delta (\vec{v}^A - \vec{u}^A) = \frac{1}{m_0} \vec{F}^{\text{ext}} \quad (175)$$

Noting that $\vec{v}_c^A = \vec{v}^A - \vec{u}^A$, we are able to write:

$$\frac{\partial}{\partial t} \vec{v}_c^A + (\vec{v}_c^A + 2\vec{u}^A) \cdot \vec{\nabla} \vec{v}_c^A - \nu \Delta \vec{v}_c^A = \frac{1}{m_0} \vec{F}^{\text{ext}} \quad (176)$$

If we set $\vec{u}^A = \vec{\nabla} p(\vec{r}, t) = 0$ and $\rho \vec{F}^{\text{ext}} = -\vec{\nabla} p(\vec{r}, t)$, where $p(\vec{r}, t)$ is the pressure and $\rho \equiv m_0 \rho(\vec{r}, t)$ defines the mass density, we get the Navier-Stokes equation of a fluid:

$$\rho \frac{\partial}{\partial t} \vec{v}_c^A + \rho \left(\vec{v}_c^A \cdot \vec{\nabla} \right) \vec{v}_c^A - \nu \rho \Delta \vec{v}_c^A + \vec{\nabla} p(\vec{r}, t) = 0 \quad (177)$$

For eq. (177), various manifestly covariant formulations have been proposed [68–71]. To derive the relativistic version of eq. (177) from first principles, one typically starts with relativistic Boltzmann equations [72–74]. Because of the close similarity between eq. (177) and eq. (69), it seems at least plausible, that it might be possible to formulate the theory of Fritsche and Haugk in a relativistic setting.

9.3 Problems of standard quantum mechanics in curved space-times, which might be absent in the model of Fritsche and Haugk

It is known, that in curved space-times there exists an ambiguity, if one wants to derive quantum mechanical operators. Usually, we are advised by the correspondence principle to make the replacement $p_i \rightarrow \hat{p}_i \equiv -i\hbar \frac{\partial}{\partial q_i}$ for the momentum operator, with q_i as the generalised coordinate of the particle. In Cartesian coordinates, the Hamilton function of a free particle is $H_{\text{free}} \equiv \frac{p_x^2 + p_y^2 + p_z^2}{2m_0}$. Using the replacement rule of the correspondence principle, we can get the free quantum mechanical Hamilton operator:

$$\hat{H}_{\text{free}} = -\frac{\hbar}{2m_0} \left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2} \right) = -\frac{\hbar}{2m_0} \Delta \quad (178)$$

We may convert its Laplacian in cylindrical coordinates ρ, φ, z and derive the free Hamiltonian in the new coordinate system:

$$\hat{H}_{\text{free}} = -\frac{\hbar}{2m_0} \left(\frac{\partial^2}{\partial \rho^2} + \frac{\partial}{\rho \partial \rho} + \frac{\partial^2}{\rho^2 \partial \varphi^2} + \frac{\partial^2}{\partial z^2} \right). \quad (179)$$

Now, we want to derive this operator in cylindrical coordinates from the beginning. We start with the classical Lagrange function for a free particle:

$$\mathcal{L}_{\text{free}}(q_i, \dot{q}_i) \equiv \frac{m_0}{2} (\dot{\rho}^2 + \rho^2 \dot{\varphi}^2 + \dot{z}^2) \quad (180)$$

If we insert the particle's momentum $p_\rho = \frac{\partial \mathcal{L}_{\text{free}}}{\partial \dot{\rho}} = m_0 \dot{\rho}$, $p_\varphi = \frac{\partial \mathcal{L}_{\text{free}}}{\partial \dot{\varphi}} = m_0 \rho^2 \dot{\varphi}$, $p_z = \frac{\partial \mathcal{L}_{\text{free}}}{\partial \dot{z}} = m_0 \dot{z}$, we get the following expression for $H_{\text{free}}(p_i, q_i, \dot{q}_i) \equiv \sum p_i \dot{q}_i - \mathcal{L}_{\text{free}}$:

$$H_{\text{free}}(p_i, q_i, \dot{q}_i) = p_\rho \dot{\rho} + p_\varphi \dot{\varphi} + p_z \dot{z} - \mathcal{L}_{\text{free}} = \frac{1}{2m_0} \left(p_\rho^2 + \frac{p_\varphi^2}{\rho^2} + p_z^2 \right) \quad (181)$$

After using the replacement rules defined by correspondence principle, we arrive at:

$$\hat{H}_{\text{free}} = -\frac{\hbar}{2m_0} \left(\frac{\partial^2}{\partial \rho^2} + \frac{\partial}{\rho \partial \rho} + \frac{\partial^2}{\rho^2 \partial \varphi^2} + \frac{\partial^2}{\partial z^2} \right) \quad (182)$$

Unfortunately, eq. (182) is different from eq. (179) and wrong. A similar problem is present in the path integral formalism in similar form. (Although a technique, which at least seems to overcome some of

those problems with path integrals was recently developed: [75]) Generally speaking, the above mentioned paradox implies, that the naive rules of the correspondence principle do only work, if applied on Cartesian coordinates. In the curved space-times of general relativity, there is no such coordinate system. Thereby, one cannot give, for example, an unambiguous expression for the probability density of an electron near the singularity of a black hole.

Unlike in ordinary quantum mechanics, in stochastic mechanics the quantum mechanical operators are not simply defined as replacement rules. Instead, their expressions are derived completely from classical equations (see [19] at pp. 290). It might well be, that it is possible to extend the theory of Fritsche and Haugk to special and general relativity via a derivation of its classical diffusion equations (69) in covariant form from relativistic Boltzmann equations. Then, as in the Cartesian case, one might get information about the form of the quantum mechanical operators, but for curved space-times. As a result, we could be able to understand the behaviour of quantum objects near space-time singularities.

Acknowledgements: The author wants to thank prof. Edward Nelson from Princeton University, prof. Lothar Fritsche from Technical University Clausthal and phd. Klaus Hornberger from Ludwig-Maximilians University Munich for helpful discussions and suggestions.

References

- [1] A. Einstein, B. Podolski, N. Rosen, *Phys. Rev.* 47, pp. 777 (1935)
- [2] D. Bohm, *Quantum Theory* (Prentice-Hall, Englewood Cliffs, NJ 1951), pp. 614-619
- [3] D. Bohm, Y. Aharonov, *Phys. Rev.* 108, pp. 1070 (1957)
- [4] J. S. Bell, On The Einstein Podolsky Rosen Paradox, in [5]
- [5] J. S. Bell, *Speakable and unspeakable in Quantum mechanics* (Cambridge University Press 1987)
- [6] J. F. Clauser, R. A. Holt, M. A. Horne, A. Shimony, *Phys. Rev. Lett.* 23, pp. 880 (1969)
- [7] J. S. Bell, The theory of local be-ables, in [5]
- [8] E. Nelson, Field theory and the future of stochastic mechanics, in *Stochastic Processes in Classical and Quantum Systems*, Proc. Ascona, Switzerland 1985, eds: S. Albeverio, G. Casati, D. Merlini, *Lecture Notes in Physics* 262 (Springer, Berlin 1988), pp. 438
- [9] E. Nelson, The locality problem in Stochastic Mechanics, in *New techniques and ideas in quantum measurement theory*, eds: D. Greenberger, *Ann. N. Y. Acad. Sci.* 480, pp. 533 (1986)
- [10] E. Nelson, Stochastic mechanics and random fields, in *Ecole d'Ete de Probabilites de Saint Flour XV-XVII*, *Lecture Notes in Mathematics* 1362 (Springer, Berlin 1985-87), pp. 427
- [11] N. D. Mermin, *Am. J. Phys.* 49, pp. 940 (1981)
- [12] W. G. Faris, Probability in quantum mechanics, Appendix to D. Wick, *The Infamous Boundary: Seven Decades of Controversy in Quantum Physics* (Birkhauser, Boston 1995)
- [13] J. P. Jarret, *Noûs* 18, pp. 569 (1984)
- [14] P. Suppes, M. Zanotti, On the Determinism of Hidden Variable Theories with Strict Correlation and Conditional Statistical Independence of Observables, In P. Suppes (Eds.), *Logic and probability in quantum mechanics* (Dordrecht: Reidel 1976), pp. 445-455
- [15] J. Conway, S. Kochen, *Found. Phys* 36, pp. 1441 (2006), arXiv:quant-ph/0604079v1
- [16] S. Kochen, E. Specker, *J. of Mathematics. and Mechanics.* 17, pp. 59 (1967)
- [17] E. Nelson, *Phys. Rev.* 150, pp. 1079 (1966)
- [18] E. Nelson, *Quantum Fluctuations* (Princeton University Press, Princeton NJ 1985)

- [19] L. Fritsche, M. Haugk, *Ann. Phys. (Leipzig)* 12, pp. 371 (2003)
- [20] A. Aspect, J. Dalibard, G. Roger, *Phys. Rev. Lett.* vol. 49, pp. 1804 (1982)
- [21] H. Rauch, *Neutron Interferometry* (Oxford University Press, Oxford 2000)
- [22] H. Rauch, *Reality in Neutron Interference Experiments*, *Found. Phys.* 23, pp. 7 (1993)
- [23] A. Go for the Belle Collaboration, *J. Mod. Opt.* 51, pp. 991 (2004), arXiv:quant-ph/0310192
- [24] H. Sakai, T. Saito, T. Ikeda, K. Itoh, T. Kawabata, H. Kuboki, Y. Maeda, N. Matsui, C. Rangacharyulu, M. Sasano, Y. Satou, K. Sekiguchi, K. Suda, A. Tamii, T. Uesaka, and K. Yako, *Phys. Rev. Lett.* 97, 150405 (2006)
- [25] B. G. Englert, M. O. Scully, G. Süssmann, H. Walther, *Z. Naturforsch* 47a, pp. 1175 (1992)
- [26] E. Nelson, The mystery of stochastic mechanics, conference talk, available at <http://www.math.princeton.edu/~nelson/papers/talk.pdf>
- [27] T. Dankel, *Arch. Ration. Mech. Anal.* 37, pp. 192 (1971)
- [28] W. G. Faris, A stochastic picture of spin, *Stochastic Processes in Quantum Theory and Statistical Physics*, eds. S. Albeverio, Ph. Combe, M. Sirugue-Collin (Springer, Berlin 1982), pp. 154
- [29] W. G. Faris, Spin correlations in stochastic mechanics, *Found. Phys.* 12, pp. 1 (1982)
- [30] L. Fritsche, M. Haugk, *Anschauliche Quantenmechanik*, unpublished manuscript
- [31] J. M. Levy-Leblond, *Comm. Math. Phys.* 6, pp. 286 (1967)
- [32] O. Stern, W. Gerlach, *Z. Phys.* 9, pp. 349 (1922)
- [33] T. E. Phipps, O. Stern, *Z. Phys.* 73, pp. 185 (1932)
- [34] R. Penrose, *The Road To Reality, A complete guide to the laws of the universe* (Vintage Books, London, 2005), pp. 630 figure 25.1 a)
- [35] H. Goldstein, *Classical Mechanics* (Addison-Wesley, 2nd. edition 1980), pp. 143-158
- [36] R. Kaltenbaek, R. Prevedel, M. Aspelmeyer, A. Zeilinger, High-fidelity entanglement swapping with fully independent sources, arXiv:0809.3991
- [37] J. D. Franson, *Phys. Rev. Lett.* 62, 2205 (1989)
- [38] C. Simon and J.P. Poizat, *Phys. Rev. Lett.* 94, 030502 (2005)
- [39] C. Gneiting, K Hornberger, Bell test for the free motion of material particles, arxiv:0807.1448
- [40] H. De Raedt, K. De Raedt, K. Michielsen, K. Keimpema, S. Miyashita, *J. Comp. Theor. Nanosci.* 4, pp. 957 (2007), arXiv:0712.3781v2
- [41] E. Nelson, space time chance, popular talk, available at <http://www.math.princeton.edu/~nelson/papers/cti.html>
- [42] L. Ballentine, *Quantum Mechanics, A modern development* (World Scientific 1998), pp. 608
- [43] D. Greenberger, M. Horne, A. Shimony, A. Zeilinger, Bell's theorem without inequalities, *Am. J. Phys.* 58, pp. 1131 (1990)
- [44] J. S. Bell, Introduction to the hidden variable question, in [5]
- [45] J. F. Clauser, M. A. Horne, *Phys. Rev. D* 10, pp. 526 (1974)
- [46] H. P. Stapp, *Found. Phys.* 10, pp. 767 (1980)
- [47] A. Fine, *J. Math Phys*, 23, pp. 1306 (1982)

- [48] D. Dürr, *Bohmsche Mechanik als Grundlage der Quantenmechanik* (Springer, Berlin 2001)
- [49] A. Khrennikov, Bell's inequality: Physics meets Probability, arxiv:0709.3909
- [50] R. Ursin, F. Tiefenbacher, T. Schmitt-Manderbach, H. Weier, T. Scheidl, M. Lindenthal, B. Blauensteiner, T. Jennewein, J. Perdigues, P. Trojek, B. Oemer, M. Fuerst, M. Meyenburg, J. Rarity, Z. Sodnik, C. Barbieri, H. Weinfurter, A. Zeilinger, *Nature Physics* 3, pp. 481 (2007)
- [51] L. Ballentine, *Rev. Mod. Phys.* 42, pp. 358 (1970)
- [52] M. Redhead, *Incompleteness, nonlocality and realism* (Clarendon Press, Oxford 1987)
- [53] R. Hakim, *J. Math. Phys.* 9, pp. 1805 (1968)
- [54] R. M. Dudley, *Ark. Fys. Math* 6, pp. 241 (1965)
- [55] R. M. Dudley, *Rocky Mount. J. Math.* 4, pp. 401 (1974)
- [56] F. Guerra, P. Ruggiero, *Lett. al. Nuovo Cimento* 23, pp. 529 (1978)
- [57] W. Lehr, J. L. Park, *J. Math. Phys.* 18, pp. 1235 (1977)
- [58] N. Cufaro-Petroni, C. Dewdney, P. Holland, T. Kyprianidis, J. P. Vigièr, *Phys. Rev. D* 32, pp. 1375 (1985)
- [59] M. Serva, *Ann. Inst. Henri Poincaré* 49, pp. 415 (1988)
- [60] M. Pavon, *J. Math. Phys.* 42, pp. 4846 (2001), arXiv:quant-ph/0108139v1
- [61] P. Garbaczewski, *Phys. Lett. A* 162, pp. 129 (1992)
- [62] B. Gaveau, T. Jacobson, M. Kac, L.S. Schulman, *Phys. Rev. Lett.* 53, pp. 419 (1984)
- [63] G. F. de Angelis, G. J. Lasinio, M. Serva, N. Zanghi, *J. Phys. A: Math. Gen.* 19, pp. 865 (1986)
- [64] D. G. C. McKeon, G. N. Ord, *Phys. Rev. Lett.* 69, pp. 3(1992)
- [65] J. Dunkel, P. Hänggi, *Phys. Rev. E* 74, 051106 (2006); Erratum: E74, 069902 (2006), arXiv:cond-mat/0607082v4
- [66] J. Dunkel, P. Talkner, P. Hänggi, *Phys. Rev. D* 75, 043001 (2007), arXiv:cond-mat/0608023v3
- [67] D. Cubero, J. Casado-Pascual, J. Dunkel, P. Talkner, P. Hänggi, *Phys. Rev. Lett.* 99, 170601 (2007), arXiv:0705.3328v2
- [68] I. Müller, *Z. Phys.* 198, pp. 329 (1967)
- [69] W. Israel, J. M. Stewart, *Ann. Phys.* 118, pp. 341 (1979)
- [70] I. Liu, I. Müller and T. Ruggeri, *Ann. Phys.* 169, pp. 191 (1986)
- [71] I. Müller, *Speeds of Propagation in Classical and Relativistic Extended Thermodynamics*, *Liv. Rev. Rel* 2 (1999), URL cited on 30.12.2008: <http://www.livingreviews.org/lrr-1999-1>
- [72] S. R. DeGroot, W. A. van Leeuwen, Ch. G. van Weert, *Relativistic Kinetic Theory. Principles and Applications* (North Holland, Amsterdam 1980)
- [73] A. Muronga, *Phys. Rev. C* 76, 014909 (2007), arxiv:nucl-th/0611090v3
- [74] A. Muronga, *Phys. Rev. C* 76, 014910 (2007), arxiv:nucl-th/0611091v3
- [75] H. Kleinert, *Path Integrals in Quantum Mechanics, Statistics, Polymer Physics, and Financial Markets*, chapter 10, 4th edition (World Scientific, Singapore, 2004), also available at <http://users.physik.fu-berlin.de/~kleinert/b5/psfiles/pthic10.pdf>