# A Kinetic Model for Grain Growth

R. Henseler<sup>\*</sup>

M. Herrmann<sup> $\dagger$ </sup>

nann<sup>†</sup> B. Niethammer<sup>‡</sup> J. J. L. Velázquez<sup>§</sup>

October 27, 2018

#### Abstract

We provide a well–posedness analysis of a kinetic model for grain growth introduced by Fradkov which is based on the *von Neumann–Mullins law*. The model consists of an infinite number of transport equations with a tri-diagonal coupling modelling topological changes in the grain configuration. Self–consistency of this kinetic model is achieved by introducing a coupling weight which leads to a nonlinear and nonlocal system of equations.

We prove existence of solutions by approximation with finite dimensional systems. Key ingredients in passing to the limit are suitable super–solutions, a bound from below on the total mass, and a tightness estimate which ensures that no mass is transported to infinity in finite time.

Keywords: grain growth, kinetic model, infinite-dimensional system

AMS Subject Classification: 35F25, 35R15, 74A50

# 1 Introduction

Many technologically useful materials are poly-crystalline aggregates, composed of a huge number of crystallites, called grains, separated by so-called grain boundaries. Typically such materials undergo an aging process leading to coarsening of the grain structure and therefore inducing changes in mechanical, electrical, optical, and magnetic properties of the material. For details we refer to the review articles by Fradkov and Udler [FU94] and by Thompson [Tho01].

Different approaches for modelling grain growth in two space dimensions are established in the literature. In Monte–Carlo models, compare [ASGS84a, ASGS84b], the kinetics of the boundary motion are simulated by employing a Monte–Carlo technique for moving discrete lattice points describing the microstructure. An attractive feature of this model is the simple handling of topological events like grain boundary flipping and grain disappearance.

Boundary tracking models based on partial differential equations as discussed in [KL01, MNT04] offer an alternative to Monte–Carlo models as they deal with quantities of lower dimension. They can be further reduced to so–called vertex models where movement of grain boundaries is projected onto the triple–junctions, see [KNN89, HNO03]. In both cases, however, topological changes require extra treatment.

In the sequel we focus on a kinetic mean-field type models [Fra88, Mar87, Fly93] based on the *von* Neumann-Mullins law. Such models consider time-dependent distribution functions for the grain areas and the number of sides per grain. Grain areas change according to the *von Neumann-Mullins law*, topological changes are modelled by collision-type operators. Fradkov[Fra88] was the first to develop a

<sup>\*</sup>Universität Bonn, Germany, henseler@uni-bonn.de

 $<sup>^{\</sup>dagger} University \ of \ Oxford, England, \texttt{michael.herrmann@maths.ox.ac.uk}$ 

<sup>&</sup>lt;sup>‡</sup>University of Oxford, England, niethammer@maths.ox.ac.uk

<sup>§</sup>Instituto de Ciencias Mathemáticas (CSIC-UAM-UC3M-UCM), Spain, jj\_velazquez@mat.ucm.es

model of this type which consists of an infinite–dimensional system of transport equations with a nonlocal weight, making the equations nonlinear. In this article we establish a rigorous well–posedness theory for this model.

# 2 The model

In this section we present a derivation of Fradkov's kinetic model for grain growth which is based on the *von Neumann–Mullins law* for the change of grain areas and the so–called 'gas' approximation to describe topological changes in a 2D network of grains. Starting point is the isotropic mean–curvature flow for the grain boundaries with equilibrium of forces at triple junctions.

# 2.1 Networks of grains with triple junctions

Motion by mean curvature and equilibrium of forces at triple junctions Mean curvature flow coupled with equilibrium of forces at triple junctions is a widely accepted model for two-dimensional grain growth [BR93, KL01, MNT04]. For simplicity our objects are 1-periodic spatial networks whose curves meet in triple junctions (Fig. 1). We restrict ourselves in the following to the case of isotropic surface energies, such that the curves move according to the isotropic mean curvature flow. Moreover, we assume that the mobility of the triple junctions is infinite compared to the mobility of the grain boundaries, and this leads to equilibrium of forces at triple junctions. In the isotropic case this condition, also known as *Herring condition*, just means that the curves meet in an angle of  $2\pi/3$ . The Herring condition also arises as the natural boundary condition in the interpretation of the mean curvature flow of networks as  $L^2$ -gradient flow of the surface energy, see [TC94, HNO03].



Figure 1: Cartoon of a 2D network of grains with triple junctions indicated by bullets. The Herring condition implies all angles to equal  $2\pi/3$ .

**Von Neumann–Mullins law** Under the assumptions stated above (isotropic surface energy, equal mobility of grain boundaries, and infinite mobility of triple junctions), one can derive a law of motion for the area a(t) of a single grain with n edges [Mul56], known as the *von Neumann–Mullins law*:

$$\frac{\mathrm{d}}{\mathrm{d}t}a\left(t\right) = M\sigma\frac{\pi}{3}\left(n-6\right)\tag{1}$$

Here M denotes the mobility of the grain boundaries and  $\sigma$  the surface tension. The proof uses a direct geometric computation involving motion by curvature of the grain boundaries and the prescribed jumps of the outer normal by  $2\pi/3$  at triple junctions.

The von Neumann–Mullins law implies that grains with less than six edges shrink, those with more than six grow, and such with exactly six edges retain their area (possibly not their shape).

**Topological changes** The evolution by mean curvature is well–defined until two vertices on a grain boundary collide, after which topological rearrangements may take place. This happens when either an edge or a whole grain vanishes. In the first case an unstable fourfold vertex is produced, which immediately splits up again, usually in such a way that two new vertices are connected by a new edge. In this case, two neighbouring grains decrease their topological class (i.e. the number of edges), whereas



Figure 2: Neighbour switching

the two other grains increase it (Fig. 2). The second case causing topological rearrangements is grain vanishing. Each grain vanishing is accompanied by disappearance of two vertices and three edges. Due to the *von Neumann–Mullins law* we only take grains with topological class  $2 \le n \le 5$  into account. Grains with n = 2 and n = 3 vanish in a single possible way. For n = 4 we observe two topologically



Figure 3: Grain vanishing

distinguishable possibilities and for n = 5 even five possible local configurations (Fig. 3). For further details on the resulting topological classes we refer to the review article by Fradkov and Udler [FU94].

It is unclear by which mechanism a specific topological configuration is selected within switching or after vanishing events. A natural idea is to compute all possible local configurations and select the one that minimises energy locally in the best way, see [HNO03].

### 2.2 Kinetic model for networks of grains

Our next aim is to derive the kinetic model for large networks of grains with triple junctions. This kinetic model comprises the same essential features as the gradient flow dynamics, but differs in some aspects.

**One-particle distribution** Following [Fra88, FU94] we introduce a number density  $f_n(a, t)$  that measures the number of grains with topological class  $n \ge 2$  and area  $a \ge 0$  at time  $t \ge 0$ . Using the *von* Neumann-Mullins law (1) we can describe the evolution of f by transport equations

$$\partial_t f_n(a,t) + (n-6) \partial_a f_n(a,t) = 0 \tag{2}$$

as long as no topological rearrangements take place. Furthermore we choose the following boundary conditions

$$f_n(0,t) = 0 \qquad \text{for} \quad n > 6,$$

ensuring that no additional mass is transported from the negative half-axis to the positive one. This means no additional grains can be created.

**'Collision' operator** To model topological changes we introduce a collision term  $(\hat{J}f)_n$  on the r.h.s. of (2) which couples the equations for different topological classes. We define *topological fluxes*  $\eta_n^+$  and  $\eta_n^-$  denoting the flux from class n to n+1 and from n to n-1, respectively, so that

$$(\tilde{J}f)_n = \eta_{n-1}^+ + \eta_{n+1}^- - \eta_n^+ - \eta_n^-$$

Next we state the so-called 'gas' approximation of the collision term by Fradkov [Fra88] that takes into account the following transitions between topological classes.

- Switching events cause simultaneously both a transition from n to (n-1) (grains that contained the vanishing edge) and a transition from n to (n+1) (grains that contain the created edge).
- Vanishing of a neighbouring grain corresponds to a transition from n to (n-1).

The restriction to these elementary events for topology changes is a simplification of the real dynamics in networks of grains. In particular, we ignore that the topological class of a grain is lowered by two if the neighbouring annihilated grain was a lens with topological class n = 2. Fradkov and Udler argue [FU94] that such an event takes place only very rarely as the number of lenses itself is already very small. In addition, we ignore that the annihilation of a grain with topology class n = 5 causes other grains to increase their topology class, compare Figure 3.

**Transition rates** We make a further simplification by assuming constitutive relations for the transitions rates, that in particular ignore all neighbour correlations. More precisely, in what follows we assume that the topological fluxes are given by

$$\eta_n^+ = \Gamma\beta n f_n, \quad \eta_n^- = \Gamma \left(\beta + 1\right) n f_n, \tag{3}$$

where the coupling weight  $\Gamma$  describes the intensity of topological changes and depends in a self-consistent way on the complete state of the system, see (6) below. The parameter  $\beta$  in (3) measures the ratio between switching events and vanishing events. In principle one could allow for arbitrary values of  $\beta$  but our analysis requires  $0 < \beta < 2$ . Moreover, [FUK88] provide numerical evidence that experimental data correspond to  $0.45 \leq \frac{\beta}{1+\beta} \leq 0.6$ , that means  $0.8 \leq \beta \leq 1.5$ .

According to (3) the collision terms are given by  $\tilde{J}f = \Gamma(f)Jf$  with

$$(Jf)_{2} = 3 (\beta + 1) f_{3} - 2\beta f_{2},$$
  

$$(Jf)_{n} = (\beta + 1) (n + 1) f_{n+1} - (2\beta + 1) n f_{n} + \beta (n - 1) f_{n-1} \quad \text{for} \quad n > 2.$$
(4)

Notice that this definition ensures the zero balance property, that is

$$\sum_{n \ge 2} (Jf)_n (a, t) = 0 \qquad \text{for all } a, t \ge 0.$$
(5)

This identity reflects that for each  $a_0 > 0$  the number of grains with area  $a_0$  does not change due to neither switching nor vanishing events.

**Evolution equation** The coupling weight  $\Gamma$  which makes the equations nonlinear (and nonlocal in the grain area variable a) is chosen as

$$\Gamma(f) = \Gamma_N(f) / \Gamma_D(f)$$

with

$$\Gamma_N(f(t)) = \sum_{n \ge 2} (n-6)^2 f_n(0,t), \quad \Gamma_D(f(t)) = \sum_{n \ge 2} n \int_0^\infty f_n(a,t) \, \mathrm{d}a - 2\,(\beta+1) \int_0^\infty f_2(a,t) \, \mathrm{d}a. \tag{6}$$

We will see in §2.3 below, that this choice of  $\Gamma$  guarantees consistency as the solution satisfies the polyhedral formula, compare (9), and conserves the total area covered by the grains under the evolution.

The kinetic model we consider in this paper is thus given by

$$\partial_t f_n(a,t) + (n-6)\partial_a f_n(a,t) = \Gamma(f(t))(Jf)_n(a,t).$$
(7)

These equations (7) are basically the same as in the work of Fradkov [Fra88, FUK88, FU94]. The coupling term  $(Jf)_2$  differs and we do not neglect  $\int f_2 da$  within  $\Gamma_D(f)$ . We refer moreover to [BKLT06], which presents some formal analysis and numerical simulations for a similar kinetic model. This model relies on different expressions for  $\eta_n^{\pm}$  but comprises the same essential features as (7).

Finally, we mention the following implication of Definition (6). As long as no vanishing events take place, that means as long as no small grains with topology class  $2 \le n \le 5$  do exist, switching events do likewise not occur, and (7) reduces to a system of uncoupled transport equations. This is in contrast to the gradient flow dynamics in which edge switching events occur independently of grain vanishing.

# 2.3 Qualitative properties of the kinetic model

We summarise the most important properties of the kinetic model. In order to simplify the presentation we argue by means of formal analysis but mention that all results will be proven rigorously within §3.

**Decreasing number of grains** Since equations (7) reflect a coarsening process, it is clear that the total number of grains N(f(t)) with

$$N(f) = \sum_{n \ge 2} \int_{0}^{\infty} f_n(a) \,\mathrm{d}a \tag{8}$$

should decrease in time. This is also satisfied for solutions to our model as grains with topology class n < 6 can shrink to area zero so that they are annihilated. More precisely, the evolution equations imply

$$\frac{\mathrm{d}}{\mathrm{d}t}N(f(t)) = \sum_{n\geq 2} (n-6) f_n(0,t) = \sum_{n=2}^5 (n-6) f_n(0,t) \le 0.$$

**Polyhedral formula** In this section we motivate the choice of  $\Gamma$  in (6). It is essential that our kinetic model reflects all the properties which are satisfied by a grain configuration which covers the complete area and where edges only meet in triple junctions. Hence we need to ensure that Euler's polyhedral formula is satisfied. For a finite network of grains Poincaré's version of the polyhedral formula reads

$$V + F - E = \chi(g).$$

with V, E, and F being the number of vertices, edges, and facets, respectively. Moreover, g is the genus of the surface and  $\chi(g) = 2 - 2g$  the corresponding Euler characteristic. We can encode that grain boundaries only meet in triple junctions by setting V = 2/3E, and the polyhedral formula reduces to

$$3F - E = 3\chi(g).$$

In what follows we set  $\chi(g) = 0$  because the network of grains is usually considered on a two-dimensional torus. In the kinetic model the *normalised* numbers of facets and edges are given by

$$F(f) = N(f) = \sum_{n \ge 2} \int_{0}^{\infty} f_n(a) \, \mathrm{d}a, \quad E(f) = \frac{1}{2} \sum_{n \ge 2} n \int_{0}^{\infty} f_n(a) \, \mathrm{d}a,$$

respectively, and consequently we require each solution of (7) to satisfy

$$P(f(t)) = 0 \tag{9}$$

for all times  $t \ge 0$ , where the *polyhedral defect* is given by

$$P(f) = \sum_{n \ge 2} (n-6) \int_{0}^{\infty} f_n(a) \, \mathrm{d}a.$$
(10)

The main observation is that our choice of  $\Gamma$  guarantees (9). Indeed, due to (5) we find

$$\frac{\mathrm{d}}{\mathrm{d}t}P(f(t)) = -\sum_{n\geq 2} (n-6)^2 \int_0^\infty \partial_a f_n(a,t) \,\mathrm{d}a + \sum_{n\geq 2} (n-6) \int_0^\infty \Gamma(f(t)) \,(Jf)_n(a,t) \,\mathrm{d}a$$
$$= \sum_{n\geq 2} (n-6)^2 f_n(0,t) + \Gamma(f(t)) \sum_{n\geq 2} n \int_0^\infty (Jf)_n(a,t) \,\mathrm{d}a,$$

and a simple calculation shows

$$\begin{split} \sum_{n\geq 2} n\left(Jf\right)_n &= \beta \sum_{n\geq 3} n\left(n-1\right) f_{n-1} + \left(\beta+1\right) \sum_{n\geq 2} n\left(n+1\right) f_{n+1} - \beta \sum_{n\geq 2} n^2 f_n - \left(\beta+1\right) \sum_{n\geq 3} n^2 f_n \\ &= 2\left(\beta+1\right) f_2 - \sum_{n\geq 2} n f_n, \end{split}$$

which implies

$$\frac{\mathrm{d}}{\mathrm{d}t}P(f(t)) = \Gamma_N(f) - \Gamma(f)\Gamma_D(f) = 0$$

thanks to (6). Hence, the polyhedral formula (9) is satisfied for all t > 0 if it is satisfied by the initial data.

**Conservation of area** As a consequence of the polyhedral formula we obtain that the total covered area A(f(t)) with

$$A(f) = \sum_{n \ge 2} \int_{0}^{\infty} a f_n(a) \,\mathrm{d}a \tag{11}$$

is a conserved quantity. This follows from

$$\frac{\mathrm{d}}{\mathrm{d}t}A(f(t)) = -\sum_{n\geq 2} (n-6) \int_{0}^{\infty} a\partial_a f_n(a,t) \,\mathrm{d}a + \Gamma(f(t)) \int_{0}^{\infty} a \sum_{n\geq 2} (Jf)_n(a,t) \,\mathrm{d}a$$
$$= \sum_{n\geq 2} (n-6) \int_{0}^{\infty} f_n(a,t) \,\mathrm{d}a = P(f(t)) = 0,$$

where we used an integration by parts and the zero balance property (5).

#### 2.4 Statement of the main result

Our main result in this paper concerns the existence of mild solutions to the following initial and boundary value problem, provided that the initial data satisfy certain assumptions.

**Problem 2.1.** For fixed  $0 < T < \infty$ , and given initial data  $g = g_n(a)$  we seek mild solutions  $f = f_n(a, t)$  to the following infinite system of coupled transport equations

$$\partial_t f_n(a,t) + (n-6) \partial_a f_n(a,t) = \Gamma(f(t)) (Jf)_n(a,t)$$

with initial and boundary conditions

$$f_n(a,0) = g_n(a)$$
 for  $n \ge 2$ ,  
 $f_n(0,t) = 0$  for  $n > 6$ ,

where  $a \ge 0, t \in [0, T], n \ge 2$ .

A first assumption we have to make concerns the choice of  $\beta$ . In what follows we always suppose that  $\beta$  is a fixed constant with  $0 < \beta < 2$ , where the upper bound is necessary in order to ensure that  $\Gamma_D(f)$  is non-negative for all  $f \ge 0$ . Further necessary assumptions regard the initial data.

Assumption 2.2. Suppose that

1. g is non-negative with  $g_n(0) = 0$  for all n > 6,

2. 
$$\sum_{n \ge 2} (1+n) ||g_n||_{0,1} < \infty$$
 with  $||g_n||_{0,1} = \int_0^\infty (1+a)g_n(a) da$ ,

3. g fulfils the polyhedral formula P(g) = 0.

Moreover, suppose that g is sufficiently regular and decays sufficiently fast in n- and a-direction,

According to the discussion in §2, the first three assumptions on the initial data appear very naturally. Our regularity and decay assumptions, however, are needed for technical reasons and can probably be weakened at the price of more analytical effort. The precise statement of these assumptions appears below, but we mention that we mainly assume all functions  $g_n$  to be equi-continuous and to decay exponentially with respect to a and n.

Our main result can be summarised as follows.

**Theorem 2.3.** For any initial data that satisfy Assumption 2.2 there exists a unique mild solution f to Problem (2.1) for all  $0 \le t < \infty$ . Moreover, this solution conserves the area with non-increasing number of grains, and all states f(t) satisfy Assumption 2.2.

The details of the proof are presented within §3 and rely on the following key ideas. In §3.1 we introduce an approximate system of coupled transport equations by neglecting all topological classes with  $n > n_0$ .

In order to construct mild solutions for the approximate system we solve the transport equations explicitly, and apply *Duhamel's Principle* (or *Variation of Constants*) to the collision operator. This is discussed in §3.2.

Freezing the coupling weight  $\Gamma$  in the approximate equations we can derive a comparison principle that implies both, the non-negativity of solutions and the existence of an appropriate super-solution. This will be done in §3.3.

In  $\S3.4$  we establish the short time existence and uniqueness for admissible solutions to the approximate system. To this end we propose a suitable iteration scheme, and make use of Banach's Contraction Principle.

The decay behaviour with respect to a and n is investigated in §3.5. We derive several tightness estimates, and as a consequence we obtain long-time existence for the approximate system together with estimates that are uniform in  $n_0$ .

In §3.6 we show the existence of mild solutions to Problem 2.1 by passing to the limit  $n_0 \to \infty$ . The Arzela–Ascoli theorem provides that solutions to the approximate problem converge to some reasonable limit, and the tightness estimates ensure that this limit provides an admissible solution. Finally, we sketch how both, the uniqueness of solutions and the continuous dependence on the initial data, can be obtained.

### 2.5 Further considerations

Here we point out some open questions and directions for future research.

**Stationary solutions** Nontrivial stationary solutions to (7), i.e.  $f^s \neq 0$ , are characterised by  $f_n^s(a) = 0$  for  $n \neq 6$  and  $0 < a < \infty$  (cf. [Hen07], §6.1), but the component function  $f_6^s(a)$  can be anything. This is a further difference to the gradient flow dynamics as there not all networks of hexagons are stationary. However, such nontrivial stationary solutions to the kinetic model are expected to be unstable for the following reasons: Slightly perturbed data lead to a positive  $\Gamma(f(t))$  for some times t and are therefore affected by the coupling operator  $(Jf)_n(a,t)$ . This leads to a decrease of the total number of grains. We therefore expect stationary solutions to be unstable, and thus we suspect that  $f \to 0$  weakly for  $t \to \infty$ .

**Self-similar scaling** For a coarsening process as considered here, one usually expects to find self-similarity under dynamic scaling. The natural rescaling  $\varphi_n(\xi, t) = t^2 f_n(a, t), \xi = a/t$ , yields the following equation for self-similar solutions

$$(n-6-\xi)\partial_{\xi}\varphi_n = \Gamma(\varphi)(J\varphi)_n + 2\varphi_n, \quad n \ge 2, \quad \xi \ge 0.$$
(12)

The natural boundary conditions are  $\varphi_n(0) = 0$  for n > 6, so that the solution depends on the values of  $\varphi_2(0), \ldots, \varphi_5(0)$ . Note that the coupling weight still depends on the complete solution. A starting point for future analysis is the following observation. We can integrate (12) with respect to  $\xi$  to obtain

$$(6-n)\varphi_n(0) = \Gamma(\varphi)(J\phi)_n + \phi_n, \quad n \ge 2.$$

This is a two-point iteration scheme for  $\phi_n = \int \varphi_n(\xi) d\xi$ .

Lewis' law A natural question concerning grain growth is to ask whether there are correlations between the topological class and the area of a grain. Lewis [Lew43] observed a linear relationship examining cellular structures arising in biology, and Rivier and Lissowski [RL82] derived *Lewis' law* by maximum entropy arguments applied to cell distributions. In common with Flyvbjerg [Fly93] this so-called *Lewis' law* reads

$$\left\langle \xi \right\rangle_n = b\left(n-6\right) + c \tag{13}$$

in our model. Here  $\langle \xi \rangle_n = \int \xi \varphi_n(\xi) d\xi / \int \varphi_n(\xi) d\xi$  denotes the mean grain size in the topological class n. However, it is unclear if this phenomenological law is really applicable for grain growth. Formal computations suggest in our model that (13) is valid asymptotically for large n with  $b = 1/(\Gamma + 1)$  and  $c = b((2\beta + 1) - 6\Gamma)$  (cf. [Hen07], §6.4). Similar results are achieved by Flyvbjerg [Fly93].

# **3** Proof of the main result

#### 3.1 The approximate system

The approximate system for (7) results from the original equations by neglecting all number densities belonging to topological classes with  $n > n_0$ . More precisely, we choose the parameter  $n_0$  with  $6 < n_0 <$ 

 $\infty$ , and modify the coupling operator accordingly. The approximate coupling operator J splits into its gain and loss part, that is

$$J = J_{+} - J_{-}, \tag{14}$$

which now are given by

$$(J_{+}f)_{2} = 3 (\beta + 1) f_{3},$$
  

$$(J_{+}f)_{n} = (\beta + 1) (n + 1) f_{n+1} + \beta (n - 1) f_{n-1} \text{ for } 2 < n < n_{0},$$
  

$$(J_{+}f)_{n_{0}} = \beta (n_{0} - 1) f_{n_{0} - 1},$$
(15)

and

$$(J_{-}f)_{2} = 2\beta f_{2},$$
  

$$(J_{-}f)_{n} = (2\beta + 1) n f_{n} \quad \text{for } 2 < n < n_{0},$$
  

$$(J_{-}f)_{n_{0}} = (\beta + 1) n_{0} f_{n_{0}}.$$
(16)

Notice that for  $n < n_0$  the term  $(Jf)_n$  is defined as in the original model. For the sake of consistency we must moreover adapt the formula for  $\Gamma$ . In what follows we use the approximate coupling weight

$$\Gamma(f) = \Gamma_N(f) / \Gamma_D(f), \quad \Gamma_N(f) = \sum_{n=2}^5 (n-6)^2 f_n(0), \quad \Gamma_D(f) = -\sum_{n=2}^{n_0} \int_0^\infty n(Jf)_n(a) \, \mathrm{d}a. \tag{17}$$

Analogously to §2 we define the total area A(f), the number of grains N(f), and the polyhedral defect P(f) by

$$N(f) = \sum_{n=2}^{n_0} \int_0^\infty f_n(a) \, \mathrm{d}a, \quad A(f) = \sum_{n=2}^{n_0} \int_0^\infty a f_n(a) \, \mathrm{d}a, \quad P(f) = \sum_{n=2}^{n_0} (n-6) \int_0^\infty f_n(a) \, \mathrm{d}a,$$

and for convenience we introduce in addition

$$M(f) = \sum_{n=2}^{n_0} n \int_0^\infty f_n(a) \, \mathrm{d}a, \quad R(f) = 2(\beta+1) \int_0^\infty f_2(a) \, \mathrm{d}a - n_0 \beta \int_0^\infty f_{n_0}(a) \, \mathrm{d}a.$$

**Remark 3.1.** 1. Definitions (15) and (16) imply

$$\sum_{n=2}^{n_0} (Jf)_n = 0, \quad \sum_{n=2}^{n_0} n(Jf)_n = 2(\beta+1)f_2 - n_0\beta f_{n_0} - \sum_{n=2}^{n_0} nf_n, \tag{18}$$

and we infer that

$$A(Jf) = N(Jf) = 0, \qquad P(Jf) = M(Jf) = -\Gamma_D(f).$$
 (19)

2. The polyhedral formula P(f) = 0 implies

$$\Gamma_D(f) = -R(f) + 6N(f),$$

and if f is in addition non–negative we have, thanks to  $0<\beta<2,$ 

$$\Gamma_D(f) \ge (6 - 2(\beta + 1))N(f) > 0.$$
(20)

The approximate problem we aim to solve can now be stated as follows.

**Problem 3.2.** For fixed  $n_0 > 6$  and  $0 < T < \infty$ , and given initial data  $g = g_n(a)$  we seek (mild) solutions  $f = f_n(a, t)$  to

$$\partial_t f_n(a,t) + (n-6) \partial_a f_n(a,t) = \Gamma(f(t)) (Jf)_n(a,t)$$
(21)

with initial and boundary conditions

$$\begin{array}{rcl}
f_n(a,0) &=& g_n(a) & \quad for & 2 \le n \le n_0, \\
f_n(0,t) &=& 0 & \quad for & 7 \le n \le n_0,
\end{array}$$
(22)

where  $a \ge 0, t \in [0, T], 2 \le n \le n_0$ , and J and  $\Gamma$  are defined as in (15)–(17).

# 3.2 Transport equations and notion of mild solutions

We start with some basic facts about solutions to transport equations in the upper-right space-time quadrant  $a \ge 0$  and  $t \ge 0$ . For this reason let us consider the following system of transport equations.

**Problem 3.3.** Let  $a \ge 0$ ,  $2 \le n \le n_0$  and  $t \in [0, T]$ . For fixed initial data  $g = g_n(a)$  and given right hand side  $h = h_n(a, t)$  we seek (mild) solutions  $f = f_n(a, t)$  to

$$\partial_t f_n + (n-6)\partial_a f_n = h_n \tag{23}$$

with initial and boundary conditions as in the approximate problem, see (22).

The homogeneous problem with h = 0 can be solved explicitly by the method of characteristics. This means, the general solution to the homogeneous problem is given by  $f(t) = \mathcal{T}(t)g$ , where the group of transport operators is defined by

$$\left(\mathcal{T}(t)g\right)_n = \mathcal{T}_{n-6}(t)g_n$$

with

$$(\mathcal{T}_{n-6}(t)g_n)(a) = g_n(a - (n-6)t)$$

for all  $n \leq 6$ , whereas n > 6 corresponds to

$$(\mathcal{T}_{n-6}(t)g_n)(a) = \begin{cases} g_n(a - (n-6)t) & \text{for } a \ge (n-6)t, \\ 0 & \text{for } a < (n-6)t. \end{cases}$$

Recall that  $n \leq 6$  implies the transport velocity n-6 to be non-positive, so there is no contribution from the boundary in this case.

For non-vanishing right hand side h the solutions can be constructed by means of Duhamel's Principle. More precisely, for given  $h = h_n(a, t)$  the unique mild solution to Problem 3.3 is given by

$$f(t) = \mathcal{T}(t)g + \int_{0}^{t} \mathcal{T}(t-s)h(s) \,\mathrm{d}s.$$
(24)

Moreover, the mild solution depends continuously on the data via

$$||f(t)||_{\infty} \le ||g||_{\infty} + \int_{0}^{t} ||h(s)||_{\infty} \,\mathrm{d}s, \qquad ||f(t)||_{0,1} \le C_{n_0,t} \Big( ||g||_{0,1} + \int_{0}^{t} ||h(s)||_{0,1} \,\mathrm{d}s \Big), \tag{25}$$

where  $||g||_{0,1} := \sum_{n=2}^{n_0} \int_0^\infty (1+a) |g_n(a)| da < \infty.$ 

In the sequel we make use of the following regularity results for mild solutions, that can be derived directly from the representation formula given in (24).

**Definition 3.4.** 1. A state  $g = g_n(a)$  is called

- (a) C<sup>0</sup>-regular, if g is continuous (w.r.t. to a) and  $0 = g_n(0)$  for all  $7 \le n \le n_0$ ,
- (b)  $L^1$ -regular, if  $||g||_{0,1} < \infty$ .
- 2. The right hand side  $h = h_n(a, t)$  is called
  - (a)  $C^0$ -regular, if h is continuous (w.r.t. to (a, t)),
  - (b)  $L^1$ -regular, if  $\int_0^T ||h(t)||_{0,1} dt < \infty$ .

Moreover, we say the solution  $f = f_n(a, t)$  is *regular*, if f is a regular right hand side and f(t) is a regular state for all  $t \in [0, T]$ , where 'regular' means either  $C^0$ - or  $L^1$ -regular.

**Lemma 3.5.** The mild solution f to Problem 3.3 given in (24) has the following properties.

- 1. If the data are  $C^0$ -regular, then f is  $C^0$ -regular. In particular we have  $f_n(0,t) = 0$  for all t and  $7 \le n \le n_0$ .
- 2. If the data are  $L^1$ -regular, then f is  $L^1$ -regular.

Finally, we collect some properties of mild solutions to be used in  $\S3.4$ .

**Remark 3.6.** If the data g and h are compactly supported in  $\{a : 0 \le a \le a_0\}$ , then the mild solution f(t) is compactly supported in  $\{a : 0 \le a \le a_0 + (n_0 - 6)t\}$ 

**Lemma 3.7.** Suppose that the data (g,h) for Problem 3.3 are  $C^0$ - and  $L^1$ -regular. Then the mild solution f satisfies

$$N(f(t)) = N(g) + \int_{0}^{t} N(h(s)) \,\mathrm{d}s + \sum_{n=2}^{5} (n-6) \int_{0}^{t} f_{n}(0,s) \,\mathrm{d}s,$$
  

$$P(f(t)) = P(g) + \int_{0}^{t} P(h(s)) \,\mathrm{d}s + \int_{0}^{t} \Gamma_{D}(f(s)) \,\mathrm{d}s,$$
  

$$A(f(t)) = A(g) + \int_{0}^{t} A(h(s)) \,\mathrm{d}s + \int_{0}^{t} P(f(s)) \,\mathrm{d}s.$$
(26)

In particular, N(f(t)), P(f(t)), and A(f(t)) are continuously differentiable with respect to t.

*Proof.* For classical solutions the differential counterparts of all assertion follow directly from the differential equation (23). For mild solutions we approximate with classical solutions. In fact, we can approximate the data (g, h) by some regular data  $(\tilde{g}, \tilde{h})$  that satisfy the boundary condition and are both differentiable and  $L^1$ -regular. The corresponding solution  $\tilde{f}$  to Problem 3.3 is then a classical solution and satisfies (26). Finally,  $\tilde{f} - f$  is a mild solution to Problem 3.3 with data  $(\tilde{g} - g, \tilde{h} - h)$ , and using (25) we infer that f satisfies (26).

# 3.3 Auxiliary problem with prescribed coupling weight $\Gamma$

In this section we consider an auxiliary problem that results from the approximate system 3.2 by prescribing the coupling weight  $\Gamma$  as a function of time, and study both the existence and qualitative properties of solutions in the space of bounded and continuous functions. In particular, we will derive comparison results for such solutions. Below in §3.4 we apply our results in the context of the approximate problem, and show that each solutions to the approximate problem must be non–negative and bounded from above by some appropriately chosen super–solution.

The auxiliary problem can be stated as follows.

**Problem 3.8.** Let  $0 < T < \infty$  be arbitrary,  $\Gamma \in C([0, T])$  be some non-negative weight function, and g be some initial data. Then we seek (mild) solutions f to

$$\partial_t f_n + (n-6) \,\partial_a f_n = \Gamma(t) (Jf)_n \tag{27}$$

with initial and boundary conditions (22).

The state space and solution space for Problem 3.8 are given by

$$Y_{\text{aux}} = BC([0, \infty); X_{\text{aux}})$$
 and  $\mathcal{Y}_{\text{aux}} = C([0, T]; Y_{\text{aux}}),$ 

respectively, where  $X_{\text{aux}} \cong \mathbb{R}^{n_0-1}$  denotes the set of all tuples  $(x_2, ..., x_{n_0})$ .

The key idea for arriving at comparison results for the auxiliary problem is to split the exchange operator into its loss and gain part according to (14). More precisely, we regard each solution to the approximate problem (27) as mild solution to

$$\partial_t f_n + (n-6) \,\partial_a f_n + \Gamma(t) J_- f(t) = h(t), \tag{28}$$

where the right hand side is given by

$$h(t) = \Gamma(t)J_+f(t). \tag{29}$$

Doing so we benefit from the following two observations. The loss operator  $J_{-}$  is diagonal and can hence easily be incorporated into the homogeneous problem. The resulting solution operators take care of the transport as before, but describe additionally the exponential relaxation of  $f_n$  to 0 along each characteristic line with local rate proportional to  $\Gamma(t)$ . In particular, these solution operators preserve the non-negativity of the initial data for all times. On the other side, the gain operator  $J_{+}$  preserves the cone of non-negative functions, so that the non-negativity of solutions turns out to be a direct consequence of Duhamel's principle with respect to the gain operator.

#### Exponential relaxation along characteristics

The mild solutions to the homogeneous problem corresponding to (28) are described by a two-parameter family of linear operators defined by

$$\mathcal{T}_{\Gamma}^{-}(s,t): Y_{\text{aux}} \to Y_{\text{aux}}, \qquad \left(\mathcal{T}_{\Gamma}^{-}(s,t)f\right)_{n} := \exp\left(-u_{n} \int_{s}^{t} \Gamma(\tilde{s}) \,\mathrm{d}\tilde{s}\right) \mathcal{T}_{n-6}(t-s)f_{n}, \qquad (30)$$

where s and t are two times with  $0 \le s \le t \le T$ , and  $u_n$  is the relaxation constant appearing in the loss operator. More precisely, according to (15) and (16) we have

$$u_2 = 2\beta, \qquad u_{n_0} = (\beta + 1)n_0, \qquad \text{and} \qquad u_n = (2\beta + 1)n \quad \text{for} \quad 2 < n < n_0.$$
 (31)

With (30), each mild solution to (28) satisfies

$$f(t) = \mathcal{T}_{\Gamma}^{-}(0,t)g + \int_{0}^{t} \mathcal{T}_{\Gamma}^{-}(s,t)h(s) \,\mathrm{d}s,$$

with h as in (29). As outlined above, the introduction of  $\mathcal{T}_{\Gamma}^{-}(s,t)$  is motivated by technical reasons but does not change the auxiliary problem. More precisely, we find the following equivalence.

**Remark 3.9.** A mild solution  $f \in \mathcal{Y}_{aux}$  to Problem 3.8, that is

$$f(t) = \mathcal{T}(t)g + \int_{0}^{t} \mathcal{T}(t-s)\Gamma(s)(Jf(s)) \,\mathrm{d}s,$$

also satisfies

$$f(t) = \mathcal{T}_{\Gamma}^{-}(0,t)g + \int_{0}^{t} \mathcal{T}_{\Gamma}^{-}(s,t)\Gamma(s)(J_{+}f(s)) \,\mathrm{d}s$$

and vice versa.

#### Super-solution to the auxiliary problem

In order to establish an existence result for mild solutions we start by identifying a suitable super-solution  $\phi$  to Problem 3.8 that does not depend on t and a, and provides suitable a priori bounds, see Lemma 3.12.

**Remark 3.10.** Let  $\phi \in X_{n_0}$  be defined by

$$\phi_n(a) = \frac{1}{n\beta} \left(\frac{\beta}{1+\beta}\right)^n.$$

Then,  $\phi$  is a solution to  $J\phi = 0$ . Moreover,  $\overline{f}$  with  $\overline{f}_n(a,t) = \phi_n$  for all  $t \ge 0$ ,  $a \ge 0$  and  $2 \le n \le n_0$  provides a solution to the differential equations (27).

*Proof.* For  $n \geq 3$  we find

$$j_n := (\beta+1) n\phi_n - \beta(n-1)\phi_{n-1} = \left(\frac{\beta+1}{\beta}\right) \left(\frac{\beta}{\beta+1}\right)^n - \left(\frac{\beta}{\beta+1}\right)^{n-1} = 0,$$

and this implies  $(J\phi)_n = j_{n+1} - j_n = 0$  for all  $2 < n < n_0$ . Moreover, we have  $(J\phi)_2 = j_3 = 0$  and  $(J\phi)_{n_0} = -j_{n_0} = 0$ .

We next show that a suitably chosen multiple of  $\overline{f}$  provides an upper bound for each solution to the approximate problem. For this reason we introduce a special norm for tuples  $(x_2, ..., x_{n_0}) \in X_{aux}$  by

$$|x|_{\beta} := \sup_{2 \le n < n_0 + 1} \frac{|x_n|}{\phi_n}.$$
(32)

This norm can easily be extended to states  $f \in Y_{aux}$  via  $||f||_{\beta} := \sup_{a \ge 0} |f(a)|_{\beta}$ , and writing

$$||f||_{\beta} = \sup_{2 \le n < n_0 + 1} \lfloor f \rfloor_{n, \beta}, \qquad \lfloor f \rfloor_{n, \beta} := \sup_{a \ge 0} \frac{|f_n(a)|}{\phi_n}$$
(33)

we infer that

$$||f_n||_{\infty} \le ||f||_{\beta} \phi_n \le C_{\beta} ||f||_{\beta}.$$

Notice that the norms  $||\cdot||_{\beta}$  and  $||\cdot||_{\infty}$  are equivalent on  $Y_{\text{aux}}$  since  $n_0 < \infty$ . However, we prefer working with  $||\cdot||_{\beta}$  because only this norm gives rise to estimates that are independent of  $n_0$ .

**Remark 3.11.** For all  $f \in Y_{aux}$  and  $2 \le n \le n_0$  we have

$$||(J_+f)_n||_{\infty} \le u_n \phi_n ||f||_{\beta}$$

with  $u_n$  as in (31).

*Proof.* It is sufficient to consider  $||f||_{\beta} = 1$ . For  $2 < n < n_0$  we find

$$||(J_+f)_n||_{\infty} \le (\beta+1)(n+1)||f_{n+1}||_{\infty} + \beta(n-1)||f_{n-1}||_{\infty}$$
$$\le \frac{1+\beta}{\beta} \left(\frac{\beta}{1+\beta}\right)^{n+1} + \left(\frac{\beta}{\beta+1}\right)^{n-1} = \left(\frac{\beta}{\beta+1}\right)^n \left(\frac{2\beta+1}{\beta}\right) = u_n \phi_n.$$

Finally, the estimates

$$||(J_+f)_{n_0}||_{\infty} = \beta(n_0 - 1)||f_{n_0 - 1}||_{\infty} \le \left(\frac{\beta}{\beta + 1}\right)^{n_0 - 1} = \left(\frac{\beta}{\beta + 1}\right)^{n_0} \left(\frac{\beta + 1}{\beta}\right) = u_{n_0}\phi_{n_0},$$

and

$$||(J_{+}f)_{2}||_{\infty} = (1+\beta)3||f_{3}||_{\infty} \le \frac{1+\beta}{\beta} \left(\frac{\beta}{\beta+1}\right)^{3} = \left(\frac{\beta}{\beta+1}\right)^{2} = u_{2}\phi_{2}$$

complete the proof.

### Existence of solutions to the auxiliary problem

Our existence and uniqueness results for the auxiliary problem are based on Banach's contraction principle applied to the iteration operator

$$\mathcal{I}_{\text{aux}}: \mathcal{Y}_{\text{aux}} \to \mathcal{Y}_{\text{aux}}, \qquad (\mathcal{I}_{\text{aux}}f)(t) = \mathcal{T}_{\Gamma}^{-}(t)g + \int_{0}^{t} \mathcal{T}_{\Gamma}^{-}(s,t)\Gamma(s)(J_{+}f(s)) \,\mathrm{d}s.$$

We start with deriving some a priori estimates for  $\mathcal{I}_{aux}$ .

**Lemma 3.12.** For each  $f \in \mathcal{Y}_{aux}$  we have

$$\sup_{0 \le t \le T} ||\mathcal{I}_{\text{aux}} f(t)||_{\beta} \le \max\{||g||_{\beta}, \sup_{0 \le t \le T} ||f(t)||_{\beta}\}.$$

*Proof.* Let  $u_n$  be as in (31) and  $G(t) = \int_0^t \Gamma(s) \, ds$ . Definition (30) and Remark 3.11 imply

$$\left|\left|\left(\mathcal{I}_{\mathrm{aux}}f\right)_{n}(t)\right|\right|_{\infty} \leq \phi_{n}\left(\exp\left(-u_{n}G(t)\right)\left|\left|g\right|\right|_{\beta} + \int_{0}^{t}\exp\left(-u_{n}G(t) + u_{n}G(s)\right)u_{n}\Gamma(s)\left|\left|f(s)\right|\right|_{\beta} \mathrm{d}s\right),$$

and we conclude

$$\left| \left( \mathcal{I}_{\mathrm{aux}} f \right)(t) \right|_{n,\beta} \leq \exp\left( -u_n G(t) \right) \left( ||g||_{\beta} + \mu(t) \int_0^t \exp\left( u_n G(s) \right) u_n \Gamma(s) \, \mathrm{d}s \right)$$
$$= \exp\left( -u_n G(t) \right) \left( ||g||_{\beta} + \mu(t) (\exp\left( u_n G(t) \right) - 1) \right), \tag{34}$$

with  $\mu(t) = \sup_{0 \le s \le t} ||f(s)||_{\beta}$ , which yields the desired estimate.

We mention that all results obtained so far in this section immediately apply to the case  $n_0 = \infty$ , provided that the initial data satisfy  $||g||_{\beta} < \infty$ .

**Corollary 3.13.** For arbitrary  $0 < T < \infty$  and any initial data  $g \in Y_{\text{aux}}$  there exists a unique mild solution  $f \in \mathcal{Y}_{\text{aux}}$  to Problem 3.8 which satisfies  $||f(t)||_{\beta} \leq ||g||_{\beta}$  for all  $t \in [0, T]$ . Moreover, this solutions is

- 1. non-negative provided that the initial data are non-negative, and
- 2.  $C^1$ -regular ( $L^1$ -regular) if the initial data are  $C^1$ -regular ( $L^1$ -regular).

*Proof.* The existence and uniqueness of a mild solution is a direct consequence of Banach's contraction principle and Lemma 3.12 (the Lipschitz constant of  $\mathcal{I}_{aux}$  with respect to  $|| \cdot ||_{\beta}$  can be read off from (34) with  $g \equiv 0$ , and is given by  $1 - \exp(-u_{n_0}G(t))$ ). Moreover, the remaining assertions are satisfied as  $\mathcal{I}_{aux}$  respects both the non-negativity and  $L^1$ -regularity.

## 3.4 Solutions to the approximate system

In this section we are going to establish local existence and uniqueness results for the approximate problem. To this end we consider the solution space  $\mathcal{Y}_{app} = C([0, T]; Y_{app})$ , where the state space given by

$$Y_{\text{app}} = Y_{\text{aux}} \cap \{ f = f_n(a) : ||f_n||_{0,1} < \infty \text{ for all } 2 \le n \le n_0 \}.$$

Of course, the set of all *admissible* states is only a proper subset of  $Y_{app}$  as any reasonable solution to the approximate problem must be non-negative and area conserving, and has to satisfy the boundary conditions and the polyhedral formula.

**Definition 3.14.** The state  $f \in Y_{app}$  is called *admissible*, if it is non-negative with A(f) > 0, and satisfies the polyhedral formula P(f) = 0 as well as the boundary conditions  $f_n(0) = 0$  for  $n \ge 7$ .

#### Iteration scheme for the approximate system

We construct solutions to the approximate system as fixed points to the following iteration operator  $\mathcal{I}_{app} : \mathcal{Y}_{app} \to \mathcal{Y}_{app}$ . For given  $f \in \mathcal{Y}_{app}$  let  $\tilde{f} = \mathcal{I}_{app} f$  be the mild solution to

$$\partial_t \tilde{f}_n + (n-6)\partial_a \tilde{f}_n = |\Gamma(f)| (Jf)_n \tag{35}$$

with initial and boundary conditions (22).

**Lemma 3.15.** For all initial data  $g \in Y_{app}$  there exist a time T > 0 and a constant  $C_0$  (both depending on  $n_0$  and g) such that the operator  $\mathcal{I}_{app}$  has the following properties.

- 1. For given  $f \in \mathcal{Y}_{aux}$  the function  $\tilde{f} = \mathcal{I}_{app} f \in \mathcal{Y}_{aux}$  is a mild solution to (35) that satisfies the initial and boundary conditions (22).
- 2. For  $C^1$ -regular data g and f the function  $\mathcal{I}_{app}f$  is  $C^1$ -regular.
- 3. The set  $\widetilde{\mathcal{Y}}_{app} = BC([0, T]; \widetilde{Y}_{app})$  with

$$\widetilde{Y}_{\text{app}} = \left\{ f \in Y_{\text{app}} : \Gamma_D(f) \ge C_0^{-1}, \quad ||f||_{\infty} \le C_0 \right\}.$$
(36)

is invariant under the action of  $\mathcal{I}_{app}$ , and the operator  $f \mapsto |\Gamma(f)| Jf$  is Lipschitz-continuous on  $\widetilde{Y}_{app}$ .

*Proof.* At first let  $C_0$  and T be arbitrary but fixed, and suppose that  $f(t) \in \tilde{Y}_{app}$  for all  $t \in [0, T]$ . For the remainder of this proof  $\tilde{C}$  always denotes a constant depending only on  $n_0$  and  $\beta$ , but the value of  $\tilde{C}$  may change from line to line. From (35) and (25) we conclude that

$$||\tilde{f}(t)||_{\infty} \le ||g||_{\infty} + \int_{0}^{t} \frac{|\Gamma_{N}(f(s))|}{\Gamma_{D}(f(s))} ||Jf(s)||_{\infty} \,\mathrm{d}s \le ||g||_{\infty} + \tilde{C} \int_{0}^{t} \frac{||f(s)||_{\infty}^{2}}{\Gamma_{D}(f(s))} \,\mathrm{d}s,$$

and due to  $f\in \widetilde{Y}_{\mathrm{app}}$  we have

$$||\hat{f}(t)||_{\infty} \le ||g||_{\infty} + \hat{C} C_0^{-3} T.$$
(37)

Moreover, since  $\Gamma_D$  is linear in f we find

$$\left|\Gamma_D(\tilde{f}(t)) - \Gamma_D(\mathcal{T}(t)g)\right| \le \int_0^t \frac{|\Gamma_N(f(s))|}{\Gamma_D(f(s))} \left|\Gamma_D(\mathcal{T}(t-s)Jf(s))\right| \,\mathrm{d}s \le \tilde{C} \,C_0^{-3} \,T.$$
(38)

In order to prove the invariance of  $\tilde{Y}_{app}$  under the action of  $\mathcal{I}_{app}$ , we choose  $C_0$  sufficiently large such that

$$2||g||_{\infty} < C_0, \quad \Gamma_D(g) > 3C_0^{-1},$$

In addition, we choose T sufficiently small with

$$\Gamma_D(\mathcal{T}(t)g) \ge 2C_0^{-1} \text{ for all } 0 \le t \le T, \qquad \tilde{C} C_0^3 T < ||g||_{\infty}, \qquad \tilde{C} C_0^3 T < C_0^{-1}.$$

and thanks to (37) and (38) we thus find  $\tilde{f} \in \mathcal{Y}_{app}$ . Finally, the claimed regularity results are provided by Lemma 3.5, and the proof of the Lipschitz-continuity is straightforward.

#### Local existence and uniqueness

We prove the existence of *admissible* solutions to the approximate system by combining the following arguments: (1) Banach's contraction principle implies the existence of a unique fixed point f for  $\mathcal{I}_{app}$ . (2) Our results concerning the auxiliary problem provide the desired non-negativity. (3) The polyhedral formula is a consequence of the equation itself, and implies the conservation of area.

**Lemma 3.16.** Let T and  $C_0$  be as in Lemma 3.15, and suppose that the initial data g are admissible. Then there exists a unique mild solution  $f \in \mathcal{Y}_{app}$  to the approximate system on the time interval [0, T]. Moreover,

- 1. all states f(t) are admissible (in the sense of Definition 3.14) and fulfil  $||f(t)||_{\beta} \leq ||g||_{\beta}$ ,
- 2. f conserves the total area and the number of grains is non-increasing.

*Proof.* Banach's contraction principle provides the existence of a fixed point  $f \in \mathcal{Y}_{app}$  with  $\mathcal{I}_{app}f = f$ . Then we consider the auxiliary problem with prescribed coupling weight  $|\Gamma(f(t))|$ , and both the positivity and boundedness of f are consequences of Corollary 3.13. In particular, we find  $\Gamma(f(t)) \ge 0$  and this shows that f is indeed a solution to the approximate system. In order to prove the polyhedral formula and the conservation of area we use basically the same computations as in §2. More precisely, Lemma 3.7 implies

$$\frac{\mathrm{d}}{\mathrm{d}t}P(f(t)) - \sum_{n=2}^{n_0} (n-6)^2 f(0,t) = \Gamma(f(t))P(Jf(t)),$$

and with (17) and (19) we infer that

$$\frac{\mathrm{d}}{\mathrm{d}t}P(f(t)) = \Gamma_N(f(t)) - \frac{\Gamma_N(f(t))}{\Gamma_D(f(t))}\Gamma_D(f(t)) = 0.$$

This implies P(f(t)) = 0 for all t due to P(g) = 0. Similarly, we find

$$\frac{\mathrm{d}}{\mathrm{d}t}A(f(t)) = P(f(t)) = 0.$$

Finally, exploiting Lemma 3.7 once again, gives

$$\frac{\mathrm{d}}{\mathrm{d}t}N(f(t)) = \sum_{n=2}^{5} (n-6)f_n(0,t) + \Gamma(f(t))N(Jf(t)) \le 0$$

where we used N(Jf) = 0 and  $f \ge 0$ .

**Remark 3.17.** The proofs of Lemma 3.15 and Lemma 3.16 imply that the local solution does exist as long as  $\Gamma_D(f(t))$  can be bounded from below by a positive constant  $d_0$ . In fact, for non-negative f both  $||f(t)||_{\infty}$  and  $\Gamma_N(f(t))$  are bounded by the super-solution  $\overline{f}_n(a,t) = ||g||_{\beta}\phi_n$ , and thus the constant  $C_0$  from (36) can be chosen to depend on  $d_0$  and g only. Consequently, each lower estimate for  $\inf_{0 \le t \le T} \Gamma_D(f(t))$  will imply the existence and uniqueness of solutions on the interval [0, T]. In what follows we derive several variants of such estimates by bounding N(f(t)) from below, respectively, which is sufficient in view of (20).

The first lower bound for the number of grains is an elementary consequence of the von Neumann– Mullins law. The main observation is that grains can grow only with speed  $n_0 - 6$ , so that for compactly supported initial data the number of grains can be bounded from below by the total area. More precisely, if the initial data g are compactly supported in  $\{a : 0 \le a \le a_0\}$  then we have

$$N(f(t)) \ge \frac{A(g)}{a_0 + t(n_0 - 6)}.$$
(39)

This bound is, however, not optimal because it holds only for compactly supported initial data and depends strongly on  $n_0$ . For this reason we finally rely on refined estimates to be derived in Lemma 3.25.

# 3.5 Tightness estimates and global existence

We introduce the notion of 'quasi-complement', that is the number of grains outside a bounding frame. By enlarging this frame in time, we show that no mass runs off at infinity in finite time, and as a consequence we establish long-time existence for the approximate system. Moreover, these tightness estimates also apply to the original problem with  $n_0 = \infty$  provided that the initial data decay sufficiently fast with respect to both a and n.

Within this subsection we use the following notations. Let f be a fixed non-negative solution to the approximate system (3.2), defined for  $0 \le t \le T < \infty$ , and set

$$\gamma = \log(1 + 1/\beta), \quad N(t) = N(f(t)), \quad \Gamma(t) = \Gamma(f(t)), \quad \overline{\Gamma}(t) = \sup_{0 \le s \le t} \Gamma(s).$$

Notice that the definition of  $\gamma$  implies  $\beta n \phi_n = \exp(-n\gamma)$  with  $\phi = (\phi_n)_{2 \le n \le n_0}$  being the super-solution from Remark 3.10.

Given  $\nu \in \mathbb{N}$  and  $\alpha \in [0, \infty)$  we define the first quasi-complement by

$$N^{\perp}(t, \alpha, \nu) = \sum_{n=\nu+1}^{n_0} \int_0^{\alpha} f_n(a, t) \, \mathrm{d}a + \sum_{n=2}^{n_0} \int_{\alpha}^{\infty} f_n(a, t) \, \mathrm{d}a$$

as the non–essential part of N(t). In order to simplify the notation we allow for  $\nu = 0$  by setting  $f_1 \equiv 0$ and  $(Jf)_1 = 0$ , so that  $N(f(t)) = N^{\perp}(t, 0, 0)$ .

In what follows we write  $N^{\perp}(t, \alpha) := N^{\perp}(t, \alpha, \lfloor \alpha \rfloor)$  if the third argument of  $N^{\perp}$  is given by rounding down the second one, and refer to

$$M^{\perp}(t,\alpha) = \sum_{n=\lfloor \alpha \rfloor + 1}^{n_0} n \int_0^{\infty} f_n(a,t) \,\mathrm{d}a + \sum_{n=2}^{n_0} \int_{\alpha}^{\infty} a f_n(a,t) \,\mathrm{d}a$$
(40)

as the second quasi-complement.

**Remark 3.18.** For all  $t \ge 0$  and  $\alpha \ge 0$  we have  $N^{\perp}(t, \alpha) \le M^{\perp}(t, \alpha)$ .

*Proof.* The desired estimate follows immediately from the definitions provided that  $\alpha \ge 1$ . For  $0 \le \alpha < 1$  we find

$$N^{\perp}(t,\alpha) = N(t) \le \sum_{n=2}^{n_0} n \int_0^\infty f_n(a,t) \,\mathrm{d}a \le M^{\perp}(t,\alpha)$$

due to  $\lfloor \alpha \rfloor = 0$ .

## Estimates for $N^{\perp}$

In order to derive suitable estimates for the first quasi-complement  $N^{\perp}$  we choose  $\nu = \lfloor \mu \rfloor$  and  $\alpha$  as functions of time. More precisely,  $\alpha$  should grow at least linearly in  $\nu$  to compensate for the transport in a along characteristic lines, and  $\nu$  should grow exponentially to control the diffusion in n.

**Lemma 3.19.** Let  $\mu, \alpha : [0, T] \to [0, \infty]$  be two smooth functions with

$$\dot{\mu}(t) \ge 0$$
 and  $\dot{\alpha}(t) \ge \max\{\lfloor \mu(t) \rfloor - 6, 0\}.$ 

Then, we have

$$N^{\perp}(t,\alpha(t),\lfloor\mu(t)\rfloor) \le N^{\perp}(0,\alpha(0),\lfloor\mu(0)\rfloor) + ||g||_{\beta}\overline{\Gamma}(t)\int_{0}^{t}\exp\left(-\gamma\lfloor\mu(s)\rfloor\right)\alpha(s) \,\mathrm{d}s \tag{41}$$

for all  $t \in [0, T]$ .

Proof. We define  $\nu(t) = \lfloor \mu(t) \rfloor$  as the integer part of  $\mu(t)$  and denote the jump of  $\nu(t)$  by  $\llbracket \nu \rrbracket(t) = \nu(t+) - \nu(t-)$ . Obviously, we can restrict ourselves to  $\nu(t) \leq n_0$ , and for simplicity we consider classical solutions. Our results then can be generalised to mild solutions by approximation arguments. At first we study the case  $\llbracket \nu \rrbracket(t) = 0$ . Differentiating  $N^{\perp}(t, \alpha(t), \nu(t))$  and using the evolution equation (21) yields

$$\frac{\mathrm{d}}{\mathrm{d}t}N^{\perp}(t,\alpha,\nu) = -\sum_{n=\nu+1}^{n_0} (n-6) f_n(\alpha,t) + \Gamma(t) \sum_{n=\nu+1}^{n_0} \int_0^{\alpha} (Jf)_n(a,t) \,\mathrm{d}a + \sum_{n=\nu+1}^{n_0} \dot{\alpha} f_n(\alpha,t) - \sum_{n=2}^{n_0} \int_{\alpha}^{\infty} (n-6) \,\partial_a f_n(a,t) \,\mathrm{d}a + \Gamma(t) \int_{\alpha}^{\infty} \sum_{n=2}^{n_0} (Jf)_n(a,t) \,\mathrm{d}a - \sum_{n=2}^{n_0} \dot{\alpha} f_n(\alpha,t) \,,$$
(42)

where  $\alpha$  and  $\nu$  are shorthand for  $\alpha(t)$  and  $\nu(t)$ , and due to (18) this implies

$$\begin{aligned} \frac{\mathrm{d}}{\mathrm{d}t} N^{\perp}\left(t,\alpha,\nu\right) &= \sum_{n=2}^{\nu} \left(n-6-\dot{\alpha}\right) f_n\left(\alpha,t\right) + \Gamma(t) \sum_{n=\nu+1}^{n_0} \int_0^\alpha \left(Jf\right)_n\left(a,t\right) \mathrm{d}a \\ &\leq \sum_{n=\nu+1}^{n_0} \Gamma(t) \int_0^\alpha \left(Jf\right)_n\left(a,t\right) \mathrm{d}a \end{aligned}$$

For  $\nu = 0$  or  $\nu = 1$  we find  $\frac{d}{dt}N^{\perp}(t, \alpha) \leq 0$  due to  $\sum_{n=2}^{n_0} (Jf)_n = 0$ , while for  $\nu > 2$  the identities (15) and (16) give

$$\sum_{n=\nu+1}^{n_0} (Jf)_n (a,t) = \beta \nu f_{\nu} (a,t) - (\beta + 1) (\nu + 1) f_{\nu+1} (a,t)$$
$$\leq \beta \nu f_{\nu} (a,t) \leq ||g||_{\beta} \beta \nu \phi_n = ||g||_{\beta} \exp(-\gamma \nu),$$

where we used that  $||f(t)||_{\beta} \leq ||g||_{\beta}$ , see Corollary 3.13. Therefore,

$$\frac{\mathrm{d}}{\mathrm{d}t}N^{\perp}\left(t,\alpha(t),\nu(t)\right) \leq \overline{\Gamma}(t)\left|\left|g\right|\right|_{\beta} \exp\left(-\gamma\nu\left(t\right)\right)\alpha\left(t\right)$$
(43)

holds for all  $\nu$ . In the case  $\llbracket \nu \rrbracket = +1$  we find

$$\begin{split} \llbracket N^{\perp}\left(t,\alpha,\lfloor\mu\rfloor\right) \rrbracket &= N^{\perp}\left(t_{+},\alpha(t_{+}),\lfloor\mu\rfloor(t_{-})+1\right) - N^{\perp}\left(t_{-},\alpha(t_{-}),\lfloor\mu\rfloor(t_{-})\right) \\ &= -\int_{0}^{\alpha(t)} f_{\lfloor\mu(t_{-})\rfloor}\left(a,t\right) \,\mathrm{d}a < 0 \end{split}$$

as an additional part in the r.h.s. of (42), so (43) still holds in a distributional sense. Finally, (41) follows by integrating (43).  $\Box$ 

Next we derive a decay result for  $N^{\perp}(t, \alpha)$  with respect to the variable  $\alpha$  which does not depend on  $n_0$  but only on the initial data g. These decay estimates turn out to be the crucial ingredient for both, the global existence proof for solutions and the passage to the limit  $n_0 \to \infty$ .

Corollary 3.20. The estimate

$$N^{\perp}(t,\alpha) \leq N^{\perp}(0,\alpha\exp\left(-t\right)) + C||g||_{\beta}\overline{\Gamma}(t)\exp\left(-\gamma\alpha\exp\left(-t\right)\right)$$

holds for all  $t \ge 0$  and  $\alpha \ge 0$ , where C depends only on  $\beta$ .

*Proof.* Let  $\alpha_0 \geq 0$  be arbitrary and consider  $\alpha(t) = \mu(t) = \alpha_0 \exp(t)$ , and let  $\nu(t) = \lfloor \mu(t) \rfloor$ . Then,  $\dot{\alpha}(t) = \mu(t) \geq \nu(t) - 6$ , and Lemma 3.19 implies

$$N^{\perp}\left(t,\alpha_{0}\exp\left(t\right)\right) \leq N^{\perp}\left(0,\alpha_{0}\right) + \left|\left|g\right|\right|_{\beta}\overline{\Gamma}(t)\,\exp\left(\gamma\right)\,\int_{0}^{t}\exp\left(-\gamma\mu\left(s\right)\right)\alpha\left(s\right)\,\mathrm{d}s,$$

where we used that  $\exp(-\gamma \lfloor \mu(s) \rfloor) \leq \exp(-\gamma \mu(s) + \gamma)$ . Replacing s by  $a = \alpha(s) = \alpha_0 \exp(s)$  we find

$$N^{\perp}(t, \alpha_{0} \exp(t)) \leq N^{\perp}(0, \alpha_{0}) + ||g||_{\beta} \overline{\Gamma}(t) \exp(\gamma) \int_{\alpha_{0}}^{\alpha_{0} \exp(t)} \exp(-\gamma a) da$$
$$\leq N^{\perp}(0, \alpha_{0}) + ||g||_{\beta} \overline{\Gamma}(t) \frac{\exp(\gamma)}{\gamma} \exp(-\gamma \alpha_{0}).$$

Since this identity holds for arbitrary t and  $\alpha_0$ , we can choose  $\alpha_0 = \alpha \exp(-t)$ , which gives the desired result.

**Remark 3.21.** The proofs of Lemma 3.19 and Corollary 3.20, and hence all estimates derived below, can be easily generalised to the original problem with  $n_0 = \infty$ , provided that  $||g||_{\beta} < \infty$ , i.e., if for  $n \to \infty$  the term  $||g_n||_{\infty}$  decays at least as fast as  $n^{-1} \exp(-\gamma n)$ .

# Estimates for $M^{\perp}$

Here we exploit the properties of the  $N^{\perp}$  and that the decay behaviour of  $M^{\perp}$  with respect to  $\alpha$  is controlled by t and the first quasi-complement of the initial data.

#### Lemma 3.22. We have

$$M^{\perp}(t,\alpha) \le C \exp\left(t\right) \left(1 + ||g||_{\beta} + ||g||_{\beta} \overline{\Gamma}(t)\right) \mathcal{N}_{0}^{\perp}(\alpha \exp\left(-t\right))$$

for all  $t \geq 0$ ,  $\alpha \geq 0$ , where  $\mathcal{N}_0^{\perp}$  depends only on the initial quasi-complement via

$$\mathcal{N}_0^{\perp}(\alpha) = (\alpha + 1) \exp\left(-\gamma\alpha\right) + \alpha N^{\perp}(0, \alpha) + \int_{\alpha}^{\infty} N^{\perp}(0, a) \,\mathrm{d}a,\tag{44}$$

and C depends only on  $\beta$ .

Proof. Notice that

$$\sum_{n=2}^{n_0} \int_{\alpha}^{\infty} a f_n(a,t) \, \mathrm{d}a = \sum_{n=2}^{n_0} \int_{\alpha}^{\infty} \int_{a}^{\infty} f_n(s,t) \, \mathrm{d}s \, \mathrm{d}a + \alpha \sum_{n=2}^{n_0} \int_{\alpha}^{\infty} f_n(a,t) \, \mathrm{d}a \tag{45}$$

via an integration by parts. Corollary 3.20 yields

$$\sum_{n=2}^{n_0} \int_{\alpha}^{\infty} f_n(a,t) \, \mathrm{d}a \le N^{\perp}(0, \alpha \exp\left(-t\right)) + C||g||_{\beta} \overline{\Gamma}(t) \, \exp\left(-\gamma \alpha \exp\left(-t\right)\right),\tag{46}$$

and this bounds the second term on the r.h.s of (45). Moreover, integrating (46) we find

$$\sum_{n=2}^{n_0} \int_{\alpha}^{\infty} \int_{a}^{\infty} f_n(s,t) \, \mathrm{d}s \, \mathrm{d}a \leq + \exp\left(t\right) \int_{\alpha \exp\left(-t\right)}^{\infty} N^{\perp}\left(0,a\right) \, \mathrm{d}a + \frac{C||g||_{\beta} \overline{\Gamma}(t) \exp\left(t\right)}{\gamma} \exp\left(-\gamma\alpha \exp\left(-t\right)\right), \tag{47}$$

and combining (46) with (47) we conclude that

$$\sum_{n=2}^{n_0} \int_{\alpha}^{\infty} a f_n(a,t) \, \mathrm{d}a \le C \exp\left(t\right) \left(1 + \left|\left|g\right|\right|_{\beta} + \left|\left|g\right|\right|_{\beta} \overline{\Gamma}(t)\right) \mathcal{N}_0^{\perp}(\alpha \exp\left(-t\right)).$$

To derive the remaining estimate we split the integration with respect to a at a = n, and obtain

$$\sum_{n=\lfloor\alpha\rfloor+1}^{n_0} n \int_0^\infty f_n(a,t) \, \mathrm{d}a \le \sum_{n=\lfloor\alpha\rfloor+1}^{n_0} n \int_0^n f_n(a,t) \, \mathrm{d}a + \sum_{n=\lfloor\alpha\rfloor+1}^{n_0} \int_n^\infty a f_n(a,t) \, \mathrm{d}a$$
$$\le \sum_{n=\lfloor\alpha\rfloor+1}^{n_0} n \int_0^n f_n(a,t) \, \mathrm{d}a + \sum_{n=2}^{n_0} \int_\alpha^\infty a f_n(a,t) \, \mathrm{d}a.$$

We have already bounded the second term on the r.h.s., while the first one can be estimated by using  $f_n(a,t) \leq ||g||_{\beta} \phi_n = ||g||_{\beta} \exp(-\gamma n)/(n\beta)$ . This gives

$$\sum_{n=\lfloor\alpha\rfloor+1}^{n_0} n \int_0^n f_n(a,t) \, \mathrm{d}a \le \frac{||g||_\beta}{\beta} \sum_{n=\lfloor\alpha\rfloor+1}^{n_0} n \exp\left(-\gamma n\right) \le C \, ||g||_\beta \int_\alpha^\infty a \exp\left(-\gamma a\right) \, \mathrm{d}a$$
$$\le C \, ||g||_\beta \int_{\alpha \exp\left(-t\right)}^\infty a \exp\left(-\gamma a\right) \, \mathrm{d}a \le C \, ||g||_\beta \mathcal{N}_0^\perp(\alpha \exp\left(-t\right))$$

and the proof is complete.

#### Positivity of numbers of grains and global existence

The initial data g are called *rapidly decreasing* if there exists two constants  $d_0$  and  $D_0$  such that

$$M^{\perp}(0,\alpha) \le D_0 \exp\left(-d_0 \alpha\right) < \infty \tag{48}$$

for all  $\alpha \geq 0$ . Notice that (48) implies that both  $N^{\perp}(0, \alpha)$  and  $\mathcal{N}_0^{\perp}(\alpha)$  decay exponentially with respect to  $\alpha$ . This follows from Remark 3.18 and Definition (44).

**Remark 3.23.** Suppose that the initial data decay exponentially with respect to both a and n, this means there constants  $\tilde{d}_0$  and  $\tilde{D}_0$  such that  $g_n(a) \leq \tilde{D}_0 \exp(-\tilde{d}_0(a+n))$ . Then there exists a suitable choice of  $d_0$  and  $D_0$  such that (48) is satisfied.

For rapidly decreasing initial data we can estimate  $M^{\perp}(t, \alpha)$  for arbitrary t and  $\alpha$  by means of Lemma 3.22.

**Corollary 3.24.** Suppose that the initial data g are rapidly decreasing. Then, for each  $t \ge 0$  there exists constants  $d_t$  and  $D_t$  such that

$$M^{\perp}(t,\alpha) \le D_t \left(1 + \overline{\Gamma}(t)\right) \exp\left(-d_t \alpha\right)$$

holds for all  $\alpha \geq 0$ . These constants depend on  $\beta$  and g, but not on  $\overline{\Gamma}(t)$  or  $n_0$ .

Recall that the lower bound for N(t) from (39) is not optimal, as it is restricted to compactly supported initial data and depends on  $n_0$ . Here we derive a better result by exploiting the tightness estimates.

**Lemma 3.25.** For rapidly decreasing initial data g and each  $t \ge 0$  there exists a constant  $C_t > 0$  such that

$$N(t) \ge C_t, \quad \overline{\Gamma}(t) \le 1/C_t.$$

In particular, this constant is independent of  $n_0$ .

*Proof.* Within this proof let  $C_t$  denote an arbitrary constant that depends only on t,  $\beta$ , and g. According to Remark 3.1 and the monotonicity of N(t) we have

$$\overline{\Gamma}(t) = \sup_{0 \le s \le t} \Gamma(s) = \sup_{0 \le s \le t} \frac{c||g||_{\beta}}{N(s)} \le \frac{c||g||_{\beta}}{N(t)}.$$
(49)

Moreover, the conservation of area implies

$$A(g) = A(f(t)) = \sum_{n=2}^{n_0} \left( \int_0^\alpha a f_n(a,t) \,\mathrm{d}a + \int_\alpha^\infty a f_n(a,t) \,\mathrm{d}a \right) \le \alpha N(t) + M^{\perp}(t,\alpha)$$

for all  $\alpha \geq 0$ , so we infer from Corollary 3.24 that

$$\alpha N(t) \ge \left(A(g) - D_t \left(1 + \overline{\Gamma}(t)\right) \exp\left(-d_t \alpha\right)\right) \ge \left(A(g) - C_t \left(1 + N(t)^{-1}\right) \exp\left(-d_t \alpha\right)\right).$$

Now we choose  $\alpha = \alpha_t$  such that  $N(t)\alpha_t = \frac{1}{2}A(g) = c_t$ , that means

$$\alpha_t = C_t \left( 1 + \ln \left( 1 + N(t)^{-1} \right) \right),$$

and hence we find

$$C_t \left( 1 + \ln \left( 1 + N(t)^{-1} \right) \right) N(t) = 1.$$

This estimate yields the existence of a constant  $C_t$  with  $N(t) > C_t$ , and thanks to (49) this provides also an corresponding upper bound for  $\overline{\Gamma}(t)$ .

**Corollary 3.26.** Suppose that the initial data g are rapidly decreasing. Then the solution from Lemma 3.16 does exist for all times  $t \ge 0$ .

*Proof.* Lemma 3.25 provides a priori estimate for both N(t) and  $\overline{\Gamma}(t)$  for arbitrary large t. Therefore, the local solution to the approximate system exists for all times, compare Remark 3.17.

#### 3.6 Passage to the limit $n_0 \to \infty$

In this section we consider a fixed final time  $0 < T < \infty$  and pass to the limit  $n_0 \to \infty$ . Consequently, from now on (i) the formulas for Jf and  $\Gamma(f) = \Gamma_N(f)/\Gamma_D(f)$  refer to (4) and (6), the definitions for the original problem with  $n_0 = \infty$ , and (ii) the number of grains N(f), the area A(f), and the polyhedral defect P(f) are given by (8), (11), and (10), respectively.

#### **Existence of solutions**

The solution space  $\mathcal{Y}$  for Problem 2.1 is given  $\mathcal{Y} = C([0, T); Y)$  for some  $0 < T < \infty$  with state space

$$Y := \left\{ f \in BC([0, \infty); X) : ||f||_{\beta} + ||f||_{\text{mom}} < \infty \right\},\$$

where  $X \cong \mathbb{R}^{\{n \ge 2\}}$  abbreviates the space of real-valued series  $(x_2, ..., x_n, ...)$ . The norm  $|| \cdot ||_{\beta}$  is defined analogously to (32) and (33) (with  $n_0 = \infty$ ), and  $|| \cdot ||_{\text{mom}}$  reads

$$||f||_{\text{mom}} := \sum_{n \ge 2} \int_{0}^{\infty} (n+a)|f_n(a)| \, \mathrm{d}a.$$

In what follows we call a state  $f \in Y$  admissible, if f is non-negative with A(f) > 0, and satisfies the polyhedral formula P(f) = 0 and the boundary conditions  $f_n(a) = 0$  for n > 6.

Our strategy for constructing mild solutions to Problem 2.1 is rather straightforward. For fixed  $0 < T < \infty$  and given admissible initial data  $g \in Y$  we consider a sequence of corresponding solutions to the approximate problem with increasing parameter  $n_0$ , and aim to show the existence of an reasonable limit in  $\mathcal{Y}$ . More precisely, within this section we consider the functions

$$g^{(n_0)} \in Y, \quad f^{(n_0)} \in \mathcal{Y}, \quad \Gamma^{(n_0)} \in C([0, T]), \quad N^{(n_0)} \in C([0, T]), \quad M^{\perp, (n_0)} \in C([0, T] \times [0, T]).$$

which are defined for  $n_0 > 6$  as follows.

1. For all  $a \ge 0$  and  $0 \le t \le T$  we have

$$g_n^{(n_0)}(a) = \begin{cases} g_n(a) & \text{for } 2 \le n \le n_0, \\ 0 & \text{for } n > n_0, \end{cases} \qquad f_n^{(n_0)}(a,t) = \begin{cases} f_n(a,t) & \text{for } 2 \le n \le n_0, \\ 0 & \text{for } n > n_0. \end{cases}$$

- 2. For each  $n_0$  the component functions  $f_2^{(n_0)}, ..., f_{n_0}^{(n_0)}$  are the unique mild solution for the approximate Problem 3.2 with initial data given by  $g_2^{(n_0)}, ..., g_{n_0}^{(n_0)}$ .
- 3.  $\Gamma^{(n_0)}$  and  $N^{(n_0)}$  are the corresponding coupling weight and number of grains, respectively.
- 4.  $M^{\perp,(n_0)}$  is the second quasi-complement from (40).

Recall that Lemma 3.15 combined with Corollary 3.26 provide that all these functions are well defined, and that each  $\Gamma^{(n_0)}$  is positive and bounded.

It is natural to suppose the initial data to be admissible, but we need a bit more for our subsequent analysis. In order to establish compactness in  $\mathcal{Y}$  and to control the tail behaviour (w.r.t. to n and a) we must assume that the initial data are moreover *regular*.

**Definition 3.27.** A state  $f \in Y$  is called *regular* if the following statements are satisfied.

1. f is equi-continuous with respect to  $\|\cdot\|_{\beta}$ , i.e., for each  $\varepsilon > 0$  there exists  $\overline{\delta} = \overline{\delta}(\varepsilon)$  such that

$$||\mathcal{S}^+_{\delta}f - f||_{\beta} \leq \varepsilon \quad \text{for all} \quad 0 \leq \delta \leq \overline{\delta},$$

where  $\mathcal{S}^+_{\delta}$  is the left-shift operator defined by  $\mathcal{S}^+_{\delta}f(a) = f(a-\delta)$  for  $a \ge \delta$  and  $\mathcal{S}^+_{\delta}f(a) = 0$  else.

2. f is rapidly decreasing w.r.t n and a, that means there exist two constants d and D such that

$$M^{\perp}(f,\alpha) \le D \exp\left(-d\alpha\right)$$

for all  $\alpha \geq 0$ , where

$$M^{\perp}(f,\alpha) := \sum_{n \geq 2} \int_{\alpha}^{\infty} a f_n(a) \, \mathrm{d}a + \sum_{n \geq \lfloor \alpha \rfloor + 1} n \int_{0}^{\infty} f_n(a) \, \mathrm{d}a$$

is defined in line with (40).

We proceed with some remarks concerning the regularity of the initial data g. (i) The first condition is apparently satisfied if g is differentiable w.r.t a with  $||\partial_a g||_{\beta} < \infty$ . (ii) Concerning the second condition recall that  $||g||_{\beta} < \infty$  already implies an exponential decay w.r.t. n. (iii) Compactly supported initial data with  $g_n(a) = 0$  for large n and a are rapidly decreasing.

Assumption 3.28. From now on we suppose the initial data  $g \in Y$  to be admissible and regular.

As a first implication we summarise some consequences of the tightness estimates from §3.5, more precisely of Corollary 3.24 and Lemma 3.25.

**Remark 3.29.** There exist constants d and D which depend only on the initial data g and T but not on  $n_0$  such that for all  $t \in [0, T]$  we have

$$N^{(n_0)}(t) \ge d, \qquad \Gamma^{(n_0)}(t) \le D, \quad M^{\perp, (n_0)}(t, \alpha) \le D \exp(-d\alpha).$$

Our second result provides compactness in the space of bounded and continuous functions.

**Lemma 3.30.** For each n the set  $\{f_n^{(n_0)} : n_0 > n\}$  is equi-continuous in  $BC([0, T] \times [0, \infty))$ .

*Proof.* Within this proof let  $\varepsilon > 0$  be fixed and suppose that  $\delta$  is sufficiently small. We start with the modulus of continuity in *a*-direction as it can be estimated independently of *n*. Since the auxiliary Problem 3.8 is linear in *f* and invariant under shifts w.r.t *a*, we immediately obtain

$$\mathcal{S}_{\delta}^{+}f^{(n_{0})}(t) - f^{(n_{0})}(t) = \mathcal{S}_{\delta}^{+}g^{(n_{0})} - g^{(n_{0})} + \int_{0}^{t} \Gamma^{(n_{0})}(s)\mathcal{T}(t-s)J\Big(\mathcal{S}_{\delta}^{+}f^{(n_{0})}(s) - f^{(n_{0})}(s)\Big) \,\mathrm{d}s,$$

and Corollary 3.13 provides

$$||\mathcal{S}_{\delta}^{+}f^{(n_{0})}(t) - f^{(n_{0})}(t)||_{\beta} \le ||\mathcal{S}_{\delta}^{+}g^{(n_{0})} - g^{(n_{0})}||_{\beta} \le \varepsilon.$$

Therefore, all function  $f_n^{(n_0)}(t)$  are equi-continuous in *a*-direction (uniformly in *n*,  $n_0$  and *t*). Now let *n* be fixed and consider  $n_0 > n$ . The first part of this proof implies

$$\left|\left|\left(\mathcal{T}(\delta)f^{(n_0)}(t)\right)_n - f^{(n_0)}_n(t)\right|\right|_{\infty} \le \varepsilon$$

for all  $0 \leq \delta \leq \overline{\delta}$ . Note that here  $\overline{\delta}$  is expected to depend also on n as the transport velocity increases with n. Moreover, by construction we have

$$f^{(n_0)}(t+\delta) = \mathcal{T}(\delta)f^{(n_0)}(t) + \int_0^\delta \Gamma^{(n_0)}(t+s)\mathcal{T}(\delta-s)Jf^{(n_0)}(t+s)\,\mathrm{d}s,$$

and since  $\Gamma^{(n_0)}(t)$  is uniformly bounded, see Remark 3.29, we infer that

$$||f_n^{(n_0)}(t+\delta) - \mathcal{T}(\delta)f_n^{(n_0)}(t)||_{\infty} \le C \int_0^{\delta} ||(Jf^{(n_0)}(t+s))_n||_{\infty} \,\mathrm{d}s \le C_n\delta,$$

with constant  $C_n$  depending on n. Finally, we have shown that

$$||f_n^{(n_0)}(t+\delta) - f_n^{(n_0)}(t)||_{\infty} \le \epsilon$$

for all sufficiently small  $\delta$ , and the proof is complete.

We are now able to prove our main result, which was already stated in  $\S2$ , Theorem 2.3.

**Lemma 3.31.** There exists a mild solution  $f \in \mathcal{Y}$  to Problem 2.1, which has the following properties.

- 1. f is non-negative with  $||f(t)||_{\beta} \leq ||g||_{\beta}$
- 2. all states f(t) are admissible, so that P(f(t)) = 0,
- 3. all states are regular in the sense of Definition 3.27,
- 4. f conserves the area with non-increasing number of grains.

Proof. Let  $\Omega = [0, T] \times [0, \infty)$ . The Arzela–Ascoli theorem implies that any equi-continuous subset of  $BC(\Omega)$  has a subsequence that converges to a bounded and continuous limit function on  $\Omega$ , and that this convergence is uniform on each compact subset of  $\Omega$ . Thanks to Lemma 3.30 there exists a sequence  $(n_k)_k$  with  $n_k \to \infty$  such that  $f_2^{(n_k)}$  converges to a limit  $f_2$  locally uniform in  $BC(\Omega)$ . Moreover, passing to a suitably chosen subsequence, still denoted by  $(n_k)_k$ , we can assume that  $f_3^{(n_k)} \to f_3$  for some  $f_3 \in BC(\Omega)$ .

Iterating this argument, and using the usual diagonal trick, we finally find a subsequence along with limit functions  $f = (f_2, ..., f_n, ...)$  such that

$$f_n^{(n_k)} \xrightarrow{k \to \infty} f_n$$
 locally uniform in  $BC(\Omega)$ .

for all  $n \ge 2$ . By construction, the limit f is non-negative and satisfies  $||f(t)||_{\beta} \le ||g||_{\beta}$  for all t. The uniform tightness estimates from Remark 3.29 imply the  $L^1$ -regularity of f, and hence  $f \in \mathcal{Y}$ , as well as  $M^{\perp}(f(t)) \le D \exp(-dt)$ . Moreover, since these tightness estimates control the tail behaviour with respect to n and a we infer that

$$N(f^{(n_k)}(t)) \xrightarrow{k \to \infty} N(f(t)), \quad \Gamma(f^{(n_k)}(t)) \xrightarrow{k \to \infty} \Gamma(f(t)),$$

and

$$P(f^{(n_k)}(t)) \xrightarrow{k \to \infty} P(f(t)), \quad A(f^{(n_k)}(t)) \xrightarrow{k \to \infty} N(f(t)).$$

From this we conclude that f is in fact a mild solution to Problem 2.1, and the remaining assertions concerning A(f(t)), N(f(t)) and P(f(t)) follow from the corresponding properties of  $f^{(n_k)}$ .

#### **Remarks on well-posedness**

To establish well-posedness for our kinetic model we must prove both, the uniqueness of solutions and the continuous dependence on the initial data. This can done by means of energy methods, if the 'energy distance' of two states f and  $\tilde{f}$  is defined via

$$\mathcal{E}(f,\tilde{f}) = \sum_{n\geq 2} n \int_0^\infty \exp\left(-a\right) \left(f_n(a) - \tilde{f}_n(a)\right)^2 \mathrm{d}a + \left(N(f) - N(\tilde{f})\right)^2$$

The main result on  $\mathcal{E}$ , which in turn implies uniqueness and continuous dependence for solutions to Problem 2.1, can be stated as follows.

**Lemma 3.32.** For all  $0 < T < \infty$  and any two mild solutions f and  $\tilde{f}$  from Lemma 3.31 we have

$$\mathcal{E}(f(t), \tilde{f}(t)) \le \exp\left(Ct\right) E(g, \tilde{g}) \tag{50}$$

for all  $0 \le t \le T$ , where the constant C depends only on T and the initial data  $g = f(0), \ \tilde{g} = \tilde{f}(0)$ .

The proof of (50) relies on careful estimates for  $\frac{d}{dt}\mathcal{E}(f(t), \tilde{f}(t))$ . Since a concise presentation would involve lengthy computations, we omit the details here and refer the reader to [Hen07].

# Acknowledgments

RH and BN acknowledge support by the Deutsche Forschungsgemeinschaft through the Priority Program 1095 Analysis, Modeling, and Simulation of Multiscale Problems within the project Homogenization of many-particle systems at the Humboldt–Universität zu Berlin. JJLV was supported through the Alexander-von-Humboldt foundation and DGES Grant MTM2007-61755.

# References

[ASGS84a] M. P. Anderson, D. J. Srolovitz, G. S. Grest, and P. Sahni, Computer simulation of grain growth-I. Kinetics, Acta metall. 32 (1984), 783–791.

- [ASGS84b] \_\_\_\_\_, Computer simulation of grain growth-II. Grain size distribution, topology, and local dynamics, Acta metall. **32** (1984), 793–802.
- [BKLT06] K. Barmak, D. Kinderlehrer, I. Livshits, and S. Ta'asan, Remarks on a multiscale approach to grain growth in polycrystals, Variational Problems in Materials Science, Progress in Nonlinear Differential Equations and Their Applications, vol. 68, Birkhäuser Basel, 2006, pp. 1–11.
- [BR93] L. Bronsard and F. Reitich, On three-phase boundary motion and the singular limit of a vector-valued Ginzburg-Landau equation, Arch. Rat. Mech. Anal. **124** (1993), 355–379.
- [Fly93] H. Flyvbjerg, Model for coarsening froths and foams, Phys. Rev. E 47 (1993), 4047–4054.
- [Fra88] V. E. Fradkov, A theoretical investigation of two-dimensional grain growth in the 'gas' approximation, Phil. Mag. Lett. 58 (1988), 271–275.
- [FU94] V. E. Fradkov and D. G. Udler, 2D normal grain growth: Topological aspects, Adv. in Phys. 43 (1994), 739–789.
- [FUK88] V. E. Fradkov, D. G. Udler, and R. E. Kris, Computer simulation of two-dimensional normal grain growth (the 'gas' approximation), Phil. Mag. Lett. 58 (1988), 277–283.
- [Hen07] R. Henseler, A kinetic model for grain growth, Doctoral thesis, Humboldt–Universität zu Berlin, 2007.
- [HNO03] R. Henseler, B. Niethammer, and F. Otto, A reduced model for simulating grain growth, Internat. Ser. Numer. Math. 147 (2003), 177–187.
- [KL01] D. Kinderlehrer and C. Liu, Evolution of grain boundaries, Math. Mod. Meth. Appl. Sc. 11 (2001), 713–729.
- [KNN89] K. Kawasaki, T. Nagai, and K. Nakashima, Vertex models for two-dimensional grain growth, Phil. Mag. B 60 (1989), 399–421.
- [Lew43] F. T. Lewis, The geometry of growth and cell division in epithelial mosaics, Am. J. Bot. 30 (1943), 766–776.
- [Mar87] M. Marder, Soap-bubble growth, Phys. Rev. A 36 (1987), 438–440.
- [MNT04] C. Mantegazza, M. Novaga, and V. M. Tortorelli, *Motion by curvature of planar networks*, Ann. Sc. Norm. Super. Pisa Cl. Sci. **3** (2004), 235–324.
- [Mul56] W. W. Mullins, Two-dimensional motion of idealized grain boundaries, J. Appl. Phys. 27 (1956), 900–904.
- [RL82] N. Rivier and A. Lissowski, On the correlation between sizes and shapes of cells in epithelial mosaics, J. Phys. A: Math. Gen. 15 (1982), 143–148.
- [TC94] J. E. Taylor and J. W. Cahn, Linking anisotropic sharp and diffusive surface motion laws via gradient flows, J. Stat. Phys. 77 (1994), 183–197.
- [Tho01] C. V. Thompson, Grain growth and evolution of other cellular structures, Solid State Physics 55 (2001), 269–314.