Invariant expansion for the trigonal band structure of graphene

R. Winkler

Department of Physics, Northern Illinois University, DeKalb, Illinois 60115, USA and Materials Science Division, Argonne National Laboratory, Argonne, Illinois 60439, USA

U. Zülicke

Institute of Fundamental Sciences and MacDiarmid Institute for Advanced Materials and Nanotechnology,

Massey University, Manawatu Campus, Private Bag 11 222, Palmerston North, New Zealand and

Centre for Theoretical Chemistry and Physics, Massey University, Albany Campus,

Private Bag 102904, North Shore MSC, Auckland 0745, New Zealand

(Dated: 14 December 2010)

We present a symmetry analysis of the trigonal band structure in graphene, elucidating the transformational properties of the underlying basis functions and the crucial role of time-reversal invariance. Group theory is used to derive an invariant expansion of the Hamiltonian for electron states near the **K** points of the graphene Brillouin zone. Besides yielding the characteristic k-linear dispersion and higher-oder corrections to it, this approach enables the systematic incorporation of all terms arising from external electric and magnetic fields, strain, and spin-orbit coupling up to any desired order. Several new contributions are found, in addition to reproducing results obtained previously within tight-binding calculations. Physical ramifications of these new terms are discussed.

I. INTRODUCTION

In recent years, tremendous interest has been focused on studying single layers of graphene, 1-3 following the first experimental realization of this material.^{4,5} To a large extent, these continuing efforts are motivated by the unique band structure of graphene in the vicinity of the Fermi edge. In most semiconductors, the band edges are characterized by a quadratic dispersion, with k-linear corrections possible only in inversion-asymmetric materials due to spin-orbit (SO) coupling.⁶ In contrast, for graphene, the dispersion $E(\mathbf{k})$ of the uppermost valence band and the lowest conduction band is *dominated* by klinear terms, ${}^7 E(\mathbf{k}) \approx \hbar v k$ with Fermi velocity v. These bands touch at the points \mathbf{K} and \mathbf{K}' at the edge of the Brillouin zone [Fig. 1(b)] so that the resulting energy surfaces resemble those of free massless fermions described by the Dirac equation [Fig. 1(c)]. The apparent analogies between a solid-state system and relativistic quantum mechanics have greatly stimulated the interest in graphene.^{1-3,8-10}

Both in early work⁷ as well as in more recent pu-



FIG. 1: (Color online) (a) Honeycomb structure of graphene. Atoms in sublattice A (B) are marked with open (closed) circles. (b) Brillouin zone and its two inequivalent corner points \mathbf{K} and \mathbf{K}' . The remaining corners are related with \mathbf{K} or \mathbf{K}' by reciprocal lattice vectors. (c) Dispersion E(k) near the \mathbf{K} point. We have $\mathbf{k} \equiv \boldsymbol{\kappa} - \mathbf{K}$.

blications $^{3,8-11}$ the electronic properties of graphene have largely been explored using tight-binding calculations. Tight-binding models also provide the usual starting point for the derivation of simplified, effective Hamiltonians^{3,9} to describe the *k*-linear dispersion in the vicinity of the Fermi energy [Fig. 1(c)]. Alternatively, first-principles methods have also been applied to graphene.¹² The importance of group theory for the characterization of the graphene band structure was already recognized in early work.^{13–17} Here we employ a group theoretical approach to graphene⁶ that was pioneered for the study of bulk semiconductors¹⁸ and. more recently, has proved to be useful also for the systematic investigation of band-structure effects in lowdimensional systems.¹⁹ Using the theory of invariants⁶ we find the Hamiltonian describing electronic degrees of freedom near the \mathbf{K} and \mathbf{K}' points in graphene in terms of a systematic expansion in orders of wave-vector difference $\mathbf{k} \ (\equiv \boldsymbol{\kappa} - \mathbf{K} \ \text{or} \ \boldsymbol{\kappa} - \mathbf{K}', \text{ respectively})$ from these special points and various external perturbations. Both the sublattice-related (orbital) pseudospin and the intrinsic spin of quasiparticles in graphene are accounted for within this scheme, and the fundamental difference in the origin of these two degrees of freedom is reflected in the transformational properties of the Bloch-state basis functions. In principle, our analysis can be used to construct all allowed terms in the Hamiltonian up to any desired order; but we limit the present discussion to all contributions up to second order as well as selected terms up to third order. Several new terms are found that have not been obtained previously.

The important role played by discrete symmetries in protecting a k-linear dispersion in graphene was pointed out in Ref. 20. Here we discuss in detail the special significance of time-reversal invariance for the trigonal band structure of graphene. It provides an additional criterion that is satisfied only by a subset of all terms allowed by the spatial symmetry of the \mathbf{K} - and \mathbf{K}' -point Bloch functions.

The remainder of this article is organized as follows. To introduce our phase conventions and other relevant background information, the following Sec. II provides a summary of the tight-binding description for a graphene sheet. The symmetry analysis for this system is performed in Sec. III, with the role of time-reversal invariance highlighted. A discussion of the terms obtained within the symmetry analysis is given in Sec. IV before we present our conclusions in Sec. V.

II. REVIEW OF TIGHT-BINDING ANALYSIS

The honeycomb structure of graphene is sketched in Fig. 1(a). For definiteness, we use the basis vectors in real space

$$\mathbf{a}_1 = a \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \qquad \mathbf{a}_2 = a \begin{pmatrix} 1/2 \\ \sqrt{3}/2 \end{pmatrix}, \qquad (1)$$

with lattice constant a. The basis vectors in reciprocal space become

$$\mathbf{b}_1 = \frac{2\pi}{a} \begin{pmatrix} 1\\ -1/\sqrt{3} \end{pmatrix}, \qquad \mathbf{b}_2 = \frac{2\pi}{a} \begin{pmatrix} 0\\ 2/\sqrt{3} \end{pmatrix}, \qquad (2)$$

and the two inequivalent corner points of the Brillouin zone are

$$\mathbf{K} = \frac{2\pi}{a} \begin{pmatrix} 2/3\\ 0 \end{pmatrix}, \quad \mathbf{K}' = \frac{2\pi}{a} \begin{pmatrix} -2/3\\ 0 \end{pmatrix}. \tag{3}$$

We consider a tight-binding Hamiltonian for the graphene π bonds formed by the carbon p_z orbitals, taking into account nearest-neighbor and second-nearest-neighbor interactions. For a given atom in the honey-comb structure, the vectors connecting nearest-neighbor atoms are (j = 1, 2, 3)

$$\boldsymbol{\tau}_{1}^{(j)} = \mathcal{R}(2j\pi/3)\,\boldsymbol{\tau}_{1}^{(3)}, \quad \boldsymbol{\tau}_{1}^{(3)} = \begin{pmatrix} a\\ 1/\sqrt{3} \end{pmatrix}, \quad (4)$$

where $\mathcal{R}(\phi)$ denotes a two-dimensional (2D) rotation by the angle ϕ . Similarly, we get the vectors connecting second-nearest-neighbor atoms (j = 1, ..., 6)

$$\boldsymbol{\tau}_2^{(j)} = \mathcal{R}(j \, \pi/3) \, \mathbf{a}_1. \tag{5}$$

Then the tight-binding Hamiltonian becomes^{7,11}

$$\tilde{\mathcal{H}}(\boldsymbol{\kappa}) = \begin{pmatrix} \tilde{\epsilon}_{2p} + t_2 f_2(\boldsymbol{\kappa}) & t_1 f_1(\boldsymbol{\kappa}) \\ t_1 f_1^*(\boldsymbol{\kappa}) & \tilde{\epsilon}_{2p} + f_2(\boldsymbol{\kappa}) \end{pmatrix}, \qquad (6)$$

where $\tilde{\epsilon}_{2p}$ is the site energy of the p_z orbitals, t_l are the transfer integrals for *l*th-nearest neighbors, and the functions $f_l(\boldsymbol{\kappa})$ are given by

$$f_l(\boldsymbol{\kappa}) \equiv \sum_j e^{i\boldsymbol{\kappa}\cdot\boldsymbol{\tau}_l^{(j)}}.$$
(7)

The particular geometry (4) gives for $f_1(\boldsymbol{\kappa})$

$$f_1(\boldsymbol{\kappa}) = e^{i\kappa_y a/\sqrt{3}} + 2e^{-i\kappa_y a/2\sqrt{3}}\cos(\kappa_x a/2), \quad (8)$$

and we have the relation

$$f_2(\kappa) = |f_1(\kappa)|^2 - 3.$$
 (9)

Thus it is possible to rewrite the Hamiltonian (6) such that it only depends on the function $f \equiv f_1$,

$$\mathcal{H}(\boldsymbol{\kappa}) = \begin{pmatrix} \epsilon_{2p} + t_2 |f|^2 & t_1 f \\ t_1 f^* & \epsilon_{2p} + t_2 |f|^2 \end{pmatrix}, \qquad (10)$$

where $\epsilon_{2p} = \tilde{\epsilon}_{2p} - 3t_2$. We note in passing that, alternatively, we could have considered a tight-binding model that takes into account only nearest-neighbor transfer integrals as well as nearest-neighbor overlap integrals.¹¹ This model gives qualitatively similar results as the Hamiltonian (10). However, extensions of this alternative tight-binding model, e.g., to include spin-orbit effects, are hindered by the fact that the matrix of overlap integrals results in a generalized eigenvalue problem.¹¹

The tight-binding wave functions corresponding to the Hamiltonian (10) are given by

$$\Psi_{\kappa n}(\mathbf{r}) = \sum_{\lambda = A, B} \psi_{\lambda n}(\kappa) \, \Phi_{\kappa \lambda}(\mathbf{r}), \qquad (11)$$

where $\psi_{\lambda n}(\boldsymbol{\kappa})$ are $\boldsymbol{\kappa}$ -dependent expansion coefficients for band n, and the corresponding basis functions (that are Bloch functions) are

$$\Phi_{\kappa\lambda}(\mathbf{r}) = \frac{e^{i\kappa\cdot\mathbf{r}}}{\sqrt{N}} \sum_{\mathbf{R}_{\lambda}} e^{-i\kappa\cdot(\mathbf{r}-\mathbf{R}_{\lambda})} \phi_{\pi}(\mathbf{r}-\mathbf{R}_{\lambda}).$$
(12)

Here $\phi_{\pi}(\mathbf{r})$ denote the carbon π orbitals, and the sum runs over the atomic positions \mathbf{R}_{λ} in sublattice λ [see Fig. 1(a)].

Diagonalization of the Hamiltonian (10) yields the energy dispersion $(n = \pm)$

$$E_n(\boldsymbol{\kappa}) = \epsilon_{2p} + n t_1 |f(\boldsymbol{\kappa})| + t_2 |f(\boldsymbol{\kappa})|^2, \qquad (13)$$

with eigenfunctions $\psi_n(\boldsymbol{\kappa}) = (\psi_{An}, \psi_{Bn})$ given by the expressions (valid for $\boldsymbol{\kappa} \neq \mathbf{K}, \mathbf{K}'$)

$$\psi_{+}(\boldsymbol{\kappa}) = \frac{1}{\sqrt{2}} \begin{pmatrix} 1\\ \hat{f}^{*}(\boldsymbol{\kappa}) \end{pmatrix}, \quad \psi_{-}(\boldsymbol{\kappa}) = \frac{1}{\sqrt{2}} \begin{pmatrix} -\hat{f}(\boldsymbol{\kappa})\\ 1 \end{pmatrix},$$
(14)

where $\hat{f}(\boldsymbol{\kappa}) \equiv f/|f|$. For $t_2 = 0$, the spectrum (13) is symmetric around ϵ_{2p} , but this electron-hole symmetry is broken when $t_2 \neq 0$.

The two bands \pm touch at the **K** and **K'** points of the Brillouin zone where f = 0. We can expand $\mathcal{H}(\boldsymbol{\kappa})$ around **K** ($\boldsymbol{\kappa} \equiv \mathbf{K} + \mathbf{k}$), which yields up to second order in k

$$\mathcal{H}_{55}^{\mathbf{K}}(\mathbf{k}) = a_{10}^{55} \mathbb{1} + a_{61}^{55} (k_x \sigma_x + k_y \sigma_y) + a_{11}^{55} (k_x^2 + k_y^2) \mathbb{1} \\ + a_{62}^{55} [(k_y^2 - k_x^2) \sigma_x + 2k_x k_y \sigma_y] + \mathcal{O}(k^3) , (15)$$

with Pauli matrices σ_i and

$$a_{10}^{55} = \epsilon_{2p},$$
 (16a)

$$a_{61}^{55} = -\frac{\sqrt{3}}{2}at_1, \tag{16b}$$

$$a_{11}^{55} = \frac{3}{4}t_2a^2, \tag{16c}$$

$$a_{62}^{55} = -\frac{1}{8}t_1a^2.$$
 (16d)

Our notation will become clear in Sec. III. The dispersion becomes

$$E_{\pm}(\mathbf{k}) = a_{10}^{55} \pm a_{61}^{55}k + a_{11}^{55}k^2 \pm a_{62}^{55}k_x (3k_y^2 - k_x^2)/k + \mathcal{O}(k^3).$$
(17)

It follows immediately from Eq. (14) that the corresponding basis functions describing the two-fold degeneracy at the **K** point can be chosen to be nonzero only on sublattice $\lambda = A$ or B,

$$\Psi_{\mathbf{K}\lambda}(\mathbf{r}) = \Phi_{\mathbf{K}\lambda}(\mathbf{r}) = \frac{e^{i\mathbf{K}\cdot\mathbf{r}}}{\sqrt{N}} \sum_{\mathbf{R}_{\lambda}} e^{-i\mathbf{K}\cdot(\mathbf{r}-\mathbf{R}_{\lambda})}\phi_{\pi}(\mathbf{r}-\mathbf{R}_{\lambda}).$$
(18)

We obtain the Hamiltonian $\mathcal{H}_{55}^{\mathbf{K}'}(\mathbf{k})$ for \mathbf{K}' from $\mathcal{H}_{55}^{\mathbf{K}}(\mathbf{k})$ by applying any symmetry element R in the point group D_{6h} of the honeycomb structure that maps **K** onto **K**',

$$\mathcal{H}_{55}^{\mathbf{K}'}(\mathbf{k}) = \mathcal{D}(R) \,\mathcal{H}_{55}^{\mathbf{K}}(R^{-1}\mathbf{k}) \,\mathcal{D}^{-1}(R).$$
(19)

Here the matrices $\mathcal{D}(R)$ map the basis functions at **K** on the basis functions at \mathbf{K}'

$$\Psi_{\mathbf{K}'\lambda}(\mathbf{r}) = \sum_{\lambda'} \mathcal{D}_{\lambda\lambda'}(R) \Psi_{\mathbf{K}\lambda'}(R^{-1}\mathbf{r}).$$
(20)

Choosing for \mathbf{K}' the basis functions $\Psi_{\mathbf{K}'\lambda}(\mathbf{r}) = \Phi_{\mathbf{K}'\lambda}(\mathbf{r})$ and assuming that, for both \mathbf{K} and \mathbf{K}' , these basis functions are ordered as (A, B), this transformation becomes particularly simple if we consider the reflection $R = R_{\mu}$ at a perpendicular plane that includes the y axis (see Fig. 1). The reflection R_{y} preserves the sublattices λ , i.e., $\mathcal{D}(R_{\eta}) = \mathbb{1}$. So we obtain

$$\mathcal{H}_{55}^{\mathbf{K}'}(\mathbf{k}) = \mathcal{H}_{55}^{\mathbf{K}}(R_y^{-1}\mathbf{k}), \qquad (21)$$

where $R_y^{-1}(k_x, k_y) = (-k_x, k_y)$. SO coupling in the ideal material can be described by adding a second-nearest-neighbor term²¹⁻²³

$$\mathcal{H}_{\rm so}(\boldsymbol{\kappa}) = \begin{pmatrix} h_{\rm so} & 0\\ 0 & -h_{\rm so} \end{pmatrix}, \qquad (22a)$$

where

$$h_{\rm so} = i\lambda_{\rm so} \sum_{j,k} \mathbf{s} \cdot (\boldsymbol{\tau}_1^{(j)} \times \boldsymbol{\tau}_1^{(k)}) e^{i\boldsymbol{\kappa} \cdot (\boldsymbol{\tau}_1^{(j)} - \boldsymbol{\tau}_1^{(k)})} .$$
(22b)

Here \mathbf{s} denotes the operator for spin angular momentum. Similarly, the tight-binding model (6) can be extended²⁴ to include SO coupling due to the gradient $e\boldsymbol{\mathcal{E}} \equiv \nabla V$ of an external potential V by adding a term

$$\mathcal{H}_R(\boldsymbol{\kappa}) = \begin{pmatrix} 0 & h_R \\ h_R^* & 0 \end{pmatrix}, \qquad (23a)$$

where

$$h_R = \lambda_R \mathbf{s} \cdot \boldsymbol{\mathcal{E}} \times \sum_j i \boldsymbol{\tau}_j e^{i \boldsymbol{\kappa} \cdot \boldsymbol{\tau}_j} .$$
 (23b)

To lowest order in **k** and \mathcal{E} , the following additional terms beyond Eq. (15) are obtained:

$$p_{21}^{55} s_z \sigma_z + r_{61}^{55} s_z (\mathcal{E}_y \sigma_x - \mathcal{E}_x \sigma_y) + r_{62}^{55} \mathcal{E}_z (s_y \sigma_x - s_x \sigma_y) + r_{63}^{55} s_z [(k_x \mathcal{E}_y + k_y \mathcal{E}_x) \sigma_x + (k_x \mathcal{E}_x - k_y \mathcal{E}_y) \sigma_y] + r_{64}^{55} \mathcal{E}_z [(s_x k_y + s_y k_x) \sigma_x + (s_x k_x - s_y k_y) \sigma_y], \quad (24)$$

where

$$p_{21}^{55} = \frac{3}{2}\lambda_{so}a^2, \tag{25a}$$

$$r_{61}^{55} = -\frac{\sqrt{3}}{2}\lambda_R a,$$
 (25b)

$$r_{62}^{55} = \frac{\sqrt{3}}{2}\lambda_R a, \qquad (25c)$$

$$r_{63}^{55} = \frac{1}{4}\lambda_R a^2, \tag{25d}$$

$$r_{64}^{55} = -\frac{1}{4}\lambda_R a^2.$$
 (25e)

While the above approach can be further extended in various ways, it becomes difficult to explore all possible terms in a systematic manner. For example, we will show below that the expansion (24) misses SO coupling terms of the form

$$r_{11}^{55}(s_xk_y - s_yk_x)\mathcal{E}_z 1 + r_{12}^{55}s_z(k_x\mathcal{E}_y - k_y\mathcal{E}_x)1 , \quad (26)$$

which illustrates the fact that qualitatively new terms may appear if a more complete model beyond Eq. (23) is used to describe Rashba spin-orbit coupling in graphene. From a different perspective, we note that SO coupling, while small for systems made of light atoms like carbon, can be expected to be underestimated by the approach outlined above that expresses all properties in terms of interatomic matrix elements. In general, the dominant effect of the SO interaction is a coupling of the p orbitals on the same atom,²⁵ which can be expected to contribute also to SO coupling in graphene, see Sec. IVC. (As usual,³ the present approach has neglected the p_x and p_u orbitals.)

III. SYMMETRY ANALYSIS

Invariant Expansion Α.

While the power expansion (15) of the tight-binding Hamiltonian (10) can be readily extended to arbitrary orders of the wave vector \mathbf{k} , it became noticeable in the above discussion that it gets more difficult within this approach to incorporate the effects of perturbations such as external electric and magnetic fields, strain, or spin-orbit coupling. In contrast, the theory of invariants⁶ enables systematic construction of an invariant expansion for the

effective Hamiltonian $\mathcal{H}(\mathcal{K})$ describing the electron states in the vicinity of \mathbf{K} and \mathbf{K}' (or indeed any other point in the Brillouin zone of any crystalline material). Here \mathcal{K} represents a general tensor operator, which can depend, e.g., on the components of the kinetic wave vector \mathbf{k} , on external electric and magnetic fields \mathcal{E} and \mathbf{B} , on strain ϵ , and on the intrinsic spin \mathbf{s} of the electrons. Keeping in mind that we want to extend the discussion from the spinless Hamiltonian (15) to a Hamiltonian that includes spin, we first present a brief review of the general theory⁶ before applying the formalism to our particular questions of interest.

The Hamiltonian $\mathcal{H}(\mathcal{K})$ is defined relative to a set of basis functions $\Phi_{\mu}(\mathbf{r})$ transforming according to a (reducible or irreducible) representation Γ of the point group \mathcal{G} . Denoting the transformation matrices by $\mathcal{D}(g)$ $(g \in \mathcal{G})$, we have

$$\Phi_{\mu}(g^{-1}\mathbf{r}) = \sum_{\mu'} \mathcal{D}_{\mu\mu'}(g) \Phi_{\mu'}(\mathbf{r}).$$
(27)

Then the invariance of $\mathcal{H}(\mathcal{K})$ under the symmetry elements $g \in \mathcal{G}$ implies

$$\mathcal{D}(g) \mathcal{H}(g^{-1}\mathcal{K}) \mathcal{D}^{-1}(g) = \mathcal{H}(\mathcal{K}).$$
(28)

As shown in Ref. 6, Eq. (28) can be used to construct $\mathcal{H}(\mathcal{K})$. This is greatly simplified if we choose basis functions $\Phi_{\mu}(\mathbf{r})$ transforming according to *irreducible* representations (IRs) of \mathcal{G} . Then $\mathcal{H}(\mathcal{K})$ can be decomposed into blocks $\mathcal{H}_{\alpha\beta}(\mathcal{K})$, where α and β denote the spaces of the n_{α} - and n_{β} -fold degenerate basis functions, which transform according to the IRs Γ_{α} and Γ_{β} of \mathcal{G} . For each block $\mathcal{H}_{\alpha\beta}(\mathcal{K})$, one can find a complete set of linearly independent $n_{\alpha} \times n_{\beta}$ -dimensional matrices $X_l^{(\kappa)}$ that transform according to the IRs Γ_{κ} (of dimension L_{κ}) contained in the product representation $\Gamma_{\alpha} \times \Gamma_{\beta}^{*}$. Likewise, \mathcal{K} can be decomposed into irreducible tensor operators $\mathcal{K}^{(\kappa',\lambda)}$ that transform according to the IRs $\Gamma_{\kappa'}$ of \mathcal{G} (Ref. 26). Then each block $\mathcal{H}_{\alpha\beta}(\mathcal{K})$ can be written as

$$\mathcal{H}_{\alpha\beta}(\mathcal{K}) = \sum_{\kappa,\lambda} a_{\kappa\lambda}^{\alpha\beta} \sum_{l=1}^{L_{\kappa}} X_l^{(\kappa)} \mathcal{K}_l^{(\kappa,\lambda)*}, \qquad (29)$$

with material-specific coefficients $a_{\kappa\lambda}^{\alpha\beta}$. By construction, each block $\mathcal{H}_{\alpha\beta}(\mathcal{K})$ is invariant under the symmetry operations in \mathcal{G} in the sense of Eq. (28).

To proceed, we need to identify the symmetry of the eigenfunctions (18) at **K** and **K'** that form the basis functions for $\mathcal{H}(\mathcal{K})$. The group of the wave vector **K** is isomorphic to the trigonal point group $\mathcal{G} = D_{3h}$ (while the point group of the honeycomb structure is D_{6h}). Projection of $\Psi_{\mathbf{K}\lambda}$ on the IRs of D_{3h} shows^{6,15} that these functions transform according to the two-dimensional IR Γ_5 .^{27,28} More specifically, under the symmetry operations of D_{3h} , the Bloch function $\Psi_{\mathbf{K},A(B)}(\mathbf{r})$ transforms like $|\rho_{-(+)}\rangle$, where $|\rho_{\mp}\rangle \equiv \frac{1}{\sqrt{2}} |\tilde{x} \mp i\tilde{y}\rangle$. Here $|\tilde{x}\rangle$ and $|\tilde{y}\rangle$ transform like the coordinate functions x and y except that

they do not change sign under inversion. Thus, from a symmetry point of view, we may identify $\{\Psi_{\mathbf{K}\lambda}(\mathbf{r})\}$ with $\{|\rho_{\mp}\rangle\}$. Note that $\Psi_{\mathbf{K}'\lambda} = \Psi^*_{\mathbf{K}\lambda}$ [i.e., time reversal does not cause extra degeneracies, see Eq. (35) below]. The role of $|\rho_{\mp}\rangle$ is thus reversed at the \mathbf{K}' point: $|\rho_{-}\rangle$ ($|\rho_{+}\rangle$) corresponds to sublattice B (A).

The TB model (10) neglects the spin degree of freedom. Thus we have obtained an ordinary IR (Γ_5 of D_{3h}), which differs qualitatively from the double-group (spinor) IRs characterizing a particle with a genuine spin-1/2 degree of freedom coupled to a particle's orbital motion. The double group \mathcal{G}_d corresponding to a group \mathcal{G} can be written as $\mathcal{G}_d = \mathcal{G} \oplus \overline{E}\mathcal{G}$, where \overline{E} is a rotation by 2π around an arbitrary axis. The presence of \overline{E} in \mathcal{G}_d reflects the well-known fact²⁹ that spin-1/2 spinors (which may be used²⁷ as basis functions for the spinor IR Γ_7 of D_{3h}) change sign when rotated by 2π . The basis functions of Γ_5 do not change sign when rotated by 2π , i.e., \bar{E} acts like the neutral element E. In principle, this could be studied experimentally in a setup similar to the neutron interference experiments in Refs. 30.31. However, such an experiment would obviously be complicated by the fact that the electrons in graphene also carry a real spin degree of freedom that is neglected in the present discussion. (See, however, Sec. IV C below.)

Choosing basis functions for Γ_5 that transform like $\{|\rho_-\rangle, |\rho_+\rangle\}$, we obtain the basis matrices listed in Table I. Similarly, \mathcal{K} can be decomposed into irreducible tensor operators $\mathcal{K}^{(\kappa,\lambda)}$ that transform according to the IRs Γ_{κ} of D_{3h} . Using the coordinate system in Fig. 1(b), we get the lowest-order tensor operators $\mathcal{K}^{(\kappa,\lambda)}$ in Table II. We note that given two irreducible tensor operators $\mathcal{K}^{(\alpha,\lambda_1)}$ and $\mathcal{K}^{(\beta,\lambda_2)}$ transforming according to Γ_{α} and Γ_{β} , we get new higher-order tensor operators transforming according to the IRs Γ_{κ} using the relation

$$\mathcal{K}_{l}^{(\kappa,\lambda)} = \sum_{l_{1},l_{2}} C_{l_{1}l_{2},l}^{\alpha\beta,\kappa} \mathcal{K}_{l_{1}}^{(\alpha,\lambda_{1})} \mathcal{K}_{l_{2}}^{(\beta,\lambda_{2})}, \qquad (30)$$

where $C_{l_1 l_2, l}^{\alpha \beta, \kappa}$ denote the coupling coefficients for D_{3h} .²⁸ We emphasize that this approach guarantees that one obtains all irreducible tensor operators up to a desired order. However, the definition of these tensor operators is not unique because for two tensor operators $\mathcal{K}^{(\kappa,\lambda_1)}$ and $\mathcal{K}^{(\kappa,\lambda_2)}$ which both transform according to Γ_{κ} their linear combination likewise transforms according to Γ_{κ} .

Combining the basis matrices and tensor operators according to Eq. (29), we exactly reproduce Eq. (15). Third-order terms can be read off from Tables I and II. Terms of yet higher orders can be constructed using Eq. (30). For \mathbf{K}' , we obtain in generalization of Eq. (21)

$$\mathcal{H}_{55}^{\mathbf{K}'}(\mathbf{\mathcal{K}}) = \mathcal{H}_{55}^{\mathbf{K}}(R_y^{-1}\mathbf{\mathcal{K}}).$$
(31)

The part of $\mathcal{H}_{55}^{\mathbf{K}}$ linear in **k** is formally equivalent to the Dirac Hamiltonian for massless (chiral) fermions.³² Also, it is related via a simple unitary transformation with the Dresselhaus³³ and the Rashba³⁴ terms in quasi-2D systems.¹⁹ All these models give rise to a dispersion

TABLE I: Symmetrized matrices for the invariant expansion of the blocks $\mathcal{H}_{\alpha\beta}$ for the point group D_{3h} .

Block	Representations	Symn	Symmetrized matrices	
\mathcal{H}_{55}	$\Gamma_5 \times \Gamma_5^*$	Γ_1 :	1	
	$= \Gamma_1 + \Gamma_2 + \Gamma_6$	Γ_2 :	σ_z	
		Γ_6 :	σ_x, σ_y	
\mathcal{H}_{77}	$\Gamma_7 \times \Gamma_7^*$	Γ_1 :	1	
	$=\Gamma_1+\Gamma_2+\Gamma_5$	Γ_2 :	σ_z	
		Γ_5 :	$\sigma_x, -\sigma_y$	
\mathcal{H}_{99}	$\Gamma_9 imes \Gamma_9^*$	Γ_1 :	1	
	$=\Gamma_1+\Gamma_2+\Gamma_3+\Gamma_4$	Γ_2 :	σ_z	
		Γ_3 :	σ_x	
		Γ_4 :	σ_y	
\mathcal{H}_{79}	$\Gamma_7 \times \Gamma_9^*$	Γ_5 :	$1, -i\sigma_z$	
	$=\Gamma_5+\Gamma_6$	Γ_6 :	σ_x,σ_y	

that is linear in the limit of small \mathbf{k} . Yet for each of these models, the transformational properties of the basis functions under the corresponding symmetry operations are qualitatively different.³⁵ The Dirac equation reflects the continuous symmetries of the Lorentz group.³⁶ In quasi-2D systems, the k-linear Dresselhaus term is intimately related with the tetrahedral symmetry of the zinc blende structure,³³ whereas the Rashba term emerges from a model with axial¹⁹ (or hexagonal³⁷) symmetry. These terms refer to electron states transforming according to spinor representations of the corresponding crystallographic point groups, and they are nonzero only as a consequence of spin-orbit coupling.¹⁹ In contrast, the Hamiltonian $\mathcal{H}_{55}^{\mathbf{K}}$ refers to the basis functions (18). It is applicable to spinless particles or particles for which the spin degree of freedom is decoupled from the orbital motion, and the group element \overline{E} does not play a role.³⁸ From a more general perspective, we see here that the symmetries of a system determine the invariant expansion (29) and the band structure $E_n(\mathbf{k})$. Yet it is, in general, not possible to follow the opposite path and infer the symmetries of a system from the Hamiltonian and the band structure $E_n(\mathbf{k})$.

B. Implications of Time Reversal Invariance

Time-reversal invariance results in additional constraints for the allowed terms in the invariant expansion (29). Time reversal corresponds to complex conjugation. Focusing here on the important case (usually⁶ denoted "case a") that we have a linear relation between the complex-conjugate basis functions Φ_{λ}^* and the original basis functions Φ_{λ} ,

$$\Phi_{\lambda}^{*}(\mathbf{r}) = \sum_{\lambda'} \mathcal{T}_{\lambda\lambda'} \Phi_{\lambda'}(\mathbf{r}), \qquad (32)$$

TABLE II: Irreducible tensor components for the point group D_{3h} . Those printed in bold give rise to invariants in $\mathcal{H}_{55}^{\mathbf{K}}(\mathcal{K})$ allowed by time-reversal invariance. (Terms proportional to k_z or ϵ_{jz} are not listed as they are irrelevant for graphene.) Notation: $\{A, B\} \equiv \frac{1}{2}(AB + BA)$.

Γ_1	1; $k_x^2 + k_y^2$; $\{k_x, 3k_y^2 - k_x^2\}$; $k_x \mathcal{E}_x + k_y \mathcal{E}_y$;
	$\epsilon_{xx}+\epsilon_{yy};\;(\epsilon_{yy}-\epsilon_{xx})k_x+2\epsilon_{xy}k_y;$
	$(\epsilon_{yy} - \epsilon_{xx})\mathcal{E}_x + 2\epsilon_{xy}\mathcal{E}_y; s_x B_x + s_y B_y; s_z B_z;$
	$(s_xk_y - s_yk_x)\mathcal{E}_z; \ s_z(k_x\mathcal{E}_y - k_y\mathcal{E}_x);$
Γ_2	$\{k_y, 3k_x^2 - k_y^2\}; B_z; k_x \mathcal{E}_y - k_y \mathcal{E}_x;$
	$(\epsilon_{xx} - \epsilon_{yy})k_y + 2\epsilon_{xy}k_y; (\epsilon_{xx} + \epsilon_{yy})B_z;$
	$(\epsilon_{xx} - \epsilon_{yy})\mathcal{E}_y + 2\epsilon_{xy}\mathcal{E}_x; \ s_z; \ s_x B_y - s_y B_x;$
	$(s_xk_x+s_yk_y)\mathcal{E}_z;\ s_z(\epsilon_{xx}+\epsilon_{yy});$
Γ_3	$B_x k_x + B_y k_y; \ \mathcal{E}_x B_x + \mathcal{E}_y B_y; \ \mathcal{E}_z B_z;$
	$(\epsilon_{yy} - \epsilon_{xx})B_x + 2\epsilon_{xy}B_y; s_xk_x + s_yk_y;$
	$s_x \mathcal{E}_x + s_y \mathcal{E}_y; \ s_z \mathcal{E}_z; \ s_x (\epsilon_{yy} - \epsilon_{xx}) + 2s_y \epsilon_{xy}$
Γ_4	$B_x k_y - B_y k_x; \ \mathcal{E}_z; \ \mathcal{E}_x B_y - \mathcal{E}_y B_x;$
	$(\epsilon_{xx} - \epsilon_{yy})B_y + 2\epsilon_{xy}B_x; \ (\epsilon_{xx} + \epsilon_{yy})\mathcal{E}_z;$
	$s_x k_y - s_y k_x; \ s_x \mathcal{E}_y - s_y \mathcal{E}_x; \ s_y (\epsilon_{xx} - \epsilon_{yy}) + 2s_x \epsilon_{xy}$
Γ_5	$B_x, B_y; B_y k_y - B_x k_x, B_x k_y + B_y k_x; k_y \mathcal{E}_z, -k_x \mathcal{E}_z;$
	$\mathcal{E}_{y}B_{y} - \mathcal{E}_{x}B_{x}, \mathcal{E}_{y}B_{x} + \mathcal{E}_{x}B_{y}; \ (\epsilon_{xx} + \epsilon_{yy})(B_{x}, B_{y});$
	$(\epsilon_{xx} - \epsilon_{yy})B_x + 2\epsilon_{xy}B_y, (\epsilon_{yy} - \epsilon_{xx})B_y + 2\epsilon_{xy}B_x;$
	$2\epsilon_{xy}\mathcal{E}_z, (\epsilon_{xx}-\epsilon_{yy})\mathcal{E}_z; s_x, s_y;$
	$s_yk_y - s_xk_x, s_xk_y + s_yk_x; s_yB_z, -s_xB_z;$
	$s_z B_y, -s_z B_x; \ s_y \mathcal{E}_y - s_x \mathcal{E}_x, s_x \mathcal{E}_y + s_y \mathcal{E}_x;$
	$(s_x, s_y)(\epsilon_{xx} + \epsilon_{yy});$
	$s_x(\epsilon_{xx} - \epsilon_{yy}) - 2s_y\epsilon_{xy}, s_y(\epsilon_{yy} - \epsilon_{xx}) - 2s_x\epsilon_{xy}$
Γ_6	$k_x, k_y; \; \{k_y + k_x, k_y - k_x\}, 2\{k_x, k_y\};$
	$\{k_x, k_x^2 + k_y^2\}, \{k_y, k_x^2 + k_y^2\}; B_z k_y, -B_z k_x;$
	$\mathcal{E}_x, \mathcal{E}_y; \ k_y \mathcal{E}_y - k_x \mathcal{E}_x, k_x \mathcal{E}_y + k_y \mathcal{E}_x;$
	$\mathcal{E}_{y}B_{z}, -\mathcal{E}_{x}B_{z}; \ \mathcal{E}_{z}B_{y}, -\mathcal{E}_{z}B_{x};$
	$\epsilon_{yy}-\epsilon_{xx}, 2\epsilon_{xy}; \ (\epsilon_{xx}+\epsilon_{yy})(k_x,k_y);$
	$(\epsilon_{xx} - \epsilon_{yy})k_x + 2\epsilon_{xy}k_y, (\epsilon_{yy} - \epsilon_{xx})k_y + 2\epsilon_{xy}k_x;$
	$2\epsilon_{xy}B_z, (\epsilon_{xx} - \epsilon_{yy})B_z;$
	$(\epsilon_{xx} - \epsilon_{yy})\mathcal{E}_x + \epsilon_{xy}\mathcal{E}_y, (\epsilon_{yy} - \epsilon_{xx})\mathcal{E}_y + \epsilon_{xy}\mathcal{E}_x;$
	$(\epsilon_{xx} + \epsilon_{yy})(\mathcal{E}_x, \mathcal{E}_y); \ s_z k_y, -s_z k_x;$
	$s_yB_y - s_xB_x, s_xB_y + s_yB_x; \ s_z{\cal E}_y, -s_z{\cal E}_x;$
	$s_y \mathcal{E}_z, -s_x \mathcal{E}_z; \ s_z (k_x \mathcal{E}_y + k_y \mathcal{E}_x), s_z (k_x \mathcal{E}_x - k_y \mathcal{E}_y);$
	$(s_xk_y+s_yk_x){\cal E}_z,(s_xk_x-s_yk_y){\cal E}_z;$
	$2s \epsilon s (\epsilon - \epsilon)$

time-reversal invariance implies

$$\mathcal{T}^{-1}\mathcal{H}(\zeta \mathcal{K})\mathcal{T} = \mathcal{H}^*(\mathcal{K}) = \mathcal{H}^t(\mathcal{K}).$$
(33)

Here, * denotes complex conjugation and t transposition. The prefactor ζ depends on the behavior of \mathcal{K} under time reversal. **k**, **B**, and **s** are odd under time reversal so that then $\zeta = -1$, while \mathcal{E} and ϵ have $\zeta = +1$.

The general analysis needs to be modified due to the fact that the valleys **K** and **K'** are inequivalent points in the star { κ } that characterizes the IRs of the space group for these values of κ . Therefore, the eigenstates at **K** and **K'** need to be combined in order to use the general relation (32). It turns out that the bands at **K** and **K'** belong to the case denoted " a_2 " in Ref. 6, i.e., the functions $\theta \Psi_{\mathbf{K}\lambda} = \Psi^*_{\mathbf{K}\lambda}$ at **K** and $\Psi_{\mathbf{K}'\lambda}$ at **K'** are

linearly related via a unitary matrix \mathcal{T}

$$\theta \Psi_{\mathbf{K},\lambda} = \Psi_{\mathbf{K}\lambda}^* = \sum_{\lambda'} \mathcal{T}_{\lambda\lambda'} \Psi_{\mathbf{K}'\lambda'}, \qquad (34)$$

where θ denotes the time-reversal operator. But the crystallographic point group D_{6h} also contains elements Rthat likewise establish linear relations between $\Psi_{\mathbf{K}\lambda}$ and $\Psi_{\mathbf{K}'\lambda}$, see Eq. (20). Combining Eqs. (20) and (34), we obtain the linear relation

$$\theta \Psi_{\mathbf{K},\lambda} = \Psi_{\mathbf{K}\lambda}^* = \sum_{\lambda',\lambda''} \mathcal{T}_{\lambda\lambda'} \mathcal{D}_{\lambda'\lambda''}(R) \Psi_{\mathbf{K}\lambda''}.$$
 (35)

Obviously, the matrix \mathcal{T} depends on the symmetry element R. For $R = R_y$ (which preserves the sublattices A, B as discussed above) we get $\mathcal{T} = \mathbb{1}$. If we choose R as inversion [which flips the sublattices A, B, thus $\mathcal{D}(R) = \sigma_x$] we get $\mathcal{T} = \sigma_x$ (as in Ref. 20). However, in any case we have $\mathcal{T}^2 = +1$, as expected for spinless particles.²⁹

Combining the general relation (33) with (35) we obtain the condition^{6,20}

$$\mathcal{T}^{-1}\mathcal{H}(R^{-1}\mathcal{K})\mathcal{T} = \mathcal{H}^*(\zeta \mathcal{K}) = \mathcal{H}^t(\zeta \mathcal{K}).$$
(36)

It provides a general criterion for determining which terms in the expansion (29) are allowed by time-reversal invariance and which terms are forbidden.

IV. DISCUSSION OF INVARIANT EXPANSION

Tables I and II are the main result of this work. In the following, we discuss some associated physical consequences. In Sec. IV A, we focus on the orbital motion of the electrons, including the effect of perturbing electric and magnetic fields \mathcal{E} and **B**. In Sec. IV B, we consider the effect of strain. Finally, the effect of spin-orbit coupling is discussed in Sec. IV C.

We use different letters for the prefactors in the invariant expansion (29) to classify the terms according to their relevance, though such a scheme cannot be rigorous when mixed effects are considered. Coefficients $a_{\kappa\lambda}^{\alpha\beta}$ refer to invariants characterizing the orbital motion in the absence of fields; $r_{\kappa\lambda}^{\alpha\beta}$ denote the prefactors of orbital or spin-dependent Rashba-like invariants proportional to an external electric field; $z_{\kappa\lambda}^{\alpha\beta}$ denote prefactors of Zeeman-like terms proportional to B; coefficients $b_{\kappa\lambda}^{\alpha\beta}$ characterize the Bir-Pikus strain Hamiltonian for graphene; and coefficients $p_{\kappa\lambda}^{\alpha\beta}$ characterize the intrinsic (Pauli) spin-orbit coupling.

A. Orbital Motion

Equation (36) implies that all terms in Eq. (15) are allowed by time-reversal invariance, yet, e.g., the thirdorder invariant $k_y(3k_x^2 - k_y^2)\sigma_z$ is forbidden. The tensor operators that give rise to invariants in $\mathcal{H}_{55}^{\mathbf{K}}(\mathcal{K})$ allowed by time reversal have been printed in bold in Table II.

The existence of k-linear terms in $\mathcal{H}_{55}^{\mathbf{K}}$ is intimately related to the behavior of $\mathcal{H}_{55}^{\mathbf{K}}$ under time reversal characterized by Eq. (36).^{6,20} For comparison, consider the Γ point $\mathbf{k} = 0$ of a material with point group D_{3h} , where Tables I and II are valid, too. Yet the k-linear terms in $\mathcal{H}_{55}^{\mathbf{K}}$ are then forbidden because, in this case (case a_1 in Ref. 6), the matrices $\{X_l^{(\kappa)}\}$ can be classified as even or odd under time reversal with $\{X_l^{(6)}\}$ even.

To linear order of \mathcal{E}_z and in the absence of other perturbations, an external perpendicular electric field \mathcal{E}_z cannot couple to the planar orbital motion in graphene (to all orders in k). The lowest-order invariant involving in-plane electric fields reads

$$r_{21}^{55}(k_x \mathcal{E}_y - k_y \mathcal{E}_x)\sigma_z \quad . \tag{37}$$

A linear Zeeman splitting in a parallel magnetic field $\mathbf{B}_{\parallel} = (B_x, B_y, 0)$ (and independent of k) is likewise forbidden by symmetry. However, this result is not a consequence of the planar geometry of graphene. Indeed, it follows from the character tables of D_{3h} that an in-plane orbital magnetic moment is absent in all systems characterized by this point group, e.g., also in graphite. On the other hand, the allowed invariant²⁰

$$z_{21}^{55}B_z\sigma_z$$
 (38)

implies that electrons in graphene have an orbital magnetic moment in z direction. We have here an interesting difference between the Dirac-like orbital motion of Bloch electrons in graphene and truly relativistic systems. For neutrinos, which are almost-massless Dirac fermions, the magnetic moment is proportional to their mass.³⁹ The spin magnetic moment of Rashba and 2D Dresselhaus electrons has both a z and an in-plane component, though generally these are different.¹⁹

In a field $B_z > 0$ we can utilize the usual¹⁹ ladder operators $a^{\pm} = l_B(k_x \pm ik_y)/\sqrt{2}$ with magnetic length $l_B = \sqrt{\hbar/|eB_z|}$ to obtain up to linear order of **k** and B_z $[\sigma_{\pm} \equiv \frac{1}{2}(\sigma_x \pm i\sigma_y)]$

$$\mathcal{H}_{55}^{\rm L\pm} = a_{10}^{55} \mathbb{1} + \frac{\sqrt{2} \, a_{61}^{55}}{l_{\rm B}} \left(a^+ \sigma_{\mp} + a^- \sigma_{\pm} \right) \pm z_{21}^{55} \, B_z \, \sigma_z \,, \ (39)$$

where the upper (lower) sign applies to **K** (**K**'). Both \mathcal{H}_{55}^{L+} and \mathcal{H}_{55}^{L-} have the same Landau spectrum

$$E_{n\pm} = a_{10}^{55} \pm \frac{\sqrt{2} a_{61}^{55}}{l_{\rm B}} \sqrt{n + \left(\frac{z_{21}^{55}}{a_{61}^{55}}\right)^2 \left|\frac{\hbar B_z}{2e}\right|}, (40a)$$

$$E_0 = a_{10}^{55} - z_{21}^{55} |B_z| , \qquad (40b)$$

with positive integers n. Finding $E_0 \neq a_{10}^{55}$ signals broken particle-hole symmetry. For $z_{21}^{55} = 0$, Eq. (40) is identical to the spectrum of 2D massless Dirac fermions in a magnetic field B_z (Ref. 40). The spectrum (40) is obtained also for the Rashba³⁴ and 2D Dresselhaus¹⁹ models in the limit of an infinite effective electron mass. This result illustrates the fact that the spectrum (40) for $B_z > 0$ and $z_{21}^{55} = 0$ is determined by the dispersion (17) at B = 0 but does not depend on the transformational properties of the underlying basis functions (which are different for these models).

B. Strain-induced effects

To incorporate the effect of strain ϵ into the invariant expansion (29), we follow the general theory developed in Ref. 6. A small homogeneous strain for a 2D sheet such as graphene is defined by the symmetric strain tensor⁴¹

$$\epsilon_{ij} = \frac{1}{2} \left(\frac{\partial u_i}{\partial r_j} + \frac{\partial u_j}{\partial r_i} + \frac{\partial u_z}{\partial r_i} \frac{\partial u_z}{\partial r_j} \right), \tag{41}$$

where $\mathbf{u}(\mathbf{r})$ is the displacement vector at point \mathbf{r} due to strain and $i, j \in \{x, y\}$. Note that, while the position vector \mathbf{r} is two-dimensional, the displacement $\mathbf{u}(\mathbf{r})$ can have three nonzero components. Nevertheless, to lowest order, the components ϵ_{jz} and ϵ_{zz} of the general (three-dimensional) strain tensor vanish in a 2D sheet like graphene;⁴¹ only the in-plane components of $\boldsymbol{\epsilon}$ are finite. Recent experimental studies have mapped⁴² and engineered⁴³ strain in single-layer graphene.

Quite generally,⁶ the components ϵ_{ij} of the strain tensor transform like the symmetrized products $\{k_i, k_j\}$. Thus to lowest order, we get the irreducible tensor components proportional to ϵ_{ij} listed in Table II. Previous work^{44,45} has already identified the terms

$$b_{11}^{55}(\epsilon_{xx} + \epsilon_{yy})\mathbb{1} + b_{61}^{55}[(\epsilon_{yy} - \epsilon_{xx})\sigma_x + 2\epsilon_{xy}\sigma_y], \quad (42)$$

where the second term can be interpreted as arising from a geometry-related fictitious vector potential.^{46,47} From that viewpoint, the term

$$b_{12}^{55}[(\epsilon_{yy} - \epsilon_{xx})k_x + 2\epsilon_{xy}k_y]\mathbb{1}$$

$$\tag{43}$$

has a straightforward interpretation in terms of the same type of gauge-field correction to the quadratic-dispersion contribution $a_{11}^{55}(k_x^2 + k_y^2)\mathbb{1}$. Possibilities to use straininduced pseudo-magnetic fields to manipulate electronic transport in graphene have attracted significant attention recently.^{48–50} Furthermore, an isotropic renormalization of the electron velocity as embodied in the term

$$b_{62}^{55}(\epsilon_{xx} + \epsilon_{yy})(k_x\sigma_x + k_y\sigma_y) \tag{44}$$

was discussed in conjunction with smooth rippling of the graphene sheet.⁵¹ Our results suggest that even an anisotropic velocity renormalization can be engineered using strain, based on the contribution

$$b_{63}^{55}\{[(\epsilon_{xx}-\epsilon_{yy})k_x+2\epsilon_{xy}k_y]\sigma_x+[(\epsilon_{yy}-\epsilon_{xx})k_y+2\epsilon_{xy}k_x]\sigma_y\}.$$
(45)

This mechanism for creating an anisotropic dispersion provides an alternative to the previously suggested⁵² periodic modulation of graphene sheets. Several terms involve strain in combination with external fields. A strain-dependent renormalization of orbital Zeeman coupling is given by

$$b_{21}^{55}(\epsilon_{xx} + \epsilon_{yy})B_z\sigma_z \quad . \tag{46}$$

In-plane electric fields coupled with strain generate a gap via the contribution

$$b_{22}^{55}[(\epsilon_{xx} - \epsilon_{yy})\mathcal{E}_y + 2\epsilon_{xy}\mathcal{E}_x]\sigma_z \quad . \tag{47}$$

Both the orbital g-factor renormalization and the gap size could vary randomly in space, as strain is associated with certain types of disorder such as ripples.⁵³

Our symmetry analysis has yielded all the terms that can be generated from Eqs. (15) and (37) by replacing

$$k_x \to k_x + \alpha (\epsilon_{yy} - \epsilon_{xx}), \qquad k_y \to k_y + 2\alpha \, \epsilon_{xy}, \quad (48)$$

thus resembling a minimal coupling to a strain-related geometric gauge field with coupling constant α . This result reflects the fact that both the wave vector components k_x, k_y and the strain tensor components $\epsilon_{xx} - \epsilon_{yy}, 2\epsilon_{xy}$ transform according to Γ_6 (see Table II) so that these terms may be combined as in Eq. (48) to form a new (alternative) tensor operator. However, the construction of higher-order tensor operators using Eq. (30) requires that we include not only the weighted sum of k_x, k_y and $\epsilon_{xx} - \epsilon_{yy}, 2\epsilon_{xy}$ as one tensor operator, as in Eq. (48). Rather, their (weighted) difference also constitutes a linearly independent tensor operator. So even if the latter has been defined such that the prefactor of the corresponding linear-order invariant vanishes, it is not guaranteed that the prefactors of higher-order invariants constructed via Eq. (30) will vanish as well. Thus the simple replacement (48) may be insufficient to account for strain effects in higher orders.

C. Spin-Orbit Coupling

As discussed in Ref. 6, spin-orbit (SO) coupling can be incorporated in the invariant expansion (29) in two equivalent ways. In the first approach, the components of the pseudovector **s** enter the general tensor \mathcal{K} in much the same way as \mathbf{k} , \mathbf{B} , $\boldsymbol{\mathcal{E}}$, and $\boldsymbol{\epsilon}$. In Table II, we have listed the resulting lowest-order irreducible tensor operators. With spin taken into account in this way, the basis functions of the Hamiltonian $\mathcal{H}_{55}^{\mathbf{K}}(\mathcal{K})$ transform according to the direct product $\Gamma_5 \times \mathcal{D}_{1/2}$ of the representation Γ_5 according to which the coordinate functions (18) transform and the representation $\mathcal{D}_{1/2}$ of SU(2) according to which the spin functions $|\uparrow\rangle$ and $|\downarrow\rangle$ transform. In other words, the basis functions are constructed from the eigenfunctions without \mathcal{H}_{so} , and \mathcal{H}_{so} is introduced as a perturbation (similar to \mathcal{E} , **B**, and ϵ). One can thus ascertain at once which of the coefficients $a_{\kappa\lambda}^{55}$ entering $\mathcal{H}_{55}^{\mathbf{K}}(\mathcal{K})$ are relativistically small. In lowest order, we get the terms given in Eqs. (24) and (26), as well as several

invariants where strain couples to the intrinsic spin degree of freedom. As noted above, terms shown in Eq. (26) were previously omitted.

Alternatively, we can construct $\mathcal{H}(\mathcal{K})$ by directly using the double-group representations of D_{3h} contained in the product representation $\Gamma_5 \times \mathcal{D}_{1/2}$. Decomposing $\Gamma_5 \times \mathcal{D}_{1/2}$ into IRs gives rise to spinors that transform according to the double-group IRs Γ_7 (with representative basis functions $\{|\rho_- \uparrow\rangle, |\rho_+ \downarrow\rangle\}$) and Γ_9 ($\{|\rho_- \downarrow\rangle, |\rho_+ \uparrow\rangle\}$) of D_{3h} . The corresponding basis matrices are also listed in Table I which are again combined with the tensor operators in Table II. Thus we obtain the 4×4 Hamiltonian

$$\mathcal{H} = \begin{pmatrix} \mathcal{H}_{77} & \mathcal{H}_{79} \\ \mathcal{H}_{97} & \mathcal{H}_{99} \end{pmatrix}, \tag{49}$$

where each block $\mathcal{H}_{\alpha\beta}$ is given by an invariant expansion of the form (29). Obviously, both approaches are related by a unitary transformation.

The most important consequence of SO coupling, which can be inferred directly from Eq. (49), is the opening of a gap Δ between the bands Γ_7 and Γ_9 , so that we get $a_{10}^{77} = \epsilon_{2p} + \frac{1}{2}\Delta$ and $a_{10}^{99} = \epsilon_{2p} - \frac{1}{2}\Delta$ (Refs. 16,54). The origin of this gap can be understood as follows. We may replace the spinless basis functions $|\rho_{\mp}\rangle$ by the unitarily equivalent basis functions $|\tilde{x}\rangle = \frac{1}{\sqrt{2}}(|\rho_{-}\rangle + |\rho_{+}\rangle)$ and $|\tilde{y}\rangle = \frac{i}{\sqrt{2}}(|\rho_{-}\rangle - |\rho_{+}\rangle)$. The latter basis functions have equal magnitudes on both sublattices A and B. Then we have $\Delta = \frac{\hbar}{2m_{0}^{2}c^{2}}\langle \tilde{x}|[\nabla V \times \mathbf{p}]_{z}|\tilde{y}\rangle$, where V is the microscopic crystal potential of graphene. From a symmetry point of view, the gap Δ is thus analogous to the gap that separates the topmost valence band in semiconductors such as Ge and GaAs from the spin-split-off valence band.¹⁹ Of course, in graphene we would have $\Delta = 0$ if the basis functions $|\tilde{x}\rangle$ and $|\tilde{y}\rangle$ were comprised of pure π (p_z) orbitals. However, SO coupling induces a mixing of the π (p_z) and σ (p_x, p_y) orbitals in graphene that contributes to Δ in second order of SO coupling.^{15,55,56} A second contribution to Δ (linear in SO coupling) stems from an SO-induced coupling between the atomic p_z and higher atomic orbitals such as d states.^{15,16} These mechanisms refer to SO matrix elements for states localized on the same atom. As SO coupling originates from the steep gradients of the Coulomb potentials in the atomic cores, these terms generally provide the dominant effect,²⁵ though we see that such mechanisms are less effective in graphene. A third contribution to Δ is due to the second-nearest-neighbor coupling of the p_z orbitals discussed in Ref. 23. Recent first-principles calculations⁵⁷ found $\Delta \approx 24 \ \mu eV$, with this gap arising almost entirely from contributions from d and higher orbitals. The sign of Δ and, thus, the order of the bands Γ_7 and Γ_9 cannot be inferred from our analysis.

Unlike the case of inversion-asymmetric crystal structures such as zinc blende and wurtzite, SO coupling in inversion-symmetric graphene does not give rise to spin splitting. Hence, with spin taken into account, we get a two-fold spin degeneracy for all bands throughout the Brillouin zone.⁶ Accordingly, the Hamiltonian (49) preserves the two-fold spin degeneracy (for $B = \mathcal{E} = 0$). An external field \mathcal{E} breaks the spatial inversion symmetry. Tables I and II show that, to lowest order in k, the resulting spin splitting is due to the invariants given in Eqs. (24) and (26).

As is the case in other materials,⁶ strain mediates a coupling between intrinsic-spin and orbital dynamics. The lowest-order contribution

$$p_{22}^{55}(\epsilon_{xx} + \epsilon_{yy})s_z\sigma_z \tag{50}$$

constitutes a renormalization of the intrinsic SO coupling $p_{21}^{55}s_z\sigma_z$. The existence of this term enables strain engineering of spin splitting in graphene. It also implies that disorder associated with strain (such as ripples) gives rise to a spatially random SO coupling that should have implications for spin relaxation in graphene beyond previously considered mechanisms.⁵⁸

V. CONCLUSIONS

We have performed a detailed symmetry analysis of the trigonal band structure of graphene. A systematic invariant expansion of the envelope-function Hamiltonians describing electron states near the **K** and **K'** points is presented, including effects due to external electric and magnetic fields, strain, and spin-orbit coupling. Our results include all terms up to second order as well as selected terms up to third order and they include several previously unnoticed terms. Examples for the latter are shown in Eqs. (26), (37), (45)–(47), and (50). We have also highlighted the peculiar role played by time-reversal invariance in determining graphene's band structure.

It should be noted that, in principle, our analysis based on the invariant expansion (29) could be extended to multilayer graphene. However, as discussed, e.g., in Ref. 59, N-layer graphene systems consisting of an even (odd, N > 1) number of sheets have the point group $D_{3d}(D_{3h})$, and the **K** points have the point group $D_3(C_{3h})$. The proper symmetry analysis for multilayer graphene needs to be based on these symmetries and, hence, will differ qualitatively from the one presented in this work.

Before closing, we comment on the significance of the different transformational properties characterizing the wave functions of electrons in graphene and massless Dirac fermions, respectively. Our group-theoretical analysis suggests to divide the electronic properties of these systems into two categories: (I) those that emerge from the linear dispersion (17) but that are independent of the transformational properties of the basis functions, and (II) those that *do* reflect these transformational properties.³⁵ Clearly, the experimentally verified^{60,61} Landau spectrum (40) belongs to category (I). Similarly, *Zitterbewegung*-like effects,⁶² i.e., phenomena arising due to the interference between electron states from neighboring bands,⁶³ generally belong to category (II). In contrast, the magnetic moment belongs to category (II). Similarly,

Kramers' degeneracy only holds for particle states transforming according to spinor IRs. In this context, it is not important whether the symmetry group is discrete or continuous, as in both cases we can distinguish ordinary and spinor IRs.

Acknowledgments

The authors appreciate stimulating discussions with C.-S. Chu, D. Culcer, E. I. Rashba, A. I. Signal, and L.-

- ¹ A. K. Geim and K. S. Novoselov, Nature Mater. **6**, 183 (2007).
- ² C. W. J. Beenakker, Rev. Mod. Phys. **80**, 1337 (2008).
- ³ A. H. Castro Neto, F. Guinea, N. M. R. Peres, K. S. Novoselov, and A. K. Geim, Rev. Mod. Phys. 81, 109 (2009).
- ⁴ K. S. Novoselov, A. K. Geim, S. V. Morozov, D. Jiang, Y. Zhang, S. V. Dubonos, I. V. Grigorieva, and A. A. Firsov, Science **306**, 666 (2004).
- ⁵ K. S. Novoselov, D. Jiang, F. Schedin, T. J. Booth, V. V. Khotkevich, S. V. Morozov, and A. K. Geim, Proc. Natl. Acad. Sci. USA **102**, 10451 (2005).
- ⁶ G. L. Bir and G. E. Pikus, Symmetry and Strain-Induced Effects in Semiconductors (Wiley, New York, 1974).
- ⁷ P. R. Wallace, Phys. Rev. **71**, 622 (1947).
- ⁸ G. W. Semenoff, Phys. Rev. Lett. **53**, 2449 (1984).
- ⁹ D. P. DiVincenzo and E. J. Mele, Phys. Rev. B **29**, 1685 (1984).
- ¹⁰ F. D. M. Haldane, Phys. Rev. Lett. **61**, 2015 (1988).
- ¹¹ R. Saito, G. Dresselhaus, and M. S. Dresselhaus, *Physical Properties of Carbon Nanotubes* (Imperial College, London, 1998).
- 12 G. S. Painter and D. E. Ellis, Phys. Rev. B 1, 4747 (1970).
- ¹³ J. L. Carter Jr., Ph.D. thesis, Cornell University (1953).
- ¹⁴ W. M. Lomer, Proc. R. Soc. Lond. A **227**, 330 (1955).
- ¹⁵ J. C. Slonczewski, Ph.D. thesis, Rutgers University (1955).
- ¹⁶ J. C. Slonczewski and P. R. Weiss, Phys. Rev. **109**, 272 (1958).
- ¹⁷ F. Bassani and G. Pastori Parravicini, Nuovo Cim. B 50, 95 (1967).
- ¹⁸ J. M. Luttinger, Phys. Rev. **102**, 1030 (1956).
- R. Winkler, Spin-Orbit Coupling Effects in Two-Dimensional Electron and Hole Systems (Springer, Berlin, 2003).
 I. M. E. China and M. A. H. Maruelian, Dhana China, China and M. A. L. Maruelian, China and M. A. L. Maruelian, China and China a
- ²⁰ J. L. Mañes, F. Guinea, and M. A. H. Vozmediano, Phys. Rev. B **75**, 155424 (2007).
- ²¹ T. Ando, J. Phys. Soc. Jpn. **69**, 1757 (2000).
- ²² A. De Martino, R. Egger, K. Hallberg, and C. A. Balseiro, Phys. Rev. Lett. 88, 206402 (2002).
- ²³ C. L. Kane and E. J. Mele, Phys. Rev. Lett. **95**, 226801 (2005).
- ²⁴ M. Zarea and N. Sandler, Phys. Rev. B **79**, 165442 (2009).
- ²⁵ D. J. Chadi, Phys. Rev. B **16**, 790 (1977).
- ²⁶ If the IRs { Γ_{κ} } contained in the product $\Gamma_{\alpha} \times \Gamma_{\beta}$ are only a subset of all IRs { $\Gamma_{\kappa'}$ } of \mathcal{G} , a tensor operator $\mathcal{K}^{(\kappa',\lambda)}$ transforming according to an IR $\Gamma_{\kappa'} \notin {\Gamma_{\kappa}}$ is forbidden by symmetry to occur in $\mathcal{H}_{\alpha\beta}(\mathcal{K})$. For example, in $\mathcal{H}_{55}^{\mathbf{K}}(\mathcal{K})$ all tensor operators transforming according to Γ_3 , Γ_4 , and Γ_5

Y. Wang. This work is supported by the Marsden Fund Council (contract MAU0702) from New-Zealand Government funding, administered by the Royal Society of New Zealand. We thank the Kavli Institute for Theoretical Physics China at the Chinese Academy of Sciences for hospitality and support during the final stages of writing this article. Work at Argonne was supported by DOE BES under Contract No. DE-AC02-06CH11357. U. Z. gratefully acknowledges hospitality at the Aspen Center for Physics during the 2008 Summer program.

are forbidden by symmetry. Yet these tensor operators are allowed in $\mathcal{H}_{77}^{K}(\mathcal{K})$, $\mathcal{H}_{99}^{B}(\mathcal{K})$, and/or $\mathcal{H}_{79}^{K}(\mathcal{K})$, see Table I.

- ²⁷ G. F. Koster, J. O. Dimmock, R. G. Wheeler, and H. Statz, *Properties of the Thirty-Two Point Groups* (MIT, Cambridge, MA, 1963).
- ²⁸ We define the IRs of D_{3h} via the characters listed in Table 65 of Ref. 27. This requires that the basis functions for Γ_5 and Γ_6 of D_{3h} are reversed, see, e.g., Ref. 6. To match the coordinate system in our Fig. 1, we have recalculated the coupling coefficients of D_{3h} listed in Table 67 of Ref. 27 for a coordinate system, where the in-plane symmetry axes are rotated by 30° relative to those defined in Fig. 3 of Ref. 27.
- ²⁹ J. J. Sakurai, *Modern Quantum Mechanics* (Addison-Wesley, Redwood City, 1985).
- ³⁰ H. Rauch, A. Zeilinger, G. Badurek, and A. Wilfing, Phys. Lett. A **54**, 425 (1975).
- ³¹ S. A. Werner, R. Colella, A. W. Overhauser, and C. F. Eagen, Phys. Rev. Lett. **35**, 1053 (1975).
- ³² J. J. Sakurai, *Advanced Quantum Mechanics* (Addison-Wesley, Reading, MA, 1967).
- ³³ G. Dresselhaus, Phys. Rev. **100**, 580 (1955).
- ³⁴ Y. A. Bychkov and E. I. Rashba, J. Phys. C: Solid State Phys. **17**, 6039 (1984).
- ³⁵ For a Hamiltonian like Eq. (29) or the Dirac Hamiltonian the full wave function is the product of the solution of this Hamiltonian times the corresponding basis functions.
- ³⁶ The (restricted) homogenous Lorentz group is often denoted SO(3, 1). To account for spin, we need its universal covering group, which is the special linear group SL(2).
- ³⁷ E. I. Rashba, Sov. Phys.–Solid State **2**, 1109 (1960).
- ³⁸ A similar situation occurs, e.g., at the K points of wurtzite materials,⁶ yet in the absence of a center of inversion.
- ³⁹ K. Zuber, Neutrino Physics (Taylor and Francis, Bristol, 2003).
- ⁴⁰ J. W. McClure, Phys. Rev. **104**, 666 (1956).
- ⁴¹ L. D. Landau and E. M. Lifshitz, *Theory of Elasticity* (Pergamon, Oxford, 1987), 3rd ed.
- ⁴² M. L. Teague, A. P. Lai, J. Velasco, C. R. Hughes, A. D. Beyer, M. W. Bockrath, C. N. Lau, and N.-C. Yeh, Nano Lett. 9, 2542 (2009).
- ⁴³ W. Bao, F. Miao, Z. Chen, H. Zhang, W. Jang, C. Dames, and C. N. Lau, Nature Nanotech. 4, 562 (2009).
- ⁴⁴ H. Suzuura and T. Ando, Phys. Rev. B **65**, 235412 (2002).
- ⁴⁵ J. L. Mañes, Phys. Rev. B **76**, 045430 (2007).
- ⁴⁶ C. L. Kane and E. J. Mele, Phys. Rev. Lett. **78**, 1932 (1997).
- ⁴⁷ M. A. H. Vozmediano, M. I. Katsnelson, and F. Guinea,

- ⁴⁸ M. M. Fogler, F. Guinea, and M. I. Katsnelson, Phys. Rev. Lett. **101**, 226804 (2008).
- ⁴⁹ V. M. Pereira and A. H. Castro Neto, Phys. Rev. Lett. 103, 046801 (2009).
- ⁵⁰ F. Guinea, M. I. Katsnelson, and A. K. Geim, Nature Phys.
 6, 30 (2010).
- ⁵¹ F. de Juan, A. Cortijo, and M. A. H. Vozmediano, Phys. Rev. B **76**, 165409 (2007).
- ⁵² C.-H. Park, L. Yang, Y.-W. Son, M. L. Cohen, and S. G. Louie, Nature Phys. 4, 213 (2008).
- ⁵³ F. Guinea, B. Horovitz, and P. Le Doussal, Phys. Rev. B 77, 205421 (2008).
- ⁵⁴ G. Dresselhaus and M. S. Dresselhaus, Phys. Rev. 140, A401 (1965).
- ⁵⁵ H. Min, J. É. Hill, N. A. Sinitsyn, B. R. Sahu, L. Kleinman, and A. H. MacDonald, Phys. Rev. B **74**, 165310 (2006).

- ⁵⁶ D. Huertas-Hernando, F. Guinea, and A. Brataas, Phys. Rev. B **74**, 155426 (2006).
- ⁵⁷ M. Gmitra, S. Konschuh, C. Ertler, C. Ambrosch-Draxl, and J. Fabian, Phys. Rev. B 80, 235431 (2009).
- ⁵⁸ D. Huertas-Hernando, F. Guinea, and A. Brataas, Phys. Rev. Lett. **103**, 146801 (2009).
- ⁵⁹ L. M. Malard, M. H. D. Guimarães, D. L. Mafra, M. S. C. Mazzoni, and A. Jorio, Phys. Rev. B **79**, 125426 (2009).
- ⁶⁰ K. S. Novoselov, A. K. Geim, S. V. Morozov, D. Jiang, M. I. Katsnelson, I. V. Grigorieva, S. V. Dubonos, and A. A. Firsov, Nature **438**, 197 (2005).
- ⁶¹ Y. Zhang, Y.-W. Tan, H. L. Stormer, and P. Kim, Nature 438, 201 (2005).
- ⁶² R. Winkler, U. Zülicke, and J. Bolte, Phys. Rev. B 75, 205314 (2007).
- ⁶³ D. Culcer, Int. J. Mod. Phys. B **22**, 4765 (2008).