A PERIOD MAP FOR GENERALIZED DEFORMATIONS

DOMENICO FIORENZA AND MARCO MANETTI

ABSTRACT. For every compact Kähler manifold we give a canonical extension of Griffith's period map to generalized deformations, intended as solutions of Maurer-Cartan equation in the algebra of polyvector fields. Our construction involves the notion of Cartan homotopy and a canonical L_{∞} structure on mapping cones of morphisms of differential graded Lie algebras.

INTRODUCTION

Let X be a compact Kähler manifold and denote by $H^*(X, \mathbb{C})$ the graded vector space of its De Rham cohomology. The goal of this paper is to define a natural transformation

$$\Phi \colon \operatorname{Def}_X \to \operatorname{Aut}_{H^*(X,\mathbb{C})}$$

from infinitesimal deformations of X to automorphisms of $H^*(X, \mathbb{C})$. More precisely, for every local Artinian \mathbb{C} -algebra A and every deformation of X over Spec(A) we define in a functorial way a canonical morphism of schemes

$$\operatorname{Spec}(A) \to GL(H^*(X,\mathbb{C})) = \prod_n GL(H^n(X,\mathbb{C})).$$

Our construction will be carried out by using the interplay of Cartan homotopies and L_{∞} -morphisms and it is compatible with classical constructions of the theory of infinitesimal variations of Hodge structures. In particular:

(1) Via the natural isomorphism $H_X \simeq \bigoplus_{p,q} H^q(\Omega_X^p)$ induced by Dolbeault's theorem and the $\partial \overline{\partial}$ -lemma, the differential of Φ

$$d\Phi \colon H^1(T_X) \to \operatorname{Hom}^0(H^*(X,\mathbb{C}),H^*(X,\mathbb{C}))$$

is identified with the contraction operator: $d\Phi(\xi) = i_{\xi}$, where $i_{\xi}(\omega) = \xi \lrcorner \omega$.

(2) The contraction

$$i: H^2(T_X) \to \operatorname{Hom}^1(H^*(X, \mathbb{C}), H^*(X, \mathbb{C}))$$

is a morphism of obstruction theories. In particular, since $\operatorname{Aut}_{H^*(X,\mathbb{C})}$ is smooth, every obstruction to deformation of X is contained in the kernel of i.

(3) For every m let $H^*(F^m) \subseteq H^*(X, \mathbb{C})$ be the subspace of cohomology classes of closed (p, q)-forms, with $p \ge m$. Then the composition of Φ with the natural projection

$$GL(H^*(X,\mathbb{C})) \to \operatorname{Grass}(H^*(X,\mathbb{C})) = \prod_n \operatorname{Grass}(H^n(X,\mathbb{C})), \qquad f \mapsto f(H^*(F^m)),$$

is the classical period map.

Date: July 25, 2008.

¹⁹⁹¹ Mathematics Subject Classification. 14D07, 17B70, 13D10.

Key words and phrases. Differential graded Lie algebras, symmetric coalgebras, L_{∞} -algebras, functors of Artin rings, Kähler manifolds, period map.

We will define and study the morphism Φ using the framework of L_{∞} -algebras. It is however useful to give also a more geometric definition in the following way.

Denote by $A_X = \bigoplus_i A_X^i$ the space of complex valued differential forms on X, by $d = \partial + \overline{\partial} \colon A_X^i \to A_X^{i+1}$ the De Rham differential and by $\partial A_X \subseteq A_X$ the subspace of ∂ -exact forms. A small variation of the almost complex structure is determined by a form $\xi \in A_X^{0,1}(T_X)$: according to Newlander-Niremberg theorem, the integrability condition of ξ is equivalent to $(d + l_\xi)^2 = 0$, where $l_\xi \colon A_X^i \to A_X^{i+1}$ is the holomorphic Lie derivative, defined by the formula

$$\boldsymbol{l}_{\boldsymbol{\xi}}(\boldsymbol{\omega}) = \partial(\boldsymbol{\xi} \lrcorner \boldsymbol{\omega}) + \boldsymbol{\xi} \lrcorner \partial \boldsymbol{\omega}.$$

Notice that $l_{\xi}(\ker \partial) \subseteq \partial A_X$.

Assume therefore $(d + \mathbf{l}_{\xi})^2 = 0$; according to $\partial \overline{\partial}$ -lemma, the complex $(\partial A_X, d)$ is acyclic and then, if ξ is sufficiently small, the complex $(\partial A_X, d + \mathbf{l}_{\xi})$ is still acyclic.

In order to define the automorphism $\Phi_{\xi} \colon H^*(X, \mathbb{C}) \to H^*(X, \mathbb{C})$, let $[\omega] \in H^*(X, \mathbb{C})$ and choose a *d*-closed form $\omega_0 \in A_X$ representing $[\omega]$ and such that $\partial \omega_0 = 0$. Since

$$(d + \mathbf{l}_{\xi})\omega_0 = \partial(\xi \lrcorner \omega_0) \in \partial A_X$$
 and $(d + \mathbf{l}_{\xi})^2 \omega_0 = 0$

there exists $\beta \in A_X$ such that $(d + l_{\xi})\omega_0 = (d + l_{\xi})\partial\beta$. If i_{ξ} is the contraction, then it is not difficult to prove that $d(e^{i_{\xi}}(\omega_0 - \partial\beta)) = 0$ and the cohomology class of $e^{i_{\xi}}(\omega_0 - \partial\beta)$ does not depend on the choice of β and ω_0 , allowing to define $\Phi_{\xi}([\omega])$ as the cohomology class of $e^{i_{\xi}}(\omega_0 - \partial\beta)$.

Equivalently, for every d-closed form $\omega \in A_X$ and every small variation of the complex structure ξ we have

$$\Phi_{\xi}([\omega]) = [e^{i_{\xi}}(\omega - d\gamma - \partial\beta)], \quad \text{where} \quad \partial\omega = \partial d\gamma, \quad \boldsymbol{l}_{\xi}(\omega - d\gamma) = (d + \boldsymbol{l}_{\xi})\partial\beta$$

Moreover, as direct consequence of the L_{∞} approach, we will see that Φ_{ξ} is invariant under the gauge action, where two integrable small variation of the almost complex structure ξ_1, ξ_2 are gauge equivalent if and only if they give isomorphic deformations of X.

Our construction generalizes in a completely straightforward way to generalized deformations of X, defined as the solutions, up to gauge equivalence, of the Maurer-Cartan equation in the differential graded Lie algebra

$$\operatorname{Poly}_X = \bigoplus_i \operatorname{Poly}_X^i, \qquad \operatorname{Poly}_X^i = \bigoplus_{b-a=i-1} A_X^{0,b}(\wedge^a T_X),$$

endowed with the opposite of Dolbeault differential and the Schouthen-Nijenhuys bracket.

Putting together all these fact, at the end we get, for every m, a commutative diagram of morphism of functors of Artin rings

$$\begin{array}{c|c} \widetilde{\operatorname{Def}}_{X} & \xrightarrow{\Phi} \operatorname{Aut}_{H^{*}(X,\mathbb{C})} \\ & i & & & \\ i & & & & \\ \operatorname{Def}_{X} & \xrightarrow{p} \operatorname{Grass}_{H^{*}(F^{m}),H^{*}(X,\mathbb{C})} \end{array}$$

where Def_X is the functor of generalized deformations of X, i is the natural inclusion, $Grass_{H^*(F^m),H^*(X,\mathbb{C})}$ is the Grassmann functor with base point $H^*(F^m)$, π is the smooth morphism defined as $\pi(f) = f(H^*(F^m))$ and p is the classical m^{th} period map.

In view of this result is natural to candidate the composition $\pi\Phi$ as period map for generalized deformations.

Example 0.1. Our definition is compatible with yet existing notion of period map for generalized deformations of Calabi-Yau manifolds used in some mirror symmetry constructions.

In fact, if X is a Calabi-Yau manifold with volume element Ω , then by Tian-Todorov Lemma , every generalized deformation over $\operatorname{Spec}(A)$ is represented by an element $\xi \in \operatorname{Poly}_X^1 \otimes \mathfrak{m}_A$ such that

$$D\xi + \frac{1}{2}[\xi,\xi] = 0, \qquad \partial(\xi \lrcorner \Omega) = 0.$$

Under these assumptions our recipe gives

$$\Phi_{\xi}([\Omega]) = [e^{i_{\xi}}(\Omega)]$$

and therefore we recover the construction of [2].

Keywords and general notation.

We assume that the reader is familiar with the notion and main properties of differential graded Lie algebras and L_{∞} -algebras (we refer to [9, 16, 17, 18, 22] as introduction of such structures); however the basic definitions are recalled in this paper in order to fix notation and terminology. For the whole paper, \mathbb{K} is a field of characteristic 0; every vector space is intended over \mathbb{K} . **Art** is the category of local Artinian \mathbb{K} -algebras with residue field \mathbb{K} . For $A \in \mathbf{Art}$ we denote by \mathfrak{m}_A the maximal ideal of A. By abuse of notation, if $F: \mathbf{Art} \to \mathbf{Set}$ is a functor, we write $\xi \in F$ to mean $\xi \in F(A)$ for some fixed $A \in \mathbf{Art}$.

1. Deformation functors associated with DGLA morphisms

We recall from [23] that, to any morphism of differential graded Lie algebras over a field K of characteristic 0, $\chi: L \to M$, are naturally associated two functors of Artin rings MC_{χ} , $Def_{\chi}: Art \to Set$, in the following way:

$$\operatorname{MC}_{\chi}(A) = \left\{ (x, e^{a}) \in (L^{1} \otimes \mathfrak{m}_{A}) \times \exp(M^{0} \otimes \mathfrak{m}_{A}) \mid dx + \frac{1}{2}[x, x] = 0, \ e^{a} * \chi(x) = 0 \right\},$$
$$\operatorname{Def}_{\chi}(A) = \frac{\operatorname{MC}_{\chi}(A)}{\operatorname{gauge equivalence}},$$

where two solutions of the Maurer-Cartan equation are gauge equivalent if they belong to the same orbit of the gauge action

$$(\exp(L^0 \otimes \mathfrak{m}_A) \times \exp(dM^{-1} \otimes \mathfrak{m}_A)) \times \mathrm{MC}_{\chi}(A) \xrightarrow{*} \mathrm{MC}_{\chi}(A)$$

given by the formula

$$(e^{l}, e^{dm}) * (x, e^{a}) = (e^{l} * x, e^{dm} e^{a} e^{-\chi(l)}) = (e^{l} * x, e^{dm \bullet a \bullet (-\chi(l))})$$

The • in the rightmost term in the above formula is the Baker-Campbell-Hausdorff multiplication; namely $e^{x}e^{y} = e^{x \cdot y}$. Note that if L = 0 and the differential on M is trivial, then

$$\operatorname{Def}_{\chi}(A) = \exp(M^0 \otimes \mathfrak{m}_A).$$

It has been shown in [7] that the suspended cone of χ , i.e., the differential complex (C_{χ}, μ_1) given by the graded vector space

$$C_{\chi} = \bigoplus_{i} C_{\chi}^{i}, \qquad C_{\chi}^{i} = L^{i} \oplus M^{i-1},$$

endowed with the differential

$$\mu_1(l,m) = (dl, \chi(l) - dm), \qquad l \in L, m \in M,$$

carries a natural compatible L_{∞} -algebra structure, which we shall denote $\widetilde{C}(\chi)$, such that the associated deformation functor $\operatorname{Def}_{\widetilde{C}(\chi)}$ is naturally isomorphic to $\operatorname{Def}_{\chi}$. More precisely, the map $(l,m) \mapsto (l,e^m)$ induces a natural isomorphism $\operatorname{MC}_{\widetilde{C}(\chi)} \xrightarrow{\sim} \operatorname{MC}_{\chi}$, and homotopy equalence on $\operatorname{MC}_{\widetilde{C}(\chi)}$ is identified with gauge equivalence on MC_{χ} .

The higher brackets

$$\mu_n \colon \bigwedge^n C_{\chi} \to C_{\chi}[2-n], \qquad n \ge 2,$$

defining the L_{∞} -algebra structure $\widetilde{C}(\chi)$ have been explicitly described in [7]. Namely, one has

$$\mu_2((l_1, m_1) \land (l_2, m_2)) = \left([l_1, l_2], \frac{1}{2} [m_1, \chi(l_2)] + \frac{(-1)^{\deg(l_1)}}{2} [\chi(l_1), m_2] \right)$$

and for $n \geq 3$

$$\mu_n((l_1, m_1) \wedge \dots \wedge (l_n, m_n)) = \pm \frac{B_{n-1}}{(n-1)!} \sum_{\sigma \in S_n} \varepsilon(\sigma)[m_{\sigma(1)}, [\dots, [m_{\sigma(n-1)}, \chi(l_{\sigma(n)})] \cdots]]$$

Here the B_n 's are the Bernoulli numbers, ε is the Koszul sign and we refer to [7] for the exact determination of the overall \pm sign in the above formulas (it will not be needed in the present paper). Note that the projection on the first factor $\pi_1 : \tilde{C}(\chi) \to L$ is a linear L_{∞} -morphism.

By the functoriality of $\chi \mapsto \widetilde{C}(\chi)$ if



is a a commutative diagram of morphisms of differential graded Lie algebras, then $(x, e^a) \mapsto (f_L(x), e^{f_M(a)})$ is a natural transformation of Maurer-Cartan functors inducing a natural transformation

$$\operatorname{Def}_{\chi_1} \to \operatorname{Def}_{\chi_2}$$

Moreover, if f_L and f_M are quasi-isomorphisms, then $\operatorname{Def}_{\chi_1} \xrightarrow{\sim} \operatorname{Def}_{\chi_2}$ is an isomorphism.

2. An example from Kähler geometry

Let X be a compact Kähler manifold. Consider the DGLA $\operatorname{Hom}^*(A_X, A_X)$ of graded endomorphisms of the De Rham complex and their subDGLAs

$$L = \{ f \in \operatorname{Hom}^*(A_X, A_X) \mid f(\ker \partial) \subseteq \partial A_X \},\$$

$$M = \{ f \in \operatorname{Hom}^*(A_X, A_X) \mid f(\ker \partial) \subseteq \ker \partial \text{ and } f(\partial A_X) \subseteq \partial A_X \}.$$

Then we have a commutative diagram of morphisms of DGLAs, where the vertical arrows are the inclusions

$$0 \longleftarrow L \longrightarrow L$$

$$\rho \downarrow \qquad \eta \downarrow \qquad \downarrow \chi$$

$$\operatorname{Hom}^* \left(\frac{\ker \partial}{\partial A_X}, \frac{\ker \partial}{\partial A_X} \right) \longleftarrow M \longrightarrow \operatorname{Hom}^*(A_X, A_X)$$

By the $\partial \overline{\partial}$ -lemma, we have quasi-isomorphisms

$$(A_X, d) \longleftarrow (\ker \partial, d) \longrightarrow \left(\frac{\ker \partial}{\partial A_X}, 0\right) \cong (H^*(X, \mathbb{C}), 0)$$

Hence the horizontal arrows in the above commutative diagram are quasi-isomorphisms and we get the following isomorphisms of deformation functors

$$\operatorname{Def}_{\chi} \simeq \operatorname{Def}_{\eta} \simeq \operatorname{Def}_{\rho} = \operatorname{Aut}_{H^*(X,\mathbb{C})}.$$

The isomorphism ψ : $\operatorname{Def}_{\chi} \simeq \operatorname{Aut}_{H^*(X,\mathbb{C})}$ is explicitly described as follows: given a Maurer-Cartan element $(\alpha, e^a) \in \operatorname{MC}_{\chi}$ and a cohomology class $[\omega] \in H^*(X, \mathbb{C})$,

$$\psi_a([\omega]) = [e^{\tilde{a}}\omega_0 - \partial\beta_0],$$

where $(\tilde{\alpha}, e^{\tilde{a}}) \in \mathrm{MC}_{\eta} \subseteq \mathrm{MC}_{\chi}$ is gauge-equivalent to (α, e^{a}) , the differential form ω_{0} is a ∂ -closed representative for the cohomology class $[\omega]$, and $\beta_{0} \in A_{X}$ is such that $d(e^{\tilde{a}}\omega_{0} - \partial\beta_{0}) = 0$. The cohomology class $[e^{\tilde{a}}\omega_{0} - \partial\beta_{0}]$ is independent of the choices of $(\tilde{\alpha}, e^{\tilde{a}}), \omega_{0}$ and β_{0} . Since $e^{\tilde{a}}$ is an automorphism of ∂A_{X} , we can write

$$\psi_a([\omega]) = [e^a(\omega_0 - \partial\beta_0)]$$

for any $(\tilde{\alpha}, e^{\tilde{a}})$ and ω_0 as above and any $\beta \in A_X$ such that $de^{\tilde{a}}(\omega_0 - \partial \beta) = 0$.

Remark 2.1. Since $\chi: L \hookrightarrow \operatorname{Hom}^*(A_X, A_X)$ is injective, the projection on the second factor $C_{\chi} \to \operatorname{Hom}^{*-1}(A_X, A_X)$ induces an identification $H^*(C_{\chi}) \simeq H^{*-1}(\operatorname{coker} \chi)$ and so in particular

$$H^1(C_{\chi}) \xrightarrow{\sim} H^0(\operatorname{Hom}^*(\ker \partial, A_X/\partial A_X)) = \operatorname{Hom}^0(H^*(\ker \partial), H^*(A_X/\partial A_X)).$$

Hence, the differential of ψ is naturally identified with the linear isomorphism

$$\operatorname{Hom}^{0}(H^{*}(\ker \partial), H^{*}(A_{X}/\partial A_{X})) \simeq \operatorname{Hom}^{0}(H^{*}(X, \mathbb{C}); H^{*}(X, \mathbb{C}))$$

induced by the $\partial \overline{\partial}$ -lemma and the De Rham isomorphism.

For later use, we give a more explicit description of the map ψ by writing a map $\tilde{\psi} \colon \mathrm{MC}_{\chi} \to \mathrm{Aut}_{H^*(X,\mathbb{C})}$ inducing it. To define the map $\tilde{\psi}$ we need a few preliminary remarks.

Lemma 2.2. If $(\alpha, e^a) \in MC_{\chi}$, then in the associative algebra $Hom^*(A_X, A_X)$ we have the equality

$$e^{-a}de^a = d + \alpha.$$

Proof. By the definition of the gauge action in the DGLA $\operatorname{Hom}^*(A_X, A_X)$, one has for every $x \in \operatorname{Hom}^0(A_X, A_X)$, $y \in \operatorname{Hom}^1(A_X, A_X)$ the formula

$$e^x * y = e^x(d+y)e^{-x} - d$$

In particular $e^{-a}de^a = d + e^{-a} * 0$.

Corollary 2.3. If $(\alpha, e^a) \in MC_{\chi}$, then the graded subspaces

$$e^a(\ker \partial), \quad e^a(\partial A_X)$$

are subcomplexes of (A_X, d) . Moreover, the map

$$e^a \colon \left(\frac{\ker\partial}{\partial A_X}, 0\right) \to \left(\frac{e^a(\ker\partial)}{e^a(\partial A_X)}, 0\right)$$

is an isomorphism of complexes and the natural maps

$$(A_X, d) \longleftarrow (e^a(\ker \partial), d) \longrightarrow \left(\frac{e^a(\ker \partial)}{e^a(\partial A_X)}, 0\right)$$

are quasiisomorphisms.

Proof. Both $(\partial A_X, d)$ and $(\ker \partial, d)$ are subcomplexes of (A_X, d) . Since $de^a(v) = e^a(dv + \alpha(v))$, with $\alpha(\ker \partial) \subseteq \partial A_X$, we have $de^a(\partial A_X) \subseteq e^a(\partial A_X)$ and $de^a(\ker \partial) \subseteq e^a(\ker \partial)$. The induced differential on the quotient space $\frac{e^a(\ker \partial)}{e^a(\partial A_X)}$ is trivial since, by the $\partial\overline{\partial}$ -lemma, $d(\ker \partial) \subseteq \partial A_X$. Again by $\partial\overline{\partial}$ -lemma the complex $(\partial A_X, d)$ is acyclic and therefore the morphisms of complexes

$$(A_X, d) \longleftarrow (\ker \partial, d) \longrightarrow \left(\frac{\ker \partial}{\partial A_X}, 0\right)$$

are quasiisomorphisms. Since every infinitesimal perturbations of an acyclic complex is still acyclic, the complex $(e^a(\partial A_X), d)$ is acyclic and therefore the morphisms of complexes

$$(A_X, d) \longleftarrow (e^a(\ker \partial), d) \longrightarrow \left(\frac{e^a(\ker \partial)}{e^a(\partial A_X)}, 0\right)$$

are quasiisomorphisms.

Definition 2.4. The isomorphism

$$\psi_a \colon H^*(X, \mathbb{C}) \to H^*(X, \mathbb{C})$$

associated to a Maurer-Cartan element (α, e^a) via the natural map $\mathrm{MC}_{\chi} \to \mathrm{Def}_{\rho} \simeq \mathrm{Aut}^0(H^*(X;\mathbb{C}))$ is obtained by the De Rham isomorphism $H^*(X,\mathbb{C}) = H^*(A_X,d)$ and the chain of quasi-isomorphisms

More explicitly,

$$\widetilde{\psi}_a([\omega]) = [e^a(\omega_0 - \partial\beta)]$$

for any ∂ -closed representative ω_0 of the cohomology class $[\omega]$, and any $\beta \in A_X$ such that $de^a(\omega_0 - \partial \beta) = 0$.

Proposition 2.5. The natural transformation $\widetilde{\psi} \colon \mathrm{MC}_{\chi} \to \mathrm{Aut}_{H^*(X,\mathbb{C})}$ is gauge invariant and therefore factors to $\psi \colon \mathrm{Def}_{\chi} \to \mathrm{Aut}_{H^*(X,\mathbb{C})}$.

Proof. To show that $\widetilde{\psi}_{a \bullet (-l)} = \widetilde{\psi}_a$, we note that, since $l(\ker \partial) \subseteq \partial A_X$, we have $e^{-l}(\ker \partial) = \ker \partial$, $e^{-l}(\partial A_X) = \partial A_X$, and

$$e^{-l} \colon \left(\frac{\ker \partial}{\partial A_X}, 0\right) \to \left(\frac{\ker \partial}{\partial A_X}, 0\right)$$

is the identity. To prove that $\tilde{\psi}_{dm\bullet a} = \tilde{\psi}_a$, notice that we have a commutative diagram of morphism of complexes

which, since dm is homotopy equivalent to zero, induces the commutative diagram of isomorphisms

$$\begin{array}{rcccc}
H^*(X;\mathbb{C}) &\longleftrightarrow & \frac{\operatorname{ker} \partial}{\partial A_X} \\
& & \downarrow^{\operatorname{Id}} & & \downarrow^{e^{dm}} \\
H^*(X;\mathbb{C}) &\longleftrightarrow & \frac{e^{dm}(\operatorname{ker} \partial)}{e^{dm}(\partial A_X)}
\end{array}$$

Finally, gauge invariance of $\tilde{\psi}$, together with the explicit formulae for $\tilde{\psi}([\omega])$ and $\psi([\omega])$ written above immediately imply that $\tilde{\psi}$ induces ψ .

3. Morphisms of deformation functors associated to Cartan homotopies

In this section we formalize, under the notion of *Cartan homotopy*, a set of standard identities that often arise in algebra and geometry [5, Appendix B], and show how to any Cartan homotopy can be canonically associated a natural transformation of deformation functors.

Let L and M be two differential graded Lie algebras. For a given linear map $i \in$ Hom⁻¹(L, M), let $l: L \to M$ be the map defined as

$$a \mapsto \boldsymbol{l}_a = d\boldsymbol{i}_a + \boldsymbol{i}_{da}.$$

Definition 3.1. The map i is called a *Cartan homotopy* for l if, for every $a, b \in L$, we have:

$$i_{[a,b]} = [i_a, l_b], \qquad [i_a, i_b] = 0.$$

When the second equation is replaced by the weaker condition

$$\sum_{\sigma\in S_3}\pm [oldsymbol{i}_{x_{\sigma(1)}},[oldsymbol{i}_{x_{\sigma(2)}},oldsymbol{l}_{x_{\sigma(3)}}]=0,$$

where \pm is the Koszul sign, we shall say that *i* is a *weak Cartan homotopy*.

It is straightforward to show that the condition $i_{[a,b]} = [i_a, l_b]$ implies that l is a morphism of differential graded Lie algebras.

Example 3.2. The name Cartan homotopy has a clear origin in differential geometry. Namely, let M be a differential manifold, $\mathcal{X}(M)$ be the Lie algebra of vector fields on M, and $\mathcal{E}nd^*(\Omega^*(M))$ be the Lie algebra of endomorphisms of the de Rham algebra of M. The Lie algebra $\mathcal{X}(M)$ can be seen as a DGLA concentrated in degree zero, and the graded Lie algebra $\mathcal{E}nd^*(\Omega^*(M))$ has a degree one differential given by $[d_{dR}, -]$, where $d_{\rm dR}$ is the de Rham differential. Then the contraction

$$i: \mathcal{X}(M) \to \mathcal{E}nd^*(\Omega^*(M))[-1]$$

is a Cartan homotopy and its differential is the Lie derivative

$$[d, i] = \mathcal{L} \colon \mathcal{X}(M) \to \mathcal{E}nd^*(\Omega^*(M)).$$

In fact, by classical Cartan's homotopy formulas [1, Section 2.4], for any two vector fields X and Y on M, we have

- (1) $\mathcal{L}_X = d_{\mathrm{dR}} \boldsymbol{i}_X + \boldsymbol{i}_X d_{\mathrm{dR}} = [d_{\mathrm{dR}}, \boldsymbol{i}_X];$ (2) $\boldsymbol{i}_{[X,Y]} = \mathcal{L}_X \boldsymbol{i}_Y \boldsymbol{i}_Y \mathcal{L}_X = [\mathcal{L}_X, \boldsymbol{i}_Y] = [\boldsymbol{i}_X, \mathcal{L}_Y];$
- (3) $[i_X, i_Y] = 0.$

Note that the first Cartan formula above actually states that $[d, i] = \mathcal{L}$. Indeed $\mathcal{X}(M)$ is concentrated in degree zero and then its differential is trivial.

Example 3.3. Let *L* be a DGLA and endowe the complex L[-1] with the trivial bracket. Then the identity map

$$\mathrm{Id} \in \mathrm{Hom}^{-1}(L[-1], L)$$

is a weak Cartan homotopy for the trivial map 0. In fact $d_{L[-1]} = -d_L$ and then $d_L(a) + d_{L[-1]}(a) = 0$.

Remark 3.4. The composition of a (weak) Cartan homotopy with a morphism of DGLAs is a (weak) Cartan homotopy. If $i: L \to M[-1]$ is a Cartan homotopy and Ω is a differential graded-commutative algebra, then its natural extension

$$\mathbf{i} \otimes \mathrm{Id} \colon L \otimes \Omega \to (M \otimes \Omega)[-1], \qquad a \otimes \omega \mapsto \mathbf{i}_a \otimes \omega,$$

is a (weak) Cartan homotopy.

Remark 3.5. By definition, l is the differential of i in the complex Hom^{*}(L, M) and so i is a homotopy between l and the trivial map. Then the map $l: L \to M$ is a null-homotopic morphism of DGLAs and

$$i: L \to (\operatorname{Coker} l)[-1], \quad i: \ker l \to M[-1]$$

are morphisms of differential graded vector spaces.

Proposition 3.6. Let $l: L \to M$ be a DGLA morphism, and let $\pi_1: \widetilde{C}(l) \to L$ be the projection on the first factor. A weak Cartan homotopy $i: L \to M[-1]$ for l is the datum of a linear L_{∞} -morphisms $L \to \widetilde{C}(l)$ lifting the identity of L. In particular, if $i: L \to M[-1]$ is a weak Cartan homotopy for l, then the map $a \mapsto (a, e^{i_a})$ induces a natural transformation of Maurer-Cartan functors $MC_L \to MC_l$, and consequently a natural transformation of deformation functors $Def_L \to Def_l$.

Proof. A linear map $\tilde{i}: L \to \tilde{C}(l)$ lifting the identity of L has the form $\tilde{i}(a) = (a, i_a)$, with $i: L \to M[1]$. By the explicit expression for the higher brackets

$$\mu_n \colon \bigwedge^n C_{\chi} \to C_{\chi}[2-n], \qquad n \ge 2,$$

defining the L_{∞} -algebra structure $\widetilde{C}(\chi)$, it is straightforward to check that \tilde{i} is a morphism of complexes commuting with every bracket if and only if i is a weak Cartan homotopy. Indeed, $\tilde{i}(dx) = \mu_1(\tilde{i}(x))$ is the identity $i_{da} = (-d)i_a + l_a$; the identity $\tilde{i}([x,y]) = \mu_2(\tilde{i}(x) \wedge \tilde{i}(y))$ is

$$m{i}_{[a,b]} = rac{1}{2} [m{i}_a,m{l}_b] + rac{(-1)^{ ext{deg}(a)}}{2} [m{l}_a,m{i}_b] = [m{i}_a,m{l}_b],$$

and $\mu_n(\tilde{i}(x_1) \wedge \cdots \wedge \tilde{i}(x_n)) = 0$, for any x_1, x_2, \ldots, x_n and any $n \ge 3$, if and only if

$$\sum_{\sigma \in S_3} \pm [\boldsymbol{i}_{x_{\sigma(1)}}, [\boldsymbol{i}_{x_{\sigma(2)}}, \boldsymbol{l}_{x_{\sigma(3)}}] = 0$$

for any x_1, x_2, x_3 .

Since the L_{∞} -morphism i is linear, the map $l \mapsto (l, i_l)$ is a morphism of Maurer-Cartan functors $\mathrm{MC}_L \to \mathrm{MC}_{\widetilde{C}(l)}$. To conclude the proof, compose this morphism with the isomorphism $\mathrm{MC}_{\widetilde{C}(l)} \xrightarrow{\sim} \mathrm{MC}_l$ given by $(l, m) \mapsto (l, e^m)$.

Corollary 3.7. Let $i: N \to M[-1]$ be a Cartan homotopy for $l: N \to M$, let L be a subDGLA of M such that $l(N) \subseteq L$, and let $\chi: L \hookrightarrow M$ be the inclusion. Then the linear map

$$\Phi: N \to C(\chi), \qquad \Phi(a) = (\boldsymbol{l}_a, \boldsymbol{i}_a)$$

is a linear L_{∞} -morphism. In particular, the map $a \mapsto (l_a, e^{i_a})$ induces a natural transformation of Maurer-Cartan functors $MC_N \to MC_{\chi}$, and consequently a natural transformation of deformation functors $\mathrm{Def}_N \to \mathrm{Def}_{\chi}$.

Proof. We have a commutative diagram of differential graded Lie algebras



inducing an L_{∞} -morbism $\widetilde{C}(\boldsymbol{l}) \to \widetilde{C}(\chi)$. Composing this morphism with the L_{∞} -morphism $\tilde{i}: N \to \widetilde{C}(l)$ given by Proposition 3.6, one gets the L_{∞} -morphism Φ .

4. Polyvector fields and generalized periods

The notion of Cartan homotopy generalizes immediately to sheaves of DGLAs. In this section we give another example of Cartan homotopy which we will use later.

Let X be a complex manifold and denote by:

- T_{X,C} = T^{1,0}_X ⊕ T^{0,1}_X the complexified differential tangent bundle.
 T_X ≃ T^{1,0}_X the holomorphic tangent bundle.
 A^{p,q}_X the sheaf of differentiable (p,q)-forms and by A^{p,q}_X(E) the sheaf of (p,q)forms with values in a holomorphic vector bundle E.
- $A_X^{p,q}$ and $A_X^{p,q}(E)$ the vector spaces of global sections of $\mathcal{A}_X^{p,q}$ and $\mathcal{A}_X^{p,q}(E)$ respectively.

The direct sum

$$\mathcal{A}_X = \mathop{\oplus}\limits_i \mathcal{A}^i_X, \quad ext{where} \quad \mathcal{A}^i_X = \mathop{\oplus}\limits_{p+q=i} \mathcal{A}^{p,q}_X,$$

endowed with the wedge product \wedge , is a sheaf of graded algebras; we denote by $\mathcal{H}om^{a,b}(\mathcal{A}_X, \mathcal{A}_X)$ the sheaf of its \mathbb{C} -linear endomorphisms of \mathcal{A}_X of bidegree (a, b). Notice that ∂ and $\overline{\partial}$ are global sections of $\mathcal{H}om^{1,0}(\mathcal{A}_X, \mathcal{A}_X)$ and $\mathcal{H}om^{0,1}(\mathcal{A}_X, \mathcal{A}_X)$ respectively. The direct sum $\mathcal{H}om^*(\mathcal{A}_X, \mathcal{A}_X) = \bigoplus_k \bigoplus_{a+b=k} \mathcal{H}om^{a,b}(\mathcal{A}_X, \mathcal{A}_X)$ is a sheaf of graded associative algebras, and so a sheaf of differential graded Lie algebras with the natural bracket

$$[f,g] = fg - (-1)^{\deg(f)\deg(g)}gf$$

and differential $[d, -] = [\partial + \overline{\partial}, -]$. For any integer (a, b) with $a \leq 0$ and $b \geq 0$, let $\mathcal{G}erst_X^{a,b}$ be the sheaf

$$\mathcal{G}erst_X^{a,b} = \mathcal{A}_X^{0,b}(\wedge^{-a}T_X)$$

The direct sum $\mathcal{G}erst_X^* = \bigoplus_k \bigoplus_{a+b=k} \mathcal{G}erst_X^{a,b}$ is a sheaf of differential Gerstenhaber algebras, with the wedge product

$$\wedge \colon \mathcal{G}erst_X^{a_1,b_1} \otimes \mathcal{G}erst_X^{a_2,b_2} \to \mathcal{G}erst_X^{a_1+a_2,b_1+b_2}$$

as graded commutative product, the degree 1 differential

$$\overline{\partial} \colon \mathcal{G}erst_X^{a,b} \to \mathcal{G}erst_X^{a,b+1}$$

defined in local coordinates by the formula

$$\overline{\partial}\left(\phi\frac{\partial}{\partial z_{I}}\right) = \overline{\partial}(\phi)\frac{\partial}{\partial z_{I}}, \qquad \phi \in \mathcal{A}_{X}^{0,*},$$

and the degree 1 bracket

$$[\,,\,]_{\mathcal{G}}\colon \mathcal{G}erst_X^{a_1,b_1}\otimes \mathcal{G}erst_X^{a_2,b_2} \to \mathcal{G}erst_X^{a_1+a_2+1,b_1+b_2}$$

defined in local coordinates by the formula

$$\left[fd\overline{z}_{I}\frac{\partial}{\partial z_{H}},gd\overline{z}_{J}\frac{\partial}{\partial z_{K}}\right]_{G}=d\overline{z}_{I}\wedge d\overline{z}_{J}\left[f\frac{\partial}{\partial z_{H}},g\frac{\partial}{\partial z_{K}}\right]_{SN}.$$

Here $[,]_{SN}$ denotes the Schouten-Nijenhuis bracket on $\mathcal{A}^0_X(\wedge^*T_X)$, i.e., the odd graded Lie bracket obtained extending the usual Lie bracket on $\mathcal{A}^0_X(T_X)$ by imposing

 $[\xi, f]_{\rm SN} = \xi(f), \qquad \xi \in \mathcal{A}^0(T_X), \quad f \in \mathcal{A}^0_X;$

and the odd graded Poisson identity

$$[\xi, \eta_1 \land \eta_2]_{\rm SN} = [\xi, \eta_1]_{\rm SN} \land \eta_2 + (-1)^{\deg(\xi)(\deg(\eta_1) - 1)} \eta_1 \land [\xi, \eta_2]_{\rm SN}.$$

The contraction of differential forms with vector fields is used to define an injective morphisms of sheaves of bigraded vector spaces: the *contraction map*

$$i: \mathcal{G}erst_X^{a,b} \to \mathcal{H}om^{a,b}(\mathcal{A}_X, \mathcal{A}_X), \qquad \xi \mapsto i_{\xi}, \quad i_{\xi}(\omega) = \xi \lrcorner \omega$$

The contraction map is actually a morphism of sheaves of bigraded associative algebras:

$$oldsymbol{i}_{\xi\wedge\eta}=oldsymbol{i}_{\xi}oldsymbol{i}_{\eta}$$

In particular, since $(\mathcal{G}erst_X^*, \wedge)$ is a graded commutative algebra, we obtain

$$[oldsymbol{i}_{\xi},oldsymbol{i}_{\eta}]=0, \qquad orall \xi,\eta\in \mathcal{G}erst^*_X.$$

Note that iterated contractions give a symmetric map

$$\mathbf{i}^{(n)}: \bigodot^{n} \mathcal{G}erst_{X}^{*} \to \mathcal{H}om^{*}(\mathcal{A}_{X}, \mathcal{A}_{X})$$

$$\xi_{1} \odot \xi_{2} \odot \cdots \odot \xi_{n} \mapsto \mathbf{i}_{\xi_{1}}\mathbf{i}_{\xi_{2}} \cdots \mathbf{i}_{\xi_{n}}$$

Since $\mathcal{G}erst^*_X$ is a sheaf of differential graded Gerstenhaber algebras, its desuspension

$$\mathcal{P}oly_X^* = \mathcal{G}erst[-1]_X^*$$

is a sheaf of differential graded Lie algebras. Note that, due to the shift, the differential D in $\mathcal{P}oly_X^*$ is $-\overline{\partial}$, i.e., in local coordinates

$$D: \mathcal{P}oly^k_X \to \mathcal{P}oly^{k+1}_X$$

is given by the formula

$$D\left(\phi\frac{\partial}{\partial z_I}\right) = -\overline{\partial}(\phi)\frac{\partial}{\partial z_I}, \qquad \phi \in \mathcal{A}_X^{0,*},$$

The contraction map $i: \mathcal{G}erst_X^* \to \mathcal{H}om^*(\mathcal{A}_X, \mathcal{A}_X)$ can be seen as a linear map

$$i: \mathcal{P}oly_X^* \to \mathcal{H}om^*(\mathcal{A}_X, \mathcal{A}_X)[-1].$$

More in general, via the decalage isomorphism, the iterated contraction is a graded antisymmetric map

$$i^{(n)}: \bigwedge^n \mathcal{P}oly_X^* \to \mathcal{H}om^*(\mathcal{A}_X, \mathcal{A}_X)[-n].$$

Lemma 4.1. In the notation above, for every $\xi, \eta \in \mathcal{P}oly_X^*$ we have

$$oldsymbol{i}_{D\xi}=-[\overline{\partial},oldsymbol{i}_{\xi}], \qquad oldsymbol{i}_{[\xi,\eta]}=[oldsymbol{i}_{\xi},[\partial,oldsymbol{i}_{\eta}]], \qquad [oldsymbol{i}_{\xi},oldsymbol{i}_{\eta}]=0.$$

Proof. The third equation has been proved above, and the first equation is completely straightforward: it just expresses the Leibniz rule for $\overline{\partial}$. To prove the second equation, let

$$\Phi(\xi,\eta) = \boldsymbol{i}_{[\xi,\eta]} - [\boldsymbol{i}_{\xi}, [\partial, \boldsymbol{i}_{\eta}]].$$

Then, using $\mathbf{i}_{\xi \wedge \eta} = \mathbf{i}_{\xi} \mathbf{i}_{\eta}$, the (shifted) odd Poisson identity $[\xi, \eta_1 \wedge \eta_2] = [\xi, \eta_1] \wedge \eta_2 + (-1)^{(\deg(\xi)-1)\deg(\eta_1)}\eta_1 \wedge [\xi, \eta_2]$ and the third equation $[\mathbf{i}_{\xi}, \mathbf{i}_{\eta}] = 0$, one finds

$$\Phi(\xi,\eta_1 \wedge \eta_2) = \Phi(\xi,\eta_1) i_{\eta_2} + (-1)^{(\deg(\xi)-1) \deg(\eta_1)} i_{\eta_1} \Phi(\xi,\eta_2)$$

and

$$\Phi(\xi_1 \wedge \xi_2, \eta) = \mathbf{i}_{\xi_1} \Phi(\xi_2, \eta) + (-1)^{(\deg(\xi_2)(\deg(\eta) - 1))} \Phi(\xi_1, \eta) \mathbf{i}_{\eta_2}$$

Therefore, to prove $\Phi(\xi,\eta) = 0$ for any ξ,η one just needs to prove $\Phi(\xi,\eta) = 0$ for $\xi,\eta \in \mathcal{A}_X^0 \cup \{d\overline{z}_i\} \cup \{\partial/\partial z_j\}$, where z_1,\ldots,z_n are local holomorphic coordinates. This is straightforward and it is left to the reader: see also Lemma 7 of [21] and Lemma 7.21 of [22].

Corollary 4.2. The contraction map $i: \operatorname{Poly}_X^* \to \operatorname{Hom}^*(\mathcal{A}_X, \mathcal{A}_X)[-1]$ is a Cartan homotopy and the induced morphism l of sheaves of differential graded Lie algebras is the holomorphic Lie derivative

$$\boldsymbol{l} \colon \mathcal{P}oly_X^* \to \mathcal{H}om^*(\mathcal{A}_X, \mathcal{A}_X), \qquad \boldsymbol{\xi} \mapsto \boldsymbol{l}_{\boldsymbol{\xi}} = [\partial, \boldsymbol{i}_{\boldsymbol{\xi}}], \quad \boldsymbol{l}_{\boldsymbol{\xi}}(\omega) = \partial(\boldsymbol{\xi} \lrcorner \omega) + (-1)^{\deg(\boldsymbol{\xi})} \boldsymbol{\xi} \lrcorner \partial \omega.$$

Moreover, *l* is an injective morphism of sheaves.

Proof. As in Section 3, let $l_{\xi} = [d, i_{\xi}] + i_{D\xi}$. Since $i_{D\xi} = -[\overline{\partial}, i_{\xi}]$, we find $l_{\xi} = [\partial, i_{\xi}]$. The identity $i_{[\xi,\eta]} = [i_{\xi}, [\partial, i_{\eta}]]$ then reads $i_{[\xi,\eta]} = [i_{\xi}, l_{\eta}]$ and this, together with $[i_{\xi}, i_{\eta}] = 0$, tells us that the contraction map i is a Cartan homotopy and, consequently, the holomorphic Lie derivative $l_{\xi} = [\partial, i_{\xi}]$ is the induced morphism of sheaves of differential graded Lie algebras. Injectivity of l is easily checked in local coordinates.

The Corollary 4.2, applied to global sections shows that the contraction map

$$i: \operatorname{Poly}_X = \bigoplus_{i,j} A^{0,i}(\wedge^{-j}T_X) \to \operatorname{Hom}^*(\mathcal{A}_X, \mathcal{A}_X)[-1]$$

is a Cartan homotopy, as well as its restriction to the Kodaira-Spencer DGLA $KS_X = \bigoplus_i A^{0,i}(T_X)$.

Remark 4.3. The composition of the inclusion $KS_X \hookrightarrow \operatorname{Poly}_X$ with the iterated contraction $i^{(n)} \colon \wedge^n \operatorname{Poly}_X \to \operatorname{Hom}^*(\mathcal{A}_X, \mathcal{A}_X)[-n]$ induces in cohomology a graded antisymmetric map $\bigwedge^n H^*(KS_X) \to \operatorname{Hom}^*(H^*(X, \mathbb{C}), H^*(X, \mathbb{C}))[-n]$. In particular, from the isomorphism of graded vector spaces $H^1(KS_X^*) \simeq H^1(T_X)[-1]$ and the decalage isomorphism, the iterated contraction gives a symmetric morphism

$$i^{(n)}$$
: $\bigcirc^{n} H^{1}(T_{X}) \to \operatorname{Hom}^{0}(H^{*}(X, \mathbb{C}), H^{*}(X, \mathbb{C})).$

It is well known [12] and easy to prove that the image of $i^{(n)}$ consists of self-adjoint operators with respect the cup product on $H^*(X, \mathbb{C})$.

Under the identification $H^*(X, \mathbb{C}) = \bigoplus_{p,q} H^q(X, \Omega_X^p)$, when dim X = n the morphism $i^{(n)}$ reduces to the Yukawa coupling

$$i^{(n)}$$
: $\bigodot^n H^1(T_X) \to \bigodot^2 H^n(X, \mathcal{O}_X).$

Now, as in Section 2, consider the DGLA

 $L = \{ f \in \operatorname{Hom}^*(A_X, A_X) \mid f(\ker \partial) \subseteq \partial A_X \},\$

and let $\chi: L \hookrightarrow \operatorname{Hom}^*(A_X, A_X)$ be the inclusion. Since, $l(\operatorname{Poly}_X) \subseteq L$, by Corollary 3.7, we have a natural transformation of deformation functors $\operatorname{Def}_{\operatorname{Poly}_X} \to \operatorname{Def}_{\chi}$ induced, at the Maurer-Cartan level, by the map $\xi \mapsto (l_{\xi}, e^{i_{\xi}})$.

The functor $\operatorname{Def}_{\operatorname{Poly}_X}$ is called the functor of generalized deformations of X, see [2], and we will denote it by $\widetilde{\operatorname{Def}}_X$. We have shown in Section 2 that there exists a natural isomorphism $\psi \colon \operatorname{Def}_{\chi} \to \operatorname{Aut}_{H^*(X,\mathbb{C})}$. By these considerations, we obtain:

Theorem 4.4. The linear map

$$\operatorname{Poly}_X \to \widetilde{C}(\chi), \qquad \xi \mapsto (\boldsymbol{l}_{\xi}, \boldsymbol{i}_{\xi})$$

is a linear L_{∞} -morphism and induces a natural transformation of functors

$$\Phi \colon \mathrm{Def}_X \to \mathrm{Aut}_{H^*(X,\mathbb{C})},$$

given at the level of Maurer-Cartan functors by the map $\xi \mapsto \psi_{i_{\varepsilon}}$.

Proposition 4.5. Via the natural identifications $H^1(\operatorname{Poly}_X) = \bigoplus_{i\geq 0} H^i(\wedge^i T_X)$ and $H^*(X,\mathbb{C}) = \bigoplus_{p,q} H^q(X,\Omega_X^p)$ given by the Dolbeault's theorem and the $\partial\overline{\partial}$ -lemma, the differential of Φ ,

$$d\Phi \colon H^1(\operatorname{Poly}_X) \to \operatorname{Hom}^0(H^*(X,\mathbb{C}),H^*(X,\mathbb{C}))$$

is identified with the contraction

$$\left(\oplus_{i\geq 0}H^{i}(\wedge^{i}T_{X})\right)\otimes\left(\oplus_{p,q}H^{q}(X,\Omega_{X}^{p})\right)\to\oplus_{i,p,q}H^{q+i}(X,\Omega_{X}^{p-i})$$

Proof. By Lemma 4.1 we have a commutative diagram of differential complexes

where we have used the fact that on ker ∂ and on $A_X/\partial A_X$ the differentials ad_d and $\operatorname{ad}_{\overline{\partial}}$ coincide. Using the identification $H^*_{\overline{\partial}}(A_X) = H^*(X, \mathbb{C})$ coming from Dolbeault's theorem and the $\partial\overline{\partial}$ -lemma, and by Remark 2.1, the above commutative diagram induces the commutative diagram in cohomology

Since, by Theorem 4.4, the differential of Φ is $d\Phi = d\psi \circ i$, this ends the proof.

As a corollary of Theorem 4.4, the linear map $\xi \mapsto (\boldsymbol{l}_{\xi}, \boldsymbol{i}_{\xi})$ induces a morphism of obstruction spaces $H^2(\text{Poly}) \to \text{Hom}^1(H^*(X, \mathbb{C}), H^*(X, \mathbb{C}))$ commuting with Φ and obstruction maps. The same argument of Proposition 4.5 shows that this morphism is naturally identified with the contraction

$$\left(\oplus_{i\geq 0}H^{i+1}(\wedge^{i}T_{X})\right)\otimes\left(\oplus_{p,q}H^{q}(X,\Omega_{X}^{p})\right)\to\oplus_{i,p,q}H^{q+i+1}(X,\Omega_{X}^{p-i})$$

Since the deformation functor $\operatorname{Aut}_{H^*(X,\mathbb{C})}$ is smooth, we obtain the following version of the so-called Kodaira principle (ambient cohomology annihilates obstruction):

Proposition 4.6. The obstructions to extended deformations of a compact Kähler manifold X are contained in the subspace

$$\bigoplus_{i\geq 0} \bigcap_{p,q} \ker \left(H^{i+1}(\wedge^{i}T_{X}) \xrightarrow{\boldsymbol{i}} \operatorname{Hom} \left(H^{q}(X, \Omega_{X}^{p}), H^{q+i+1}(X, \Omega_{X}^{p-i}) \right) \right)$$

of $H^2(\operatorname{Poly}_X)$.

As an immediate corollary we recover the fact that extended deformations of compact Calabi-Yau manifolds are unobstructed [2]. Indeed, if X is an n-dimensional compact Calabi-Yau manifold, then for any $i \ge 0$ the contraction pairing

$$H^{i+1}(\wedge^{i}T_{X}) \otimes H^{0}(X,\Omega^{n}_{X}) \to H^{i+1}(X,\Omega^{n-i}_{X})$$

is nondegenerate.

5. Restriction to classical deformations

Let (X, \mathcal{O}_X) be a complex manifold. It is well known that the infinitesimal deformations of the complex structure of X are governed by the Kodaira-Spencer DGLA of X. More precisely, there is a natural isomorphism of deformation functors

$$\operatorname{Def}_{KS_X} \xrightarrow{\sim} \operatorname{Def}_X$$

mapping a Maurer-Cartan element $\xi \in A_X^{0,1}(T_X)$ to the complex manifold (X, \mathcal{O}_{ξ}) , where the structure sheaf \mathcal{O}_{ξ} is defined by

$$\mathcal{O}_{\xi} = \ker\{\overline{\partial}_{\xi} \colon \mathcal{A}_X^{0,0} \to \mathcal{A}_X^{0,1}\} = \{f \in \mathcal{A}_X^0 \mid (\overline{\partial} + \boldsymbol{l}_{\xi})f = 0\},\$$

see [5], [11], [16, Ex. 3.4.1] or [14]. The above equations have to be intended as identities among functors of Artin rings; namely, they mean that for any local Artin algebra (B, \mathfrak{m}_B) the Kuranishi data $\xi \in \mathrm{MC}_{KS_X}(B) \subseteq A_X^{0,1}(T_X) \otimes \mathfrak{m}_B$ are mapped to the family (X, \mathcal{O}_{ξ}) of complex manifolds over $\mathrm{Spec}(B)$, whose structure sheaf \mathcal{O}_{ξ} is defined by

$$\mathcal{O}_{\xi} = \ker\{\overline{\partial}_{\xi} \colon \mathcal{A}^{0,0}_X \otimes B \to \mathcal{A}^{0,1}_X \otimes B\} = \{f \in \mathcal{A}^0_X \otimes B \mid (\overline{\partial} + \boldsymbol{l}_{\xi})f = 0\}.$$

Let now

$$A_X = F_{\xi}^0 \supseteq F_{\xi}^1 \supseteq \cdots$$

be the Hodge filtration of differential forms on the complex manifold (X, \mathcal{O}_{ξ}) , i.e. for every $m \geq 0$ F_{ξ}^m is the complex of global sections of the differential ideal sheaf $\mathcal{F}_{\xi}^m \subseteq \mathcal{A}_X$ generated by $(d\mathcal{O}_{\xi})^m$. Again, here we are writing \mathcal{A}_X for the functor of Artin rings defined by $B \mapsto \mathcal{A}_X \otimes B$. If X is a compact Kähler manifold, the cohomology of (F_{ξ}^m, d) naturally embeds into the cohomology of (\mathcal{A}_X, d) . Since the dimension of $H^*(F_{\xi}^m, d)$ is independent of ξ , one can look at $H^*(F_{\xi}^m, d)$ as a different linear embedding of $H^*(F^m, d)$ into $H^*(X; \mathbb{C})$, hence $\xi \mapsto H^*(F_{\xi}^m, d)$ is a map

$$\operatorname{Def}_X \to \operatorname{Grass}_{H^*(F^m), H^*(X;\mathbb{C})},$$

called the m^{th} period map.

The inclusion of DGLAs $KS_X \hookrightarrow \operatorname{Poly}_X$ induces an embedding of deformation functors $\operatorname{Def}_X \to \widetilde{\operatorname{Def}}_X$. Hence, the restriction of Φ to Def_X is a natural transformation

$$\Phi \colon \operatorname{Def}_X \to \operatorname{Aut}_{H^*(X,\mathbb{C})}$$
.

Theorem 5.1. For any $m \ge 0$, the map $\Phi \colon \operatorname{Def}_X \to \operatorname{Aut}_{H^*(X,\mathbb{C})}$ lifts the m^{th} period map $\operatorname{Def}_X \to \operatorname{Grass}_{H^*(F^m),H^*(X;\mathbb{C})}$.

Proof. Let ξ be a Maurer-Cartan element in KS_X . Then $(\boldsymbol{l}_{\xi}, e^{\boldsymbol{i}_{\xi}}) \in \mathrm{MC}_{\chi}$ is a Maurer-Cartan element with \boldsymbol{l}_{ξ} of bidegree (0, 1). Let $[\omega]$ be an element in $H^*(F^m)$. To compute $\psi_{\xi}[\omega]$ we pick a ∂ -closed representative, for the class $[\omega]$, which we can assume to be ω , and then we take the cohomology class of a *d*-closed representative of $e^{\boldsymbol{i}_{\xi}}\omega$ in $e^{\boldsymbol{i}_{\xi}}(\ker \partial)/e^{\boldsymbol{i}_{\xi}}(\partial A_X)$, i.e., we have $\psi_{\xi}[\omega] = [e^{\boldsymbol{i}_{\xi}}(\omega - \partial\beta)]$ for any $\beta \in A_X$ such that $de^{\boldsymbol{i}_{\xi}}(\omega - \partial\beta) = 0$. For such a β we have

$$0 = e^{-i_{\xi}} de^{i_{\xi}} (\omega - \partial \beta) = -\overline{\partial} \partial \beta + l_{\xi} (\omega) - l_{\xi} (\partial \beta).$$

and so

$$(\overline{\partial} + \boldsymbol{l}_{\xi})\partial\beta = \boldsymbol{l}_{\xi}(\omega).$$

Write $\eta_{\leq m}$ and $\eta_{\geq m}$ for the components of a differential form η in $\bigoplus_{i \leq m} A_X^{i,*}$ and in $\bigoplus_{i \geq m} A_X^{i,*}$, respectively. Since both \boldsymbol{l}_{ξ} and $(\overline{\partial} + \boldsymbol{l}_{\xi})$ are homogeneous of bidegree (0, 1), we have

$$\begin{split} \boldsymbol{l}_{\xi}(\omega) &= (\boldsymbol{l}_{\xi}(\omega))_{\geq m} = ((\overline{\partial} + \boldsymbol{l}_{\xi})\partial\beta)_{\geq m} = (\overline{\partial} + \boldsymbol{l}_{\xi})\partial(\beta_{\geq m-1}).\\ \text{Hence } \psi_{\boldsymbol{i}_{\xi}}[\omega] &= [e^{\boldsymbol{i}_{\xi}}(\omega - \partial(\beta_{\geq m-1}))] \in H^{*}(e^{\boldsymbol{i}_{\xi}}F^{m}), \text{ and so} \end{split}$$

$$\Phi_{\xi}(H^*(F^m)) = H^*(e^{\mathbf{i}_{\xi}}F^m).$$

On the other hand, the period of the infinitesimal deformation $\mathcal{O}_{\xi} = \ker(\overline{\partial} + \mathbf{l}_{\xi})$ is $H^*(F^m_{\xi}) \subseteq H^*(A_X)$, where F^m_{ξ} is the complex of global sections of the differential ideal sheaf $\mathcal{F}^m_{\xi} \subseteq \mathcal{A}_X$ generated by $(d\mathcal{O}_{\xi})^m$. Since $e^{\mathbf{i}_{\xi}}$ is the identity on $\mathcal{A}^{0,0}_X$, by Lemma 2.2 we can write

$$e^{-\boldsymbol{i}_{\xi}}(d\mathcal{O}_{\xi}) = e^{-\boldsymbol{i}_{\xi}}de^{\boldsymbol{i}_{\xi}}\mathcal{O}_{\xi} = (\partial + \overline{\partial} + \boldsymbol{l}_{\xi})\mathcal{O}_{\xi} = \partial\mathcal{O}_{\xi} \subseteq \partial\mathcal{A}_{X}^{0,0} \subseteq \mathcal{A}_{X}^{1,0}.$$

Since $e^{i_{\xi}} \colon \mathcal{A}_X \to \mathcal{A}_X$ is a morphism of sheaves of differential graded commutative algebras, we get $e^{-i_{\xi}}(\mathcal{F}^m_{\xi}) \subseteq \mathcal{F}^m$ and then, by rank considerations, $e^{i_{\xi}}(\mathcal{F}^m) = \mathcal{F}^m_{\xi}$. Hence

$$\Phi_{\xi}(H^*(F^m)) = H^*(\mathcal{F}^m_{\xi})$$

Remark 5.2. It has been shown in [8] that also the m^{th} period map $\text{Def}_X \to \text{Grass}_{H^*(F^m),H^*(X;\mathbb{C})}$ is induced by an L_{∞} -morphism. Namely, let $\chi_m \colon L_m \hookrightarrow \text{Hom}^*(A_X, A_X)$ the inclusion of the subalgebra

$$L_m = \{ f \in \operatorname{Hom}^*(A_X, A_X) \, | \, f(F^m) \subseteq F^m \}$$

in the DGLA of endomorphisms of A_X . Then $\operatorname{Def}_{\chi_m} \simeq \operatorname{Grass}_{H^*(F^m),H^*(X)}$ and the map $\xi \mapsto (\boldsymbol{l}_{\xi}, \boldsymbol{i}_{\xi})$ is an L_{∞} -morphism between KS_X and C_{χ_m} inducing the m^{th} period map.

References

- R. Abraham, J.E. Marsden: Foundations of mechanics. Second edition, Benjamin/Cummings Publishing Co., Inc., Advanced Book Program, Reading, Mass. (1978).
- S. Barannikov, M. Kontsevich: Frobenius manifolds and formality of Lie algebras of polyvector fields. Internat. Math. Res. Notices 4 (1998) 201-215.
- [3] M. Bershadsky, S. Ceccotti, H. Ooguri and C. Vafa: Kodaira-Spencer theory of gravity and exact results for quantum string amplitudes. Comm. Math. Phys. 165 (1994) 311-427.
- [4] F. Catanese: Moduli of algebraic surfaces. Theory of moduli (Montecatini Terme, 1985), 1-83, Lecture Notes in Math. 1337, Springer (1988).
- [5] H. Clemens: Geometry of formal Kuranishi theory. Adv. Math. 198 (2005), no. 1, 311-365.
- [6] B. Fantechi, M. Manetti: Obstruction calculus for functors of Artin rings, I. J. Algebra 202 (1998) 541-576.
- [7] D. Fiorenza, M. Manetti: L_{∞} structures on mapping cones. Algebra Number Theory 1 (2007) 301-330; arXiv:math.QA/0601312.

- [8] D. Fiorenza, M. Manetti: L_{∞} algebras, Cartan homotopies and period maps. Preprint arXiv:math/0605297.
- [9] K. Fukaya: Deformation theory, homological algebra and mirror symmetry. Geometry and physics of branes (Como, 2001), Ser. High Energy Phys. Cosmol. Gravit., IOP Bristol (2003) 121-209. Electronic version available at http://www.math.kyoto-u.ac.jp/%7Efukaya/fukaya.html (may 2006).
- [10] W.M. Goldman, J.J. Millson: The deformation theory of representations of fundamental groups of compact Kähler manifolds. Inst. Hautes Études Sci. Publ. Math. 67 (1988) 43-96.
- [11] W.M. Goldman, J.J. Millson: The homotopy invariance of the Kuranishi space. Illinois J. Math. Math. 34 (1990) 337-367.
- [12] M. Green: Infinitesimal methods in Hodge theory. Algebraic cycles and Hodge theory (Torino, 1993), Lecture Notes in Math. 1594, Springer (1994) 1-92.
- [13] Ph. Griffiths: Periods of integrals on algebraic manifolds. II. Local study of the period mapping. Amer. J. Math. 90 (1968) 805-865.
- [14] D. Iacono: Differential Graded Lie Algebras and Deformations of Holomorphic Maps. PhD Thesis, Roma (2006). arXiv:math.AG/0701091.
- [15] K. Kodaira: Complex manifold and deformation of complex structures. Springer-Verlag (1986).
- [16] M. Kontsevich: Deformation quantization of Poisson manifolds, I. Letters in Mathematical Physics 66 (2003) 157-216; arXiv:q-alg/9709040.
- [17] T. Lada, M. Markl: Strongly homotopy Lie algebras. Comm. Algebra 23 (1995) 2147-2161; arXiv:hep-th/9406095.
- [18] T. Lada, J. Stasheff: Introduction to sh Lie algebras for physicists. Int. J. Theor. Phys. 32 (1993) 1087-1104; arXiv:hep-th/9209099.
- [19] M. Manetti: Deformation theory via differential graded Lie algebras. In Seminari di Geometria Algebrica 1998-1999 Scuola Normale Superiore (1999).
- [20] M. Manetti: Extended deformation functors. Int. Math. Res. Not. 14 (2002) 719-756; arXiv:math.AG/9910071.
- [21] M. Manetti: Cohomological constraint to deformations of compact Kähler manifolds. Adv. Math. 186 (2004) 125-142; arXiv:math.AG/0105175.
- [22] M. Manetti: Lectures on deformations on complex manifolds. Rend. Mat. Appl. (7) 24 (2004) 1-183; arXiv:math.AG/0507286.
- [23] M. Manetti: Lie description of higher obstructions to deforming submanifolds. Ann. Sc. Norm. Super. Pisa Cl. Sci. 6 (2007) 631-659. arXiv:math.AG/0507287
- [24] A. Newlander, L. Nirenberg: Complex analytic coordinates in almost complex manifolds. Ann. of Math. (2) 65 (1957) 391-404.
- [25] M. Schlessinger: Functors of Artin rings. Trans. Amer. Math. Soc. 130 (1968) 208-222.
- [26] A.N. Todorov: The Weil-Peterson geometry of the moduli space of $SU(n \ge 3)$ (Calabi-Yau) manifolds I. Commun. Math. Phys. **126** (1989) 325-346.
- [27] C. Voisin: *Théorie de Hodge et géométrie algébrique complexe I, II.* Société Mathématique de France, Paris (2002).

DIPARTIMENTO DI MATEMATICA "GUIDO CASTELNUOVO", UNIVERSITÀ DI ROMA "LA SAPIENZA", P.LE ALDO MORO 5, I-00185 ROMA ITALY.

E-mail address: fiorenza@mat.uniroma1.it *URL*: www.mat.uniroma1.it/~fiorenza/

E-mail address: manetti@mat.uniroma1.it *URL*: www.mat.uniroma1.it/people/manetti/