

μ K cooling of single nuclear spins in a room temperature diamond

P. Neumann¹, V. Jacques^{1,*}, J. Beck¹, M. Markham², D. Twitchen²,
J. Meijer³, G. Balasubramanian¹, F. Jelezko¹ and J. Wrachtrup^{1,*}

¹*Physikalisches Institut, Universität Stuttgart, 70550 Stuttgart, Germany*

²*Element Six Ltd., King's Ride Park, Ascot, Berkshire SL5 8BP, UK*

³*Central laboratory of ion beam and radionuclides, Ruhr Universität Bochum, Germany*

(Dated: September 1, 2022)

We report a method to efficiently polarized single nuclear spins in diamond, which is based on optical pumping of a single NV color center and mediated by a level-anti crossing in its excited state. A nuclear spin polarization higher than 98% is achieved at room temperature for the ¹⁵N nuclear spin associated to the NV center, corresponding to μ K effective nuclear spin temperature. We then demonstrate simultaneous deterministic initialization of two nuclear spins (¹³C and ¹⁵N) in close vicinity to NV defect, which provide efficient initialization of a three qubit quantum register by including the electron spin. This result appears as an important step toward the development of diamond-based quantum information processing.

PACS numbers:

Nuclear spins are attractive candidates for solid-state quantum information processing because of their extremely long coherence time resulting from a strong isolation from electronic and vibrational mechanisms that usually lead to decoherence in solid-state systems [1, 2]. However, their weak interaction with the environment also makes difficult to isolate and manipulate nuclear spins at the single quantum system level. Indeed, conventional nuclear magnetic resonance techniques require the use of macroscopic nuclear spin ensembles to detect measurable signals for quantum computation [3, 4, 5].

For quantum dots, nuclear spins are present in too high abundance to be addressed independantly. Furthermore, such system require a high fidelity polarization of the nuclear spin bath in order to achieve long electron spin coherence times [6, 7].

Recently, readout of individual ¹³C nuclear spins in diamond has been demonstrated [8, 9]. In such experiments, single nuclear spins are coupled by hyperfine interaction to the electron spin of a single NV color center in diamond, which can be optically polarized and read-out with long coherence time, even at room temperature [10, 11]. By coherently mapping nuclear spin states onto the electron spin of a single NV color center, spectacular solid-state quantum physics experiments have been performed, ranging from the implementation of a nuclear-spin-based quantum register [12], a conditional two-qubit CNOT gate [8], and very recently the generation of Bell and GHZ states with extraordinarily long coherence times [13].

However, most of these experiments [8, 13] were performed without any deterministic polarization of the nuclear spin state. This random initialization unavoidably decreases the success rate of all local operations as $1/2^N$ where N is the number of qubits. Deterministic polarization of the nuclear spins thus appears as a crucial step toward the development of diamond based quantum in-

formation processing.

In Ref. [12], a high degree of polarization over a single nuclear spin in diamond has been demonstrated, using a combination of selective microwave excitation and controlled Larmor precession of the nuclear spin state. In this letter, we demonstrate a new method to polarize single nuclei in diamond which is simply based on optical pumping. A nuclear spin polarization higher than 98% is achieved, corresponding to μ K effective nuclear spin temperature. A nuclear spin polarization mechanism is given to account for such experimental observation, based on a level-anti crossing (LAC) in the excited state of the NV color center. We then demonstrate simultaneous initialization of two nuclear spins in close vicinity to NV defect, which provide efficient initialization of a three qubit quantum register by including the electron spin.

The NV defect center in diamond consists of a substitutional nitrogen atom (N) associated with a vacancy (V) in an adjacent lattice site of the diamond crystalline matrix (Fig. 1-(a)). For the negatively charged NV color center, which is addressed in this study, the ground state is a spin triplet state ³A, with a zero-field splitting $D_{gs} = 2.87$ GHz between spin sublevels $m_s = 0$ and $m_s = \pm 1$. The excited state ³E is also a spin triplet, associated with a broadband photoluminescence emission with zero phonon line at 1.945 eV, which allows optical detection of single NV defects. Symmetry considerations of the NV center shows that a strong electron spin polarization into $m_s = 0$ sublevel is achieved after a few optical excitation-emission cycles. This effect results from spin-selective non-radiative inter-system crossing to a metastable state lying between the ground and excited triplet state [14, 15]. Furthermore, the photoluminescence intensity is higher when the $m_s = 0$ state is populated, allowing optical detection of spin-rotation of a single NV center at room temperature by optically detected magnetic resonance (ODMR) [16].

Recent electron-spin resonance (ESR) experiments have provided new insights into the structure of the excited-state 3E , demonstrating that the excited-state zero-field splitting D_{es} is around 1.42 GHz with the same orientation as in the ground state and that the isotropic g-factor is similar to the ground state g-factor ($g \approx 2$) [17, 18]. Using such values, an excited state LAC between the $m_s = 0$ and $m_s = -1$ sublevels is expected for a magnetic field magnitude on the order of 500 G [17, 18, 19]. In the following, we use this excited-state LAC to efficiently polarize the nuclear spin associated to the nitrogen atom of the NV color center.

We investigate ultra-pure synthetic type IIa diamond crystals prepared using a microwave assisted CVD process. Ensemble EPR experiments have shown that such grown diamond crystals have a single substitutional nitrogen concentration below 1 ppb ($< 1.7 \times 10^{14} \text{ cm}^{-3}$). Furthermore, the initial NV colour center concentration is smaller than 10^{10} cm^{-3} , as determined using confocal photoluminescence microscopy.

Single NV color centers were then artificially created by implanting 7 MeV isotopically pure ${}^{15}\text{N}$ atoms and by annealing the sample for two hours in vacuum at 800 °C [20, 21]. ESR measurements on single NV defects produced by implantation show that all NV defects contain implanted ${}^{15}\text{N}$ atoms. This indicates absence of intrinsic ${}^{14}\text{N}$ atoms in the track of vacancies created by implanted high energy nitrogen ions. Since nitrogen is the major paramagnetic center, we conclude that the sample is free of electron paramagnetic impurities and spin dynamics of NV color centers is mainly governed by nuclear spins present in the diamond lattice, which are mainly ${}^{13}\text{C}$ and ${}^{15}\text{N}$.

NV centers are optically addressed at room temperature using a confocal microscope and their unicity is verified through photon antibunching measurements. The implanted NV color centers are associated to ${}^{15}\text{N}$ isotope which is a $I = 1/2$ nucleus. The energy splitting resulting from hyperfine interaction between this nuclear spin and the electron spin is about 3 MHz in the ground state (see Fig. 1-(b)). ODMR spectra are recorded by applying microwaves, using a copper microwire (20 μm diameter) close to the NV center, and by monitoring the photoluminescence intensity. In addition, a magnetic field (B) is applied along the NV symmetry axis, which corresponds to a [111] crystal axis.

As shown in Fig. 1-(c) (upper trace), two ESR lines are evidenced in ODMR spectra recorded at small magnetic field magnitude, each line being associated to a given orientation of the nuclear spin $|\uparrow\rangle$ at frequency ν_\uparrow and $|\downarrow\rangle$ at frequency ν_\downarrow . By increasing the magnetic field magnitude up to 500 G, which correspond to the excited state LAC, the ESR line at frequency ν_\downarrow disappears, giving an evidence of strong polarization of the nuclear spin in state $|0, \uparrow\rangle$ (see Fig. 1-(c), middle trace).




FIG. 1: (a)-Atomic structure of the Nitrogen-Vacancy (NV) centre in diamond. (b)-Simplified energy-level diagram showing the ground state hyperfine structure associated with ${}^{15}\text{N}$ nuclear spin states $|\uparrow\rangle$ and $|\downarrow\rangle$. The levels associated to electron spin state $m_s = +1$ are not shown. Optical transitions (green arrow) are used to polarize and read-out the electron spin state. (c)-ODMR spectra recorded for different magnitudes of a magnetic field applied along the NV symmetry axis ([111] crystal axis). Away from the excited-state LAC (upper and lower trace), two ESR lines at frequencies ν_\uparrow and ν_\downarrow are evidenced, each related to a given orientation of the nuclear spin. Around the LAC (middle trace), the ESR line at ν_\downarrow vanishes, demonstrating a strong nuclear spin polarization in state $|\uparrow\rangle$. Solid lines are fits of the experimental data using lorentzian functions. (d)- and (e)- Experimental pulse sequence (see main text) used to measure Rabi nutations of the electron spin by applying selective microwave pulse (MW) at frequency ν_\uparrow (blue points) or ν_\downarrow (red points). The experiment is performed for $B = 40$ G (d) and $B = 500$ G (e). Solid lines are data fitting using cosine functions.

The nuclear-spin polarization is measured as :

$$\mathcal{P} = \frac{\mathcal{I}(\nu_\uparrow) - \mathcal{I}(\nu_\downarrow)}{\mathcal{I}(\nu_\uparrow) + \mathcal{I}(\nu_\downarrow)} \quad (1)$$

where $\mathcal{I}(\nu_\uparrow)$ (resp. $\mathcal{I}(\nu_\downarrow)$) is the integral of the ESR peak at frequency ν_\uparrow (resp. ν_\downarrow). By fitting each ESR line with lorentzian functions, we infer a polarization

$\mathcal{P} = 0.98 \pm 0.01$. Owing the nuclear-Zeeman splitting between states $|0, \uparrow\rangle$ and $|0, \downarrow\rangle$ (≈ 200 KHz at $B = 500$ G), such polarization corresponds to a μK effective nuclear spin temperature.

In order to confirm this observation, selective Rabi nutations are performed for each ESR line. For such measurement, the NV center is first initialized into the ground state $m_s = 0$ sublevel using an optical pulse of duration $3 \mu\text{s}$ at wavelength $\lambda = 532$ nm. After a time delay of $1 \mu\text{s}$, which ensures that the NV center has relaxed to the ground state, a selective microwave pulse at frequency ν_\uparrow or ν_\downarrow is applied. The spin-state read-out is realized by measuring the photoluminescence intensity and Rabi nutation can then be observed by varying the duration of the microwave pulse (see Fig. 1-(d)). At low magnetic field amplitude, the contrast of the Rabi nutation is almost equal for each resonance line, demonstrating that the two states $|0, \uparrow\rangle$ and $|0, \downarrow\rangle$ are populated with similar probabilities. Around the excited-state LAC, the contrast associated to state $|0, \downarrow\rangle$ vanished whereas the one associated to state $|0, \uparrow\rangle$ becomes twice higher (see Fig. 1-(e)). This result constitutes another demonstration of strong nuclear spin polarization in state $|0, \uparrow\rangle$.

We now propose a mechanism to account for the observed nuclear-spin polarization. Assuming that the magnetic field B is perfectly aligned along the NV-axis (z -axis) and neglecting the nuclear Zeeman splitting, the excited-state Hamiltonian is given by

$$H = D_{es} \hat{S}_z^2 + g_e \mu_B B \hat{S}_z + \mathcal{A} \hat{S}_z \hat{I} \quad (2)$$

where \hat{S} and \hat{I} are the electron and nuclear spin operators, D_{es} is the excited-state zero-field splitting [23], g_e the electron g-factor, μ_B the Bohr magneton and \mathcal{A} the hyperfine coupling, assumed isotropic. Note that the hyperfine interaction \mathcal{A} is expected to be much stronger in the excited state than in the ground state because the electronic wave function is localized on the nitrogen in the excited-state whereas it is almost entirely on the three carbon dangling orbitals in the ground state [22]. Recent ESR experiments have indeed demonstrated that hyperfine interaction with ^{15}N nuclear spin is $\mathcal{A} \approx 60$ MHz in the excited-state [17]. In the following, we use this value as a fixed parameter.

We first restrict the study to $m_s = 0$ and $m_s = -1$ sublevels. In the basis $\{|0, \uparrow\rangle; |0, \downarrow\rangle; |-1, \uparrow\rangle; |-1, \downarrow\rangle\}$ and by choosing the origin of energy at level $|0, \uparrow\rangle$, the Hamiltonian described by Eq. (2) can be written as

$$H = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & a & 0 \\ 0 & a & \epsilon_{-1}^\uparrow - b & 0 \\ 0 & 0 & 0 & \epsilon_{-1}^\downarrow - b \end{pmatrix} \text{ with } \begin{cases} \epsilon_{-1}^{\uparrow\downarrow} = D_{es} \mp \mathcal{A}/2 \\ b = -g_e \mu_B B \\ a = \mathcal{A}/\sqrt{2} \end{cases}.$$

The eigen states of such Hamiltonian are $|0, \uparrow\rangle$, $|-1, \downarrow\rangle$,




FIG. 2: Nuclear spin polarization mechanism. (a)-Eigen states of the excited-state Hamiltonian described by Eq. (2) showing a LAC at $B_{\text{LAC}} = 497$ G. This simulation is performed using $D_{es} = 1420$ MHz and $\mathcal{A} = 60$ MHz. The insert shows the evolution of parameters α , β and $4\alpha^2\beta^2$ as a function of the magnetic field magnitude. (b)-Away from the LAC, no nuclear spin polarization is achieved as optical transitions ($^3\text{A} \rightarrow ^3\text{E}$) are fully nuclear-spin conserving (black arrows). (c)-At the LAC, precession at frequency Ω between excited-state sublevels $|0, \downarrow\rangle$ and $|-1, \uparrow\rangle$ can lead to nuclear-spin flip, which can be transferred to the ground state through non-radiative inter-system crossing (blue arrow). The result is a strong nuclear-spin polarization in state $|0, \uparrow\rangle$ after a few optical excitation-emission cycles.

$|+\rangle$ and $|-\rangle$ with

$$|+\rangle = \alpha |0, \downarrow\rangle + \beta |-1, \uparrow\rangle \quad (3)$$

$$|-\rangle = \beta |0, \downarrow\rangle - \alpha |-1, \uparrow\rangle \quad (4)$$

The position of the associated eigen energies as well as the values of parameters α and β are represented as a function of the magnetic field magnitude in Fig. 2-(a). This energy diagram shows an excited-state LAC occurring at $B_{\text{LAC}} = 497$ G, corresponding to $b = \epsilon_{-1}^\uparrow$.

At low magnetic field magnitudes, $\alpha \approx 1$ and $\beta \approx 0$. In such regime, optical transitions from the ground state to the excited state are fully nuclear spin-conserving as no state mixing in the excited state is occurring (see Fig. 2-(b)). As a result, the nuclear spin is not polarized.

By increasing the magnetic field magnitude, α and β begin to balance. The transition from $|0, \uparrow\rangle$ to the excited state remains nuclear spin conserving whereas the transition from $|0, \downarrow\rangle$ yields to $(\alpha|+\rangle + \beta|-\rangle)$ in the excited state. This superposition state then starts to precess between $\alpha|+\rangle + \beta|-\rangle = |0, \downarrow\rangle$ and $\alpha|+\rangle - \beta|-\rangle =$




FIG. 3: (a)-Nuclear spin polarization \mathcal{P} as a function of the magnetic field amplitude. For each point the orientation of the magnetic field is kept along the NV symmetry axis and \mathcal{P} is estimated by recording ODMR spectra and using Eq. (1). Fitting the data with Eq. (9) using k_{eq}^0/Γ as single adjustable parameter leads to $k_{eq}^0/\Gamma = 0.009 \pm 0.001$ (red solid line). (b)-Nuclear spin polarization \mathcal{P} as a function of the magnetic field angle, its magnitude being fixed at $B = 472$ G. The solid line is a guide for eyes.

$(\alpha^2 - \beta^2) |0, \downarrow\rangle + 2\alpha\beta | -1, \uparrow\rangle$ at frequency

$$\Omega = \frac{4\pi\sqrt{(b - \epsilon_{-1}^\dagger)^2 + a^2}}{h}, \quad (5)$$

where h is the Planck constant (Fig. 2-(c)).

The maximum probability $p_{max}(B)$ to find the nuclear spin flipped from $|\downarrow\rangle$ to $|\uparrow\rangle$ within this precession is given by $p_{max}(B) = 4\alpha^2\beta^2$, which follows a Lorentzian magnetic field dependence as shown in the insert of Fig. 2-(a). The nuclear spin-flip has then a probability to be transferred to the ground state sublevel $|0, \uparrow\rangle$ by non-radiative inter-system crossing through the metastable singlet state responsible for electron spin polarization of the NV defect [14]. Every subsequent excitation and decay cycles increase spin polarization in state $|0, \uparrow\rangle$. We notice that at the excited-state LAC, m_s is still a good quantum number in the ground state. Indeed, the ground state LAC is known to be around $B = 1024$ G [19].

As long as the precession frequency Ω is on the same order or faster than the excited state decay rate $1/\tau$, the average probability to flip the nuclear spin in the excited-state is $p_-(B) = p_{max}(B)/2$. Using $\tau \approx 12$ ns and $\mathcal{A} \approx 60$ MHz [17], this requirement is fulfilled even at the exact position of the LAC where Ω is minimal. Note that all afore mentioned arguments hold as well for the crossing of the levels $m_s = 0$ and $m_s = +1$ when the sign of the magnetic field is reversed, leading to the probability $p_+(B) = p_-(-B)$ for a spin flip from $|\uparrow\rangle$ to $|\downarrow\rangle$ [24].

Starting with a mixture of $c|0, \uparrow\rangle$ and $d|0, \downarrow\rangle$ in the ground state, the nuclear spin polarization \mathcal{P} defined by

Eq. (1) can be written as

$$\mathcal{P} = \frac{c^2 - d^2}{c^2 + d^2}.$$

As rates of polarisation k_{\pm} and depolarization k_{eq} we assume

$$k_- = d^2 p_- \Gamma = (1 - \mathcal{P}) p_- \Gamma / 2 \quad (6)$$

$$k_+ = -c^2 p_+ \Gamma = -(1 + \mathcal{P}) p_+ \Gamma / 2 \quad (7)$$

$$k_{eq} = -k_{eq}^0 \mathcal{P}. \quad (8)$$

Here Γ is the rate of nuclear-spin conserving intersystem crossing from the excited state to the ground state sublevel $|0, \uparrow\rangle$ and k_{eq}^0 takes into account all forces that are driving the nuclear spin polarization to its equilibrium state. For such simple model, Γ depends on the intensity of the laser which drives the optical transitions. For the steady state, $k_+ + k_- + k_{eq} = 0$, leading to

$$\mathcal{P}(B) = \frac{p_- - p_+}{\frac{2k_{eq}^0}{\Gamma} + p_- + p_+}. \quad (9)$$

The evolution of nuclear-spin polarization as a function of the magnetic field magnitude is depicted on Fig. 3-(a). The data are well fitted using the function defined by Eq. (9) and taking k_{eq}^0/Γ as single adjustable parameter. The dependence of \mathcal{P} with the magnetic field magnitude is broad, demonstrating that even small state mixing in the excited-state leads to nuclear-spin polarization




FIG. 4: ODMR spectrum recorded for a single NV center with a ^{13}C on the first coordination shell for a magnetic field magnitude $B = 60$ G (upper trace). The four ESR lines are associated to the electron-spin transition $m_s = 0 \rightarrow -1$, each line being related to a given orientation of ^{13}C and ^{15}N nuclear spins. By increasing the magnetic field magnitude up to the excited-state LAC, only one ESR line is observed, giving an evidence of strong polarization of both nuclear spins. Red solid lines are data fitting using Lorentzian functions and black dot lines are guides for eyes. Note that the width of the ESR line is bigger at the LAC only because the microwave power was higher for that experiment. Such width is however smaller than the 3 MHz hyperfine splitting related to ^{15}N nuclear spin.

through optical pumping. Note that by saturating the optical transition, the speed of the polarization process is just limited by the parameter Γ which is on the order of 4 MHz according to the metastable state lifetime.

As depicted in Fig. 3-(b), the nuclear spin polarization is however very sensitive to the magnetic field alignment along the NV-axis. Inferring the evolution of the nuclear spin polarization as a function of the magnetic field angle would require more sophisticated numerical simulations of the NV color center spin-dynamics, as all spin-states are partially mixed in that case.

As a final experiment, we demonstrate that the nuclear-spin polarization method can be extended to more than one nuclear spin. Fig. 4 (upper trace) shows the ODMR spectra of a single NV center with a ^{13}C on the first coordination shell, leading to a hyperfine splitting around 130 MHz in the ground state. In such spectrum, four ESR lines are observed, each being associated to a given orientation of ^{13}C and ^{15}N nuclear spins. By increasing the magnetic field up to the excited-state LAC, we observe an efficient polarization of both nuclear spins as only one ESR line remains (see Fig. 4 lower trace), corresponding to a deterministic initialization of two nuclear spins (^{13}C and ^{15}N) in close vicinity to NV defect. From the data we infer a polarization $\mathcal{P} = 0.90 \pm 0.01$ for the double nuclear spin system.

Summarizing, we have demonstrated a new method to strongly polarize single nuclear spins in diamond at room temperature, which is experimentally simple to implement as it is only based on optical pumping without any use of microwave excitation. Furthermore the method could be extended for an arbitrary number of nuclear spins. Such robust control of nuclear spin states will be a key ingredient for further scaling up of nuclear-spin based quantum register in diamond [12, 13].

The authors are grateful to R. Kolesov, L. Rogers and N. B. Manson for many fruitful discussions. We acknowledge financial support by the European Union (QAP, EQUIND, and NEDQIT) and Deutsche Forschungsgemeinschaft (SFB/TR21). V. J. is supported by the Hum-

boldt Foundation.

-
- * Electronic address: v.jacques@physik.uni-stuttgart.de
- [1] N. A. Gershenfeld and I. L. Chuang, *Science* **275**, 350 (1997).
 - [2] B. E. Kane, *Nature* **393**, 133 (1998).
 - [3] L. M. K. Vandersypen *et al.*, *Nature* **414**, 883 (2001).
 - [4] H. G. Krojanski and D. Suter, *Phys. Rev. Lett.* **93**, 090501 (2004).
 - [5] L. M. K. Vandersypen and I. L. Chuang, *Rev. Mod. Phys.* **76**, 1037 (2004).
 - [6] C. W. Lai, P. Maletinsky, A. Badolato, and A. Imamoglu, *Phys. Rev. Lett.* **96**, 167403 (2006).
 - [7] P. Maletinsky, A. Badolato and A. Imamoglu, *Phys. Rev. Lett.* **99**, 056804 (2007).
 - [8] F. Jelezko, T. Gaebel, I. Popa, M. Domhan, A. Gruber, and J. Wrachtrup, *Phys. Rev. Lett.* **93**, 130501 (2004).
 - [9] L. Childress *et al.*, *Science* **314**, 281 (2006).
 - [10] F. Jelezko, T. Gaebel, I. Popa, A. Gruber and J. Wrachtrup, *Phys. Rev. Lett.* **92**, 076401 (2004).
 - [11] T. Gaebel *et al.*, *Nat. Phys.* **2**, 408 (2006).
 - [12] M. V. Gurudev Dutt *et al.*, *Science* **316**, 1312 (2007).
 - [13] P. Neumann *et al.*, *Science* **320**, 1326 (2008).
 - [14] N. B. Manson, J. P. Harrison and M. J. Sellars, *Phys. Rev. B* **74**, 104303 (2006).
 - [15] L. J. Rogers, S. Armstrong, M. J. Sellars and N. B. Manson, *preprint arXiv:0806.0895*.
 - [16] A. Gruber *et al.*, *Science* **276**, 2012 (1997).
 - [17] G. D. Fuchs *et al.*, *preprint arXiv:0806.1939*.
 - [18] P. Neumann *et al.*, *preprint arXiv:0807.2379*.
 - [19] R. J. Epstein, F. M. Mendoza, Y. K. Kato and D. D. Awschalom, *Nat. Phys.* **1**, 94 (2005).
 - [20] J. Meijer *et al.*, *Appl. Phys. Lett.* **87**, 261909 (2005).
 - [21] J. R. Rabreau *et al.*, *Appl. Phys. Lett.* **88**, 023113 (2006).
 - [22] S. Felton *et al.*, *Phys. Rev. B* **77**, 081201(R) (2008).
 - [23] It was recently demonstrated that the fine-structure tensor \underline{D}_{es} is coaxial with the NV-axis and is axially symmetric without any significant E-parameter [18].
 - [24] As the polarized nuclear spin is pointing in the opposite direction for a reversed magnetic field, the ODMR-spectrum still shows the same line polarized but now belonging to the opposite spin orientation (data not shown).