

Contraction in L^1 and large time behavior for a system arising in chemical reactions and molecular motors

Michel Chipot ¹, Danielle Hilhorst ²,
David Kinderlehrer ³, Michał Olech ⁴

Abstract

We prove a contraction in L^1 property for the solutions of a nonlinear reaction–diffusion system whose special cases include intercellular transport as well as reversible chemical reactions. Assuming the existence of stationary solutions we show that the solutions stabilize as t tends to infinity. Moreover, in the special case of linear reaction terms, we prove the existence and the uniqueness (up to a multiplicative constant) of the stationary solution.

Key words: weakly coupled system, molecular motor, transport, parabolic systems, contraction property.

AMS subject classification: 34D23, 35K45, 35K50, 35K55, 35K57, 92C37, 92C45.

1 Introduction

We start with two specific reaction-diffusion systems. The first one describes a reversible reaction and the other one a molecular motor. We first consider the reversible chemical reaction (see also Bothe [4], Bothe and Hilhorst [5], Desvillettes and Fellner [10] and Érdi and Tóth [11]). It involves a reaction-diffusion system of the form

$$\begin{aligned} u_t &= d_1 \Delta u - \alpha k(r_A(u) - r_B(v)) & \text{in } \Omega \times (0, T), \quad \Omega \subset \mathbb{R}^d, \\ v_t &= d_2 \Delta v + \beta k(r_A(u) - r_B(v)) & \text{in } \Omega \times (0, T), \quad \Omega \subset \mathbb{R}^d, \end{aligned} \quad (1.1)$$

¹Angewandte Mathematik, Universitaet Zurich, CH-8057 Zurich

E-mail: m.m.chipot@math.unizh.ch

²Laboratoire de Mathématiques, CNRS and Université de Paris-Sud, 91405 Orsay Cédex, France; E-mail: Danielle.Hilhorst@math.u-psud.fr

³Center for Nonlinear Analysis and Department of Mathematical Sciences, Carnegie Mellon University, Pittsburgh, PA 15213; E-mail: davidk@andrew.cmu.edu

⁴Instytut Matematyczny Uniwersytetu Wrocławskiego, pl. Grunwaldzki 2/4, 50-384 Wrocław, Polska; Laboratoire de Mathématiques, CNRS Université de Paris-Sud, 91405 Orsay Cédex, France; E-mail: olech@math.uni.wroc.pl.

The preparation of this article has been partially supported by a KBN/MNiI grant 1 P03A 008 30 and a Marie Curie Transfer of Knowledge Fellowship of the European Community's Sixth Framework Programme under the contracts MTKD-CT-2004-013389, DMS 0305794, DMS 0405343, and DMS 0806703. We would also like to mention the support of the Swiss National Science Foundation under the contract # 20-113287/1.

together with the homogeneous Neumann boundary conditions, where $d_1, d_2, \alpha, \beta, k$ and T are positive constants and where Ω is a bounded subset of \mathbb{R}^d with smooth boundary. Such systems describe, with a suitable choice of the functions r_A and r_B , chemical reactions for two mobile species. For example, functions $r_A(u) = u^k, r_B(v) = v^m$ correspond to a reversible reaction $kA \rightleftharpoons mB$. Reactions of the type $q_1A_1 + \dots + q_kA_k \rightleftharpoons q_1B_1 + \dots + q_mB_m$ can also be described by similar systems with more complicated reactions terms.

Another model problem is a system in $d = 1$ space dimension and n unknown variables $u_1, \dots, u_n, n > 1$, for intercellular transport, namely

$$\begin{aligned} \frac{\partial u_i}{\partial t} &= \frac{\partial}{\partial x} \left(\sigma \frac{\partial u_i}{\partial x} + u_i \psi'_i \right) \\ &\quad + \sum_{j=1}^n a_{ij} u_j \quad \text{in } Q_T = [0, 1] \times (0, T) \\ \sigma \frac{\partial u_i}{\partial x} + u_i \psi'_i &= 0 \quad \text{on } \partial Q_T = \{0, 1\} \times (0, T), \end{aligned}$$

where

$$\begin{aligned} a_{ii} &\leq 0, \quad a_{ij} \geq 0 \quad \text{for all } i \in \{1, \dots, n\}, i \neq j, \\ \sum_{i=1}^n a_{ij} &= 0 \quad \text{for all } i, j \in \{1, \dots, n\}. \end{aligned} \tag{1.2}$$

It models transport via motor proteins in the eukaryotic cell where chemical energy is transduced into directed motion. A derivation of the system from a mass transport viewpoint is given in [7]. For an analysis of the steady state solutions and for further references we refer to [6], [12], [13], and [20].

In this paper we study the corresponding system in higher space dimension, namely

$$\begin{aligned} \frac{\partial u_i}{\partial t} &= \operatorname{div}(\sigma_i \nabla u_i + u_i \nabla \psi_i) \\ &\quad + \alpha_i \left(\sum_{j=1}^n \lambda_{ij} r_j(u_j(x, t), x) \right) \quad \text{in } Q_T, \end{aligned} \tag{1.3a}$$

where $i \in \{1, \dots, n\}$, and $u_i(x, t) : Q_T \rightarrow \mathbb{R}^+$, with $Q_T = \Omega \times (0, T)$, Ω an open bounded subset of \mathbb{R}^d with smooth boundary, and T some positive constant. We supplement this system with the Robin (no-flux) boundary conditions

$$\sigma_i \frac{\partial u_i}{\partial \nu} + u_i \frac{\partial \psi_i}{\partial \nu} = 0, \quad i \in \{1, \dots, n\}, \quad \text{on } \partial \Omega \times (0, T), \tag{1.3b}$$

where ν is the outward normal vector to $\partial\Omega$, and the initial conditions

$$u_1(x, 0) = u_{0,1}(x), \dots, u_n(x, 0) = u_{0,n}(x), \quad x \in \Omega. \quad (1.3c)$$

We assume that the following hypotheses hold

1. The constants σ_i and $\alpha_i \in \mathbb{R}$, where $i \in \{1, \dots, n\}$, are strictly positive;
2. For $i, j \in \{1, \dots, n\}$, $\lambda_{ii} \leq 0$, $\lambda_{ij} \geq 0$ if $i \neq j$, $\sum_{k=1}^n \lambda_{kj} = 0$;
3. for all $i \in \{1, \dots, n\}$, the smooth functions r_i are nondecreasing with respect to the first variable; $r_i(0, x) = 0$ and we assume that the functions ψ_i are smooth as well;
4. $u_i(\cdot, 0) = u_{0i} \in C(\overline{\Omega})$, $u_{0i} \geq 0$.

In the linear case of the molecular motors, it amounts to choosing

$$r_i(s, x) = s, \quad \lambda_{ij} = a_{ij} \quad \text{and} \quad \alpha_i = 1 \quad \text{for all } i, j \in \{1, \dots, n\}. \quad (1.4)$$

We denote by Problem (P) the system (1.3a) together with the boundary and initial conditions (1.3b), (1.3c), and admit without proof that Problem (P) possesses a unique smooth and bounded solution on each time interval $(0, T]$. An essential idea for proving the existence of a solution would be to apply the Comparison principle Theorem 2.2 below to deduce that any solution of Problem (P) has to be nonnegative and bounded from above by a stationary solution.

Finally, we note that because of the boundary conditions (1.3b) the quantity

$$\sum_{i=1}^n \frac{1}{\alpha_i} \int_{\Omega} u_i(x, t) \, dx \quad (1.5)$$

is conserved in time.

The organization of this paper is as follows. In Section 2 we prove a comparison principle for Problem (P). The main idea, which permits to show that Problem (P) is cooperative, is a change of functions which transforms the Robin boundary conditions into the homogeneous Neumann boundary conditions. In Section 3 we establish a contraction in L^1 property for the corresponding semigroup solution. Let us point out the similarity with an old result due to Crandall and Tartar [8] where they proved in a scalar case that in the presence of a conservation of the integral property such as (1.5), a comparison principle such as Theorem 2.2 is equivalent to a contraction in L^1 property such as the inequality (3.4) below. As far as we know such an abstract result is not known in the case of systems.

Section 4 deals with the large time behavior of the solutions. Supposing the existence of a stationary solution, we construct a continuum of

stationary solutions and prove that the solutions stabilize as t tends to infinity. Let us mention a result by Perthame [19] who proved the stabilization in the case of the two component one-dimensional molecular motor problem. Finally in Section 5, show the existence and uniqueness (up to a multiplicative constant) of the stationary solution of the molecular motor problem.

Acknowledgment: The authors acknowledge the preliminary master thesis work of Aude Brisset about the corresponding two component system. They are grateful to the professors Piotr Biler, Stuart Hastings, Annick Lesne and Hiroshi Matano for very fruitful discussions.

2 Comparison principle

First, we remark that the system of equations (1.3a) is cooperative. However, since nothing is known about the sign of the coefficients $\frac{\partial \psi_i}{\partial \nu}$ in the Robin boundary conditions (1.3b), we cannot decide whether the Problem (P) is cooperative. This leads us to perform a change of variables which transforms the Robin boundary conditions into the homogeneous Neumann boundary conditions.

2.1 The change of unknown functions

Performing the change of variables

$$w_i(x, t) = u_i(x, t) e^{\psi_i(x)/\sigma_i}, \quad i \in \{1, \dots, n\}, \quad (2.1)$$

we deduce from (1.3) that $\vec{w} := (w_1, \dots, w_n)$ satisfies the parabolic problem

$$\begin{aligned} \frac{\partial w_i}{\partial t} &= \sigma_i e^{\psi_i(x)/\sigma_i} \operatorname{div} \left(e^{-\psi_i(x)/\sigma_i} \nabla w_i \right) \\ &+ \alpha_i e^{\psi_i(x)/\sigma_i} \left(\sum_{j=1}^n \lambda_{ij} r_j(w_j(x, t) e^{-\psi_j(x)/\sigma_j}, x) \right) \quad \text{in } Q_T, \end{aligned} \quad (2.2)$$

together with the homogeneous Neumann boundary conditions

$$\frac{\partial w_i}{\partial \nu} = 0, \quad i \in \{1, \dots, n\}, \quad \text{on } \partial\Omega, \quad (2.3)$$

and the initial conditions

$$w_i(x, 0) = u_{0,i}(x) e^{\psi_i(x)/\sigma_i}, \quad i \in \{1, \dots, n\}, \quad x \in \Omega. \quad (2.4)$$

In the following, we denote by Problem P_N — the problem (2.2), (2.3), (2.4). To begin with we define the operators

$$\begin{aligned} \mathcal{L}_i(w_i) &= \frac{\partial w_i}{\partial t} - \sigma_i e^{\psi_i(x)/\sigma_i} \operatorname{div}(e^{-\psi_i(x)/\sigma_i} \nabla w_i) \\ &- \alpha_i e^{\psi_i(x)/\sigma_i} \left(\sum_{j=1}^n \lambda_{ij} r_j(w_j(x, t) e^{-\psi_j(x)/\sigma_j}, x) \right) \quad \text{in } Q_T. \end{aligned} \quad (2.5)$$

We say that $(\underline{w}_1, \dots, \underline{w}_n)$ is a subsolution of Problem P_N if

$$\begin{aligned} \mathcal{L}_i(\underline{w}_i) &\leq 0 \quad \text{in } Q_T, \\ \frac{\partial \underline{w}_i}{\partial \nu} &\leq 0 \quad \text{on } \partial\Omega \times (0, T), \\ \underline{w}_i(x, 0) &\leq w_i(x, 0), \quad x \in \Omega \end{aligned} \quad (2.6)$$

for all $i \in \{1, \dots, n\}$. We define similarly a supersolution $(\overline{w}_1, \dots, \overline{w}_n)$ of Problem P_N by the inequalities

$$\begin{aligned} \mathcal{L}_i(\overline{w}_i) &\geq 0 \quad \text{in } Q_T, \\ \frac{\partial \overline{w}_i}{\partial \nu} &\geq 0 \quad \text{on } \partial\Omega \times (0, T), \\ \overline{w}_i(x, 0) &\geq w_i(x, 0), \quad x \in \Omega. \end{aligned} \quad (2.7)$$

The following comparison theorem holds ([2], [21]).

Theorem 2.1. *Let $(\underline{w}_1, \dots, \underline{w}_n)$ and $(\overline{w}_1, \dots, \overline{w}_n)$, be a sub- and a super-solution, respectively, for the operators \mathcal{L}_j defined by (2.5) with $j \in \{1, \dots, n\}$, which means that (2.6) and (2.7) hold for $i \in \{1, \dots, n\}$. Then $\underline{w}_i \leq \overline{w}_i$ in Q_T . Moreover, for all $i \in \{1, \dots, n\}$ such that $\underline{w}_i \leq \overline{w}_i$ and $\underline{w}_i \not\equiv \overline{w}_i$ on $\{t = 0\} \times \Omega$ then $\underline{w}_i < \overline{w}_i$ in Q_T .* ■

This comparison theorem immediately translates into a comparison theorem for solutions of the original Problem (P). For all $i \in \{1, \dots, n\}$, we define the operators

$$\begin{aligned} L_i(u_i) &= (u_i)_t - \operatorname{div}(\sigma_i \nabla u_i + u_i \nabla \psi_i) \\ &- \alpha_i \left(\sum_{j=1}^n \lambda_{ij} r_j(u_j, x) \right) \quad \text{in } Q_T. \end{aligned} \quad (2.8)$$

The following result holds.

Theorem 2.2. *Let $(\underline{u}_1, \dots, \underline{u}_n)$ and $(\overline{u}_1, \dots, \overline{u}_n)$, be a sub- and a super-solution, respectively, for the operators L_j , defined by (2.8) with $j \in \{1, \dots, n\}$. Then $\underline{u}_i \leq \overline{u}_i$ in Q_T . Moreover, for all $i \in \{1, \dots, n\}$ such that $\underline{u}_i \leq \overline{u}_i$ and $\underline{u}_i \not\equiv \overline{u}_i$ on $\{t = 0\} \times \Omega$ then $\underline{u}_i < \overline{u}_i$ in Q_T .* ■

Next we state two immediate corollaries of Theorem 2.2.

Corollary 2.3. (*uniqueness*) If (u_1^1, \dots, u_n^1) and (u_1^2, \dots, u_n^2) are solutions of Problem (P) with the same initial condition $(u_{0,1}, \dots, u_{0,n}) \in (C(\overline{\Omega}))^n$, then for all $i \in \{1, \dots, n\}$, $u_i^1 = u_i^2$. ■

Corollary 2.4. (*positivity*) If (u_1, \dots, u_n) is the solution of Problem (P) with the nonnegative initial condition $(u_{0,1}, \dots, u_{0,n}) \in (C(\overline{\Omega}))^n$, then for all $i \in \{1, \dots, n\}$, $u_i \geq 0$. Moreover, for all $i \in \{1, \dots, n\}$, such that $u_{0,i} \geq 0$ and $u_{0,i} \neq 0$, $u_i > 0$ in Ω . ■

3 Contraction property

The purpose of this section is to show a contraction in $(L^1(\Omega))^n$ property for the solutions of Problem (P) with the initial conditions belonging to $(L^\infty(\Omega))^n$. The main steps of the proof rely upon arguments due to [3] and [18].

We first introduce some notation. We suppose that the functions (u_1^1, \dots, u_n^1) and (u_1^2, \dots, u_n^2) are the solutions of Problem (P) with the initial conditions $(u_{0,1}^1, \dots, u_{0,n}^1)$ and $(u_{0,1}^2, \dots, u_{0,n}^2)$, respectively. Define

$$(U_1, \dots, U_n) := (u_1^1 - u_1^2, \dots, u_n^1 - u_n^2). \quad (3.1)$$

Then

$$\begin{aligned} (U_i)_t &= \operatorname{div}(\sigma_i \nabla U_i + U_i \nabla \psi_i) \\ &+ \alpha_i \sum_{j=1}^n \lambda_{ij} (r_j(u_j^1(x, t), x) - r_j(u_j^2(x, t), x)) \quad \text{in } Q_T, \\ \sigma_i \frac{\partial U_i}{\partial \nu} + U_i \frac{\partial \psi_i}{\partial \nu} &= 0 \quad \text{on } \partial\Omega \times (0, T), \\ U_i(x, 0) &= U_{0,i}(x) \quad \text{for } x \in \Omega, \end{aligned} \quad (3.2)$$

together with

$$U_{0,i} = u_{0,i}^1 - u_{0,i}^2, \quad (3.3)$$

for each $i \in \{1, \dots, n\}$.

Next we prove the following contraction in L^1 property.

Theorem 3.1. For all $t > 0$,

$$\begin{aligned} \frac{1}{\alpha_1} \|U_1(\cdot, t)\|_{L^1(\Omega)} + \dots + \frac{1}{\alpha_n} \|U_n(\cdot, t)\|_{L^1(\Omega)} \\ \leq \frac{1}{\alpha_1} \|U_{0,1}(\cdot)\|_{L^1(\Omega)} + \dots + \frac{1}{\alpha_n} \|U_{0,n}(\cdot)\|_{L^1(\Omega)}, \end{aligned} \quad (3.4)$$

where U_i and $U_{0,i}$, $i \in \{1, \dots, n\}$, are defined by (3.1) and (3.3), respectively.

Proof Dividing each partial differential equation of (3.2) by α_i and summing them up, we obtain

$$\begin{aligned}
\frac{d}{dt} \left(\sum_{i=1}^n \frac{1}{\alpha_i} U_i \right) &= \sum_{i=1}^n \frac{1}{\alpha_i} \operatorname{div} (\sigma_i \nabla U_i + U_i \nabla \psi_i) \\
&\quad + \sum_{i=1}^n \sum_{j=1}^n \lambda_{ij} \left(r_j(u_j^1(x, t), x) - r_j(u_j^2(x, t), x) \right) \\
&= \sum_{i=1}^n \frac{1}{\alpha_i} \operatorname{div} (\sigma_i \nabla U_i + U_i \nabla \psi_i) \\
&\quad + \sum_{j=1}^n \left\{ \left(r_j(u_j^1(x, t), x) - r_j(u_j^2(x, t), x) \right) \sum_{i=1}^n \lambda_{ij} \right\} \\
&= \sum_{i=1}^n \frac{1}{\alpha_i} \operatorname{div} (\sigma_i \nabla U_i + U_i \nabla \psi_i),
\end{aligned}$$

where we have used Hypothesis 2.

This, together with the boundary conditions (1.3b), implies the conservation in time of the quantity

$$\frac{d}{dt} \sum_{i=1}^n \frac{1}{\alpha_i} \int_{\Omega} U_i(x, t) dx = 0. \quad (3.5)$$

Let us look closer at the nonlinear term in (3.2). We can write, for fixed index i

$$\begin{aligned}
&\sum_{j=1}^n \lambda_{ij} \left(r_j(u_j^1(x, t), x) - r_j(u_j^2(x, t), x) \right) \\
&= \sum_{j=1}^n \lambda_{ij} U_j \int_0^1 \frac{\partial}{\partial u} r_j(\theta u_j^1 + (1 - \theta) u_j^2, x) d\theta = \sum_{j=1}^n A_{ij} U_j.
\end{aligned}$$

Freezing the functions u_i^k for $i \in \{1, \dots, n\}$, $k \in \{1, 2\}$, we deduce that the functions U_1, \dots, U_n satisfy a system of the form

$$(U_i)_t = \operatorname{div} (\sigma_i \nabla U_i + U_i \nabla \psi_i) + \sum_{j=1}^n A_{ij} U_j \quad \text{in } Q_T, \quad (3.6)$$

with the boundary and initial conditions

$$\begin{aligned}
\sigma_i \frac{\partial U_i}{\partial \nu} + U_i \frac{\partial \psi_i}{\partial \nu} &= 0 \quad \text{on } \partial\Omega \times (0, T), \\
U_i(x, 0) &= U_{0,i}(x), \quad x \in \Omega.
\end{aligned} \quad (3.7)$$

for $i \in \{1, \dots, n\}$, where A_{ij} are functions of space and time.

In order to make the notation more concise, we write

$$\begin{aligned}\vec{U}_0 &= (U_{0,1}, \dots, U_{0,n}), \\ \vec{U} &= (U_1, \dots, U_n), \\ \vec{U}_0^\pm &= (U_{0,1}^\pm, \dots, U_{0,n}^\pm), \\ \vec{U}^\pm &= (U_1^\pm, \dots, U_n^\pm),\end{aligned}$$

where $s^+ = \max\{s, 0\}$, $s^- = \max\{-s, 0\}$. By (3.6), (3.7) and Corollary 2.3 we can write \vec{U} in the form

$$(\vec{U})(x, t) = \mathcal{S}(t) \vec{U}_0(x) = (\mathcal{S}_1(t) \vec{U}_0, \dots, \mathcal{S}_n(t) \vec{U}_0)(x)$$

with some operator $\mathcal{S}(t)$. We set

$$(W_1, \dots, W_n) = -(U_1 e^{\psi_1(x)/\sigma_1}, \dots, U_n e^{\psi_n(x)/\sigma_n}),$$

and $\tilde{A}_{ij} = A_{ij} e^{\psi_i(x)/\sigma_i} e^{-\psi_j(x)/\sigma_j}$. Then, the system of equations (3.6) can be expressed in the form

$$(W_i)_t = \sigma_i e^{\psi_i(x)/\sigma_i} \operatorname{div} \left(e^{-\psi_i(x)/\sigma_i} \nabla W_i \right) + \sum_{j=1}^n \tilde{A}_{ij} W_j \quad \text{in } Q_T, \quad (3.8)$$

with the boundary and initial conditions

$$\frac{\partial W_i}{\partial \nu} = 0 \quad \text{on } \partial\Omega \times (0, T), \quad (3.9)$$

$$W_i(x, 0) = -U_{0,i} e^{\psi_i(x)/\sigma_i}, \quad x \in \Omega, \quad (3.10)$$

for $i \in \{1, \dots, n\}$.

Next we show that the solutions W_i of the problem (3.8)–(3.10) with nonpositive initial conditions are nonpositive in $\overline{\Omega}$ for all $t \in (0, T)$. To that purpose we consider the auxiliary problem

$$(W_i)_t - \vartheta_i(x) \operatorname{div} \left(\zeta_i(x) \nabla W_i \right) - \sum_{j=1}^n \gamma_{ij} W_j \leq 0 \quad \text{in } Q_T, \quad (3.11)$$

$$\frac{\partial W_i}{\partial \nu} \leq 0 \quad \text{on } \partial\Omega \times (0, T), \quad (3.12)$$

$$W_i(x, 0) = W_{0,i}(x) \leq 0 \quad x \in \Omega, \quad (3.13)$$

for $i \in \{1, \dots, n\}$. We assume that $\vartheta_i(x)$ and $\zeta_i(x)$ are nonnegative in $\overline{\Omega}$ and that the coefficients γ_{ij} satisfy the same assumptions as the coefficients λ_{ij} in Problem (P). The following result holds.

Lemma 3.2. *Let (W_1, \dots, W_n) be a smooth and bounded solution of the problem (3.11) – (3.13) with nonpositive initial conditions $W_{0,i}$ on a time interval $[0, T]$. Then $W_i(x, t) \leq 0$ in $\overline{\Omega} \times (0, T]$. Moreover, for each $i \in \{1, \dots, n\}$ such that $W_{0,i} \leq 0$ and $W_{0,i} \not\equiv 0$, $W_i < 0$ in $\overline{\Omega} \times (0, T]$.*

Proof The result of Lemma 3.2 follows from the fact that the system (3.11), (3.12), (3.13), with the inequalities $\{\leq\}$ replaced by the equalities $\{=\}$, is a cooperative system. However, for the sake of completeness, we present a proof below. We first remark that, in view of [21, Remark (i), p. 191], one can always satisfy the condition

$$\sum_{j=1}^n \gamma_{ij} \leq 0 \text{ for all } i \in \{1, \dots, n\}, \quad (3.14)$$

for the matrix of coefficients $(\gamma_{ij})_{i,j=1}^n$ by performing the change of variables $\overline{W}_i = W_i e^{-ct}$ for all $i \in \{1, \dots, n\}$ and $c > 0$ large enough. Thanks to the regularity of each W_i , we can apply Theorem 15, p. 191 from [21] to conclude that $W_i - M \leq 0$ in $\overline{\Omega} \times [0, T]$ for some $M > 0$ and all $i \in \{1, \dots, n\}$. In fact, we can deduce that $W_i - M < 0$ in $\overline{\Omega} \times (0, T)$.

Indeed, if for some $k \in \{1, \dots, n\}$, $W_k = M$ in an interior point $(\tilde{x}, \tilde{t}) \in \Omega \times (0, T)$, then Theorem 15, p. 191 in [21] implies that $W_k \equiv M$ for all $0 \leq t < \tilde{t}$, which is impossible since $W_k(x, 0) \leq 0$. If the maximum M of W_k is attained at a boundary point $P \in \partial\Omega \times (0, T)$ then either there exists an open ball $K \subset \Omega \times (0, T)$ such that $P \in \partial K$ and $W_k - M < 0$ in K , and the last part of Theorem 15, p. 191 in [21] contradicts the boundary inequality (3.12), or for all open balls $K \subset \Omega \times (0, T)$ such that $P \in \partial K$ there exists a point $(\tilde{x}, \tilde{t}) \in K$ such that $W_k(\tilde{x}, \tilde{t}) = M$, and we proceed as in the case before.

Hence, there exists $\overline{M} > 0$, such that $W_i \leq \overline{M} < M$ in $\overline{\Omega} \times [0, T]$ for all $i \in \{1, \dots, n\}$. Then we can repeat the reasoning for all $M > 0$ until $M = 0$. Indeed, if this would not be the case, we find the least real number $\overline{M} > 0$, with $W_i \leq \overline{M} \leq \widetilde{M}$ in $\overline{\Omega} \times [0, T]$, which leads again to the existence of a real number $0 \leq \widehat{M} < \overline{M}$ with the same property. This contradicts the fact that \overline{M} was defined as the least such real number. ■

Since the functions u_i^1 , u_i^2 are bounded on $\overline{\Omega} \times [0, T]$, it follows that the functions W_i are bounded on $\overline{\Omega} \times [0, T]$ for all $i \in \{1, \dots, n\}$. Then we are in a position to apply Lemma 3.2 with $\vartheta_i(x) = e^{\psi_i/\sigma_i}$, $\zeta_i(x) = \sigma_i e^{-\psi_i/\sigma_i}$ and $\gamma_{ij} = \widetilde{A}_{ij}$ for $i, j \in \{1, \dots, n\}$. We deduce that the solutions W_i of the problem (3.8)–(3.10) with nonpositive initial conditions are nonpositive in $\overline{\Omega}$ for all $t \in (0, T)$.

Next we remark that the above reasoning can be applied either with \vec{U}_0 replaced by U_0^+ or with \vec{U}_0 replaced by U_0^- . This permits to show that $\mathcal{S}_i(t)\vec{U}_0^+, \mathcal{S}_i(t)\vec{U}_0^- \geq 0$ and that

$$\mathcal{S}_i(t)\vec{U}_0^\pm > 0 \quad \text{if} \quad \vec{U}_0^\pm \neq 0. \quad (3.15)$$

We easily compute

$$\begin{aligned} & \sum_{i=1}^n \frac{1}{\alpha_i} \|U_i(\cdot, t)\|_{L^1(\Omega)} - \sum_{i=1}^n \frac{1}{\alpha_i} \|U_{0,i}(\cdot)\|_{L^1(\Omega)} \\ &= \sum_{i=1}^n \frac{1}{\alpha_i} \|\mathcal{S}_i(t)\vec{U}_0^+ - \mathcal{S}_i(t)\vec{U}_0^-\|_{L^1(\Omega)} - \sum_{i=1}^n \frac{1}{\alpha_i} \|U_{0,i}(\cdot)\|_{L^1(\Omega)} \\ &= \sum_{i=1}^n \int_{\Omega} \frac{1}{\alpha_i} \left\{ \max \{ \mathcal{S}_i(t)\vec{U}_0^+, \mathcal{S}_i(t)\vec{U}_0^- \} \right. \\ & \quad \left. - \frac{1}{\alpha_i} \min \{ \mathcal{S}_i(t)\vec{U}_0^+, \mathcal{S}_i(t)\vec{U}_0^- \} \right\} dx - \sum_{i=1}^n \frac{1}{\alpha_i} \int_{\Omega} \{U_{i,0}^+ + U_{i,0}^-\} dx \\ &= \sum_{i=1}^n \int_{\Omega} \frac{1}{\alpha_i} (\mathcal{S}_i(t)\vec{U}_0^+ + \mathcal{S}_i(t)\vec{U}_0^-) dx - \sum_{i=1}^n \frac{1}{\alpha_i} \int_{\Omega} \{U_{i,0}^+ + U_{i,0}^-\} dx \\ & \quad - 2 \sum_{i=1}^n \int_{\Omega} \frac{1}{\alpha_i} \min \{ \mathcal{S}_i(t)\vec{U}_0^+, \mathcal{S}_i(t)\vec{U}_0^- \} dx \\ &= -2 \sum_{i=1}^n \int_{\Omega} \frac{1}{\alpha_i} \min \{ \mathcal{S}_i(t)\vec{U}_0^+, \mathcal{S}_i(t)\vec{U}_0^- \} dx \leq 0, \end{aligned} \quad (3.16)$$

$$\begin{aligned} & - \frac{1}{\alpha_i} \min \{ \mathcal{S}_i(t)\vec{U}_0^+, \mathcal{S}_i(t)\vec{U}_0^- \} \} dx - \sum_{i=1}^n \frac{1}{\alpha_i} \int_{\Omega} \{U_{i,0}^+ + U_{i,0}^-\} dx \\ &= \sum_{i=1}^n \int_{\Omega} \frac{1}{\alpha_i} (\mathcal{S}_i(t)\vec{U}_0^+ + \mathcal{S}_i(t)\vec{U}_0^-) dx - \sum_{i=1}^n \frac{1}{\alpha_i} \int_{\Omega} \{U_{i,0}^+ + U_{i,0}^-\} dx \\ & \quad - 2 \sum_{i=1}^n \int_{\Omega} \frac{1}{\alpha_i} \min \{ \mathcal{S}_i(t)\vec{U}_0^+, \mathcal{S}_i(t)\vec{U}_0^- \} dx \\ &= -2 \sum_{i=1}^n \int_{\Omega} \frac{1}{\alpha_i} \min \{ \mathcal{S}_i(t)\vec{U}_0^+, \mathcal{S}_i(t)\vec{U}_0^- \} dx \leq 0, \end{aligned} \quad (3.17)$$

which completes the proof of (3.4). ■

Corollary 3.3. *Let $(u_{0,1}^1, \dots, u_{0,n}^1), (u_{0,1}^2, \dots, u_{0,n}^2) \in (C(\overline{\Omega}))^n$ be as in Theorem 3.1. Moreover, let us assume that for at least one index $k \in \{1, \dots, n\}$ the difference $u_{0,k}^1 - u_{0,k}^2$ changes the sign. Then, the inequality (3.4) is strict for all $t > 0$, so that solution satisfies a strict contraction property.*

4 Large time behavior of solutions

In this section we assume the existence and uniqueness of a positive solution $\vec{v} = (v_1, \dots, v_n) \in (C(\overline{\Omega}) \cap C^2(\Omega))^n$ of the elliptic problem

$$\operatorname{div}(\sigma_i \nabla v_i + v_i \nabla \psi_i) + \alpha_i \left(\sum_{j=1}^n \lambda_{ij} r_j(v_j(x), x) \right) = 0 \quad \text{in } \Omega, \quad (4.1)$$

$$\sigma_i \frac{\partial v_i}{\partial \nu} + v_i \frac{\partial \psi_i}{\partial \nu} = 0 \quad \text{on } \partial\Omega, \quad (4.2)$$

$$\sum_{i=1}^n \frac{1}{\alpha_i} \int_{\Omega} v_i(x) \, dx = 1, \quad (4.3)$$

for $i \in \{1, \dots, n\}$.

Definition 4.1. *We say that a vector function $\vec{v} = (v_1, \dots, v_n) \in (C(\overline{\Omega}))^n$ is nonnegative (resp. positive) if $v_i(x) \geq 0$ (resp. $v_i(x) > 0$) for all $x \in \overline{\Omega}$ and all $i \in \{1, \dots, n\}$.*

Next we introduce the semigroup notation for the unique solution of Problem (P), namely

$$\vec{u}(t) = \mathcal{T}(t) \vec{u}_0 = \left(\mathcal{T}_1(t) \vec{u}_0, \dots, \mathcal{T}_n(t) \vec{u}_0 \right),$$

with the initial data $\vec{u}_0 \in (C(\overline{\Omega}))^n$. The method of the proof is based upon an idea of Osher and Ralston [18]. It mainly exploits the contraction properties for the nonlinear semigroup $\mathcal{T}(t)$ given by Theorem 3.1 and Corollary 3.3. A similar reasoning was developed in other contexts by Bertsch and Hilhorst [3], Hilhorst and Hulshof [14] and Hilhorst and Peletier [15].

We suppose there exists a set $\mathcal{H} \subset (C(\overline{\Omega}) \cap C^2(\Omega))^n$ of positive stationary solutions with the following property which we denote by \mathcal{S} :

For each $\vec{f} = (f_1, \dots, f_n) \in (C(\overline{\Omega}) \cap C^2(\Omega))^n$ either $\vec{f} \in \mathcal{H}$ or there exists $(\xi_1, \dots, \xi_n) \in \mathcal{H}$, such that $f_i - \xi_i$ changes the sign for at least one index $i \in \{1, \dots, n\}$.

One can prove that a set \mathcal{H} satisfying Property \mathcal{S} exists in at least two cases:

- i)* In the case of the system (1.1) where the Robin boundary conditions reduce to the homogeneous Neumann boundary conditions,

the set \mathcal{H} is given by

$$\mathcal{H} = \left\{ (a, b) : a > 0, b = r_B^{-1}(r_A(a)) \right. \\ \left. \text{and } \frac{a}{\alpha} + \frac{b}{\beta} = \int_{\Omega} \left(\frac{u}{\alpha} + \frac{v}{\beta} \right) dx \right\}.$$

For more details we refer to [5].

ii) In the case of the molecular motor with a linear n -component system the set \mathcal{H} is given by

$$\mathcal{H} = \{c\vec{v} : c \in \mathbb{R}^+\},$$

where \vec{v} is a unique solution of the elliptic problem (4.1)–(4.3).

Proposition 4.2. *The continuum \mathcal{H} is such that for each*

$$\vec{f} = (f_1, \dots, f_n) \in (C(\overline{\Omega}) \cap C^2(\Omega))^n$$

either $\vec{f} \in \mathcal{H}$, or there exists $(\xi_1, \dots, \xi_n) \in \mathcal{H}$ such that $f_i - \xi_i$ changes the sign for at least one index $i \in \{1, \dots, n\}$.

Proof

- i) In the case of system (1.1) the proof is rather obvious since the continuum \mathcal{H} is composed of constant pairs.
- ii) In the case of the molecular motor, let us assume that $\vec{f} \notin \mathcal{H}$. Then there does not exist any positive constant c such that $c\vec{v} = \vec{f}$. In particular, there exists an index $i \in \{1, \dots, n\}$ such that v_i is not proportional to f_i , or in other words $cv_i \neq f_i$ for all $c > 0$. Without loss of generality we can assume that the first coordinate has this property. Let $x_0 \in \Omega$ be arbitrary. Since v_1 is strictly positive in $\overline{\Omega}$, we can define

$$c_0 = \frac{f_1(x_0)}{v_1(x_0)},$$

so that

$$(f_1 - c_0 v_1)(x_0) = 0.$$

Let $\mathcal{Z} = \{x \in \overline{\Omega} : (f_1 - c_0 v_1)(x) = 0\}$. From the continuity of f_1 and v_1 , \mathcal{Z} is closed as a subset of Ω . If there exist $x_1, x_2 \in \mathcal{Z}^c$, such that $(f_1 - c_0 v_1)(x_1)$ and $(f_1 - c_0 v_1)(x_2)$ are of different signs, then the proof is complete. Now suppose that $(f_1 - c_0 v_1)(x)$ is positive for all $x \in \mathcal{Z}^c$. In particular

$$(f_1 - c_0 v_1)(\tilde{x}) = d > 0$$

for some fixed $\tilde{x} \in \mathcal{Z}^c$. Then choosing $\varepsilon = \frac{d}{2v_1(\tilde{x})}$ we see that

$$(f_1 - (c_0 + \varepsilon)v_1)(\tilde{x}) = \frac{d}{2} > 0.$$

However

$$(f_1 - (c_0 + \varepsilon)v_1)(x_0) < 0.$$

We proceed similarly when $(f_1 - c_0 v_1)(x)$ is negative for all $x \in \mathcal{Z}^c$. ■

In the sequel we suppose that the initial data $\vec{u}_0 = (u_{0,1}, \dots, u_{0,n})$ from $(C(\overline{\Omega}))^n$ also satisfy the following property:

$$\text{There exists } \vec{h} \in \mathcal{H} \text{ such that } 0 \leq \vec{u}_0 \leq \vec{h} \text{ in } \overline{\Omega}, \quad (4.4)$$

and remark that this property is satisfied in both the cases (i) and (ii).

Proposition 4.3. *Let $\vec{u}_0 = (u_{0,1}, \dots, u_{0,n}) \in (C(\overline{\Omega}))^n$ satisfy the property (4.4). Then the solution (u_1, \dots, u_n) of Problem (P) is such that $0 \leq \vec{u}(t) \leq \vec{h}$ for all $t > 0$.*

Proof We remark that $\vec{0}$ is a subsolution of Problem (P) and that \vec{h} is a supersolution, and apply Theorem 2.2. ■

Next we prove the main result of this section. To that purpose we first define the norm $\|\cdot\|_1$ by

$$\|\vec{f}\|_1 = \sum_{i=1}^n \frac{1}{\alpha_i} \|f_i\|_{L^1(\Omega)}.$$

Note that this norm is equivalent to the usual product norm in the space $(L^1(\Omega))^n$.

Theorem 4.4. *For all nonnegative $\vec{u}_0 = (u_{0,1}, \dots, u_{0,n}) \in (C(\overline{\Omega}))^n$ there exists $\vec{f} = (f_1, \dots, f_n) \in \mathcal{H}$, such that*

$$\lim_{t \rightarrow \infty} \|\mathcal{T}(t)\vec{u} - \vec{f}\|_1 = 0.$$

Proof

The proof consists of several steps. To begin with we define the ω -limit set

$$\omega(\vec{u}_0) = \left\{ \vec{g} \in (L^1(\Omega))^n : \text{there exists a sequence } t_k \rightarrow \infty \text{ as } k \rightarrow \infty, \text{ such that } \lim_{k \rightarrow \infty} \|\mathcal{T}(t_k)\vec{u}_0 - \vec{g}\|_1 = 0 \right\}, \quad (4.5)$$

The organization of the proof is as follows. First we show that $\omega(\vec{u}_0)$ is not empty. In the second step we define the Lyapunov functional

$$\mathcal{V}(\vec{\xi}) = \|\vec{\xi} - \vec{w}\|_{\mathbf{1}},$$

where \vec{w} is a stationary solution and check that it is constant on $\omega(\vec{u}_0)$. We then deduce that $\omega(\vec{u}_0) \subset \mathcal{H}$, and finally prove that $\omega(\vec{u}_0)$ consists of exactly one function.

Step 1. $\omega(\vec{u}_0)$ is not empty.

Let $\varepsilon > 0$ be arbitrary. Suppose that $\Omega' \subset\subset \Omega$ satisfy

$$|\Omega \setminus \Omega'| \leq \frac{\varepsilon}{2K}.$$

and set

$$K = \sum_{i=1}^n \frac{2}{\alpha_i} \|h_i\|_{C(\bar{\Omega})}, \quad (4.6)$$

where \vec{h} has been introduced in (4.4). We have already proved in Proposition 4.3 that $\mathcal{T}(t)\vec{u}_0$ is bounded in $(L^\infty(\Omega))^n$. Therefore there exist a vector function $\vec{g} \in (L^\infty(\Omega))^n$ and a sequence $\{\vec{u}(t_k)\}$ such that

$$\vec{u}(t_k) \rightharpoonup \vec{g} \text{ weakly in } (L^2(\Omega))^n, \quad (4.7)$$

as $t_k \rightarrow \infty$. Next we deduce from [16, Chap. III, Theorem 10.1] that there exists a positive constant C such that

$$|u_i(x_1, t) - u_i(x_2, t)| \leq C|x_1 - x_2|^\alpha$$

for all $x_1, x_2 \in \Omega'$ and all $t > 0$. Therefore, it follows from the Ascoli-Arzelà Theorem (see, e.g., [1, Theorem 1.33]) that $\vec{u}(t_k) \rightarrow \vec{g}$ as $t_k \rightarrow \infty$, uniformly in $\bar{\Omega}'$. We choose t_0 large enough such that for all $t_k \geq t_0$

$$\|\vec{u}(\cdot, t_k) - \vec{g}(\cdot)\|_{\mathbf{1}, \Omega'} \leq \frac{\varepsilon}{2}, \quad (4.8)$$

where $\|\cdot\|_{\mathbf{1}, \Omega'}$ corresponds to the L^1 norm in Ω' . We deduce that, in view of (4.6) and (4.7) that

$$\|\vec{u}(\cdot, t_k) - \vec{g}(\cdot)\|_{\mathbf{1}, \Omega \setminus \Omega'} \leq K|\Omega \setminus \Omega'| \leq \frac{\varepsilon}{2},$$

which together with (4.8) yields

$$\|\vec{u}(\cdot, t_k) - \vec{g}(\cdot)\|_{\mathbf{1}} \leq \varepsilon.$$

Step 2. $\omega(\vec{u}_0) \subset \mathcal{H}$.

Indeed, let $\vec{g} \in \omega(\vec{u}_0)$ and suppose $\vec{g} \notin \mathcal{H}$. According to Proposition

4.2 we can find a steady state solution $\vec{w} \in \mathcal{H}$, such that at least one component of $\vec{w} - \vec{g}$ changes the sign. Without loss of generality we can assume that it happens for the first component, namely that $f_1 - w_1$ changes the sign. We remark that, by the contraction property in Theorem 3.1, the functional

$$\mathcal{V}(\vec{\xi}) = \|\vec{\xi} - \vec{w}\|_{\mathbf{1}}$$

is a Lyapunov functional for Problem (P), where $\vec{\xi} \in (L^1(\Omega))^n$. Next we describe some of its properties.

Property (a) *The functional \mathcal{V} is constant on $\omega(\vec{u}_0)$.*

Since $\mathcal{T}(t)\vec{w} = \vec{w}$ and $\mathcal{T}(t)$ has the contraction property (3.4), the functional \mathcal{V} is nonincreasing in time along the trajectory $\mathcal{T}(t)\vec{u}_0$, which yields

$$\begin{aligned} \mathcal{V}(\mathcal{T}(t)\vec{u}_0) &= \|\mathcal{T}(t)\vec{u}_0 - \vec{w}\|_{\mathbf{1}} \\ &= \|\mathcal{T}(t)\vec{u}_0 - \mathcal{T}(t)\vec{w}\|_{\mathbf{1}} \leq \|\vec{u}_0 - \vec{w}\|_{\mathbf{1}} < \infty . \end{aligned}$$

Thus there exists a finite limit \mathcal{V}^* of $\mathcal{V}(\mathcal{T}(t)\vec{u}_0)$ as $t \rightarrow \infty$. Let $\vec{h}_1, \vec{h}_2 \in \omega(\vec{u}_0)$. We can find a sequence $t_k \rightarrow \infty$ as $k \rightarrow \infty$, such that

$$\|\mathcal{T}(t_{2k})\vec{u}_0 - \vec{h}_1\|_{\mathbf{1}} \rightarrow 0 \quad \text{and} \quad \|\mathcal{T}(t_{2k+1})\vec{u}_0 - \vec{h}_2\|_{\mathbf{1}} \rightarrow 0,$$

as k tends to ∞ . It follows that $\mathcal{V}(\vec{h}_1) = \mathcal{V}(\vec{h}_2) = \mathcal{V}^*$.

Property (b) *The ω -limit set $\omega(\vec{u}_0)$ is invariant with respect to the semigroup $\mathcal{T}(t)$, namely if $\vec{h} \in \omega(\vec{u}_0)$, then for all $t > 0$ also $\mathcal{T}(t)\vec{h} \in \omega(\vec{u}_0)$.*

Let the sequence $t_k \rightarrow \infty$ as $k \rightarrow \infty$ be such that $\|\mathcal{T}(t_k)\vec{u}_0 - \vec{h}\|_{\mathbf{1}} \rightarrow 0$. From the contraction property (3.4)

$$\begin{aligned} \|\mathcal{T}(t_k + t)\vec{u}_0 - \mathcal{T}(t)\vec{h}\|_{\mathbf{1}} &= \|\mathcal{T}(t)\mathcal{T}(t_k)\vec{u}_0 - \mathcal{T}(t)\vec{h}\|_{\mathbf{1}} \\ &\leq \|\mathcal{T}(t_k)\vec{u}_0 - \vec{h}\|_{\mathbf{1}}. \end{aligned}$$

Since the last term above tends to 0 as k tends to ∞ this shows that $\mathcal{T}(t)\vec{h} \in \omega(\vec{u}_0)$.

Now, remember that $\vec{g} \in \omega(\vec{u}_0)$ is such that $\vec{g} \notin \mathcal{H}$ and $\vec{w} \in \mathcal{H}$ is such that the first component of $\vec{w} - \vec{g}$ changes the sign in Ω . Then, Corollary 3.3 yields

$$\begin{aligned} \mathcal{V}(\mathcal{T}(t)\vec{g}) &= \|\mathcal{T}(t)\vec{g} - \vec{w}\|_{\mathbf{1}} \\ &= \|\mathcal{T}(t)\vec{g} - \mathcal{T}(t)\vec{w}\|_{\mathbf{1}} < \|\vec{g} - \vec{w}\|_{\mathbf{1}} = \mathcal{V}(\vec{g}), \end{aligned}$$

for all $t > 0$, which contradicts Property (a). Therefore $\vec{g} \in \mathcal{H}$.

Step 3. *The set $\omega(\vec{u}_0)$ contains only one element.*

Suppose that $\vec{g}_1, \vec{g}_2 \in \omega(\vec{u}_0)$. Then we can find two sequences t_k, s_k tending to ∞ as $k \rightarrow \infty$, such that $s_k \leq t_k$ and $\|\mathcal{T}(t_k) \vec{u}_0 - \vec{g}_1\|_1, \|\mathcal{T}(s_k) \vec{u}_0 - \vec{g}_2\|_1 \rightarrow 0$ as $t_k \rightarrow \infty$. Since $\omega(\vec{u}_0) \subset \mathcal{H}$, it follows that

$$\begin{aligned} \|\vec{g}_1 - \vec{g}_2\|_1 &\leq \|\mathcal{T}(t_k) \vec{u}_0 - \vec{g}_1\|_1 + \|\mathcal{T}(t_k) \vec{u}_0 - \vec{g}_2\|_1 \\ &= \|\mathcal{T}(t_k) \vec{u}_0 - \vec{g}_1\|_1 + \|\mathcal{T}(t_k - s_k) \mathcal{T}(s_k) \vec{u}_0 - \mathcal{T}(t_k - s_k) \vec{g}_2\|_1 \\ &\leq \|\mathcal{T}(t_k) \vec{u}_0 - \vec{g}_1\|_1 + \|\mathcal{T}(s_k) \vec{u}_0 - \vec{g}_2\|_1, \end{aligned}$$

which tends to 0 as $k \rightarrow \infty$. ■

5 Stationary solutions for the linear molecular motor problem

In this section we show the existence and the uniqueness (up to a multiplicative constant) of the classical stationary solution of the problem for the molecular motor. We suppose that Ω is an open bounded subset of \mathbb{R}^d with smooth boundary $\partial\Omega$.

We consider the linear system

$$\operatorname{div}(\sigma_i \nabla v_i(x) + v_i(x) \nabla \psi_i(x)) + \sum_{j=1}^n a_{ij} v_j(x) = 0 \quad \text{in } \Omega, \quad (5.1)$$

where $i \in \{1, \dots, n\}$, $n > 1$. The system (5.1) is supplemented with the Robin boundary conditions

$$\sigma_i \frac{\partial v_i}{\partial \nu} + v_i \frac{\partial \psi_i}{\partial \nu} = 0 \quad \text{on } \partial\Omega, \quad (5.2)$$

where $i \in \{1, \dots, n\}$. Thus, the problem can be written as

$$\mathcal{A}\vec{v} = 0,$$

with a linear operator \mathcal{A} in a suitable Banach space \mathcal{X} of functions on Ω , to be made precise later. Moreover, we impose the integral constraint

$$\sum_{i=1}^n \int_{\Omega} v_i(x) dx = 1. \quad (5.3)$$

The adjoint problem $\mathcal{A}^* \vec{\varphi} = 0$ to (5.1), in a dual space \mathcal{X}^* , is now

$$\sigma_i \Delta \varphi_i - \nabla \psi_i \cdot \nabla \varphi_i + \sum_{j=1}^n a_{ji} \varphi_j = 0, \quad \text{in } \Omega, \quad (5.4)$$

with the Neumann boundary conditions for each $i = 1, \dots, n$

$$\frac{\partial \varphi_i}{\partial \nu} = 0 \quad \text{on} \quad \partial\Omega. \quad (5.5)$$

Since $\sum_{j=1}^n a_{ji} = 0$, the problem (5.4) has the obvious solution

$$\vec{\varphi} = (\varphi_1, \dots, \varphi_n) = (1, \dots, 1). \quad (5.6)$$

We are going to apply the Krein-Rutman theorem on the first eigenvalues and eigenvectors of positive operators, and this will permit us to conclude that the problem (5.1)–(5.2) has a one-dimensional space of solutions. Therefore, under the additional constraint (5.3), the original problem (5.1)–(5.2) has a unique solution.

Perthame and Souganidis sketched this argument for $n > 1$ and $d = 1$ in [20].

Theorem 5.1. *Under the assumption $\sum_{j=1}^n a_{ji} = 0$, there exists a unique smooth solution \vec{v} of the system (5.1)–(5.3).*

Before proving Theorem 5.1 we recall some basic definitions as well as the Krein-Rutman theorem from [9, Ch. VIII, p. 188–191].

Definition 5.2 (Reproducing cone). *We say that a closed set K in \mathcal{X} is a cone, if it possesses the following properties:*

- i) $0 \in K$,
- ii) $u, v \in K \implies \alpha u + \beta v \in K$, for all $\alpha, \beta \geq 0$,
- iii) $v \in K$ and $-v \in K \implies v = 0$.

A cone $K \subset \mathcal{X}$ is said to be reproducing if $\mathcal{X} = K - K \equiv \{k_1 - k_2 : k_1, k_2 \in K\}$.

Definition 5.3 (Dual cone). *If K is a cone in \mathcal{X} , then the set $K^* \subset \mathcal{X}^*$ is said to be a dual cone if*

$$\langle f^*, v \rangle \geq 0,$$

for every $v \in K$.

Definition 5.4 (Strict positivity). *Let \mathcal{B} be a linear operator on \mathcal{X} . Then \mathcal{B} is said to be strongly positive if $\mathcal{B}v \in K^\circ$ for all $v \in K$ such that $v \neq 0$.*

Theorem 5.5. *Let K be a reproducing cone in a Banach space \mathcal{X} , with nonempty interior $K^\circ \neq \emptyset$, and let \mathcal{B} be a strongly positive compact operator on K in a sense of Definition 5.4. Then the spectral radius of \mathcal{B} , $r(\mathcal{B})$, is a simple eigenvalue of \mathcal{B} and \mathcal{B}^* , and their associated eigenvectors belong to K° and $(K^*)^\circ$. More precisely, there exists a unique associated eigenvector in K° (resp. $(K^*)^\circ$) of norm 1. Furthermore, all other eigenvalues are strictly less in absolute value than $r(\mathcal{B})$.*

Proof We will apply Theorem 5.5 to the space $\mathcal{X} = (C(\overline{\Omega}))^n \subset (L^1(\Omega))^n$ endowed with the usual supremum norm, and the operators

$$\begin{aligned}\mathcal{B} &= (\lambda I - \mathcal{A})^{-1} : \mathcal{X} \rightarrow \mathcal{X}, \\ \mathcal{B}^* &= (\lambda I - \mathcal{A}^*)^{-1} : \mathcal{X}^* \rightarrow \mathcal{X}^*,\end{aligned}$$

where $\lambda > 0$ is a strictly positive real number to be fixed later.

Let

$$K = \{\vec{u} \in \mathcal{X} : u_i(x) \geq 0 \text{ for each } x \in \overline{\Omega}, i = 1, \dots, n\}.$$

We remark that K is a reproducing cone, with nonempty interior

$$K^o = \{\vec{u} \in \mathcal{X} : \inf_{x \in \overline{\Omega}} u_i(x) > 0, i = 1, \dots, n\}.$$

From the standard theory [17, Theorem 2.1 and Theorem 3.1, Ch. 7] for elliptic partial differential linear systems, the boundary value problem

$$\sigma_i \Delta \varphi_i - \nabla \psi_i \cdot \nabla \varphi_i + \sum_{j=1}^n a_{ji} \varphi_j - \lambda \varphi_i = f_i \text{ in } \Omega, \quad (5.7)$$

with the homogeneous Neumann conditions (5.5) on $\partial\Omega$, for $\lambda = \tilde{\lambda} > 0$ sufficiently large, has a solution $\vec{\varphi} = (\varphi_1, \dots, \varphi_n) \in \mathcal{X}$ for each $\vec{f} = (f_1, \dots, f_n) \in \mathcal{X}$. Moreover, if $f_i(x) \geq 0$ for each $i = 1, \dots, n$, and $x \in \overline{\Omega}$, then $\varphi_i(x) \geq 0$ (in fact, $\varphi_i(x) > 0$ in Ω), which is a consequence of the maximum principle (cf. also Example 3 on p. 196–197 in [9]). Thus, the operator $\mathcal{B}^* = (\tilde{\lambda} I - \mathcal{A}^*)^{-1}$ is a strongly positive and compact operator, and by Theorem 5.5, the largest eigenvalue μ of \mathcal{B} and \mathcal{B}^* is simple.

Since

$$\begin{aligned}-\sigma_i \Delta \varphi_i + \nabla \psi_i \cdot \nabla \varphi_i - \sum_{j=1}^n a_{ji} \varphi_j + \tilde{\lambda} \varphi_i &= \tilde{\lambda} \varphi_i \text{ in } \Omega \\ \frac{\partial \varphi_i}{\partial \nu} &= 0 \text{ on } \partial\Omega,\end{aligned}$$

for all $i \in \{1, \dots, n\}$, with $\vec{\varphi} = (\varphi_1, \dots, \varphi_n) = (1, \dots, 1)$, and since $(1, \dots, 1) \in (K^*)^o$, it follows that $\frac{1}{\tilde{\lambda}} = r\left((\tilde{\lambda} I - \mathcal{A}^*)^{-1}\right)$ is a simple eigenvalue of the operator $(\tilde{\lambda} I - \mathcal{A}^*)^{-1}$. Applying again Theorem 5.5, we deduce that $\frac{1}{\tilde{\lambda}}$ is the largest eigenvalue of the operator $(\tilde{\lambda} I - \mathcal{A})^{-1}$ and that it is simple, and that there exists $\vec{v} \in K^o \subset \mathcal{X}$ such that

$$(\tilde{\lambda} I - \mathcal{A})^{-1} \vec{v} = \frac{1}{\tilde{\lambda}} \vec{v},$$

which is equivalent to

$$A\vec{v} = 0.$$

This proves the existence of the solution of the problem (5.1)–(5.3).

■

References

- [1] R. A. Adams, Sobolev Spaces, *Pure and Applied Mathematics*, vol. 65, Academic Press, New York-London, 1975.
- [2] N. D. Alikakos, P. Hess, H. Matano, Discrete order preserving semigroups and stability for periodic parabolic differential equations, *J. Differential Equations*, 1989, 2, 82, 322–341.
- [3] M. Bertsch and D. Hilhorst, A density dependent diffusion equation in population dynamics: stabilization to equilibrium, *SIAM J. Math. Anal.*, 1986, 4, 17, 863–883.
- [4] D. Bothe, Instantaneous limits of reversible chemical reactions in presence of macroscopic convection, *J. Differential Equations*, 2003, 1, 193, 27–48.
- [5] D. Bothe and D. Hilhorst, A reaction-diffusion system with fast reversible reaction, *J. Math. Anal. Appl.*, 2003, 1, 268, 125–135.
- [6] M. Chipot, S. Hastings, and D. Kinderlehrer, Transport in a molecular motor system, *M2AN Math. Model. Numer. Anal.*, 2004, 38, 6, 1011–1034.
- [7] M. Chipot, D. Kinderlehrer, and M. Kowalczyk, 2004 *A variational principle for molecular motors*, *Meccanica*, **38**, 505–518
- [8] M. G. Crandall and L. Tartar, Some relations between nonexpansive and order preserving mappings, *Proc. Amer. Math. Soc.*, 1980, 3, 78, 385–390.
- [9] R. Dautray, J.-L. Lions, *Mathematical Analysis and Numerical Methods for Science and Technology*, Volume 3: Spectral Theory and Applications, Springer-Verlag, Berlin, 1990.
- [10] L. Desvillettes and K. Fellner, Exponential decay toward equilibrium via entropy methods for reaction-diffusion equations, *J. Math. Anal. Appl.*, 2006, 1, 319, 157–176.
- [11] P. Érdi and J. Tóth, *Mathematical Models of Chemical Reactions*, *Nonlinear Science: Theory and Applications*, Princeton University Press, Princeton, NJ, 1989.

- [12] Hastings, S., Kinderlehrer, D. and McLeod, J.B. Diffusion mediated transport in multiple state systems, An elliptic-parabolic problem in combustion theory: convergence to travelling waves, *SIAM J. Math. Anal.*, 2007, 39.4, 1208-1230.
- [13] Hastings, S., Kinderlehrer, D. and McLeod, J.B. Diffusion mediated transport with a look at motor proteins, Recent advances in nonlinear analysis, (Chipot, M., Lin, C-S., and Tsai, D-H., eds) (2008) World Scientific, 95 - 112.
- [14] D. Hilhorst and J. Hulshof, An elliptic-parabolic problem in combustion theory: convergence to travelling waves, *Nonlinear Analysis TMA*, 1991, 6, 17, 519–546.
- [15] D. Hilhorst and M. A. Peletier, Convergence to travelling waves in a reaction-diffusion system arising in contaminant transport, *J. Differential Equations*, 2000, 1, 163, 89–112.
- [16] O. A. Ladyženskaja and V. A. Solonnikov and N. N. Ural'ceva, Linear and Quasilinear Equations of Parabolic Type, *Translations of Mathematical Monographs*, Vol. 23, American Mathematical Society, Providence, R.I., 1967.
- [17] O. A. Ladyženskaja and N. N. Ural'ceva, Linear and Quasilinear Elliptic Equations, Academic Press, New York, 1968.
- [18] S. Osher and J. Ralston, L^1 stability of travelling waves with applications to convective porous media flow, *Comm. Pure Appl. Math.*, 1982, 6, 35, 737–749.
- [19] B. Perthame, The general relative entropy principle—applications in Perron-Frobenius and Floquet theories and a parabolic system for biomotors, *Rend. Accad. Naz. Sci. XL Mem. Mat. Appl. (5)*, 2005, 29, 307–325.
- [20] B. Perthame, P. E. Souganidis, Asymmetric potentials and motor effect: a large deviation approach, 1–16, *Arch. Rat. Mech. Anal.*, to appear, 2008.
- [21] M. H. Protter and H. F. Weinberger, Maximum Principles in Differential Equations, Springer-Verlag, New York, 1984.