

Toward Nonlinear Quantum Fokker-Planck Equations

Roumen Tsekov
DWI, RWTH, 52056 Aachen, Germany
3 August 2008

It is demonstrated how the original semiclassical approach of Coffey et al can be improved to describe correctly the time evolution as well. As a result a new semiclassical Klein-Kramers equation for the Wigner function is derived, which remains quantum for a free quantum Brownian particle as well. It is transformed also to a semiclassical Smoluchowski equation, which leads to our semiclassical generalization of the classical Einstein law of Brownian motion. A possibility is discussed how to extend these semiclassical equations to nonlinear quantum Fokker-Planck equations.

PACS numbers: 03.65.Yz, 05.40.-a

It is well known that the classical Fokker-Planck equations are linear due to the logarithmic dependence of the Boltzmann entropy on the probability density. More complex systems, however, exchange non-Shannon information as well, which reflects in nonlinear mean-field Fokker-Planck equations [1, 2]. In many systems, including quantum ones, the Fisher information plays an important role [3]. The scope of the present paper is to generalize the existing linear semiclassical equations to nonlinear Fokker-Planck equations, accounting for the complete quantum Fisher information.

Last year Coffey et al [4] introduced heuristically the following semiclassical Klein-Kramers equation

$$\partial_t W + \frac{p}{m} \partial_x W - \partial_x V \partial_p W + \frac{\hbar^2}{24} \partial_x^3 V \partial_p^3 W = \zeta \partial_p \left(\frac{p}{m} W + k_B T \partial_p W + \frac{\hbar^2}{12 m k_B T} \partial_x^2 V \partial_p W \right) \quad (1)$$

describing the evolution of the Wigner function $W(x, p, t)$ of a quantum Brownian particle. Here $V(x)$ is an arbitrary potential and ζ is the Brownian particle friction coefficient. A novelty in Eq. (1) is the quantum term on the right hand side, which ensures that the equilibrium semiclassical Wigner distribution, known from statistical thermodynamics, is also a solution of Eq. (1). The result of Coffey et al is very important since it corrects the Caldeira-Leggett master equation, which is believed to be rigorous. Obviously this is not the case since the latter does not provide the exact quantum canonical Gibbs distribution at equilibrium. However, as we demonstrated in [5], Eq. (1) does not describe well enough the evolution since in the case of a free quantum Brownian particle ($V = 0$) it reduces to the classical Klein-Kramers equation

$$\partial_t W + \frac{p}{m} \partial_x W = \zeta \partial_p \left(\frac{p}{m} W + k_B T \partial_p W \right) \quad (2)$$

Hence, according to Coffey et al a free quantum Brownian particle behaves as a classical one, which is only true if it moves in vacuum ($\zeta = 0$) since in this case no energy exchange takes place. The problem above appears from a non-rigorous extension of an equilibrium result to the non-equilibrium case. Thus, the temperature dependence of the quantum term on the right hand side on Eq. (1) clearly shows its equilibrium origin and validity.

To extend an equilibrium expression to the non-equilibrium state is not unique. Since we are looking for a semiclassical approximation, we expect that the temperature dependence of the quantum term on the right hand side of Eq. (1) originates from the classical limit of the Wigner function W_{cl} . At equilibrium the following relation $\partial_x^2 V / k_B T = -\partial_x^2 \ln W_{cl}^{eq}$ holds between the potential V and the equilibrium classical Wigner function W_{cl}^{eq} , which is just the Maxwell-Boltzmann distribution. Hence, our proposal for an alternative of Eq. (1) is

$$\partial_t W + \frac{p}{m} \partial_x W - \partial_x V \partial_p W + \frac{\hbar^2}{24} \partial_x^3 V \partial_p^3 W = \zeta \partial_p \left[\frac{p}{m} W + k_B T \partial_p W - \frac{\hbar^2}{12m} \partial_p (W \partial_x^2 \ln W_{cl}) \right] \quad (3)$$

which provides also the correct semiclassical equilibrium distribution. However, in contrast to Eq. (1), Eq. (3) keeps the new quantum term in the case of a free quantum Brownian particle ($V = 0$) as well

$$\partial_t W + \frac{p}{m} \partial_x W = \zeta \partial_p \left[\frac{p}{m} W + k_B T \partial_p W - \frac{\hbar^2}{12m} \partial_p (W \partial_x^2 \ln W_{cl}) \right] \quad (4)$$

As seen this equation is linear only because of the semiclassical linearization of the quantum term but in general we expect the quantum Klein-Kramers equation to be nonlinear [6]. As a first step toward this direction one could replace W_{cl} by W in Eq. (4) to obtain

$$\partial_t W + \frac{p}{m} \partial_x W = \zeta \partial_p \left[\frac{p}{m} W + k_B T \partial_p W - \frac{\hbar^2}{12m} \partial_p (W \partial_x^2 \ln W) \right] \quad (5)$$

One can recognize now the Fisher information density in the last term of this nonlinear quantum Fokker-Planck equation.

In the case of strong friction one can easily derive from Eq. (4) a semiclassical Smoluchowski equation for the probability density in the coordinate space $P = \int_{-\infty}^{\infty} W dp$

$$\partial_t P = \frac{1}{\zeta} \partial_x^2 \left[(k_B T - \frac{\hbar^2}{12m} \partial_x^2 \ln P_{cl}) P \right] \quad (6)$$

In the derivation of Eq. (6) the following equality $\partial_x^2 \ln W_{cl} = \partial_x^2 \ln P_{cl}$ is employed, which follows from the known non-equilibrium Gaussian solution of Eq. (2). Equation (6) shows that the quantum effects correspond to effective increase of the temperature. This is, however, not surprising, when one realizes that the quantum term in Eq. (6) is in fact the semiclassical limit of the so-called local quantum temperature [7] given by

$$T_Q = -\frac{\hbar^2}{12mk_B} \partial_x^2 \ln P \quad (7)$$

The classical solution of Eq. (6) is the well-known Gaussian distribution

$$P_{cl} = \frac{1}{\sqrt{4\pi Dt}} \exp\left(-\frac{x^2}{4Dt}\right) \quad (8)$$

where $D = k_B T / \zeta$ is the Einstein diffusion constant. Substituting Eq. (8) in Eq. (6) leads to a semiclassical diffusion equation

$$\partial_t P = D(1 + \lambda_T^2 / 6Dt) \partial_x^2 P \quad (9)$$

where $\lambda_T = \hbar / 2\sqrt{mk_B T}$ is the de Broglie thermal wavelength. The expression in the brackets represents the relative increase of the diffusion constant by quantum effects. The solution of Eq. (9) is also a Gaussian distribution density with dispersion σ^2 satisfying the following differential equation

$$\partial_t \sigma^2 = 2(D + \lambda_T^2 / 6t) \quad (10)$$

Integrating on time yields the already known semiclassical generalization of the classical Einstein law of Brownian motion

$$\sigma^2 = 2Dt + \lambda_T^2 \ln(6Dt/\lambda_T^2)/3 \quad (11)$$

which is derived from a nonlinear quantum Smoluchowski equation [5]. Note that Eq. (11) is applicable for large times $t > \lambda_T^2/2D$ only, while at shorter times the diffusion is strongly quantum and cannot be described by the semiclassical approach [6].

How it was mentioned one would expect to obtain even a quantum Smoluchowski diffusion equation

$$\partial_t P = \partial_x^2[(D + D_Q)P] = \partial_x(D\partial_x P + P\partial_x Q/3\zeta) \quad (12)$$

via replacing P_{cl} by P in Eq. (6) to restore the quantum temperature T_Q from Eq. (7), which is included now in the quantum diffusion coefficient $D_Q = k_B T_Q / \zeta$. This equation follows also directly from Eq. (5) since for Gaussian distributions $\partial_x^2 \ln W = \partial_x^2 \ln P$. In the last expression of Eq. (12) the Bohm quantum potential $Q = -\hbar^2 \partial_x^2 \sqrt{P} / 2m\sqrt{P}$ is introduced [8]. Note that the mean value of Q is proportional to the Fisher information. The solution of Eq. (12) is also a Gaussian distribution density with dispersion given by

$$\sigma^2 - \lambda_T^2 \ln(1 + 3\sigma^2/\lambda_T^2)/3 = 2Dt \quad (13)$$

Substituting σ^2 in the logarithmic term by the classical expression $2Dt$ yields an improved semiclassical result $\sigma^2 = 2Dt + \lambda_T^2 \ln(1 + 6Dt/\lambda_T^2)/3$, which is always positive and recovers Eq. (11) at large time. At short time Eq. (13) reduces to $\sigma^2 = \hbar\sqrt{t/3mb}$, which is close to the exact expression $\sigma^2 = \hbar\sqrt{t/mb}$ for the purely quantum diffusion [6]. This indicates, however, that Eq. (12) is approximate, since it is derived from semiclassical considerations. As shown in [5], the factor 3 originates from an integral over the reciprocal temperature $\beta = 1/k_B T$, which cancels at zero temperature. A more general analysis of thermo-quantum diffusion is given in [9] and the correct nonlinear quantum Smoluchowski equation reads

$$\partial_t P = D \partial_x (\partial_x P + P \partial_x \int_0^\beta Q d\beta) \quad (14)$$

The integral in Eq. (14) accounts for the interplay between the thermal Shannon and the quantum Fisher information. Hence, the total entropy $-\ln P - \beta Q + \int_0^\beta \beta dQ$ is not simply a superposition of the Shannon and Fisher components. The last integral represents the thermo-quantum entropy [9], which originates from the temperature dependence of the quantum potential.

The author is thankful to Professors Aurelian Isar and Joachim Messer for interesting and stimulating discussions.

- [1] T D Frank 2005 *Nonlinear Fokker-Planck Equations* Springer, Berlin
- [2] P H Chavanis 2006 *Comp. Rend. Phys. (Paris)* **7** 318
- [3] B R Frieden 1998 *Physics from Fisher Information* University Press, Cambridge
- [4] W T Coffey, Yu P Kalmykov, S V Titov and B P Mulligan 2007 *J. Phys. A: Math. Theor.* **40** F91, 12505
- [5] R Tsekov 2007 *J. Phys. A: Math. Theor.* **40** 10945
- [6] R Tsekov 2008 *Int. J. Theor. Phys.* [in press](#) ([arXiv:0711.1442](#))
- [7] S Sonego 1991 *Found. Phys.* **21** 1135
- [8] M G Ancona G J Iafrate 1989 *Phys. Rev. B* **39** 9536
- [9] R Tsekov 2008 [arXiv:0803.4409](#)