

On the theory of magnetic field dependence of heat conductivity in dielectric in isotropic model

L.A.Maksimov and T.V.Khabarova*
Kurchatov Institute, Moscow 123182, Russia

Phonon polarization in a magnetic field is analyzed in isotropic model. It is shown, that at presence of spin-phonon interaction phonon possess circular polarization which causes the appearance of heat flux component perpendicular both to temperature gradient and magnetic field.

Recently [1], [2] a novel phenomenon was found. It is the detection in dielectric of a heat flow perpendicular both to magnetic field B and temperature gradient ∇T – so called Phonon Hall Effect (PHE). Such flow is caused by spin-phonon interaction (SPI) of phonons and paramagnetic ions Tb^{+3} . The theory of PHE was considered in works [3] and [4] (the critic of work [3] see in [4]).

However not only SPI leads to magnetic field dependence of dielectric heat conductivity. As it is known, in molecular gases magnetic field affects the transport phenomena because of the molecular rotary moments \mathbf{M} , precessed in a magnetic field, dependence of collision cross-section (Senftleben-Beenakker Effect [5]).

We expect, that the phenomenon similar to PHE in $Tb_3Ga_5O_{12}$ can be found out in molecular crystals where is a component, consisting of the molecules with rotary degrees of freedom (RDF). Typical representatives of such substances are criocrystals [6]. To magnetic field dependence of heat conductivity can lead a RDF-dependent phonon scattering on molecules. The suitable theory is easy to construct via generalizing Senftleben-Beenakker theory on a case of heat transfer by phonons. In approach where RDF are considered quasi-classically such theory is considered in [7]. In this paper we have another approach to this problem, and we show that in crystals with RDF there is a mechanism close to SPI in ionic crystals which leads to PHE through renormalization of acoustic waves because of lattice vibrations interaction with molecular rotation. From symmetry considerations describing such interaction Hamiltonian can be written down in similar to SPI form:

$$H_1 = -g \sum_n (\mathbf{M}_n, [\mathbf{u}_n \times \mathbf{p}_n]). \quad (1)$$

Here \mathbf{M}_n is the rotary moment of the molecule in a cell n , $[\mathbf{u}_n \times \mathbf{p}_n]$ – total orbital moment of oscillating atoms in a cell. Speaking generally, molecular crystals have a complex structure containing of many particles in an elementary cell. But at low temperatures heat is transferred by long-wave elastic waves, when all particles of a cell oscillate with the same amplitude \mathbf{u}_n and velocity \mathbf{v}_n . Therefore instead of a multicomponent crystal it is possible to consider a crystal with one atom in a cell with mass m , equal to total mass of particles in a cell and one rotary moment. The value of g in case of SPI was estimated in numerous works on spin-lattice relaxation [4]

and [8]. In case of acoustic vibrations interaction with rotary degrees of freedom factor g we shall consider as small phenomenological parameter having the same order as for SPI when the molecule has uncompensated electronic moment, as in O_2 . We shall use system of units where $k_B = 1$, $\hbar = 1$. For short, interaction (1) shall be also called SPI in the case of crystals with rotary degrees of freedom. Almost identical form of Hamiltonian describing phonon interaction with internal degrees of freedom in ionic and molecular crystals (components of operator \mathbf{M} are replaced with Pauli matrices in ionic crystals with quasi-doublet structure of the bottom levels) leads to qualitative similarity of transverse thermal conductivity theories in both cases. But from the physical point of view interaction mechanisms are far from each other.

Usual dielectric heat conductivity weakly depends on detailed properties of acoustic vibrations. As it is shown in [4], PHE is caused by occurrence of phonon elliptic polarization at presence of a magnetic field and SPI. But character of acoustic branches polarization essentially depends on symmetry properties of a dynamic matrix. Unlike the general case considered in [4], where all three vibration modes are nondegenerate almost at all directions of wave vector \mathbf{k} , in present paper the model of vibrations in isotropic medium where transverse acoustic modes are degenerate will be considered at all \mathbf{k} . Though this model is realized only when atoms with a large interaction radius interact, in this model dynamic matrix has rather simple form. In isotropic medium model we shall find a heat flow in a $[B \times \nabla T]$ direction as for crystals with rotary degrees of freedom, and for ionic crystals with quasi-doublet structure. The first result is interesting as a prediction of PHE presence in molecular crystals, and the second is correction of work [3].

In an external magnetic field the average value of the rotary moment $\langle \mathbf{M} \rangle = \langle \mathbf{M}_n \rangle$ is nonzero, and crystal elastic vibrations are determined by Hamiltonian renormalized taking into account (1):

$$H = \sum_n h_n, \quad (2)$$

where

$$h_n = \frac{1}{2m} p_n^2 - \frac{1}{2} \sum_{n'} m D_{nn'}^{ab} u_n^a u_{n'}^b - g \langle \mathbf{M} \rangle [\mathbf{u}_n \times \mathbf{p}_n]. \quad (3)$$

It is essential, that the effective coupling constant $g \langle \mathbf{M} \rangle$

is proportional to magnetization of particles and it disappears when someone switches off the magnetic field.

Using (3) it is possible to write the quantum energy continuity equation and to deduce the density of a heat flow formula ([3], [4], [9])

$$j_H^\gamma = \frac{m}{2V} \sum_{nn'} r^\gamma D_{nn'}^{ab} u_n^a v_{n'}^b. \quad (4)$$

We emphasize, that in this expression term v_n^b is a velocity of this ion, but not an impulse of an ion p_n^b divided by mass

$$v_n^a = \partial_t u_n^a = \partial H / \partial p_n^a = p_n^a / m - e_{abc} g \langle M^b \rangle u_n^c. \quad (5)$$

Let's turn in (4) to impulse representation using

$$\sum_r r^\gamma D_r^{ab} \exp(ikr) = i \nabla_k^\gamma D_k^{ab}.$$

We find

$$j_H^\gamma = i \frac{m}{2V} \sum_{k s k' s'} (\nabla_k^\gamma D_k^{ab}) u_{ks}^a v_{-ks'}^b. \quad (6)$$

(s is the mode number) Let's introduce decomposition of displacement vectors and velocity vectors for normal vibration modes

$$\begin{aligned} u_{ks}^a &= \sqrt{\frac{1}{2m\omega_{ks}}} (e_{ks}^a a_{ks} + e_{-ks}^{a*} a_{-ks}^+), \\ v_{ks}^a &= (-i\omega_{ks}) \sqrt{\frac{1}{2m\omega_{ks}}} (e_{ks}^a a_{ks} - e_{-ks}^{a*} a_{-ks}^+). \end{aligned} \quad (7)$$

It is very important, that usual form has a velocity decomposition instead of an impulse $\mathbf{p}_i = m\mathbf{v}_i - gm(\mathbf{u}_i \times \mathbf{M})$ as authors of [3] supposed. We substitute u and v in (6) for (7) and average by state, which is diagonal on phonon numbers. Then we discard anomalous averages $\langle a_{ks} a_{-ks'} \rangle$ and $\langle a_{-ks'}^+ a_{ks}^+ \rangle$ and we come to heat flow in a $[B \times \nabla T]$ direction:

$$\begin{aligned} \langle j_H^y \rangle &= \frac{1}{4V} \sum_{k s s'} \left(\sqrt{\frac{\omega_{ks}}{\omega_{ks'}}} + \sqrt{\frac{\omega_{ks'}}{\omega_{ks}}} \right) \\ &\quad \times \left(\nabla_k^y D_k^{\alpha\beta} \right) e_{ks}^{\alpha*} e_{ks'}^\beta \langle a_{ks}^+ a_{ks'} \rangle. \end{aligned} \quad (8)$$

In isotropic body PHE is characterized by one factor (as a Hall constant). Without loss of generality we choose magnetization on Z -axis, a temperature gradient – on X , and a heat flow – on Y . Using this expression in [4] a PHE amount for acoustic vibrations of a general view has been calculated. In present paper we consider an elastic vibrations model in isotropic body where the dynamic matrix in zeroth-order approximation D_k^{ab} has a form

$$D_k^{ab} = c_0^2 \delta^{ab} k^2 + w k_a k_b, \quad (9)$$

with degenerate branches $\omega_{+,-}^2 = \omega_0^2 = c_0^2 k^2$ and longitudinal branch $\omega_{\parallel}^2 = \omega_0^2 + \lambda$, $\lambda = w k^2$. Here the third

independent term like $\delta^{ab} k_a^2$, which in a cubic crystal removes degeneration is discarded.

For isotropic body model considering orthogonality of polarization vectors $e_{ks}^{a*} e_{ks'}^a = \delta_{ss'}$, we have

$$\begin{aligned} \langle j_H^y \rangle &= \frac{1}{4V} \sum_{k s s'} \left(\sqrt{\frac{\omega_{ks}}{\omega_{ks'}}} + \sqrt{\frac{\omega_{ks'}}{\omega_{ks}}} \right) [\delta_{ss'} 2c_0^2 k^y + \\ &\quad + w((k_a e_{ks}^{a*}) e_{ks'}^y + e_{ks}^{y*} (k_b e_{ks'}^b))] \langle a_{ks}^+ a_{ks'} \rangle. \end{aligned} \quad (10)$$

We see that (4), (8), (10) do not contain factor g evidently. The heat flow magnitude changes only because of renormalization of phonon spectrum and polarization. Everyone can verify that diagonal terms in (10) do not lead to heat flow in $[B \times \nabla T]$ direction. PHE is described by off-diagonal terms in (10):

$$\langle j_H^y \rangle = \frac{w}{2V} \sum_{k s s'} \left(\sqrt{\frac{\omega_{ks}}{\omega_{ks'}}} + \sqrt{\frac{\omega_{ks'}}{\omega_{ks}}} \right) \text{Re} \{ e_{ks}^{y*}(\mathbf{k} e_{ks'}) \langle a_{ks}^+ a_{ks'} \rangle \} \quad (11)$$

Particularly, transverse modes correlated motion gives

$$\begin{aligned} \langle j_{\perp}^y \rangle &= \frac{w}{2V} \sum_k \text{Re} \{ e_{k-}^{y*}(\mathbf{k} e_{k+}) \langle a_{k-}^+ a_{k+} \rangle + \\ &\quad + e_{k+}^{y*}(\mathbf{k} e_{k-}) \langle a_{k+}^+ a_{k-} \rangle \}. \end{aligned} \quad (12)$$

Expression (12) essentially depends on phonon polarization vectors. First of all it is necessary to establish their form in consideration of SPI. Hamiltonian (3) leads to the dispersion equation

$$\omega_{ks}^2 e_{ks}^a = \tilde{D}_k^{ab} e_{ks}^b, \quad (13)$$

where

$$\tilde{D}_k^{ab} = D_k^{ab} + i D_{1k}^{ab}, \quad D_{1k}^{ab} = e_{abc} G^c, \quad G^c = 2\omega g \langle M^c \rangle \quad (14)$$

$$\tilde{D}_k^{ab}(\mathbf{G}) = \left(\tilde{D}_k^{ba}(-\mathbf{G}) \right)^*, \quad (\mathbf{e}_{ks}^* \mathbf{e}_{ks'}) = \delta_{ss'}. \quad (15)$$

SPI contribution to dynamic matrix is an imaginary antisymmetric tensor. In model (9) dispersion equation takes a form

$$W e^a = \lambda \hat{k}^a (\hat{k} e) + i [e \times \mathbf{G}]^a \quad (16)$$

Here $W = \omega^2 - \omega_0^2$, $\lambda = w k^2$, $\hat{k} = \mathbf{k}/k = (\sin \theta \cos \varphi, \sin \theta \sin \varphi, \cos \theta)$ and $\mathbf{G} = (0, 0, G)$, $(\mathbf{G} \hat{k}) = G \cos \theta$, $Q = G \sin \theta$. Let's use orthonormal basis vectors

$$\begin{aligned} 1) \hat{k}, \quad 2) \hat{m} &= (\mathbf{G} - (\mathbf{G} \hat{k}) \hat{k}) / Q = \\ &= (-\cos \theta \cos \varphi, -\cos \theta \sin \varphi, \sin \theta), \\ 3) \hat{n} &= [\hat{k} \times \hat{m}] = Q^{-1} [\hat{k} \times \mathbf{G}] = (\sin \varphi, -\cos \varphi, 0), \end{aligned}$$

From (16) we find vector \mathbf{e} components

$$\mathbf{e} = \xi \left[\frac{Q}{\lambda - W} \hat{\mathbf{k}} + \frac{(\mathbf{G}\hat{\mathbf{k}})}{W} \hat{\mathbf{m}} - i\hat{\mathbf{n}} \right] \quad (17)$$

and eigenvalue equation

$$\frac{Q^2 W}{\lambda - W} + W^2 - (\mathbf{G}\hat{\mathbf{k}})^2 = 0. \quad (18)$$

Dispersion equation (13) determines polarization vectors accurate within phase. We set phase of parameter ξ equal to zero. Parameter absolute value is set by normalization:

$$\xi^{-2} = \left(\frac{Q}{\lambda - W} \right)^2 + \left(\frac{(\mathbf{G}\hat{\mathbf{k}})}{W} \right)^2 + 1. \quad (19)$$

At inversion ($\hat{\mathbf{k}}, \hat{\mathbf{m}}, \hat{\mathbf{n}} \rightarrow -\hat{\mathbf{k}}, \hat{\mathbf{m}}, -\hat{\mathbf{n}}$) polarization vector (17) changes its sign as a polar vector. Complexity of vector (17) means that it is elliptically polarized.

Expressions (17), (18), (19) are correct with any G . Then we assume phonon and internal degrees of freedom of ions (molecules) interaction to be weak $G \ll \lambda$. In this case we get one longitudinal mode $W_{\parallel} = \lambda$, $\mathbf{e} = \hat{\mathbf{k}}$ from (18).

For transverse modes in zeroth-order approximation W is equal to zero, and in consideration of G is equal to

$$W_{\eta} = \frac{1}{2} \left(- (Q^2/\lambda) + \eta \sqrt{(Q^2/\lambda)^2 + 4 (\mathbf{G}\hat{\mathbf{k}})^2} \right), \quad \eta = \pm 1 \quad (20)$$

and describes their splitting

$$\Delta = \omega_+ - \omega_- = \frac{1}{\omega_0} \sqrt{(Q^2/2\lambda)^2 + (\mathbf{G}\hat{\mathbf{k}})^2}. \quad (21)$$

Splitting is minimal on equator ($|\cos \theta| < G/\lambda$, $\Delta = Q^2/(2\lambda\omega_0)$). For other $\hat{\mathbf{k}}$ directions we have

$$W_{\eta} = \eta \left| \mathbf{G}\hat{\mathbf{k}} \right|, \quad \Delta = G |\cos \theta| / \omega_0. \quad (22)$$

Real and imaginary parts of each \mathbf{e}_{η} in zeroth-order approximation mutually perpendicular and equal to each other.

$$\mathbf{e}_{\eta} \simeq \frac{1}{\sqrt{2}} [\eta (\text{sign} \cos \theta) \hat{\mathbf{m}} - i\hat{\mathbf{n}}]. \quad (23)$$

This means transverse phonons have circular polarization [3].

Polarization vector (23) direction changes stepwise crossing equator. However, if we change transverse mode numeration and write $W_{\eta} = \eta \mathbf{G}\hat{\mathbf{k}}$ instead of (22) then projection of (17) on the $\hat{\mathbf{m}}$ -axis will be constant. It always happens when crossing levels. The magnitude of (12) does not depend on numeration.

In linear approximation we find

$$(\mathbf{e}_{\eta}\hat{\mathbf{k}}) = \frac{\xi_{\eta} Q}{\lambda} \left(1 + \frac{W_{\eta}}{\lambda} \right). \quad (24)$$

We also need following expressions:

$$\xi_{\eta} \simeq \frac{1}{\sqrt{2}} \left(1 - \frac{Q^2}{4\lambda W_{\eta}} \right), \quad (25)$$

$$e_{\eta}^y = \xi_{\eta} \left[\frac{Q}{\lambda - W_{\eta}} (\sin \theta \sin \varphi) + \frac{(\mathbf{G}\hat{\mathbf{k}})}{W_{\eta}} (-\cos \theta \sin \varphi) + i \cos \varphi \right]. \quad (26)$$

Expression (12) contains projection of polarization vector onto wave vector direction ($\mathbf{e}_{\eta}\hat{\mathbf{k}}$) which is linear on SPI. So, off-diagonal density matrix $\langle a_{ks}^+ a_{ks'} \rangle$ is quite enough to calculate in zeroth-order approximation on SPI.

As is known, the nonequilibrium part of density matrix (diagonal on modes) in tau-approximation is equal

$$\begin{aligned} f_p &= \langle a_p^+ a_p \rangle - N_p = -\frac{1}{\Omega_{pp}} (\mathbf{c}_p \nabla) N_p = \\ &= -\frac{1}{\Omega_{pp} T^2} N_p (1 + N_p) (\omega_p \mathbf{c}_p) \nabla T. \end{aligned} \quad (27)$$

Here and afterwards, we use $p = ks$ for short, $1/\Omega_{pp}$ is a p -phonon relaxation time. Furthermore, in obvious cases we shall leave out an index \mathbf{k} .

The corresponding energy flow is parallel to ∇T and leads to heat conductivity coefficient

$$\chi^{xx} \simeq T^3 (c\Omega)^{-1}, \quad (28)$$

where c, Ω – average values of c_p, Ω_{pp} .

Off-diagonal part of density matrix can be expressed by f_p constructing generalized Boltzmann equation. Phonon scattering has many channels: anharmonicity, resonant scattering with molecular multiplet structure excitation, scattering with rotary degrees of freedom contribution, impurity scattering. In all cases the answer has similar structure. In [4] it is shown, that

$$\langle a_p^+ a_q \rangle = \frac{i J_{pq}}{(\omega_p - \omega_q)}, \quad (29)$$

where J_{pq} is a Hermitian matrix which differs mainly from usual collision integral in that there is product of the phonon scattering amplitudes instead of a square of the scattering amplitude module (Fermi's golden rule). In tau-approximation this expression has a form of

$$\langle a_p^+ a_q \rangle = i \frac{\Omega_{qp}(\omega_p) f_p + \Omega_{qp}(\omega_q) f_q}{2(\omega_p - \omega_q)}, \quad (30)$$

where $\Omega_{qp}^*(\omega_p) = \Omega_{pq}(\omega_p)$ and $\langle a_p^+ a_q \rangle^* = \langle a_q^+ a_p \rangle$ are Hermitian.

The effective relaxation frequencies form depends on scattering mechanism. In [4] corresponding expressions for resonant scattering and anharmonicity are shown. In case of potential impurity scattering

$$\Omega_{qp}(\omega_p) = 2\pi \frac{N_{im}}{N^2} \sum_g A_{qg} A_{gp} \delta(\omega_p - \omega_g). \quad (31)$$

It is essential, that the transition amplitude $A_{pq} = e_p^{a*} A_{pq}^{ab} e_q^b$ is proportional to polarization vectors. It guarantees the absence of a stream (12) dependence on general phases of phonon modes (phases of parameter ξ in (17)). The A_{qg} polarization-dependence is automatically transferred on relaxation frequencies $\Omega_{qp} = e_q^{a*} \Omega_{qp}^{ab} e_p^b$, where tensor $\Omega_{qp}^{ab} = (\Omega_{pq}^{ba})^*$ does not depend on external polarization vectors. In isotropic body model we can neglect mode splitting for transverse modes in numerator of (30)

$$\langle a_{k+}^+ a_{k-} \rangle = i\Omega_{-+}(\omega_0) \frac{f_{\perp}}{\Delta} \quad (32)$$

When one of modes is longitudinal from (30) we have

$$\langle a_3^+ a_{\eta} \rangle = \frac{i}{2\varepsilon} [\Omega_{3\eta}(\omega_3) f_3 + \Omega_{\eta 3}(\omega_0) f_{\perp}]. \quad (33)$$

where $\varepsilon = \omega_3 - \omega_0 = \sqrt{\omega_0^2 + \lambda^2} - \omega_0$.

Let's use the formulas above for calculation of transverse heat conductivity coefficient \varkappa^{yx} . Using (32) in (12) we find

$$\langle j_{\perp}^y \rangle = \frac{w}{2V} \sum_k \text{Re}\{-e_{k-}^{y*}(\mathbf{k}e_{k+}) i\Omega_{-+}^*(\omega_0) \frac{f_{\perp}}{\Delta} + e_{k+}^{y*}(\mathbf{k}e_{k-}) i\Omega_{-+}(\omega_0) \frac{f_{\perp}}{\Delta}\}. \quad (34)$$

In (27) function f_{\perp} is proportional to $e^x \sim \cos\varphi$, and nonzero result is given only by imaginary part of (26): $e_{k\eta}^{y*} = (\dots) - i\xi_{\eta} \cos\varphi$. From (26), (24) and (34) we have

$$\langle j_{\perp}^y \rangle = \frac{w}{2V} \sum_k \frac{f_{\perp}}{\Delta} \frac{kQ}{\lambda} \xi_{-} \xi_{+} \cos\varphi \times \text{Re}\left\{-\left(1 + \frac{W_{+}}{\lambda}\right) \Omega_{-+}^* + \left(1 + \frac{W_{-}}{\lambda}\right) \Omega_{-+}\right\}. \quad (35)$$

We see, that only second-order terms in (24) give the contribution to PHE. Using concrete form of parameters λ , Δ , Q , ξ_{η} , W_{η} , G we find

$$\langle j_{\perp}^y \rangle = -\frac{g\langle M \rangle w}{V} \sum_k f_{\perp} \frac{\omega_0^2}{w^2 k^3} \sin\theta \cos\varphi \text{Re} \Omega_{-+}. \quad (36)$$

Then we integrate using (27)

$$\varkappa^{yx} = -\frac{g\langle M \rangle w}{V} \sum_k \frac{\omega_0^3 c}{w^2 k^3} (\sin\theta \cos\varphi)^2 \frac{\text{Re} \Omega_{-+}}{\Omega_{pp} T^2} \times N_p (1 + N_p). \quad (37)$$

Consequently, accepting $w \simeq c^2$ and $\left\langle \frac{\text{Re} \Omega_{-+}}{\Omega_{pp}} \right\rangle \simeq 1$, we find accurate within numerical coefficient

$$\varkappa^{yx} = -\frac{g\langle M \rangle}{V} \sum_k \frac{c^2}{T^2} N_p (1 + N_p) \simeq -\frac{g\langle M \rangle T}{c} \quad (38)$$

Dividing it on longitudinal heat conductivity coefficient $\varkappa^{xx} \simeq T^3 (c\Omega)^{-1}$ we find a Hall angle

$$\frac{\varkappa^{yx}}{\varkappa^{xx}} \simeq \langle M \rangle \frac{g\Omega}{T^2}.$$

The second channel of Hall heat conductivity is caused by occurrence in temperature gradient presence a correlated transverse and longitudinal phonon motion which is characterized by off-diagonal density matrix component (33):

$$\langle a_1^+ a_3 \rangle = -\frac{i}{2\varepsilon} [\Omega_{3\eta}(\omega_0) f_1 + \Omega_{\eta 3}(\omega_3) f_3]. \quad (39)$$

Here $1 = (\mathbf{k}, \perp)$, $3 = (\mathbf{k}, \parallel)$. This density matrix (and its Hermitian conjugation) forms Hall heat flow (see (11))

$$\langle j_H^y \rangle = \frac{w}{2V} \sum_k k \left(\sqrt{\frac{\omega_1}{\omega_3}} + \sqrt{\frac{\omega_3}{\omega_1}} \right) \text{Re}[e_1^{y*}(\hat{k}\mathbf{e}_3) \langle a_1^+ a_3 \rangle + e_3^{y*}(\hat{k}\mathbf{e}_1) \langle a_3^+ a_1 \rangle] \quad (40)$$

The matrix (33) is proportional to (27) and $\cos\varphi$. Besides, $e_{k\eta}^{y*} = -i\xi_{\eta} \cos\varphi$ and $e_3^{y*} = \sin\theta \sin\varphi$. So, only the first term in square brackets gives a contribution to integral. In this case $(\hat{k}\mathbf{e}_3) \simeq 1$ and integration element in (40) does not contain series expansion parameter of SPI.

$$\langle j_H^y \rangle = -\frac{w}{2V} \frac{1}{\sqrt{2}} \sum_k \left(\sqrt{\frac{\omega_1}{\omega_3}} + \sqrt{\frac{\omega_3}{\omega_1}} \right) \cos\varphi \frac{k}{2\varepsilon} (f_1 \text{Re} \Omega_{31} + f_3 \text{Re} \Omega_{13}(\omega_3)) \quad (41)$$

Using (27) and integrating we estimate longitudinal mode contribution to Hall heat conductivity

$$\varkappa_{\parallel}^{yx} \sim \frac{T^2 \text{Re} \Omega_{31}}{(c_{\parallel} - c_0) \Omega_{pp}} \sim \frac{T^2}{c_0} \quad (42)$$

In this case the Hall angle has an order of magnitude of

$$\frac{\varkappa_{\parallel}^{yx}}{\varkappa^{xx}} \sim \frac{\Omega}{T} \quad (43)$$

In conclusion we compare the estimations of heat conductivity coefficients caused by transverse and longitudinal mode correlation $\varkappa_{\parallel}^{yx}$ (42), transverse modes \varkappa_{\perp}^{yx} (38), and estimation \varkappa_{KM}^{yx} for ion crystal with nondegenerate modes [4]

$$\varkappa_{\parallel}^{yx} \sim \frac{T^2}{c_0}, \quad \varkappa_{\perp}^{yx} \sim \langle M \rangle \frac{gT}{c_0}, \quad \varkappa_{KM}^{yx} \simeq \langle \sigma \rangle \frac{gT}{c}. \quad (44)$$

We see, that estimations for transverse modes in isotropic model and cubic crystal coincide accurate within average polarization caused by rotary moment substitution for pseudospin polarization. It is essential, that in both cases frequency dependence is absent as it should be for heat flow component in $[B \times \nabla T]$. direction. The longitudinal mode (42) contribution has qualitative other form. It does not depend not only on frequency but on coupling constant too, as it is in common Hall Effect. At high temperatures ($T \gg g$) channel $\varkappa_{\parallel}^{yx}$ plays a main role. If there were crystals where even approximately there were acoustic vibrations with ellipticity close to circular, then PHE experimental observation would become rather simple problem.

The first theoretical work devoted to PHE [3] contained a mistake at using relation (5) between particle velocity and their impulse at presence of SPI. Nevertheless it is interesting to compare a \varkappa^{yx} estimation in [3] with our formulas (42), (38), as in [3] calculations were performed in isotropic model like in present paper. If to not pay attention to numerical coefficients which are obvious excess of accuracy, they received a result

$$\varkappa_{sheng}^{yx} = \frac{gT}{c_0},$$

which coincide with (38) and main result of [4]. This concurrence is caused not by of our theory equivalence, and only trivial consequence of similar dimension of results and that all of them are received in linear approach on g . So, it is necessary to emphasize, that the basic result of the this work (and [4]) is not the estimation of \varkappa^{yx} , and

the establishment of that fact, that PHE is caused by joint action of two equally important factors - 1) phonon elliptic polarization caused by SPI, and 2) induced by a temperature gradient the correlated movement of two phonon modes with off-diagonal density matrix formation.

* Electronic address: frau_sych@mail.ru

- [1] C.Strohm, G.L.J.A.Rikken, and P.Wyder, Phys.Rev.Lett. 95, 155901 (2005)
- [2] A.V. Inyushkin, A.N. Taldenkov, JETP Lett., (in press).
- [3] L. Sheng, D. N. Sheng, and C. S. Ting, Phys.Rev.Lett. 96, 155901 (2006).
- [4] Yu. Kagan and L. A. Maksimov, arXiv:0707.2565 (July 2007)
- [5] L.J.F. Hermans, P.H. Fortuin, H.F.P. Knaap and J.J.M. Beenakker, Phys.Lett. A, 25, 81 (1967); L.L. Gorelik, V.G. Nikolaevskii, V.V. Sinitsyn, JETP Lett., 4, 11, 307 (1966); Yu. Kagan, L.A. Maksimov, JETP, 51, 1893 (1966).
- [6] Cryocrystals, edited by B. I. Verkin and A. F. Prikhotko (Naukova Dumka, Kiev, 1983).
- [7] L. A. Maksimov and T.V. Khabarova, arXiv:0308.1234 (March 2008)
- [8] Spin-Lattice Relaxation in Ionic Solids, edited by A.A. Manenkov and R. Orbach, Harper and Row, New York,1966; A.A. Abragam, B. Bleaney, Electron Paramagnetic Resonance of Transition Ions. Clarendon Press, Oxford, 1970; H. Capellmann, S. Lipinski, Z. Phys. B - Cond. Mat., 83, 199, (1991); A. S. Ioselevich, H. Capellmann, Phys. Rev. B, 51, 11446 (1995).
- [9] R.J. Hardy, Phys. Rev, 132, 168 (1963).