

Proofs On Arnold Conjectures

Renyi Ma

Department of Mathematical Sciences

Tsinghua University

Beijing, 100084

People's Republic of China

rma@math.tsinghua.edu.cn

Abstract

In this article, we give proofs on the Arnold Lagrangian intersection conjecture on the cotangent bundles, Arnold-Givental Lagrangian intersection conjecture, the Arnold fixed point conjecture.

Keywords fixed points, intersection points, Arnold conjectures.

1 Introduction and results

Let M be a smooth manifold and let T^*M denote its cotangent space. Let $pr_M : T^*M \rightarrow M$ denote the natural projection. The canonical or Liouville 1-form λ_M on T^*M is

$$\lambda_M(V) = \beta(d(pr_M)(V)) \quad \text{for } V \in T_\beta(T^*M).$$

The standard symplectic form on T^*M is the 2-form $\omega_M = -d\lambda_M$. If $q = (q_1, \dots, q_n)$ are local coordinates on M then $(q, p) = (q_1, p_1, \dots, q_n, p_n)$ are local coordinates on T^*M , where (q, p) corresponds to the covector

$$p_1 dq_1 + \dots + p_n dq_n \in T_q^*M.$$

In these local coordinates we have

$$\lambda_M = -pdq = -\sum_j p_j dq_j \quad \text{and} \quad \omega_M = dq \wedge dp = \sum_j dq_j \wedge dp_j.$$

It is easy to see that $(T^*M, d\lambda_M)$ is an exact symplectic manifold. Let $\phi : T^*M \rightarrow T^*M$ be a Hamiltonian symplectomorphism (see[2]). The Arnold Lagrange intersection conjecture(see[1, 2]) in cotangent bundle is a well-known conjecture in symplectic geometry. We recall the formulation. Consider a smooth function $f : M \rightarrow \mathbb{R}$, we denote by $\text{crit}(f)$ and $\text{crit}_m(f)$ the number of critical points of f resp. Morse function. Let $\text{Crit}(M) = \min\{\text{crit}(f)\}$ and $\text{Crit}_m(M) = \min\{\text{crit}_m(f)\}$ where f runs over all smooth functions $M \rightarrow \mathbb{R}$ resp. Morse functions. Furthermore, let $\text{Int}(M, \phi)$ and $\text{Int}_s(M, \phi)$ denote the number of intersection points of $\phi(M)$ with M resp. $\phi(M)$ intersects M transversally. Finally, let

$$\text{Arn} - \text{int}(M) := \min_{\phi} \text{int}(M, \phi); \quad \text{and} \quad \text{Arn} - \text{int}_s(M) := \min_{\phi} \text{int}_s(M, \phi)$$

where ϕ runs over all Hamiltonian symplectomorphisms $T^*M \rightarrow T^*M$, resp. ϕ runs over all Hamiltonian symplectomorphisms $T^*M \rightarrow T^*M$ such that $\phi(M)$ intersects M transversally. The Arnold Lagrange intersection conjecture claims that $\text{Arn} - \text{int}(M) \geq \text{Crit}(M)$ and $\text{Arn} - \text{int}_s(M) \geq \text{Crit}_m(M)$. It is well known and easy to see that $\text{Arn} - \text{int}(M) \leq \text{Crit}(M)$ and $\text{Arn} - \text{int}_s(M) \leq \text{Crit}_m(M)$. Thus, in fact, the Arnold conjecture claims the equality $\text{Arn} - \text{int}(M) = \text{Crit}(M)$ and $\text{Arn} - \text{int}_s(M) = \text{Crit}_m(M)$.

Theorem 1.1 *Let (T^*M, ω_M) ($\omega_M = d\lambda_M$) be the cotangent bundles of close manifold M . Then,*

$$\text{Arn} - \text{int}(M) := \text{Crit}(M); \quad \text{Arn} - \text{int}_s(M) := \text{Crit}_m(M),$$

i.e., the Arnold Lagrange intersection conjecture in cotangent bundle holds.

Theorem 1.1 in the stable case was proved by Hofer[8], the other methods provided for example in [9, 4], for the complete reference, see [3].

Now we generalize the above definition to the exact Lagrangian submanifolds in the exact symplectic manifolds. Let (V', ω') be an exact symplectic manifold with exact symplectic form $\omega' = d\alpha'$. Let $W' \subset V'$ a close submanifold, we call W' an exact Lagrange submanifold if $\alpha'|_{W'}$ an exact form,

i.e., $\alpha'|W' = df$. Let $\phi'_t : V' \rightarrow V'$ be a Hamiltonian isotopy with compact support in V' such that $\phi'_0 = Id$ and $\phi'_1 = \phi'$ (see[2]). Let $Int(W', V', \phi')$ denote the number of intersection points of $\phi'(W')$ and W' in V' . Finally, let

$$Arn - int(W', V') := \min_{\phi'} int(W', V', \phi')$$

where ϕ' runs over all Hamiltonian symplectomorphisms $V' \rightarrow V'$ as above. Similarly, one defines $Arn - int_s(W', V') := \min_{\phi'} int_s(W', V', \phi')$.

Then the generalized Arnold Lagrange intersection conjecture claims that $Arn - int(W', V') = Crit(W')$ and $Arn - int_s(W', V') = Crit_m(W')$.

Theorem 1.2 *Let (V', ω') be an exact symplectic manifold with exact symplectic form $\omega = d\alpha'$. Let $W' \subset V'$ a close exact Lagrange submanifold. Then,*

$$Arn - int(W', V') := Crit(W'); \quad Arn - int_s(W', V') := Crit_m(W'),$$

i.e., the generalized Arnold Lagrange intersection conjecture holds.

Again we generalize the above definition to the close Lagrangian submanifold in the general symplectic manifolds. Let (V', ω') be a symplectic manifold with symplectic form ω' . Let $W' \subset V'$ a close submanifold, we call W' a Lagrange submanifold if $\omega'|W' = 0$. Let $\phi'_t : V' \rightarrow V'$ be a Hamiltonian isotopy with compact support in V' such that $\phi'_0 = Id$ and $\phi'_1 = \phi'$ (see[2]). Let $Int(W', V', \phi')$ denote the number of intersection points of $\phi'(W')$ and W' in V' . Finally, let

$$Arn - int(W', V') := \min_{\phi'} int(W', V', \phi')$$

where ϕ' runs over all Hamiltonian symplectomorphisms $V' \rightarrow V'$ as above. Similarly, one defines $Arn - int_s(W', V') := \min_{\phi'} int_s(W', V', \phi')$. Then the Arnold-Givental Lagrange intersection conjecture claims that $Arn - int(W', V') = Crit(W')$ and $Arn - int_s(W', V') = Crit_m(W')$ if W' is symmetric under the anti-symplectic involution.

Theorem 1.3 *Let (V', ω') be a symplectic manifold with symplectic form ω' . Let $W' \subset V'$ a close Lagrange submanifold such that $H^1(W') = 0$. Then,*

$$Arn - int(W', V') := Crit(W'); \quad Arn - int_s(W', V') := Crit_m(W'),$$

Theorem 1.4 *Let (V', ω') be a symplectic manifold with symplectic form ω' . Let $\Phi : (V', \omega') \rightarrow (V', \omega')$ be a diffeomorphism such that $\Phi^*\omega' = -\omega'$, $\Phi^2 = Id$. Let $W' \subset V'$ be the symmetric Lagrange submanifold about Φ , i.e., $W' = \{x \in V' | \Phi(x) = x\}$. Then,*

$$Arn - int(W', V') := Crit(W'); \quad Arn - int_s(W', V') := Crit_m(W'),$$

i.e., the Arnold-Givental Lagrange intersection conjecture holds.

Now let (M, ω) be a closed symplectic manifold, and let $\phi : M \rightarrow M$ be a Hamiltonian symplectomorphism (see[2]). Furthermore, let $Fix(\phi)$ and $Fix_s(\phi)$ denote the number of fixed points of general ϕ resp transversal ϕ . Finally, let

$$Arn(M, \omega) := \min_{\phi} Fix(\phi) \text{ and } Arn_s(M, \omega) := \min_{\phi} Fix_s(\phi),$$

where ϕ runs over all Hamiltonian symplectomorphisms $M \rightarrow M$ resp. ϕ runs over all Hamiltonian symplectomorphisms $M \rightarrow M$ such that its fixed points are all nondegenerate. The famous Arnold conjecture claims that $Arn(M, \omega) = Crit(M)$ and $Arn_s(M, \omega) = Crit_m(M)$. Let $(V', \omega') = (M \times M, \omega \ominus \omega)$, then $L = \{(\sigma, \sigma) \in M \times M | \sigma \in M\}$, i.e., its diagonal of the product is Lagrangian submanifold in (V', ω') . Moreover, the map $P : (M \times M, \omega \ominus \omega) \rightarrow (M \times M, \omega \ominus \omega)$, $P(x, y) = (y, x)$, $x, y \in M$ is an anti-symplectic involution and L is its fixed points set. So, by the above theorem, we have

Theorem 1.5 *Let (M, ω) be a close symplectic manifold with symplectic form ω . Then,*

$$Arn(M, \omega) := Crit(M) \text{ and } Arn_s(M, \omega) := Crit_m(M),$$

i.e., the Arnold fixed point conjecture holds.

Theorem 1.6 *Let (V', ω') be a symplectic manifold with symplectic form ω' . Let $W' \subset V'$ a close Lagrange submanifold such that $i_* : H_1(W') \rightarrow H_1(V')$ is injective. Then,*

$$Arn - int(W', V') := Crit(W'); \quad Arn - int_s(W', V') := Crit_m(W'),$$

The proofs of these theorems does not depend on the hard analysis and relies on the analysis of the primitive one form of the symplectic form.

2 Proof of Theorem 1.1-1.2

Proof of Theorem1.1: Assumption $n \geq 2$. Consider the exact isotopy of Lagrange submanifolds in T^*M given by $\varphi_t, t \in [0, 1]$ as a hamilton isotopy of T^*M induced by hamilton vector field X_{h_t} with hamilton function h_t . Consider the Lie derivative $L_{X_{h_t}}\lambda_M$, by hamilton perturbation, we can assume that $L_{X_{h_t}}\lambda_M|(\varphi_t(M) \cap U(M)) = \lambda_M|(\varphi_t(M) \cap U(M))$, here $U(M)$ is the small neighbourhood of zero section M in T^*M . Moreover, $L_{X_{h_t}}\lambda_M = i_{X_{h_t}}d\lambda_M + di_{X_{h_t}}\lambda_M = dH_t$, here H_t is defined on T^*M . The level sets of H_t defines a foliation \mathcal{F}_t on T^*M . For t small enough, the Lagrangian submanifold $\varphi_t(M)$ in T^*M is transversal to the foliation \mathcal{F}_t except the Lagrangian submanifold M . So, we can perturb the Lagrangian isotopy φ_t such that the intersection points sets $\varphi_t(M) \cap M$ is invariant and the Lagrangian submanifold $\varphi_t(M)$ is transversal to the foliation \mathcal{F}_t except the intersection points set $\varphi_t(M) \cap M$.

This shows that the critical points of $H_1|_{\varphi_1(M)}$ are in the intersection points set $\varphi_1(M) \cap M$.

This yields Theorem1.1.

Proof of Theorem1.2: Same as the proof of Theorem1.1.

3 Proof of Theorem1.3-1.6

Proposition 3.1 *Let (V', ω') be a symplectic manifold with symplectic form ω' . Let $W' \subset V'$ a close Lagrange submanifold. Let $\varphi'_t : V' \rightarrow V'$ be a Hamiltonian isotopy with compact support in V' such that $\varphi'_0 = Id$ and $\varphi'_1 = \varphi'$. Consider the isotopy of Lagrange submanifolds in V' given by $F'_t = \varphi'_t, t \in [0, 1]$ as a C^∞ -map $F' : W' \times [0, 1] \rightarrow V'$. We assume that $F' : [0, 1] \times W' \setminus \{q_i | i = 1, \dots, k\} \rightarrow V'$ is an immersion and $F' : ([0, 1] \times W') \setminus \Sigma \rightarrow V'$ is an embedding, here Σ is the Riemann surfaces with boundaries contained in the boundaries of $[0, 1] \times W'$ (see[7]). Let $L_{n+1} = \cup_{t \in [0, 1]} \varphi'_t(W')$, then there exists a neighbourhood $U'(L_{n+1})$ of L_{n+1} in V' and a 1-form α' on $U'(L_{n+1})$ such that $\omega' = d\alpha'$.*

Proof. With loss of generality, we assume $n \geq 4$. Now we compute $F'^*\omega$ as in [6] as

$$F'^*\omega'|([0, 1] \times W') = \varphi'^*\omega|W' + i_{\dot{\varphi}}\omega \wedge dt$$

$$\begin{aligned}
&= \omega|_{W'} + d_{W'} h_t \wedge dt \\
&= d\theta'.
\end{aligned} \tag{3.1}$$

So, $\omega|_{F'([0,1] \times W' \setminus (\Sigma \cup \{q_i | i = 1, \dots, k\})}$ is exact. Let $\Sigma' \subset U'(L_{n+1})$ is a 2-cycle in W' , if necessary, by the small perturbation, we can assume that $\Sigma' \cap (\Sigma \cup \{q_i | i = 1, \dots, k\}) = \emptyset$, so we have $\int_{\Sigma'} \omega' = \int_{\Sigma'} d\theta' = 0$, so $\omega'|_{U'(L_{n+1})}$ is exact. This proves the proposition.

Proof of Theorem1.3. Since $H^1(W') = 0$, by Proposition 3.1 and Theorem1.2, it is obvious.

Proof of Theorem1.4. Since W' is the fixed points set of Φ and $\Phi^* \omega' = -\omega'$. By the proof of Proposition3.1, there exists a neighbourhood $U'(L_{n+1} \cup \Phi(L_{n+1}))$ of $L_{n+1} \cup \Phi(L_{n+1})$ in V' and a 1-form α' on $U'(L_{n+1} \cup \Phi(L_{n+1}))$ such that $\omega' = d\alpha'$. Let $\theta' = \frac{1}{2}(\alpha' - \Phi^* \alpha')$, then $\omega' = d\theta'$ and $\theta'|_{W'} = 0$. By Theorem1.2, Theorem1.4 follows.

Proof of Theorem1.6. By the proof of Proposition3.1, there exists a neighbourhood $U'(L_{n+1})$ of L_{n+1} in V' and a 1-form α' on $U'(L_{n+1})$ such that $\omega' = d\alpha'$. Then, by the condition that $i_* : H_1(W') \rightarrow H_1(V')$ is injective, we can assume that α' is exact. Then, Theorem1.2 yields Theorem1.6.

References

- [1] Arnold, V. I., First steps in symplectic topology, Russian Math. Surveys 41(1986),1-21.
- [2] Arnold, V.& Givental, A., Symplectic Geometry, in: Dynamical Systems IV, edited by V. I. Arnold and S. P. Novikov, Springer-Verlag, 1985.
- [3] Eliashberg, Y., Symplectic topology in the nineties, Differential geometry and its applications 9(1998)59-88.
- [4] Eliashberg, Y.& Gromov, M., Lagrangian Intersection Theory: Finite-Dimensional Approach, Amer. Math. Soc. Transl. 186(1998): 27-118.
- [5] Givental, A. B., Nonlinear generalization of the Maslov index, Adv. in Sov. Math., V.1, AMS, Providence, RI, 1990.
- [6] Gromov, M., Pseudoholomorphic Curves in Symplectic manifolds. Inv. Math. 82(1985), 307-347.

- [7] Hirsch, M., Differential Topology, Springer-Verlag, 1976.
- [8] Hofer, H., Lagrangian embedding and critical point theory, Ann. Inst.H Poincare, 2(1985), 407-462.
- [9] Laudenbach, F., Engouffrement symplectique et intersections Lagrangiennes, Comment. Math. Helvetici 70(1995) 558-614.