

CP violation of $\tau \rightarrow K\pi(\eta, \eta')\nu$ decays

Daiji Kimura^(a), Kang Young Lee^(b), T. Morozumi^(a), Keita Nakagawa^(a),

(a) Graduate School of Science, Hiroshima University,

Higashi-Hiroshima, 739-8526, Japan

(b) Department of Physics, Korea University, Seoul 136-713, Korea

Abstract

We study direct CP violation of $\tau \rightarrow K\pi(\eta, \eta')\nu$ decays. By studying the forward and backward asymmetry, the interference of $L = 0$ and $L = 1$ amplitudes of the hadronic system can be extracted. By including the scalar and vector mesons into the chiral Lagrangian, we compute the form factors which correspond to $L = 0$ and $L = 1$ angular momentum state of the hadronic system. We include real and imaginary parts of the one loop corrections to the self-energies of the scalar and vector mesons. The direct CP violation of the forward and backward asymmetry is computed using a two Higgs doublet model in which a new CP violating phase is introduced. We show how the CP violation of the forward and the backward asymmetry may depend on the new CP violating phase and the strong phase of the form factors.

PACS numbers: 13.35.Dx, 11.30.Er, 12.39.Fe, 12.60.Fr

I. INTRODUCTION

Two B factories, both Belle and Babar have accumulated the very large samples of τ decays. The τ lepton physics beyond the standard model, such as τ lepton number violation and time reversal violation through the measurement of electric dipole moment have been studied. The CP violation of the hadronic τ decay also has been investigated both theoretically [1] and experimentally [2]. Various angular distributions including the quantities using the τ spin polarization have been also proposed [1, 3]. Recently, Belle and Babar reported the precise measurements of the branching fractions of $\tau \rightarrow K_s \pi \nu$ [4] and $\tau^- \rightarrow K^- \pi^0 \nu$ [5]. The improved measurement of the branching fraction for $\tau \rightarrow K \eta \nu$ has been also obtained [6]. About the $\tau \rightarrow K \pi \nu$ decays, the hadronic invariant mass spectrum has been measured.

Motivated by these measurements, we study the direct CP violation of $\tau^\pm \rightarrow K^\pm P \bar{\nu}(\nu)$ with $P = \pi^0, \eta, \eta'$ and $\tau^\pm \rightarrow K_s \pi^\pm \bar{\nu}(\nu)$. Non-vanishing direct CP violation in the processes may arise with some new source of the CP violation in addition to Kobayashi Maskawa phase [7] and the strong phase shifts of the final states of hadrons.

In $\tau \rightarrow KP\nu$ decays, the hadronic system KP may have the angular momentum $L = 0$ (s wave) and $L = 1$ (p wave). The interference term of them can be extracted from the forward and backward asymmetry [8]. In the present paper, we define the asymmetry as the difference of the numbers of events for K meson scattered into the forward and backward directions with respect to the incoming τ momentum in the hadronic CM frame. By comparing the forward and backward asymmetries for CP conjugate processes, the direct CP violation can be defined. The s wave and p wave of the hadronic amplitudes are related to the scalar and vector form factors in the time like region which have their own strong phases. To evaluate them, we use a chiral Lagrangian including scalar and vector meson resonances such as $\kappa(800)$ and $K^*(890)$. We compute the both real and imaginary parts of the one loop corrections to the self-energy of the resonances and obtain the strong phase shifts. We include the pseudoscalar meson loop correction and scalar and vector meson loop correction. The latter may give an important contribution to the form factors at higher invariant mass regions above 1 GeV up to $m_\tau \sim 1.7(\text{GeV})$.

As a new physics effect, we study a two Higgs doublet model with non-minimal Yukawa couplings to the charged leptons. The two Higgs doublets contribute to the charged lepton mass through the Yukawa couplings. In the non-minimal model, the interaction of the

charged Higgs boson to the τ lepton family can be CP violating. The interaction generates the amplitude $\tau_R \rightarrow \nu_{\tau L} H^- \rightarrow \nu_{\tau L} (\bar{u}_L s_R)_{s\text{-wave}}$. The interference with the charged current interaction due to W boson exchanged diagram may lead to the direct CP violation which can be measured in the forward and backward asymmetry.

The paper is organized as follows. In section II, we show the hadronic chiral Lagrangian including scalar and vector resonances. In section III, we derive the form factors. In section IV, by fixing the finite renormalization constants, we numerically evaluate the form factors and the hadronic invariant mass spectrum. In section V, we introduce the two Higgs doublet model and present the direct CP violation. Section VI is devoted to conclusion and discussion.

II. CHIRAL LAGRANGIAN INCLUDING SCALAR AND VECTOR MESONS

In this section, we show the chiral Lagrangian with vector and scalar resonances. The following aspects are the main feature of the chiral Lagrangian.

- $U(1)_A$ breaking effect is taken into account so that we can apply the Lagrangian to τ decays into the final states including $K\eta$ and $K\eta'$.
- $SU(3)$ breaking of the vector mesons are taken into account.

About the inclusion of the scalar resonances, we followed the approach of Ref.[9]. About the vector meson sector, our Lagrangian is equivalent to the one in Ref.[10] except $SU(3)$ breaking effect for vector mesons. The chiral Lagrangian is given by,

$$\begin{aligned}
\mathcal{L} = & \frac{f^2}{4} \text{Tr} D U D U^\dagger + B \text{Tr} M (U + U^\dagger) - i g_{2p} \text{Tr} (\xi M \xi - \xi^\dagger M \xi^\dagger) \eta_0 - \frac{M_0^2}{2} \eta_0^2 \\
& + \text{Tr} D_\mu S D^\mu S - M_\sigma^2 \text{Tr} S^2 \\
& + \frac{g_1}{4} \text{Tr} (D_\mu U D^\mu U^\dagger) (\xi S \xi^\dagger) + g_2 \text{Tr} ((\xi M \xi + \xi^\dagger M \xi^\dagger) S) \\
& - \frac{1}{2} \text{Tr} F_{\mu\nu} F^{\mu\nu} + M_V^2 \text{Tr} (V_\mu - \frac{\alpha_\mu}{g})^2 + g_{1V} \text{Tr} S (V_\mu - \frac{\alpha_\mu}{g})^2, \tag{1}
\end{aligned}$$

where S and V are the scalar nonets and vector nonets respectively. (See appendix A.) U is the chiral field and is given as $U = \exp(2i\pi/f) = \xi^2$. π is $SU(3)$ octet pseudo Nambu Goldstone boson and η_0 corresponds to $U(1)_A$ pseudoscalar of which mass is denoted by M_0 . The $U(1)_A$ symmetry is broken by the mass term explicitly. The covariant derivatives for

the chiral field and the scalar field are given as,

$$D_\mu U = (\partial_\mu + iA_{L\mu})U, \quad (2a)$$

$$D_\mu S = \partial_\mu S + i[\alpha_\mu, S], \quad (2b)$$

$$\alpha_\mu = \alpha_\mu^0 + \xi^\dagger \frac{A_{L\mu}}{2} \xi, \quad (2c)$$

$$\alpha_\mu^0 = \frac{\xi^\dagger \partial_\mu \xi + \xi \partial_\mu \xi^\dagger}{2i}, \quad (2d)$$

where A_L denotes the external vector field corresponds to $SU(3)_L$. M in Eq.(1) is the chiral breaking term for the light quarks and is given by,

$$\begin{aligned} M &= \text{diag.}(m_u, m_d, m_s) \\ &= m_s \cdot \text{diag.}(\Delta, \Delta_d, 1). \end{aligned} \quad (3)$$

Δ denotes $\frac{m_u}{m_s}$ and $\Delta_d = \frac{m_d}{m_s}$. In this work, we work in the isospin limit, $m_u = m_d$. Below we explain how we determine the parameters in the Lagrangian of Eq.(1).

- B, g_1, g_2

In the isospin limit, the vacuum expectation values of the scalar fields are given as,

$$S_{01} = S_{02} = \frac{g_2 m_u}{M_\sigma^2}, \quad S_{03} = \frac{g_2 m_s}{M_\sigma^2}, \quad (4)$$

which leads to the $SU(3)$ breaking of the wave function renormalization constants and the decay constants of the pseudo Nambu Goldstone bosons,

$$Z_{ij} = 1 + g_1 \frac{S_{0i} + S_{0j}}{2f^2}, \quad (5a)$$

$$F_K = f\sqrt{Z_{13}}, \quad F_\pi = f\sqrt{Z_{11}}. \quad (5b)$$

The decay constant for η_8 is written as,

$$\begin{aligned} F_8 &= f\sqrt{\frac{Z_{11}}{3} + \frac{2Z_{33}}{3}} \\ &= F_\pi\sqrt{\frac{4}{3}R^2 - \frac{1}{3}}, \end{aligned} \quad (6)$$

where R is $\frac{F_K}{F_\pi}$. The generalized Gell-Mann Oakes Renner relation becomes,

$$m_K^2 F_K^2 = (m_u + m_s) (2B + g_2(S_{01} + S_{03})), \quad (7a)$$

$$m_\pi^2 F_\pi^2 = 2m_u (2B + 2g_2 S_{01}), \quad (7b)$$

which can be used to express g_2 , g_1 and B with Eq.(4) and Eq.(5) in terms of the physical quantities as,

$$g_2 m_s = \frac{M_\sigma F_K m_K}{\sqrt{1-\Delta}} \sqrt{\frac{1}{1+\Delta} - R^{-2} \frac{m_\pi^2}{m_K^2} \frac{1}{2\Delta}}, \quad (8a)$$

$$g_1 = 2 \frac{M_\sigma}{m_K} F_K (1 - R^{-2}) \frac{1}{\sqrt{\frac{1}{1+\Delta} - \frac{1}{2\Delta R^2} \frac{m_\pi^2}{m_K^2}}} \frac{1}{\sqrt{1-\Delta}}, \quad (8b)$$

$$B m_s = \frac{1}{4\Delta} \frac{1+\Delta}{1-\Delta} m_\pi^2 F_\pi^2 - \frac{1}{1+\Delta} \frac{\Delta}{1-\Delta} m_K^2 F_K^2. \quad (8c)$$

- η and η' mesons and octet and singlet mixing angle θ_{08}

g_{2p} in Eq.(1) leads to the η_0 and η_8 mixing. The mass matrix for η_0 and η_8 sector is diagonalized as,

$$\mathcal{L}_{08} = -\frac{1}{2} (\eta_8, \eta_0) \begin{pmatrix} M_{88}^2 & M_{08}^2 \\ M_{08}^2 & M_{00}^2 \end{pmatrix} \begin{pmatrix} \eta_8 \\ \eta_0 \end{pmatrix} = -\frac{1}{2} (\eta, \eta') \begin{pmatrix} M_\eta^2 & 0 \\ 0 & M_{\eta'}^2 \end{pmatrix} \begin{pmatrix} \eta \\ \eta' \end{pmatrix}, \quad (9a)$$

$$\begin{pmatrix} \eta \\ \eta' \end{pmatrix} = \begin{pmatrix} \cos \theta_{08} & -\sin \theta_{08} \\ \sin \theta_{08} & \cos \theta_{08} \end{pmatrix} \begin{pmatrix} \eta_8 \\ \eta_0 \end{pmatrix}. \quad (9b)$$

where $M_{08}^2 = \frac{4}{\sqrt{3}} \frac{m_s - m_u}{F_8} g_{2p}$ and we take the convention $M_{08}^2 < 0$. Beacuse the octet mass M_{88} is given by

$$M_{88}^2 = \frac{1}{3F_8^2} \left(\frac{8}{1+\Delta} M_K^2 F_K^2 - \frac{2}{\Delta} M_\pi^2 F_\pi^2 + M_\pi^2 F_\pi^2 \right), \quad (10)$$

M_{88}^2 can be determined by F_K , F_π , M_K , M_π , and Δ . With M_{88}^2 given by Eq. (10), the parameters M_{00}^2 and M_{08}^2 are also determined by using the masses of η and η' as,

$$M_{00}^2 = M_\eta^2 + M_{\eta'}^2 - M_{88}^2, \quad (11a)$$

$$M_{08}^2 = -\sqrt{M_{00}^2 M_{88}^2 - M_\eta^2 M_{\eta'}^2}. \quad (11b)$$

Therefore, one may predict the η_0 and η_8 mixing angle with the relation,

$$\theta_{08} = -\frac{1}{2} \arctan \frac{2|M_{08}^2|}{M_{00}^2 - M_{88}^2}. \quad (12)$$

The prediction of the mixing angle is rather close to the one experimentally extracted from $J/\psi \rightarrow \gamma\eta(\eta')$ decays. (See for example [11].)

$$\theta_{08\text{th}} = -22.38(\Delta = \frac{1}{24}), \quad -21.49(\Delta = \frac{1}{25}), \quad (13a)$$

$$\begin{aligned} \theta_{08\text{exp}} &= -\arctan \sqrt{\frac{\Gamma[J/\psi \rightarrow \gamma\eta]}{\Gamma[J/\psi \rightarrow \gamma\eta']}} \left(\frac{M_{J/\psi}^2 - M_{\eta'}^2}{M_{J/\psi}^2 - M_\eta^2} \right)^{\frac{3}{2}} \\ &\simeq -(22.36_{-1.21}^{+1.12}). \end{aligned} \quad (13b)$$

- Vector meson mass spectrum

The vector meson masses are given by the following formulae.

$$M_{Vij}^2 = M_V^2 + g_{1V} \frac{S_{0i} + S_{0j}}{2}. \quad (14)$$

With this formulae, the U(3) nonets vector mesons masses are given by,

$$M_\rho^2 = M_\omega^2 = M_V^2 + g_{1V} S_{01}, \quad (15a)$$

$$M_\phi^2 = M_V^2 + g_{1V} S_{03}, \quad (15b)$$

$$M_{K^*}^2 = M_V^2 + g_{1V} \frac{S_{01} + S_{03}}{2}. \quad (15c)$$

One can fix the parameter g_{1V} as,

$$g_{1V} = 2 \frac{M_\phi^2 - M_{K^*}^2}{\Delta S}, \quad (16)$$

where ΔS is the difference of the vacuum expectation values in Eq.(4),

$$\Delta S = S_{03} - S_{01} = \frac{g_2 m_s (1 - \Delta)}{M_\sigma^2}. \quad (17)$$

One can also derive the following relation by using Eq.(15),

$$M_\rho = \sqrt{2M_{K^*}^2 - M_\phi^2} \quad (18)$$

The relation leads to the prediction $M_\rho = 743$ MeV which is about -4% smaller than the measured value. In table I, we summarize the numerical values for the parameters in the chiral Lagrangian of Eq. (1).

$M_K = M_\sigma$ (MeV)	800	840	760
$g_2 m_s$ (MeV ³)	2.65×10^7	2.79×10^7	2.52×10^7
g_1 (MeV)	215	225	204
$B m_s$ (MeV ⁴)	9.24×10^8	9.24×10^8	9.24×10^8
$\Delta = \frac{m_u}{m_s}$	$\frac{1}{25}$	$\frac{1}{25}$	$\frac{1}{25}$
$\Delta S = S_{03} - S_{01}$ (MeV)	39.8	37.9	41.9
g_{1V} (MeV)	12200	12800	11600
g	5.90	5.90	5.90

TABLE I: The numerical values for the parameters in the chiral Lagrangian. We use $M_K = 494$ (MeV), $M_\pi = 135$ (MeV), $F_K = 113$ (MeV), $F_\pi = 92.2$ (MeV) and $\Gamma_{K^*} = 50.8$ (MeV) as input. g is determined with the width of K^* .

III. FORM FACTORS

The hadronic form factors relevant for the processes $\tau^+ \rightarrow \bar{\nu} K^+ P$ ($P = \pi^0, \eta, \eta'$) are,

$$\langle K^+(p_K) P(p_P) | \bar{u} \gamma_\mu s | 0 \rangle = F^{K^+P}(Q^2) q^\mu + \left(F_s^{K^+P}(Q^2) - \frac{\Delta_{KP}}{Q^2} F^{K^+P}(Q^2) \right) Q^\mu, \quad (19)$$

with $Q^\mu = (p_K + p_P)^\mu$ and $\Delta_{KP} = m_K^2 - m_P^2$. The form factor denoted by F is the vector form factor and F_s is the scalar form factor. The form factors have been computed by using the variety of the methods, Ref.[12, 13, 14, 15]. In this work, we have used the hadronic chiral Lagrangian including the vector and the scalar resonances in Eq.(1). We compute the loop corrections to the self-energy of the vector and the scalar resonances. The real part of the self-energy is divergent and we need to subtract the divergence. Corresponding to the subtractions, we have added the polynomials. Some of the coefficients of the polynomials are determined by the pole positions and the residues of the propagator for the resonances.

To compute the form factors, let us write the V-A charged current in terms of hadrons. By differentiating Eq.(1) with the external vector fields A_L , we obtain the current as,

$$\begin{aligned} \bar{q}_{jL} \gamma_\mu q_{iL} = & -i \frac{f^2}{2} (U \partial_\mu U^\dagger)_{ij} + \frac{M_V^2}{g} \left(\xi (V_\mu - \frac{\alpha_\mu^0}{g}) \xi^\dagger \right)_{ij} - i \frac{g_1}{4} \{ U \partial_\mu U^\dagger, \xi S \xi^\dagger \}_{ij} \\ & + \frac{g_{1V}}{2g} \{ S, \xi (V_\mu - \frac{\alpha_\mu^0}{g}) \xi^\dagger \}_{ij} - i (\xi [S, \partial_\mu S] \xi^\dagger)_{ij} - (\xi [S, [S, \alpha_\mu^0]] \xi^\dagger)_{ij}. \end{aligned} \quad (20)$$




FIG. 1: The Feynman diagrams contributing to the form factors

We first show the results of the form factors for $K\pi$ final state.

$$F^{K^+\pi^0}(Q^2) = \frac{1}{\sqrt{2}} \left\{ -\frac{R + R^{-1}}{2} + \frac{(\Delta S)^2}{2F_K F_\pi} + \frac{M_{K^*}^2}{2g^2 F_K F_\pi} \left(1 - \frac{M_{K^*}^2}{A_R} \right) + \frac{\Pi_{VS}^T}{2g^2 F_K F_\pi} \left(1 - \frac{2M_{K^*}^2}{A_R} \right) \right\}, \quad (21a)$$

$$F_s^{K^+\pi^0}(Q^2) = \frac{\Delta_{K\pi}}{Q^2} F^{K^+\pi^0}(Q^2) + \frac{1}{2\sqrt{2}} \left\{ R^{-1} - R + \frac{M_{K^*}^2}{g^2 F_K F_\pi} \frac{M_{K^*}^2}{A_R} \frac{\Delta_{K\pi}(B_R D_R - C_R^2)}{(A_R + Q^2 B_R) D_R - Q^2 C_R^2} + \frac{1}{\frac{Q^2 C_R^2}{A_R + Q^2 B_R} - D_R} g_{\kappa K\pi}(Q^2) \frac{\Delta S}{F_K F_\pi} + \frac{g_{\kappa K\pi}(Q^2) M_{K^*}^2 + M_{K^*}^2 \Delta_{K\pi} \Delta S}{g F_K F_\pi} \frac{C_R}{(A_R + Q^2 B_R) D_R - Q^2 C_R^2} \right\} + \frac{1}{2\sqrt{2} F_K F_\pi g^2} \left(\frac{\Delta_{K\pi}}{Q^2} (\Pi_{VS}^L - \Pi_{VS}^T) \left(1 - \frac{2M_{K^*}^2}{A_R} \right) + \frac{2\Delta_{K\pi}}{A_R} \Pi_{VS}^L \right). \quad (21b)$$

The form factors include the contribution of the Feynman diagrams shown in Fig. 1. In Fig. 1, the propagators for K^* and κ mesons are represented by the thick solid lines which include the one loop corrections to the self-energy. Let us consider the propagators for K^* and κ . They are obtained by inverting the inverse propagators for K^* and κ .

$$\begin{pmatrix} (g^{\mu\nu} A_R(Q^2) + Q^\mu Q^\nu B_R(Q^2)) & Q^\mu C_R(Q^2) \\ Q^\nu C_R(Q^2) & D_R(Q^2) \end{pmatrix} \begin{pmatrix} K_\nu \\ \kappa \end{pmatrix} = \begin{pmatrix} J^\mu \\ J \end{pmatrix}, \quad (22)$$

where J_μ and J are source terms for K^* and κ respectively. $A_R g^{\mu\nu} + B_R Q^\mu Q^\nu$ is the inverse propagator for K^* and D_R denotes the inverse propagator for the κ . C_R denotes the mixing between the K^* and κ . Inverting Eq. (22), one can obtain the propagator.

$$\begin{pmatrix} K^\mu \\ \kappa \end{pmatrix} = \begin{pmatrix} g^{\mu\nu} A_R^{-1} + \frac{Q^\mu Q^\nu}{Q^2} \left(\frac{D_R}{(A_R + Q^2 B_R) D_R - Q^2 C_R^2} - A_R^{-1} \right) & - \frac{Q^\mu C_R}{(A_R + Q^2 B_R) D_R - Q^2 C_R^2} \\ - \frac{Q^\nu C_R}{(A_R + Q^2 B_R) D_R - Q^2 C_R^2} & \frac{A_R + Q^2 B_R}{(A_R + Q^2 B_R) D_R - Q^2 C_R^2} \end{pmatrix} \begin{pmatrix} J_\nu \\ J \end{pmatrix}. \quad (23)$$

To obtain the contributions to the form factors from Feynman diagrams Fig.(1-a) ~ Fig.(1-e), we set $i = s, j = u$ in Eq. (20),

$$\begin{aligned} \overline{u_L} \gamma_\mu s_L &= -\frac{1}{\sqrt{2}} F_K \partial_\mu K^- + \frac{M_{K^*}^2}{\sqrt{2}g} (K_\mu^{*-} - ig \frac{\Delta S}{M_{K^*}^2} \partial_\mu \kappa^-) \\ &- i \frac{1}{2\sqrt{2}} \left(K^- \partial_\mu \pi^0 \left(\frac{F_\pi}{F_K} - \frac{M_{K^*}^2 + g^2 (\Delta S)^2}{2g^2 F_K F_\pi} \right) - \partial_\mu K^- \pi^0 \left(\frac{F_K}{F_\pi} - \frac{M_{K^*}^2 + g^2 (\Delta S)^2}{2g^2 F_K F_\pi} \right) \right) \\ &- i \frac{\sqrt{3}}{2\sqrt{2}} \left(K^- \partial_\mu \eta_8 \left(\frac{F_8}{F_K} - \frac{M_{K^*}^2 + g^2 (\Delta S)^2}{2g^2 F_K F_8} \right) - \partial_\mu K^- \eta_8 \left(\frac{F_K}{F_8} - \frac{M_{K^*}^2 + g^2 (\Delta S)^2}{2g^2 F_K F_8} \right) \right) \\ &- i \frac{1}{2} \left(\overline{K^0} \partial_\mu \pi^- \left(\frac{F_\pi}{F_K} - \frac{M_{K^*}^2 + g^2 (\Delta S)^2}{2g^2 F_K F_\pi} \right) - \partial_\mu \overline{K^0} \pi^- \left(\frac{F_K}{F_\pi} - \frac{M_{K^*}^2 + g^2 (\Delta S)^2}{2g^2 F_K F_\pi} \right) \right) \\ &+ \frac{g_{1V}}{4g} \left(\frac{1}{\sqrt{2}} \kappa^- (\rho_\mu + \omega_\mu) + \overline{\kappa^0} \rho_\mu^- + f^0 K_\mu^{*-} \right. \\ &\left. + \frac{1}{\sqrt{2}} K_\mu^{*-} (a^0 + \sigma) + \overline{K_\mu^{*0}} a^- + \phi_\mu \kappa^- \right). \quad (24) \end{aligned}$$

For the diagram in Fig.(1-a), the direct coupling of the charged current to $K\pi$ can be easily extracted from Eq. (24). For the other diagrams, the amplitudes are obtained by multiplying the propagators in Eq. (23) with the production amplitudes of K^* and κ and the amplitudes corresponding to their decays.

The matrix elements corresponding to Fig.(1-a) ~ Fig.(1-e) are given as,

$$\begin{aligned} \langle K^+ \pi^0 | \overline{u} \gamma_\mu s | 0 \rangle_{(1-a)} &= \frac{1}{\sqrt{2}} \left\{ Q_\mu \frac{R^{-1} - R}{2} + q_\mu \left(-\frac{R^{-1} + R}{2} + \frac{M_{K^*}^2 + g^2 (\Delta S)^2}{2g^2 F_K F_\pi} \right) \right\}, \\ \langle K^+ \pi^0 | \overline{u} \gamma_\mu s | 0 \rangle_{(1-b)} &= \frac{M_{K^*}^4}{2\sqrt{2}g^2 F_K F_\pi} \left(-q_\mu A_R^{-1} + Q_\mu \frac{\Delta_{K\pi}}{A_R} \frac{B_R D_R - C_R^2}{(A_R + Q^2 B_R) D_R - Q^2 C_R^2} \right), \\ \langle K^+ \pi^0 | \overline{u} \gamma_\mu s | 0 \rangle_{(1-c)} &= -\frac{\Delta S g_{\kappa K\pi} (Q^2)}{2\sqrt{2} F_K F_\pi} \frac{1}{D_R - \frac{Q^2 C_R^2}{A_R + Q^2 B_R}} Q_\mu, \\ \langle K^+ \pi^0 | \overline{u} \gamma_\mu s | 0 \rangle_{(1-d)} &= \frac{M_{K^*}^2}{2\sqrt{2}g F_K F_\pi} \frac{\Delta S \Delta_{K\pi} C_R}{(A_R + Q^2 B_R) D_R - Q^2 C_R^2} Q_\mu, \\ \langle K^+ \pi^0 | \overline{u} \gamma_\mu s | 0 \rangle_{(1-e)} &= \frac{M_{K^*}^2}{2\sqrt{2}g F_K F_\pi} \frac{g_{\kappa K\pi} (Q^2) C_R}{(A_R + Q^2 B_R) D_R - Q^2 C_R^2} Q_\mu. \quad (25) \end{aligned}$$




FIG. 2: The Feynman diagrams of scalar and vector mesons loop which contribute to the form factor for $\langle KP|\bar{u}\gamma^\mu s|0\rangle$. They can be written in terms of the self-energy correction function Π_{VS} .

To derive Eq. (25), we have used the production amplitudes for K^* and κ due to the vector current $\bar{u}\gamma_\mu s$,

$$\begin{aligned}\langle K^*_\rho|\bar{u}\gamma_\mu s|0\rangle &= g_{\rho\mu}\sqrt{2}\frac{M_{K^*}^2}{g}, \\ \langle \kappa|\bar{u}\gamma_\mu s|0\rangle &= Q_\mu\sqrt{2}\Delta S.\end{aligned}\quad (26)$$

We also have used the strong interaction vertices which are given as,

$$\begin{aligned}\langle K^+\pi^0|\mathcal{L}_{VPP}|K^{*+}\rangle &= \frac{M_{K^*}^2}{4gF_KF_\pi}q_\sigma, \\ \langle K^+\pi^0|\mathcal{L}_{SPP}|\kappa^+\rangle &= \frac{1}{4F_KF_\pi}g_{\kappa K\pi}(Q^2),\end{aligned}\quad (27)$$

with $q = p_K - p_\pi$. $g_{\kappa K\pi}(Q^2)$ is the strong coupling for $\kappa \rightarrow K\pi$ defined by,

$$g_{\kappa K\pi}(Q^2) = g_1\frac{m_K^2 + m_\pi^2 - Q^2}{2} - g_2(3m_u + m_s) + \Delta_{K\pi}(\Delta S).\quad (28)$$

In addition to the pseudoscalar loops, we have taken into account the vector and scalar meson loops denoted by Π_{VS} . Each contribution is given by,

$$\begin{aligned}\langle K^+\pi^0|\bar{u}\gamma^\mu s|0\rangle|_{(2-a)} &= \frac{q_\nu\Pi_{VS}^{\nu\mu}}{2\sqrt{2}F_KF_\pi g^2}, \\ \langle K^+\pi^0|\bar{u}\gamma^\mu s|0\rangle|_{(2-b)} &= -\frac{M_{K^*}^2}{A}q_\nu\Pi_{VS}^{\nu\rho}(\delta_\rho^\mu - \frac{Q_\rho Q^\mu}{M_{K^*}^2})\frac{1}{2\sqrt{2}F_KF_\pi g^2}, \\ \langle K^+\pi^0|\bar{u}\gamma^\mu s|0\rangle|_{(2-c)} &= -\frac{M_{K^*}^2}{A}q^\rho(g_{\rho\nu} - \frac{Q_\rho Q_\nu}{M_{K^*}^2})\frac{1}{2\sqrt{2}F_KF_\pi g^2}\Pi_{VS}^{\nu\mu},\end{aligned}\quad (29)$$

where Π_{VS} is identical to the self-energy function in Fig. (3-d),

$$\Pi_{VS}^{\mu\nu} = (g^{\mu\nu} - \frac{Q^\mu Q^\nu}{Q^2})\Pi_{VS}^T + \frac{Q^\mu Q^\nu}{Q^2}\Pi_{VS}^L.\quad (30)$$

By denoting the self-energy corrections as $\delta A_R, \delta B_R, C_R$ and Π_{VS} corresponding to the




FIG. 3: (3-a): The pseudoscalar meson loop corrections to the self-energy for K^* ($\delta A_R, \delta B_R$). (3-b): The self-energy for κ (δD_R). (3-c): The mixing term (C_R). (3-d): Π^{VS} .

Feynman diagrams in Fig.3, one obtains

$$\begin{aligned}
 A_R(s) &= M_{K^*}^2 - s - iM_{K^*}\Gamma_{K^*}(s) + \text{Re.}(\delta A_R(s)) + \Pi_{VS}^T + (a_0 + k_0) + (a_1 + k_1)s, \\
 sB_R(s) &= s(1 + b_0 + l_1) + i \left(\frac{\nu_{K\pi}^3}{48\pi s^2} + \frac{\nu_{K\pi}\Delta_{K\pi}^2}{16\pi s^2} \right) 3g_{K^*K\pi}^2 \\
 &+ i \left(\frac{\nu_{K\eta}^3}{48\pi s^2} + \frac{\nu_{K\eta}\Delta_{K\eta}^2}{16\pi s^2} \right) 3g_{K^*K\pi}^2 \left(\frac{F_\pi}{F_8} \right)^2 \cos^2 \theta_{08} \\
 &+ i \left(\frac{\nu_{K\eta'}^3}{48\pi s^2} + \frac{\nu_{K\eta'}\Delta_{K\eta'}^2}{16\pi s^2} \right) 3g_{K^*K\pi}^2 \left(\frac{F_\pi}{F_8} \right)^2 \sin^2 \theta_{08} + s\text{Re.}(\delta B_R(s)) + \Pi_{SV}^L - \Pi_{SV}^T, \\
 C_R(s) &= \text{Re.}(C_R(s)) + c_0 + i \frac{\nu_{K\pi}(s)}{16\pi s^2} \frac{\Delta_{K\pi}}{4F_K F_\pi} 3g_{\kappa K\pi}(s) g_{K^*K\pi} \\
 &- i \frac{\nu_{K\eta}(s)}{16\pi s^2} \frac{\Delta_{K\eta} F_\pi}{4F_K F_8^2} g_{\kappa K\eta}(s) g_{K^*K\pi} \cos^2 \theta_{08} \\
 &- i \frac{\nu_{K\eta'}(s)}{16\pi s^2} \frac{\Delta_{K\eta'} F_\pi}{4F_K F_8^2} g_{\kappa K\eta'}(s) g_{K^*K\pi} \sin^2 \theta_{08}, \\
 D_R(s) &= s - M_\kappa^2 + iM_\kappa\Gamma_\kappa(s) + \text{Re.}(\delta D_R(s)) + d_0 + d_1s + d_2s^2, \tag{31}
 \end{aligned}$$

where $s = Q^2$ and $g_{K^*K\pi} = \frac{M_{K^*}^2}{4gF_K F_\pi}$. The momentum dependent widths $\Gamma_{K^*}(s)$ and $\Gamma_\kappa(s)$ are given as,

$$\begin{aligned}
 \Gamma_{K^*}(s) &= 3 \frac{1}{48\pi M_{K^*}} \left(\frac{\nu_{K\pi}^3}{s^2} + \cos^2 \theta_{08} \frac{\nu_{K\eta}^3}{s^2} \left(\frac{F_\pi}{F_8} \right)^2 + \sin^2 \theta_{08} \frac{\nu_{K\eta'}^3}{s^2} \left(\frac{F_\pi}{F_8} \right)^2 \right) g_{K^*K\pi}^2, \\
 \Gamma_\kappa(s) &= 3 \frac{\nu_{K\pi} g_{\kappa K\pi}^2(s)}{16\pi s M_\kappa} \left(\frac{1}{4F_K F_\pi} \right)^2 \\
 &+ \left(\cos^2 \theta_{08} \frac{\nu_{K\eta} g_{\kappa K\eta}^2(s)}{16\pi s M_\kappa} + \sin^2 \theta_{08} \frac{\nu_{K\eta'} g_{\kappa K\eta'}^2(s)}{16\pi s M_\kappa} \right) \left(\frac{1}{4\sqrt{3}F_K F_8} \right)^2, \tag{32}
 \end{aligned}$$

where $\nu_{KP} = \sqrt{s^2 - 2s(m_K^2 + m_P^2) + \Delta_{KP}^2}$ ($P = \pi, \eta, \eta'$) and is related to the momentum of kaon p_K in the hadronic rest frame as $p_K = \frac{\nu_{KP}}{2\sqrt{s}}$. The real part of the self-energy corrections are divergent. We have subtracted the divergences at zero momentum as,

$$\begin{aligned}\delta A_R(s) &= \delta A(s) - \delta A(0) - sA'(0), \\ \delta B_R(s) &= \delta B(s) - \delta B(0), \\ C_R(s) &= C(s) - C(0), \\ \delta D_R(s) &= \delta D(s) - \delta D(0) - sD'(0) - \frac{s^2}{2}D''(0).\end{aligned}\quad (33)$$

Then we have added the polynomials with respect to s which coefficients are finite renormalization constants. We have added the polynomial $a_0 + a_1s$ which corresponds to the twice subtractions for δA_R in Eq. (33). For δB_R and C_R , we have added a finite constant to each denoted by b_0 and c_0 respectively. About the self-energy correction of the scalar meson $\delta D_R(s)$, we need to subtract divergences up to s^2 . Therefore, we need to add the polynomial $d_0 + d_1s + d_2s^2$ which is quadratic with respect to s .

The self-energy corrections in Eq. (33), $\delta A_R \sim C_R$, are given by,

$$\begin{aligned}\delta A_R(s) &= \\ &\frac{3g_{K^*K\pi}^2}{16\pi^2} \left\{ \left(-\Sigma_{K\pi} \left(\frac{1}{2} + \log \frac{s}{m_K m_\pi} \right) + \frac{s}{3} \left(-1 + \log \frac{s}{m_K m_\pi} \right) - 2sR_{K\pi} \right. \right. \\ &- \frac{m_K^6 + m_\pi^6 - 3m_K^2 m_\pi^2 \Sigma_{K\pi}}{3\Delta_{K\pi}^3} \log \frac{m_K}{m_\pi} + \frac{s}{18\Delta_{K\pi}^2} (5m_K^4 + 5m_\pi^4 - 22m_K^2 m_\pi^2) \\ &+ \frac{\Sigma_{K\pi}^2 + \Delta_{K\pi}^2}{4\Delta_{K\pi}} \log \frac{m_K^2}{m_\pi^2} \left. \right) + \left(\frac{F_\pi}{F_8} \right)^2 \cos^2 \theta_{08} \left(\frac{s}{18\Delta_{K\eta}^2} (5m_K^4 + 5m_\eta^4 - 22m_K^2 m_\eta^2) \right. \\ &- \Sigma_{K\eta} \left(\frac{1}{2} + \log \frac{s}{m_K m_\eta} \right) + \frac{\Sigma_{K\eta}^2 + \Delta_{K\eta}^2}{4\Delta_{K\eta}} \log \frac{m_K^2}{m_\eta^2} + \frac{s}{3} \left(-1 + \log \frac{s}{m_K m_\eta} \right) \\ &- \left. 2sR_{K\eta} - s \frac{m_K^6 + m_\eta^6 - 3m_K^2 m_\eta^2 \Sigma_{K\eta}}{3\Delta_{K\eta}^3} \log \frac{m_K}{m_\eta} \right) + \sin^2 \theta_{08} \left(\frac{F_\pi}{F_8} \right)^2 (-2sR_{K\eta'}) \\ &- \Sigma_{K\eta'} \left(\frac{1}{2} + \log \frac{s}{m_K m_{\eta'}} \right) + \frac{s}{3} \left(-1 + \log \frac{s}{m_K m_{\eta'}} \right) + \frac{\Sigma_{K\eta'}^2 + \Delta_{K\eta'}^2}{4\Delta_{K\eta'}} \log \frac{m_K^2}{m_{\eta'}^2} \\ &- \left. \left. s \frac{m_K^6 + m_{\eta'}^6 - 3m_K^2 m_{\eta'}^2 \Sigma_{K\eta'}}{3\Delta_{K\eta'}^3} \log \frac{m_K}{m_{\eta'}} + \frac{s}{18\Delta_{K\eta'}^2} (5m_K^4 + 5m_{\eta'}^4 - 22m_K^2 m_{\eta'}^2) \right) \right\},\end{aligned}\quad (34)$$

$$\begin{aligned}
\delta B_R(s) = & \frac{3g_{K^*K\pi}^2}{16\pi^2} \left\{ \left(-\frac{1}{3} \log \frac{s}{m_K m_\pi} - R'_{K\pi} + \frac{\Sigma_{K\pi}^3}{6\Delta_{K\pi}^3} \log \frac{m_K^2}{m_\pi^2} - \frac{4}{9} \frac{m_K^4 + m_K^2 m_\pi^2 + m_\pi^4}{\Delta_{K\pi}^2} \right) + \right. \\
& \cos^2 \theta_{08} \left(\frac{F_\pi}{F_8} \right)^2 \left(-\frac{1}{3} \log \frac{s}{m_K m_\eta} - R'_{K\eta} + \frac{\Sigma_{K\eta}^3}{6\Delta_{K\eta}^3} \log \frac{m_K^2}{m_\eta^2} - \frac{4}{9} \frac{m_K^4 + m_K^2 m_\eta^2 + m_\eta^4}{\Delta_{K\eta}^2} \right) + \\
& \left. \sin^2 \theta_{08} \left(\frac{F_\pi}{F_8} \right)^2 \left(-\frac{1}{3} \log \frac{s}{m_K m_{\eta'}} - R'_{K\eta'} + \frac{\Sigma_{K\eta'}^3}{6\Delta_{K\eta'}^3} \log \frac{m_K^2}{m_{\eta'}^2} - \frac{4}{9} \frac{m_K^4 + m_K^2 m_{\eta'}^2 + m_{\eta'}^4}{\Delta_{K\eta'}^2} \right) \right\}, \tag{35}
\end{aligned}$$

$$\begin{aligned}
C_R(s) = & \frac{3g_{K^*K\pi}}{64\pi^2 F_K F_\pi} \left\{ -g_{\kappa K\pi}(s)(2R_{K\pi}^1 - R_{K\pi}^0) + g_{\kappa K\pi}(0) \left(\frac{m_K^2 m_\pi^2 \log \frac{m_K^2}{m_\pi^2}}{\Delta_{K\pi}^2} - \frac{\Sigma_{K\pi}}{2\Delta_{K\pi}} \right) \right\} - \\
& \cos^2 \theta_{08} \frac{3g_{K^*K\pi} F_\pi}{64\pi^2 F_K F_8^2} \left\{ -g_{\kappa K\eta}(s)(2R_{K\eta}^1 - R_{K\eta}^0) + g_{\kappa K\eta}(0) \left(\frac{m_K^2 m_\eta^2 \log \frac{m_K^2}{m_\eta^2}}{\Delta_{K\eta}^2} - \frac{\Sigma_{K\eta}}{2\Delta_{K\eta}} \right) \right\} - \\
& \sin^2 \theta_{08} \frac{3g_{K^*K\pi} F_\pi}{64\pi^2 F_K F_8^2} \left\{ -g_{\kappa K\eta'}(s)(2R_{K\eta'}^1 - R_{K\eta'}^0) + g_{\kappa K\eta'}(0) \left(\frac{m_K^2 m_{\eta'}^2 \log \frac{m_K^2}{m_{\eta'}^2}}{\Delta_{K\eta'}^2} - \frac{\Sigma_{K\eta'}}{2\Delta_{K\eta'}} \right) \right\}, \tag{36}
\end{aligned}$$

where R_{PQ} and R'_{PQ} are defined as,

$$\begin{aligned}
R_{PQ} &= \int_0^1 dx \left(x^2 - x \left(1 + \frac{\Delta_{PQ}}{s} \right) + \frac{M_P^2}{s} \right) \log \left(x^2 - x \left(1 + \frac{\Delta_{PQ}}{s} \right) + \frac{M_P^2}{s} - i\epsilon \right), \\
R'_{PQ} &= 4R_{PQ}^2 - 4R_{PQ}^1 + R_{PQ}^0, \tag{37}
\end{aligned}$$

with

$$R_{PQ}^{(n)} = \int_0^1 dx \quad x^n \log \left(x^2 - x \left(1 + \frac{\Delta_{PQ}}{s} \right) + \frac{M_P^2}{s} - i\epsilon \right). \tag{38}$$

We give the explicit forms for $R_{PQ}^{(n)}$ ($n = 0 \sim 2$) in appendix B. The inverse propagator for

the scalar meson is,

$$\begin{aligned}
\delta D_R(s) = & \frac{3}{(4F_K F_\pi)^2} \left\{ g_{\kappa K\pi}^2(s) \bar{J}_{K\pi}(s) - g_{\kappa K\pi}^2(0) J'_{K\pi}(0) s - s^2 (2g_{\kappa K\pi}(0) g'_{\kappa K\pi}(0) J'_{K\pi}(0) \right. \\
& \left. + \frac{1}{2} g_{\kappa K\pi}(0)^2 J''_{K\pi}(0)) \right\} + \frac{\cos^2 \theta_{08}}{(4\sqrt{3} F_K F_8)^2} \left\{ g_{\kappa K\eta}^2(s) \bar{J}_{K\eta}(s) - g_{\kappa K\eta}^2(0) J'_{K\eta}(0) s \right. \\
& \left. - s^2 \left(2g_{\kappa K\eta}(0) g'_{\kappa K\eta}(0) J'_{K\eta}(0) + \frac{1}{2} g_{\kappa K\eta}(0)^2 J''_{K\eta}(0) \right) \right\} \\
& + \frac{\sin^2 \theta_{08}}{(4\sqrt{3} F_K F_8)^2} \left\{ g_{\kappa K\eta'}^2(s) \bar{J}_{K\eta'}(s) - g_{\kappa K\eta'}^2(0) J'_{K\eta'}(0) s \right. \\
& \left. - s^2 \left(2g_{\kappa K\eta'}(0) g'_{\kappa K\eta'}(0) J'_{K\eta'}(0) + \frac{1}{2} g_{\kappa K\eta'}(0)^2 J''_{K\eta'}(0) \right) \right\}, \tag{39}
\end{aligned}$$

where

$$\begin{aligned}
\bar{J}_{PQ}(s) = & \frac{\theta(s - (M_P + M_Q)^2)}{32\pi^2} \left\{ 2 + \left(\frac{\Delta_{PQ}}{s} - \frac{\Sigma_{PQ}}{\Delta_{PQ}} \right) \log \frac{M_Q^2}{M_P^2} - \frac{\nu_{PQ}}{s} \log \frac{(s + \nu_{PQ})^2 - \Delta_{PQ}^2}{(s - \nu_{PQ})^2 - \Delta_{PQ}^2} + \right. \\
& \left. i2\pi \frac{\nu_{PQ}}{s} \right\} + \frac{\theta((M_P + M_Q)^2 - s) \theta(s - (M_P - M_Q)^2)}{32\pi^2} \left\{ \left(\frac{\Delta_{PQ}}{s} - \frac{\Sigma_{PQ}}{\Delta_{PQ}} \right) \log \frac{M_Q^2}{M_P^2} \right. \\
& \left. + 2 + \frac{2\sqrt{-\nu_{PQ}^2}}{s} \left(\tan^{-1} \frac{s - \Delta_{PQ}}{\sqrt{-\nu_{PQ}^2}} + \tan^{-1} \frac{s + \Delta_{PQ}}{\sqrt{-\nu_{PQ}^2}} \right) \right\}, \tag{40}
\end{aligned}$$

with $\Sigma_{PQ} = m_P^2 + m_Q^2$ and $\nu_{PQ}^2 = s^2 - 2s\Sigma_{PQ} + \Delta_{PQ}^2 = (s - (M_P + M_Q)^2)(s - (M_P - M_Q)^2)$.

We also note,

$$\begin{aligned}
J'_{PQ}(0) = & \frac{1}{32\pi^2} \left\{ \frac{\Sigma_{PQ}}{\Delta_{PQ}^2} + 2 \frac{M_P^2 M_Q^2}{\Delta_{PQ}^3} \log \frac{M_Q^2}{M_P^2} \right\}, \\
J''_{PQ}(0) = & \frac{1}{32\pi^2} \left\{ \frac{2}{3\Delta_{PQ}^4} (3\Sigma_{PQ}^2 - 2\Delta_{PQ}^2) + 4 \frac{M_P^2 M_Q^2}{\Delta_{PQ}^5} \Sigma_{PQ} \log \frac{M_Q^2}{M_P^2} \right\}. \tag{41}
\end{aligned}$$

The self-energy corrections due to the vector and the scalar meson loop are also divergent,

$$\begin{aligned}
\Pi^{\mu\nu} = & -\frac{g_{1V}^2}{8} \sum_{VS} C_{VS} \mu^{4-d} \int \frac{d^d k}{(2\pi)^d i} \frac{g^{\mu\nu} - \frac{k^\mu k^\nu}{m_V^2}}{(k^2 - m_V^2)((Q - k)^2 - m_S^2)} \\
= & -\frac{g_{1V}^2}{8} \sum_{VS} C_{VS} (g^{\mu\nu} K_1 - Q^\mu Q^\nu K_2), \tag{42}
\end{aligned}$$

where C_{VS} are factors determined by scalar and vector mesons which contribute to the loop and are given as,

$$(C_{K^* \sigma}, C_{\omega \kappa}, C_{\rho \kappa}, C_{K^* a}, C_{\phi \kappa}, C_{K^* f^0}) = \left(\frac{1}{2}, \frac{1}{2}, \frac{3}{2}, \frac{3}{2}, 1, 1 \right). \tag{43}$$

We subtracted the divergences of K_1 and K_2 as,

$$\begin{aligned} K_{1VS} &= K_1(s) - K_1(0) - K_1'(0)s, \\ K_{2VS} &= K_2(s) - K_2(0). \end{aligned} \quad (44)$$

Using K_{1VS} and K_{2VS} , we write the self-energies Π_{VS}^T and Π_{VS}^L in Eq. (30) as,

$$\begin{aligned} \Pi_{VS}^T &= -\frac{g_{1V}^2}{8} \sum_{VS} C_{VS} K_{1VS}, \\ \frac{\Pi_{VS}^L - \Pi_{VS}^T}{s} &= \frac{g_{1V}^2}{8} \sum_{VS} C_{VS} K_{2VS}. \end{aligned} \quad (45)$$

The explicit forms for K_{iVS} ($i = 1, 2$) are given as,

$$\begin{aligned} K_{1VS} &= \frac{1}{16\pi^2} \left\{ -\frac{1}{2} R_{VS}^{(0)} - \frac{s + \Delta_{VS}}{2M_V^2} R_{VS}^{(1)} + \frac{s}{2M_V^2} R_{VS}^{(2)} - 1 + \frac{s}{12M_V^2} + \frac{\Sigma_{VS}}{8M_V^2} - \frac{\Sigma_{VS}}{2s} \right. \\ &\quad + \frac{s}{12M_V^2} \left(-\frac{3}{2} + \frac{2M_V^4 + M_V^2 M_S^2 + M_S^4}{\Delta_{VS}^2} \right) + \log \frac{M_V M_S}{s} \left(1 - \frac{\Sigma_{VS}}{4M_V^2} + \frac{s}{12M_V^2} \right) \\ &\quad \left. + \log \frac{M_V}{M_S} \left(\frac{\Sigma_{VS}}{\Delta_{VS}} - \frac{M_V^4 + M_S^4}{4\Delta_{VS} M_V^2} + \frac{2M_V^2 M_S^2}{s\Delta_{VS}} + \frac{s}{12M_V^2} \frac{M_V^6 + M_S^6 - 3M_V^2 M_S^2 \Sigma_{VS}}{\Delta_{VS}^3} \right) \right\}, \\ K_{2VS} &= \frac{1}{16\pi^2 M_V^2} \frac{1}{\Delta_{VS}^3} \left\{ \frac{M_V^6 \log \frac{M_V^2}{s}}{3} - \frac{11}{18} M_V^6 - \frac{M_S^6 \log \frac{M_S^2}{s}}{3} + \frac{M_S^6}{9} \right. \\ &\quad \left. - \frac{M_V^2 M_S^4}{2} + M_V^4 M_S^2 - M_V^2 M_S^2 \Delta_{VS} \log \frac{M_S^2}{s} \right\} - \frac{1}{16\pi^2 M_V^2} R_{VS}^{(2)}. \end{aligned} \quad (46)$$

The absorptive parts of K_{iVS} ($i = 1, 2$) are written as,

$$\begin{aligned} \text{Im}(K_{1VS}) &= \frac{\nu_{VS}}{32\pi s} \left(1 + \frac{s}{6M_V^2} + \frac{\Delta_{VS}}{2M_V^2} + \frac{\Sigma_{VS}}{6M_V^2} + \frac{\Delta_{VS}^2}{6sM_V^2} \right), \\ \text{Im}(K_{2VS}) &= \frac{\nu_{VS}}{48\pi s M_V^2} \left(1 - \frac{\Sigma_{VS}}{2s} + \frac{3\Delta_{VS}}{2s} + \frac{\Delta_{VS}^2}{s^2} \right). \end{aligned} \quad (47)$$

In A_R and B_R of Eq.(32), we have added the polynomial $l_1 s$ to $\Pi_{VS}^L - \Pi_{VS}^T$ and $k_0 + k_1 s$ to Π_{VS}^T . The polynomials corresponds to the subtraction of the divergent parts of Eq. (44).

The above derivation of the form factors for $K\pi$ final state is easily extended to $K\eta$ and $K\eta'$ case.

$$\begin{aligned} F^{K^+\eta}(Q^2) &= \cos \theta_{08} \frac{\sqrt{3}}{\sqrt{2}} \left\{ -\frac{R_8 + R_8^{-1}}{2} + \frac{(\Delta S)^2}{2F_K F_8} + \right. \\ &\quad \left. \frac{\Pi_{VS}^T}{2g^2 F_K F_8} \left(1 - \frac{2M_{K^*}^2}{A_R} \right) + \frac{M_{K^*}^2}{2g^2 F_K F_8} \left(1 - \frac{M_{K^*}^2}{A_R} \right) \right\}, \end{aligned} \quad (48a)$$

$$\begin{aligned}
F_s^{K^+\eta}(Q^2) &= \frac{\Delta_{K\eta}}{Q^2} F^{K^+\eta}(Q^2) \\
&+ \cos\theta_{08} \frac{\sqrt{3}}{2\sqrt{2}} \left\{ R_8^{-1} - R_8 + \frac{M_{K^*}^2}{g^2 F_K F_8} \frac{M_{K^*}^2}{A_R} \frac{\Delta_{K\eta}(B_R D_R - C_R^2)}{(A_R + Q^2 B_R) D_R - Q^2 C_R^2} \right. \\
&- \frac{1}{\frac{Q^2 C_R^2}{A_R + Q^2 B_R} - D_R} g_{\kappa K\eta}(Q^2) \frac{\Delta S}{3 F_K F_8} \\
&- \frac{g_{\kappa K\eta}(Q^2) M_{K^*}^2 - 3 M_{K^*}^2 \Delta_{K\eta} \Delta S}{3 g F_K F_8} \frac{C_R}{(A_R + Q^2 B_R) D - Q^2 C^2} \\
&\left. + \frac{1}{F_K F_8 g^2} \left(\frac{\Delta_{K\eta}}{s} (\Pi_{VS}^L - \Pi_{VS}^T) \left(1 - \frac{2 M_{K^*}^2}{A_R} \right) + \frac{2 \Delta_{K\eta}}{A_R} \Pi_{VS}^L \right) \right\}, \quad (48b)
\end{aligned}$$

$$\begin{aligned}
F^{K^+\eta'}(Q^2) &= \sin\theta_{08} \frac{\sqrt{3}}{\sqrt{2}} \left\{ -\frac{R_8 + R_8^{-1}}{2} + \frac{(\Delta S)^2}{2 F_K F_8} + \frac{M_{K^*}^2}{2 g^2 F_K F_8} \left(1 - \frac{M_{K^*}^2}{A_R} \right) + \right. \\
&\left. \frac{\Pi_{VS}^T}{2 g^2 F_K F_8} \left(1 - \frac{2 M_{K^*}^2}{A_R} \right) \right\}, \quad (49a)
\end{aligned}$$

$$\begin{aligned}
F_s^{K^+\eta'}(Q^2) &= \frac{\Delta_{K\eta'}}{Q^2} F^{K^+\eta'}(Q^2) \\
&+ \sin\theta_{08} \frac{\sqrt{3}}{2\sqrt{2}} \left\{ R_8^{-1} - R_8 + \frac{M_{K^*}^2}{g^2 F_K F_8} \frac{M_{K^*}^2}{A_R} \frac{\Delta_{K\eta'}(B_R D_R - C_R^2)}{(A_R + Q^2 B_R) D - Q^2 C_R^2} \right. \\
&- \frac{1}{\frac{Q^2 C_R^2}{A_R + Q^2 B_R} - D_R} g_{\kappa K\eta'}(Q^2) \frac{\Delta S}{3 F_K F_8} \\
&- \frac{g_{\kappa K\eta'}(Q^2) M_{K^*}^2 - 3 M_{K^*}^2 \Delta_{K\eta'} \Delta S}{3 g F_K F_8} \frac{C_R}{(A_R + Q^2 B_R) D_R - Q^2 C_R^2} \\
&\left. + \frac{1}{F_K F_8 g^2} \left(\frac{\Delta_{K\eta'}}{s} (\Pi_{VS}^L - \Pi_{VS}^T) \left(1 - \frac{2 M_{K^*}^2}{A_R} \right) + \frac{2 \Delta_{K\eta'}}{A_R} \Pi_{VS}^L \right) \right\}, \quad (49b)
\end{aligned}$$

with $R_8 = \frac{F_K}{F_8}$ and $\Delta_{K\eta^{(\prime)}} = m_K^2 - m_{\eta^{(\prime)}}^2$. To derive Eq. (48a) and Eq. (49a), we used the strong interaction vertices for $K^* \rightarrow K\eta$ and $K^* \rightarrow K\eta'$,

$$\begin{aligned}
\langle K^+\eta | \mathcal{L}_{VPP} | K_{\sigma}^{*+} \rangle &= \cos\theta_{08} \frac{M_{K^*}^2}{4g F_K F_8} \sqrt{3} q_{\sigma}, \\
\langle K^+\eta' | \mathcal{L}_{VPP} | K_{\sigma}^{*+} \rangle &= \sin\theta_{08} \frac{M_{K^*}^2}{4g F_K F_8} \sqrt{3} q_{\sigma}, \quad (50)
\end{aligned}$$

where $q = p_k - p_P$ ($P = \eta, \eta'$). We also use $\kappa \rightarrow K\eta^{(\prime)}$ vertices which are given by,

$$\begin{aligned}
\langle K^+\eta | \mathcal{L}_{SPP} | \kappa^+(Q) \rangle &= \frac{-\cos\theta_{08}}{4\sqrt{3} F_K F_8} g_{\kappa K\eta}(Q^2), \\
\langle K^+\eta' | \mathcal{L}_{SPP} | \kappa^+(Q) \rangle &= \frac{-\sin\theta_{08}}{4\sqrt{3} F_K F_8} g_{\kappa K\eta'}(Q^2), \\
g_{\kappa K\eta^{(\prime)}}(Q^2) &= g_1 \frac{m_K^2 + m_{\eta^{(\prime)}}^2 - Q^2}{2} - g_2 (5m_s - m_u) - 3\Delta_{K\eta^{(\prime)}}(\Delta S). \quad (51)
\end{aligned}$$

m_K	493.7	$m_{\eta'}$	957.8
m_π	135.0	M_ρ	775.5
M_{K^*}	891.7	M_τ	1777
M_ϕ	1019	Γ_τ	2.26×10^{-9}
M_ω	782.7	V_{us}	0.2257
m_η	547.5	Γ_{K^*}	50.8

TABLE II: The numerical values for masses,widths and V_{us} used in our numerical analysis. The units of mass and width are MeV.

IV. NUMERICAL ANALYSIS OF THE FORM FACTORS

In this section, we give the numerical results of the form factors. We summarize the numerical values for the masses of hadrons and width which are used for the numerical analysis in table II. We first determine the finite renormalization constants. There are ten constants. We first renormalize $\text{Re}.A_R$ and $\text{Re}.D_R$ so that each inverse propagator has zero at the on-shell mass $s = M_{K^*}^2$ and $s = M_\kappa^2$ respectively. And then we require the residues of the propagators $\text{Re}.\frac{1}{A_R}$ and $\text{Re}.\frac{1}{D_R}$ are unity on their pole masses. Because of the four conditions, we can constrain the parameters $a_0 + k_0, a_1 + k_1, d_0, d_1$ and d_2 . We show the four conditions below,

$$\begin{aligned}
a_0 + k_0 + (a_1 + k_1)M_{K^*}^2 &= -\text{Re}.\delta A_R(M_{K^*}^2) - \text{Re}.\Pi_{VS}^T(M_{K^*}^2), \\
a_1 + k_1 &= -\left.\frac{d\text{Re}.\delta A_R + \Pi_{VS}^T}{ds}\right|_{s=M_{K^*}^2}, \\
d_0 + d_1M_\kappa^2 + d_2M_\kappa^4 &= -\text{Re}.\delta D_R|_{s=M_\kappa^2}, \\
d_1 + 2M_\kappa^2d_2 &= -\left.\frac{d\text{Re}.\delta D_R}{ds}\right|_{s=M_\kappa^2}.
\end{aligned} \tag{52}$$

Moreover we set four parameters c_0, b_0, l_1, d_2 to be zeros. Below we show the numerical values of constants for $M_\kappa = 800$ MeV case.

$$\begin{aligned}
a_0 + k_0 &= -5.68 \times 10^5(\text{MeV}^2), \\
a_1 + k_1 &= 0.345, \\
d_1 &= -0.140, \\
d_0 &= -1.87 \times 10^3(\text{MeV}^2).
\end{aligned} \tag{53}$$

M_κ (MeV)	760	800	840
$a_1 + k_1$	0.33961	0.34471	0.35100
$a_0 + k_0$	-63519.7	-56822.9	-52534.2
d_1	-0.139154	-0.140467	-0.134301
d_0	6485.53	-1871.23	-16362.1
k_0	-4.16×10^5	-4.03×10^5	-3.92×10^5
k_1	0.6558	0.6561	0.6564
$\text{Br}(K\pi^0)$	0.00416079	0.00416079	0.00415823
$\text{Br}(K\eta)$	0.000162063	0.000162015	0.000162325
$\text{Br}(K\eta')$	4.14285×10^{-6}	3.87288×10^{-6}	3.94976×10^{-6}

TABLE III: The numerical values for the finite renormalization constants. They are chosen so that the the branching fractions of $\tau \rightarrow K\pi\nu$ and $\tau \rightarrow K\eta\nu$ can be reproduced.

We also show the values of the constants in Table.III for $M_\kappa = 760$ MeV and 840 MeV. Because the other parameters were set to be zero, there are only two undetermined parameters. We choose k_0 and k_1 as the parameters to be adjusted. They are fixed so that the branching fractions for $\tau \rightarrow K\pi\nu$ and $\tau \rightarrow K\eta\nu$ can be reproduced. For $(k_0, k_1) = (-4.03 \times 10^5, 0.656)$, we obtain

$$\begin{aligned}
Br(\tau^\mp \rightarrow K^\mp \pi \nu) &= 0.416 \times 10^{-2}, \\
Br(\tau^\mp \rightarrow K^\mp \eta \nu) &= 1.62 \times 10^{-4},
\end{aligned} \tag{54}$$

which are close to the experimental results [5] and [6],

$$\begin{aligned}
Br(\tau^- \rightarrow K^- \pi^0 \nu) &= 0.416 \pm 0.003 \pm 0.018 \times 10^{-2}, \\
Br(\tau^- \rightarrow K^- \eta \nu) &= (1.62 \pm 0.05 \pm 0.09) \times 10^{-4}.
\end{aligned} \tag{55}$$

The branching fraction for $\tau \rightarrow K\eta'\nu$ becomes,

$$\text{Br}(\tau \rightarrow K\eta'\nu) = 3.87 \times 10^{-6}. \tag{56}$$

We have plotted the hadronic invariant mass spectrum for $K\pi, K\eta$ and $K\eta'$ cases in Fig. 4.

The formulae can be found in [1],

$$\frac{d\text{Br}(\tau^\pm \rightarrow K^\pm P \bar{\nu}(\nu))}{d\sqrt{s}} = \frac{p_K G_F^2 |V_{us}|^2 (m_\tau^2 - s)^2}{\Gamma_\tau 2^5 \pi^3 m_\tau^3} \left(\left(\frac{2m_\tau^2}{3s} + \frac{4}{3} \right) p_K^2 |F^{KP}(s)|^2 + \frac{m_\tau^2}{2} |F_s^{KP}(s)|^2 \right), \quad (57)$$

where p_K is the momentum of kaon in the hadronic CM frame. In the hadronic invariant mass spectrum for $K^\pm \pi^0$ at low invariant mass region, K^* resonance can be seen. Just below K^* , we can see the effect of $\kappa(800)$. At the high invariant mass region, the new thresholds due to the vector and scalar channels are open and these effects can be seen in $K\pi$, $K\eta$ and $K\eta'$ cases in Fig. 4.




FIG. 4: The hadronic invariant mass spectrum $\frac{d\text{Br}}{d\sqrt{s}}$ for $K\pi^0$ (solid line), $K\eta$ (dashed line) and $K\eta'$ (short dashed line) cases. We choose $M_\kappa = 800$ MeV and the other parameters are fixed as in the corresponding columns of Table.III.

V. FORWARD AND BACKWARD ASYMMETRY AND CP VIOLATION

In this section, we compute the forward and backward asymmetry [8] and the direct CP violation for $\tau \rightarrow KP\nu$ decay. We start with the double differential rate of the unpolarized

τ decay [1],

$$\frac{d\text{Br}}{d\sqrt{s}d\cos\theta} = \frac{1}{\Gamma_\tau} \frac{G_F^2 |V_{us}|^2 (m_\tau^2 - s)^2}{2^5 \pi^3 m_\tau^3} \left\{ \left(\frac{m_\tau^2}{s} \cos^2 \theta + \sin^2 \theta \right) p_K^2 |F^{KP}(s)|^2 + \frac{m_\tau^2}{4} |F_s^{KP}|^2 - \frac{m_\tau^2}{\sqrt{s}} p_K \cos \theta \text{Re.}(F^{KP}(s)F_s^{KP}(s)^*) \right\}, \quad (58)$$

where θ is the scattering angle of kaon with respect to the incoming τ in the hadronic CM frame. The forward and backward asymmetry extracts the interference term of the vector form factor and the scalar form factor.

$$\begin{aligned} A_{\text{FB}}(s) &= \frac{\int_0^1 d\cos\theta \frac{d\text{Br}}{d\sqrt{s}d\cos\theta} - \int_{-1}^0 d\cos\theta \frac{d\text{Br}}{d\sqrt{s}d\cos\theta}}{\frac{d\text{Br}}{d\sqrt{s}}} \\ &= -\frac{\frac{p_K}{\sqrt{s}} \frac{|F_s^{KP}|}{|F^{KP}|} \cos \delta_{\text{st}}^{KP}}{\left(\frac{2m_\tau^2}{3s} + \frac{4}{3} \right) \frac{p_K^2}{m_\tau^2} + \frac{1}{2} \left| \frac{F_s^{KP}}{F^{KP}} \right|^2}, \end{aligned} \quad (59)$$

with $\delta_{\text{st}}^{KP} = \arg.\left(\frac{F_s^{KP}}{F^{KP}}\right)$. As we can see from Eq. (59), the forward and the backward asymmetry is determined by the ratio of the scalar and the vector form factors. It is also proportional to cosine of the strong phase shift δ_{st} . In Fig. 5, we show the vector and the scalar form factors and their ratio for $K\pi$ case. We also show the strong phase shifts in Fig. 6. The forward and backward asymmetries for $K\pi$, $K\eta$ and $K\eta'$ cases are shown in Fig. 7. As can be seen from Fig. 7, the forward and backward asymmetry for $K\pi$ case is large near the threshold region where the scalar contribution is also large. (See Fig. 5.) We can expect about 50% asymmetry for $\tau \rightarrow K\pi\nu$ decay.




FIG. 5: Left: The vector form factor $|F^{K\pi}|$ (thick solid line) and the scalar form factor $10 \times |F_s^{K\pi}|$ (thin solid line). Right: The ratio $\frac{|F_s^{K\pi}|}{|F^{K\pi}|}$.

By including new physics source of CP violation, we can predict the direct CP violation of the forward and backward asymmetry. To be definite, we start with non-minimal two




FIG. 6: The phase of the vector form factor: $\delta_V^{K\pi} = \arg.F^{K\pi}$ (short dashed line) and the phase of the scalar form factor $\delta_S^{K\pi} = \arg.F_s^{K\pi}$ (dashed line). The strong phase shift $\delta_{st}^{K\pi} = \delta_V^{K\pi} - \delta_S^{K\pi}$ is shown with thick solid line. $\delta_{st}^{K\eta}$ and $\delta_{st}^{K\eta'}$ are shown with long short dashed line and solid line respectively.




FIG. 7: The predictions of the forward and backward asymmetries of $\tau \rightarrow K\pi\nu$ (thick solid line) $\tau \rightarrow K\eta\nu$ (dashed line) and $\tau \rightarrow K\eta'\nu$ (short dashed line) in the standard model.

Higgs doublet model in which the two Higgs doublets have couplings to the charge leptons,

$$\begin{aligned}
-\mathcal{L} = & y_{1ij}\overline{e_{Ri}}\tilde{H}_1^\dagger L_{Lj} + y_{2ij}\overline{e_{Ri}}H_2^\dagger L_{Lj} + y_{1ij}^d\overline{d_{Ri}}\tilde{H}_1^\dagger Q_{Lj} \\
& + y_{2ij}^u\overline{u_{Ri}}\tilde{H}_2^\dagger Q_{Lj} + y_{2ij}^\nu\overline{\nu_{Ri}}\tilde{H}_2^\dagger L_{Lj} + \frac{M_{N_i}}{2}\overline{\nu_{Ri}}(\nu_{Ri})^c + \text{h.c.}, \quad (60)
\end{aligned}$$

where we assume that H_2 is coupled with neutrinos and both H_1 and H_2 are coupled with the charged leptons. The neutrino mass is given by the seesaw mechanism. However, the right-handed Majorana neutrino with the mass M_N much heavier than the electro-weak

scale does not affect on the interaction terms with the mass dimension equal to four at all. Therefore, we keep the terms which are not suppressed by a factor $\frac{1}{M_N}$ in discussion below. In two Higgs doublet model, without loss of generality, one can parametrize the Higgs fields as,

$$H_2 = e^{i\frac{\theta_{CP}}{2}} \begin{pmatrix} -\cos\beta H^+ \\ \frac{v_2+h_2-i\cos\beta A}{\sqrt{2}} \end{pmatrix}, \quad \tilde{H}_1 = i\tau_2 H_1^* = e^{-i\frac{\theta_{CP}}{2}} \begin{pmatrix} -\sin\beta H^+ \\ -\frac{v_1+h_1+i\sin\beta A}{\sqrt{2}} \end{pmatrix}, \quad (61)$$

where θ_{CP} is the relative phase of the vacuum expectation values of Higgs and its value can be determined from Higgs potential. The charged current interactions of the model are,

$$\begin{aligned} \mathcal{L} = & H^+ \bar{\nu}_{Li} l_{Rj} \left(\frac{Y_{2ji}^* e^{+i\frac{\theta}{2}}}{\cos\beta} - \delta_{ij} \frac{g \tan\beta m_j}{\sqrt{2} M_W} \right) \\ & - H^- \bar{d}_{iR} V_{ui}^* u_L \frac{g m_{di} \tan\beta}{\sqrt{2} M_W} - M_H^2 H^+ H^- \\ & - \frac{g}{\sqrt{2}} V_{ui}^* W_\mu^- \bar{d}_{Li} \gamma^\mu u_L - \frac{g}{\sqrt{2}} \bar{\nu}_{iL} \gamma_\mu l_{Li} W^{+\mu} + M_W^2 W^{+\mu} W_{-\mu} + \text{h.c.}, \end{aligned} \quad (62)$$

where the charged lepton masses $m_l = \text{diagonal}(m_e, m_\mu, m_\tau)$ are obtained by the diagonalization.

$$V_R \frac{1}{\sqrt{2}} (-y_1 v_1 e^{i\frac{\theta_{CP}}{2}} + y_2 v_2 e^{-i\frac{\theta_{CP}}{2}}) V_L^\dagger = m_l. \quad (63)$$

Eq. (63) can be used to express y_1 in terms of the charged lepton mass and the other Yukawa coupling y_2 . By introducing,

$$Y_2 = V_R y_2 V_L^\dagger, \quad (64)$$

one can obtain Eq. (62). The four fermi interactions induced by the charged Higgs exchange are,

$$\begin{aligned} \mathcal{L}_H = & \frac{G_F \tan^2\beta}{\sqrt{2} M_H^2} \left\{ \bar{\nu}_i \left(\delta_{ij} - r_{2ij} \frac{1}{\sin\beta} \right) m_{lj} (1 + \gamma_5) l_j \right\} \\ & \times \left\{ \bar{d} V_{KM}^\dagger (m_d (1 - \gamma_5) + m_u \cot^2\beta (1 + \gamma_5)) u \right\} + \text{h.c.}, \end{aligned} \quad (65)$$

where r_{2ij} denotes the non-minimal couplings of charged Higgs boson between the charged lepton l_j to the neutrino ν_i ,

$$r_{2ij} = \frac{Y_{2ji}^* e^{i\frac{\theta_{CP}}{2}}}{Y_j^{SM}}, \quad (66)$$

where Y^{SM} denotes the standard model Yukawa couplings for charged leptons, $Y_j^{SM} \frac{v}{\sqrt{2}} = m_{lj}$. The amplitude of the two Higgs doublet model is,

$$\text{Amp.}(\tau^- \rightarrow \nu_i K^- P) = -\frac{G_F}{\sqrt{2}} V_{us}^* \bar{u}_i \gamma^\mu (1 - \gamma_5) u_\tau \left\{ \delta_{i\tau} \left(q_\mu - \frac{\Delta_{KP}}{Q^2} Q_\mu \right) F^{KP} + \left(\delta_{i\tau} \left(1 - \frac{Q^2}{M_H^2} \tan^2 \beta \right) + \frac{Q^2}{M_H^2} \frac{\tan^2 \beta}{\sin \beta} r_{2i\tau} \right) F_s^{KP} Q_\mu \right\}, \quad (67)$$

where the matrix element of the scalar current is given by,

$$\langle K^- \pi^0 | \bar{s} u | 0 \rangle = \frac{Q^2}{m_s - m_u} F_s^{KP}. \quad (68)$$

Then $\tau \rightarrow KP\nu_i$ branching fraction is,

$$\frac{dBr(\tau^- \rightarrow K^- P \nu_i)}{d\sqrt{s} d \cos \theta} = \frac{1}{\Gamma} \frac{G_F^2}{2^5 \pi^3} \frac{(m_\tau^2 - s)^2}{m_\tau^3} \left\{ \left(\frac{m_\tau^2}{s} \cos^2 \theta + \sin^2 \theta \right) p_K^2 |F^{KP}|^2 \delta_{i\tau} + \frac{m_\tau^2}{4} |F_{si\tau}^{KP}|^2 - \delta_{i\tau} \frac{m_\tau^2}{\sqrt{s}} p_K \cos \theta \text{Re.}(F^{KP} F_{s\tau\tau}^{KP*}) \right\}, \quad (69)$$

where we define

$$F_{si\tau}^{KP} = \left\{ \delta_{i\tau} \left(1 - \frac{Q^2}{M_H^2} \tan^2 \beta \right) + \frac{Q^2}{M_H^2} \tan^2 \beta \frac{r_{2i\tau}}{\sin \beta} \right\} F_s^{KP} \quad (70)$$

for $i = e, \mu$ and τ . We neglect the small corrections proportional to up quark mass. For CP conjugate processes $\tau^+ \rightarrow K^+ \pi^0 \bar{\nu}_i$ is obtained by replacing r_2 in the amplitude $\tau^- \rightarrow K^- \pi^0 \nu_i$ with its complex conjugate r_2^* . Therefore the direct CP violation of the forward and backward asymmetries is given as,

$$A_{FB} - \bar{A}_{FB} = \frac{2 \frac{p_K}{\sqrt{s}} \frac{|F_s^{KP}|}{|F^{KP}|} \sin \delta_{st}}{\left(\frac{2m_\tau^2}{3s} + \frac{4}{3} \right) \frac{p_K^2}{m_\tau^2} + \frac{1}{2} \sum_i \frac{|F_{si\tau}^{KP}|^2}{|F_{\tau\tau}^{KP}|^2}} \left(\frac{Q^2}{M_H^2} \frac{\tan^2 \beta}{\sin \beta} \right) |r_{2\tau\tau}| \sin \theta_{2\tau\tau}, \quad (71)$$

where we parametrize CP violating phase of the flavor diagonal coupling as,

$$r_{2\tau\tau} = |r_{2\tau\tau}| e^{i\theta_{2\tau\tau}}. \quad (72)$$

We set the flavor off-diagonal couplings in $r_{2i\tau}$ to be zeros. We note that in the isospin limit $m_u = m_d$, the CP asymmetry of Eq. (71) for $K^\pm \pi^0$ case is identical to the direct CP violation of $\tau^- \rightarrow K_s \pi^- \nu_i$ and $\tau^+ \rightarrow K_s \pi^+ \bar{\nu}_i$. Contrary to the CP violation of the total branching ratios which is sensitive to the CP violation of the mixing of K^0 and \bar{K}^0 [16], the CP violation of the forward and backward asymmetries does not depend on the mixing induced CP violation.

The CP violation of the forward and backward asymmetry is shown for $K^\mp\pi^0$ case in Fig. 8. By taking $|r_{2\tau\tau}| = 1$, one can see that CP asymmetries can be as large as a few %. At low invariant mass region $\sqrt{s} < 900(\text{MeV})$, the direct CP violation is negative while at high invariant mass region $\sqrt{s} > 900 (\text{MeV})$, the CP violation is positive. The sign is correlated to $\sin\delta_{st}$ as can be seen in Eq. (71). From Fig. 6, we can see $\sin\delta_{st}$ also changes its sign around $\sqrt{s} = 900(\text{MeV})$. We also change the charged Higgs boson mass. For $M_H > 500 (\text{GeV})$, CP violation is suppressed to less than 1 %. In Fig. 9, we also show the CP asymmetries for $K\eta$ and $K\eta'$ cases. We note the sign of the CP violation is opposite to the sign at high invariant mass region of $K\pi$ case.




FIG. 8: CP violation for the forward and backward asymmetries of $\tau \rightarrow K\pi\nu$. The charged Higgs boson mass is changed as $M_H(\text{GeV}) = 200$ (thick solid line), 250 (dashed line), 500 (short dashed line) and 700 (solid line). The other parameters are $\tan\beta = 50$, $|r_{2\tau\tau}| = 1$ and $\theta_{2\tau\tau} = \frac{\pi}{2}$.

VI. CONCLUSION AND DISCUSSION

We have studied CP violation of $\tau \rightarrow KP\nu$ ($P = \pi^0, \eta, \eta'$) decays and $\tau \rightarrow K_s\pi\nu$. CP violation for the forward and backward asymmetries is computed using the two Higgs doublet




FIG. 9: CP violation for the forward and backward asymmetries of $\tau \rightarrow K\pi\nu$ (thick solid line), $\tau \rightarrow K\eta\nu$ (dashed line) and $\tau \rightarrow K\eta'\nu$ (short dashed line). We choose the parameters as $M_H = 300$, $\tan\beta = 50$ $|r_{2\tau\tau}| = 1$, $\theta_{2\tau\tau} = \frac{\pi}{2}$

model which has the non-minimal Yukawa couplings to the charged leptons. The effect of the CP violation is numerically studied. The direct CP violation depends on the strong phase shifts $\sin(\delta_V - \delta_s)$. To evaluate the phase shifts, we use the chiral lagrangian including the vector and the scalar resonances and take into account of the one loop corrections. We compute both real part and imaginary part of their self-energies. The divergences of the real part is subtracted properly. We have determined the finite renormalization constants so that the branching fractions for $K\pi$ and $K\eta$ modes are reproduced. We also take into account of $U(1)_A$ breaking so that η_0 and η_8 mixing can be incorporated. With those improved treatment, we can predict the hadronic invariant mass spectrum and CP violation even at high invariant mass region for $K\pi$ case as well as the same observables of $K\eta$ and $K\eta'$ cases.

The effect on the direct CP violation of the non-minimal coupling is studied. For the non-minimal Yukawa coupling as large as that of the standard model Yukawa coupling of τ lepton, we have predicted a few % CP asymmetries within the parameter region with $M_H = 200 \sim 300(\text{GeV})$ and with $\tan\beta \sim 50$.

Acknowledgments

We would like to thank K. Hayasaka, K. Inami, T. Onogi and T. Oshima for discussion and encouragement.

This work was supported by the Korea Research Foundation Grant funded by the Korean Government (MOEHRD, Basic Research Promotion Fund, KRF-2007-C00145) and the BK21 program of Ministry of Education (K.Y.L.).

The work of T. M. was supported by KAKENHI, Grant in Aid for Scientific Research on Priority Areas "New Development for Flavor Physics", No.19034008 and No.20039008, MEXT, Japan. We also thank YITP for support from international molecule type visitor program by the Yukawa International Programs for Quark-Hadron Sciences. The preliminary results of the paper were presented in BNM2006, YITP workshop; Towards the precise predictions of CP violation (YITP-W-07-21), BNM2008, New Development of flavor physics 2008, YongPyong 2008 and HQL2008.

APPENDIX A: SCALAR AND VECTOR MESON NONETS

Here we show the scalar and vector meson nonets.

$$S = \begin{pmatrix} \frac{a^0}{2} + \frac{\sigma}{2} & \frac{a^+}{\sqrt{2}} & \frac{\kappa^+}{\sqrt{2}} \\ \frac{a^-}{\sqrt{2}} & -\frac{a^0}{2} + \frac{\sigma}{2} & \frac{\kappa^0}{\sqrt{2}} \\ \frac{\kappa^-}{\sqrt{2}} & \frac{\bar{\kappa}^0}{\sqrt{2}} & f^0 \end{pmatrix}, \quad V = \begin{pmatrix} \frac{\rho}{2} + \frac{\omega}{2} & \frac{\rho^+}{\sqrt{2}} & \frac{K^{*+}}{\sqrt{2}} \\ \frac{\rho^-}{\sqrt{2}} & -\frac{\rho^0}{2} + \frac{\omega}{2} & \frac{K^{*0}}{\sqrt{2}} \\ \frac{K^{*-}}{\sqrt{2}} & \frac{K^{*0}}{\sqrt{2}} & \phi \end{pmatrix}. \quad (\text{A1})$$

APPENDIX B: FUNCTIONS $R_{PQ}^{(n)}$

The function $R_{PQ}^{(n)}$ is defined as,

$$R_{PQ}^{(n)} = \int_0^1 dx x^n \log(x^2 - (1 + \frac{\Delta_{PQ}}{s})x + \frac{M_P^2}{s} - i\epsilon). \quad (\text{B1})$$

There are two regions for s of interests depending on below the threshold, i.e., (1) $(M_P - M_Q)^2 \leq s \leq (M_P + M_Q)^2$ or the above threshold (2) $(M_P + M_Q)^2 \geq s$.

For the case (2),

$$R_{PQ}^{(n)} = \int_0^1 x^n \log((x - a)(x - b) - i\epsilon) dx, \quad (\text{B2})$$

with a and b are given as,

$$\begin{aligned} a &= \frac{1 + \frac{\Delta_{PQ}}{s}}{2} - \frac{\nu_{PQ}}{2s}, \\ b &= \frac{1 + \frac{\Delta_{PQ}}{s}}{2} + \frac{\nu_{PQ}}{2s}, \end{aligned} \quad (\text{B3})$$

with $0 \leq a, b \leq 1$. We show the real part of $R_{PQ}^{(n)}$,

$$\begin{aligned} \text{Re.}(R_{PQ}^{(0)}) &= -2 + \log(1-a)(1-b) - a \log \frac{1-a}{a} - b \log \frac{1-b}{b}, \\ \text{Re.}(R_{PQ}^{(1)}) &= \frac{a^2}{2} \log \frac{a}{1-a} + \frac{1}{2} \log(1-a) - \frac{1}{4} - \frac{a}{2} + \frac{b^2}{2} \log \frac{b}{1-b} + \frac{1}{2} \log(1-b) - \frac{1}{4} - \frac{b}{2}, \\ \text{Re.}(R_{PQ}^{(2)}) &= -\frac{6a^2 + 3a + 2}{18} + \frac{1}{3} \log(1-a) + \frac{a^3}{3} \log \frac{a}{1-a}, \\ &\quad - \frac{6b^2 + 3b + 2}{18} + \frac{1}{3} \log(1-b) + \frac{b^3}{3} \log \frac{b}{1-b}. \end{aligned} \quad (\text{B4})$$

For the case (1),

$$\begin{aligned} R_{PQ}^{(n)} &= \int_0^1 dx x^n \log((x-\beta)^2 + \alpha^2) \\ &= \int_{-\beta}^{1-\beta} dy (\beta+y)^n \log(y^2 + \alpha^2), \end{aligned} \quad (\text{B5})$$

where,

$$\begin{aligned} \alpha &= \frac{\sqrt{((M_P + M_Q)^2 - s)((s - (M_P - M_Q)^2)}}{2s} = \frac{\sqrt{-\nu_{PQ}^2}}{2s}, \\ \beta &= \frac{s + \Delta_{PQ}}{2s}. \end{aligned} \quad (\text{B6})$$

We define the indefinite integrals,

$$r_{PQ}^{(n)}(y) = \int^y dy y^n \log(y^2 + \alpha^2). \quad (\text{B7})$$

Using the integrals, one can write,

$$\begin{aligned} R_{PQ}^{(0)} &= r_{PQ}^{(0)}(1-\beta) - r_{PQ}^{(0)}(-\beta), \\ R_{PQ}^{(1)} &= r_{PQ}^{(1)}(1-\beta) - r_{PQ}^{(1)}(-\beta) + \beta R_{PQ}^{(0)}, \\ R_{PQ}^{(2)} &= r_{PQ}^{(2)}(1-\beta) - r_{PQ}^{(2)}(-\beta) + 2\beta R_{PQ}^{(1)} - \beta^2 R_{PQ}^{(0)}. \end{aligned} \quad (\text{B8})$$

The indefinite integrals are given by,

$$\begin{aligned} r_{PQ}^{(2)} &= \frac{1}{3} (y^3 \log(y^2 + \alpha^2) - \frac{2y^3}{3} + 2\alpha^2 y - 2\alpha^3 \arctan \frac{y}{\alpha}), \\ r_{PQ}^{(1)} &= \frac{1}{2} ((y^2 + \alpha^2) \log(y^2 + \alpha^2) - y^2), \\ r_{PQ}^{(0)} &= y \log(y^2 + \alpha^2) - 2y + 2\alpha \arctan \frac{y}{\alpha}. \end{aligned} \quad (\text{B9})$$

-
- [1] J. H. Kuhn. and E. Mirkes, Phys. Lett. **B398**, 407 (1997).
- [2] G. Bonvicini et al. (CLEO), Phys. Rev. Lett. **88**, 11803 (2002).
- [3] S. Y. Choi, J. Lee, and J. Song, Phys. Lett. **B437**, 191 (1998).
- [4] D. Epifanov et al.(Belle), Phys. Lett. **B654**, 65 (2007).
- [5] B. Aubert et al.(Babar), Phys. Rev. **D76**, 051104 (2007).
- [6] K. Abe et al.(Belle)(2007), hep-ex/0708.0733.
- [7] M. Kobayashi and T. Maskawa, Prog. Theor. Phys. **49**, 652 (1973).
- [8] L. Beldjoudi and T. N. Truong, Phys. Lett. **B351**, 357 (1995).
- [9] G. Ecker and J. Gasser and A. Pich and E. de Rafael, Nucl. Phys. **B321**, 311 (1989).
- [10] M. Bando, T. Kugo, S. Uehara, K. Yamawaki and T. Yanagida, Phys. Rev. Lett. **54**, 1215 (1985).
- [11] J. M. Gerard and E. Kou, Phys. Lett. **B616**, 85 (2005).
- [12] B. A. Li, Phys. Rev. **D55**, 1436 (1997).
- [13] M. Finkemeier and E. Mirkes, Z. Phys. **C72**, 619 (1996).
- [14] M. Jamin, A. Pich and J. Portoles, Phys. Lett. **B640**, 176 (2006).
- [15] B. Moussallam, Eur. Phys. J. **C53**, 401 (2008).
- [16] I. I. Bigi and A. I. Sanda, Phys. Lett. **B625**, 47 (2005).