

The Parts Determine the Whole except for n -Qubit Greenberger-Horne-Zeilinger States

Scott N. Walck* and David W. Lyons†
 Lebanon Valley College, Annville, PA 17003
 (Dated: August 5, 2008)

The generalized n -qubit Greenberger-Horne-Zeilinger (GHZ) states and their local unitary equivalents are the only pure states of n qubits that are not uniquely determined (among arbitrary states, pure or mixed) by their reduced density matrices of $n - 1$ qubits. Thus, the generalized GHZ states are the only ones containing information at the n -party level.

PACS numbers: 03.67.Mn, 03.65.Ta, 03.65.Ud

There are a number of perspectives from which to attempt an understanding of multiparty quantum entanglement. Entanglement can be viewed in terms of its ability to reject local realism and local hidden variable theories [1, 2]. From an operational point of view, entanglement can be classified based on local operations and classical communication (LOCC) and related notions [3, 4]. Quantum entanglement can be viewed as a resource for quantum information technologies [5, 6]. Finally, entanglement can be viewed in terms of the relationship between states of a quantum system and its subsystems [7, 8, 9]. Partial progress has been made in each of these perspectives, but none has emerged as a definitive way to think about entanglement. This paper deals with the last, “parts and whole” perspective on quantum entanglement, and makes a connection with the first, “rejection of local realism” view.

The “parts and whole” view of quantum entanglement asks to what extent a quantum state can be described by the (typically mixed) states of its subsystems. In particular, can an unknown state be uniquely determined, or identified, by giving the states of its subsystems (its reduced density matrices)? In [7, 8], Linden, Popescu, and Wootters proved the surprising result that almost all n -party pure states are determined by their reduced density matrices (RDMs). From the “parts and whole” perspective, the most entangled states are those that remain undetermined even after giving every possible reduced density matrix for every subsystem. In [10], the present authors proved that the only n -qubit states undetermined by their reduced density matrices *among other pure states* are the generalized n -qubit Greenberger-Horne-Zeilinger (GHZ) states,

$$\alpha |00 \cdots 0\rangle + \beta |11 \cdots 1\rangle, \quad \alpha\beta \neq 0$$

and their local unitary (LU) equivalents.

In this Letter, we prove the stronger result that the generalized n -qubit GHZ states and their LU equivalents are precisely the states that are undetermined by their reduced density matrices, among arbitrary states (pure or mixed). From the “parts and whole” perspective on quantum entanglement, the generalized GHZ states are

the most entangled quantum states. The generalized GHZ states are the only ones containing information at the n -party level.

Let D_n be the set of n -qubit density matrices. If $\rho \in D_n$ is an n -qubit density matrix, and $j \in \{1, \dots, n\}$ is a qubit label, we may form an $(n-1)$ -qubit reduced density matrix $\rho_{(j)} = \text{tr}_j \rho$ by taking the partial trace of ρ over qubit j . Let

$$\text{PTr} : D_n \rightarrow D_{n-1}^n$$

be the map $\rho \mapsto (\rho_{(1)}, \dots, \rho_{(n)})$ that associates to ρ its n -tuple of $(n-1)$ -qubit reduced density matrices. The map PTr is neither injective (one-to-one) nor surjective (onto). The failure of PTr to be injective means that multiple n -qubit states can have the same reduced density matrices. States $\rho_1 \neq \rho_2$ with $\text{PTr}(\rho_1) = \text{PTr}(\rho_2)$ require more information for their determination than is contained in their $(n-1)$ -qubit reduced density matrices.

Given a state ρ , the set $\text{PTr}^{-1}(\text{PTr}(\rho))$ contains all states with the same reduced density matrices as ρ . We define a state $\rho \in D_n$ to be *determined by its reduced density matrices* if $\text{PTr}^{-1}(\text{PTr}(\rho))$ contains only ρ , and *undetermined by its reduced density matrices* if $\text{PTr}^{-1}(\text{PTr}(\rho))$ contains more than one state.

Main result. An n -qubit pure state $|\psi\rangle$ is undetermined by its reduced density matrices (among arbitrary states, pure or mixed) if and only if $|\psi\rangle$ is LU equivalent to a generalized n -qubit GHZ state.

Proof. We will show below that if a pure n -qubit state $|\psi\rangle$ and a mixed (non-pure) n -qubit state ω have the same reduced density matrices, then there is a distinct pure n -qubit state $|\psi'\rangle \neq c|\psi\rangle$ that has the same reduced density matrices as $|\psi\rangle$. The authors showed in [10] that distinct n -qubit pure states can have the same reduced density matrices only if they are LU equivalent to generalized n -qubit GHZ states. \square

Let $|\psi\rangle$ be an n -qubit pure state and let ω be an n -qubit mixed (non-pure) state with the same RDMs as $|\psi\rangle$,

$$\rho_{(j)} := \text{tr}_j |\psi\rangle \langle \psi| = \text{tr}_j \omega,$$

where $j \in \{1, \dots, n\}$ labels a qubit. (In this paper, we use *mixed* to mean not pure. We use *arbitrary state* to mean pure or mixed.) For each j , we can Schmidt decompose $|\psi\rangle$ as

$$|\psi\rangle = \sum_{i=0}^1 \sqrt{q_i^j} |\chi_i^{(j)}\rangle |\alpha_i^j\rangle, \quad (1)$$

where $q_0^j \geq q_1^j \geq 0$ are eigenvalues of the $(n-1)$ -qubit density matrix $\rho_{(j)}$ and $|\chi_i^{(j)}\rangle$ are associated eigenvectors.

Let $|\Omega\rangle$ be a purification of ω , that is, a pure state of n qubits plus an environment E , for which

$$\omega = \text{tr}_E |\Omega\rangle \langle \Omega|.$$

Because

$$\text{tr}_j |\psi\rangle \langle \psi| = \text{tr}_{j,E} |\Omega\rangle \langle \Omega|,$$

we can write

$$|\Omega\rangle = \sum_{i=0}^1 \sqrt{q_i^j} |\chi_i^{(j)}\rangle |E_i^j\rangle, \quad (2)$$

where $|E_i^j\rangle$ is a vector in the subsystem of qubit j plus the environment, and

$$\langle E_{i'}^j | E_i^j \rangle = \delta_{i'i}. \quad (3)$$

Because

$$\text{tr}_{(j)} |\psi\rangle \langle \psi| = \text{tr}_{(j),E} |\Omega\rangle \langle \Omega|,$$

where $\text{tr}_{(j)}$ indicates a trace over all qubits *except* j , we can write

$$|\Omega\rangle = \sum_{r=0}^1 \sqrt{q_r^j} |\Omega_r^{(j)}\rangle |\alpha_r^j\rangle, \quad (4)$$

where $|\Omega_r^{(j)}\rangle$ is a vector in the subsystem of the environment plus all qubits except j , and

$$\langle \Omega_{r'}^{(j)} | \Omega_r^{(j)} \rangle = \delta_{r'r}. \quad (5)$$

Expand

$$|E_i^j\rangle = \sum_{r=0}^1 |\alpha_r^j\rangle |e_{ir}^j\rangle, \quad (6)$$

where $|e_{ir}^j\rangle$ is a vector in the environment only, which need not be normalized nor orthogonal to other environment vectors.

Plugging (6) into (2) gives

$$|\Omega\rangle = \sum_{i=0}^1 \sum_{r=0}^1 \sqrt{q_i^j} |\chi_i^{(j)}\rangle |\alpha_r^j\rangle |e_{ir}^j\rangle. \quad (7)$$

Comparing (7) with (4), we see that

$$\sqrt{q_r^j} |\Omega_r^{(j)}\rangle = \sum_{i=0}^1 \sqrt{q_i^j} |\chi_i^{(j)}\rangle |e_{ir}^j\rangle.$$

The orthonormality relations (3) and (5) give conditions on the environment vectors.

$$\sum_{r=0}^1 \langle e_{i'r}^j | e_{ir}^j \rangle = \delta_{i'i}$$

$$\sum_{i=0}^1 q_i^j \langle e_{i'r'}^j | e_{ir}^j \rangle = q_r^j \delta_{r'r}$$

Lemma 1. *If $|e_{ii^c}^j\rangle = 0$ for some qubit j and some bit $i \in \{0, 1\}$ (i^c is the bit complement of i), and $q_0^j q_1^j \neq 0$, then then $|e_{i^c i}^j\rangle = 0$.*

Proof. The first orthonormality condition above implies $\langle e_{ii}^j | e_{ii}^j \rangle = 1$, and the second then implies $|e_{i^c i}^j\rangle = 0$. \square

For a multi-index $I = (i_1 i_2 \dots i_n)$, in which each $i_j \in \{0, 1\}$, define

$$|I\rangle = |\alpha_{i_1}^1\rangle \dots |\alpha_{i_n}^n\rangle.$$

Let

$$|\psi\rangle = \sum_I c_I |I\rangle,$$

expand $|\chi_i^{(j)}\rangle$ in terms of the $|\alpha_i^j\rangle$, and substitute into (7) to give

$$|\Omega\rangle = \sum_{i=0}^1 \sum_I c_{i_1 \dots i_{j-1} i_{j+1} \dots i_n} |I\rangle |e_{ii}^j\rangle.$$

Equating coefficients of $|I\rangle$ with respect to different qubits gives

$$c_I |e_{i_j i_j}^j\rangle + c_{I_j} |e_{i_j^c i_j}^j\rangle = c_I |e_{i_k i_k}^k\rangle + c_{I_k} |e_{i_k^c i_k}^k\rangle, \quad (8)$$

where I_j is the multi-index equal to I in every slot except j , where it is complemented.

$$I_j := (i_1 \dots i_{j-1} i_j^c i_{j+1} \dots i_n)$$

We call equation (8) the *main constraint*. It holds for every multi-index I and every choice of qubits j and k .

Lemma 2. *If there are qubits j and k and a multi-index I such that $c_I = c_{I_j} = 0$ and $c_{I_k} \neq 0$, then $|e_{i_k^c i_k}^k\rangle = 0$.*

Proof. This follows directly from the main constraint. \square

Lemma 3. *If $c_I = c_{I_j} = 0$ for some multi-index I and some qubit j , then $\left|e_{i_k^c i_k}^k\right\rangle = 0$ for some k .*

Proof. If $c_{I_k} \neq 0$ for some k , then Lemma 2 applies, and $\left|e_{i_k^c i_k}^k\right\rangle = 0$. On the contrary, suppose that $c_{I_k} = 0$ for every k . In that case, every multi-index that differs from I in exactly one qubit slot has zero coefficient.

There is at least one multi-index with a nonzero coefficient. Choose a multi-index I' with a nonzero coefficient that differs from I in a minimal number of slots, say m slots. (We know that m is at least 2.) Then all multi-indexes that differ from I in $m - 1$ slots or less have zero coefficients. Choose k and l to be slots in which I' differs from I . Then we can apply Lemma 2 with I'_k playing the role of I , I'_{kl} playing the role of I_j , and I' playing the role of I_k . We conclude that $\left|e_{i_k^c i_k}^k\right\rangle = 0$. \square

Lemma 4. *If there exist I and I' that differ in slot j and agree in some slot k , such that $c_I c_{I'} \neq 0$, and $c_I c_{I'} - c_{I'_j} c_{I_j} \neq 0$, then the set*

$$\left\{\left|e_{00}^j\right\rangle, \left|e_{01}^j\right\rangle, \left|e_{10}^j\right\rangle, \left|e_{11}^j\right\rangle\right\}$$

spans at most two dimensions.

Proof. Take $c_{I'}$ times the main constraint for I minus c_I times the main constraint for I' . If I and I' differ in slot j , but agree in k , then

$$\begin{aligned} & c_{I'} c_{I'} \left(\left|e_{i_j i_j}^j\right\rangle - \left|e_{i_j^c i_j^c}^j\right\rangle \right) \\ &= -c_{I'} c_{I_j} \left|e_{i_j^c i_j}^j\right\rangle + c_I c_{I'_j} \left|e_{i_j i_j^c}^j\right\rangle + (c_I c_{I_k} - c_{I'_j} c_{I'_k}) \left|e_{i_k^c i_k}^k\right\rangle. \end{aligned}$$

If I and I' differ in slot j , but agree in k , then I and I'_j agree in slots j and k , so

$$(c_I c_{I'} - c_{I'_j} c_{I_j}) \left|e_{i_j i_j}^j\right\rangle = (c_I c_{I'_k} - c_{I'_j} c_{I_k}) \left|e_{i_k^c i_k}^k\right\rangle,$$

and I_j and I' agree in slots j and k , so

$$(c_{I_j} c_{I'_j} - c_I c_I) \left|e_{i_j i_j}^j\right\rangle = (c_{I_j} c_{I'_k} - c_I c_{I_k}) \left|e_{i_k^c i_k}^k\right\rangle.$$

Since $c_I c_{I'} \neq 0$, $\left|e_{00}^j\right\rangle - \left|e_{11}^j\right\rangle$ is a linear combination of $\left|e_{01}^j\right\rangle$, $\left|e_{10}^j\right\rangle$, and $\left|e_{i_k^c i_k}^k\right\rangle$. Since $c_I c_{I'} - c_{I'_j} c_{I_j} \neq 0$, we know that $\left|e_{01}^j\right\rangle$, $\left|e_{10}^j\right\rangle$, and $\left|e_{i_k^c i_k}^k\right\rangle$ span one dimension. \square

Proposition 1. *If an n -qubit pure state $|\psi\rangle$ and an n -qubit mixed state ω have the same reduced density matrices, then ω has rank 2.*

Proof. We show that ω has rank 2 by showing that at most 2 dimensions of the environment are used in $|\Omega\rangle$. It

is not possible that only one dimension of the environment is used, because that would make ω a pure state, contrary to our assumptions.

First we treat the case in which $q_0^j q_1^j = 0$ for some j , which is equivalent to $|\psi\rangle$ being a product of a single qubit state (in qubit j) and an $(n - 1)$ -qubit state. In this case, equation (1) becomes

$$|\psi\rangle = \left|\chi_0^{(j)}\right\rangle \left|\alpha_0^j\right\rangle,$$

and equation (2) becomes

$$|\Omega\rangle = \left|\chi_0^{(j)}\right\rangle \left|E_0^j\right\rangle.$$

This expression involves only two dimensions of the environment, so the proposition holds in this case.

From here on, we assume that $q_0^j q_1^j \neq 0$ for every $j \in \{1, \dots, n\}$. Next suppose that $\left|e_{i_i^c}^j\right\rangle = 0$ for some j and some i . Then, by Lemma 1, two of the four environment kets are zero, so the proposition holds.

Consider the case in which $c_I = 0$ for some I . If there is a qubit j for which $c_{I_j} = 0$ also, then Lemma 3 ensures that ω has rank 2. On the other hand, if $c_{I_j} \neq 0$ for every qubit j , then we can apply Lemma 4 with $j = 1$, $k = 2$, I_1 playing the role of I in the Lemma and I_2 playing the role of I' .

Finally, suppose that $c_I \neq 0$ for every multi-index I . If there are multi-indexes I and I' that disagree in some slot j and agree in some slot k , and satisfy $c_I c_{I'} - c_{I'_j} c_{I_j} \neq 0$, then Lemma 4 applies, and ω has rank 2. Otherwise, every pair of multi-indexes I and I' that disagree in some slot j and agree in some other slot satisfy $c_I c_{I'} = c_{I'_j} c_{I_j}$, or equivalently,

$$\frac{c_{i_1 \dots i_{j-1} i_j i_{j+1} \dots i_n}}{c_{i_1 \dots i_{j-1} i_j^c i_{j+1} \dots i_n}} = \frac{c_{i'_1 \dots i'_{j-1} i_j i'_{j+1} \dots i'_n}}{c_{i'_1 \dots i'_{j-1} i_j^c i'_{j+1} \dots i'_n}}.$$

For $n \geq 3$, this condition implies that $|\psi\rangle$ is the product of a 1-qubit state and an $(n - 1)$ -qubit state, contrary to our current assumptions. \square

Proposition 2. *If an n -qubit pure state $|\psi\rangle$ and an n -qubit mixed state ω have the same reduced density matrices, then there is a distinct pure state $|\psi'\rangle \neq c|\psi\rangle$ that has the same reduced density matrices as $|\psi\rangle$.*

Proof. Since ω has the same reduced density matrices as $|\psi\rangle$, Proposition 1 ensures ω has rank 2. In fact, the matrix

$$(1 - a) |\psi\rangle \langle \psi| + a \omega \tag{9}$$

has the same RDMS as $|\psi\rangle$ for all real a that give a legitimate density matrix, and consequently has rank at most two. We claim that by extending a beyond 1, we will reach a pure state with the same RDMS as $|\psi\rangle$.

Let $|\phi_1\rangle$ and $|\phi_2\rangle$ be eigenvectors of ω , so that

$$\omega = p |\phi_1\rangle \langle \phi_1| + (1-p) |\phi_2\rangle \langle \phi_2|.$$

Notice that $|\psi\rangle$ is a linear combination of $|\phi_1\rangle$ and $|\phi_2\rangle$,

$$|\psi\rangle = c_1 |\phi_1\rangle + c_2 |\phi_2\rangle.$$

If this were not the case, then expression (9), with $a = 1/2$, say, would have rank 3. Since $|\psi\rangle$ is in the span of $|\phi_1\rangle$ and $|\phi_2\rangle$, expression (9) has two real eigenvalues that sum to one for any real value of a .

The eigenvalues of a trace 1 Hermitian matrix

$$\begin{bmatrix} \frac{1}{2} + z & \bar{u} \\ u & \frac{1}{2} - z \end{bmatrix}$$

are

$$\lambda = \frac{1}{2} \pm \sqrt{|u|^2 + z^2}.$$

In particular, the lowest eigenvalue is less than or equal to the lowest diagonal element.

Now extend $|\psi\rangle$ to an orthonormal basis $\{|\psi\rangle, |\psi_2\rangle\}$ by defining

$$|\psi_2\rangle = \bar{c}_2 |\phi_1\rangle - \bar{c}_1 |\phi_2\rangle.$$

Then,

$$\begin{aligned} |\phi_1\rangle &= \bar{c}_1 |\psi\rangle + c_2 |\psi_2\rangle \\ |\phi_2\rangle &= \bar{c}_2 |\psi\rangle - c_1 |\psi_2\rangle. \end{aligned}$$

The coefficient of $|\psi\rangle \langle \psi|$ in the $\{|\psi\rangle, |\psi_2\rangle\}$ basis for expression (9) is

$$1 - a + a \left[p |c_1|^2 + (1-p) |c_2|^2 \right].$$

Since the term in square brackets is greater than zero and less than one, this coefficient will become negative for large enough a . Hence the lowest eigenvalue of expression (9), which is lower than either diagonal entry, will become negative for large enough a . Since u and z above are linear functions of a , the eigenvalues are continuous functions of a . Since the low eigenvalue of this matrix is positive for $a = 1$, and becomes negative for large a , it must pass through zero for some value of a , indicating a pure state. It is clear that this pure state is different from $|\psi\rangle$, since we can form ω as a mixture of $|\psi\rangle$ and this new pure state. \square

Conclusion. GHZ states first appeared as a way to achieve a simpler and non-statistical rejection of local realism and local hidden variable theories, as a way to provide a direct example of an “element of physical reality” for Einstein, Podolsky, and Rosen [11]. That the “parts and whole” view of quantum entanglement makes a connection with the “rejection of local realism” viewpoint is encouraging and provocative. Whether stronger connections can be made is a subject for future work.

The authors thank the National Science Foundation for their support of this work through NSF Award No. PHY-0555506.

* Electronic address: walck@lvc.edu

† Electronic address: lyons@lvc.edu

- [1] D. M. Greenberger, M. A. Horne, and A. Zeilinger, in *Bell's Theorem, Quantum Theory, and Conceptions of the Universe*, edited by M. Kafatos (Kluwer Academic, Dordrecht, The Netherlands, 1989), pp. 69–72, arXiv:0712.0921v1 [quant-ph].
- [2] J. S. Bell, *Phys.* **1**, 195 (1964), reprinted in J. S. Bell, *Speakable and Unsayable in Quantum Mechanics*, second edition (Cambridge University Press, Cambridge, England, 2004).
- [3] M. A. Nielsen, *Phys. Rev. Lett.* **83**, 436 (1999), arXiv:quant-ph/9811053.
- [4] W. Dür, G. Vidal, and J. I. Cirac, *Phys. Rev. A* **62**, 062314 (2000), arXiv:quant-ph/0005115.
- [5] C. H. Bennett, G. Brassard, C. Crépeau, R. Jozsa, A. Peres, and W. K. Wootters, *Phys. Rev. Lett.* **70**, 1895 (1993).
- [6] P. W. Shor, in *Proceedings, 35th Annual Symposium on Foundations of Computer Science* (IEEE Press, Los Alamitos, CA, 1994).
- [7] N. Linden and W. K. Wootters, *Phys. Rev. Lett.* **89**, 277906 (2002).
- [8] N. Linden, S. Popescu, and W. K. Wootters, *Phys. Rev. Lett.* **89**, 207901 (2002).
- [9] N. S. Jones and N. Linden, *Phys. Rev. A* **71**, 012324 (2005).
- [10] S. N. Walck and D. W. Lyons, *Phys. Rev. Lett.* **100**, 050501 (2008), arXiv:0707.4428 [quant-ph].
- [11] A. Einstein, B. Podolsky, and N. Rosen, *Phys. Rev.* **47**, 777 (1935).