

ENUMERATION OF $(k, 2)$ -NONCROSSING PARTITIONS

Toufik Mansour¹ and Simone Severini²

¹Department of Mathematics, University of Haifa, 31905 Haifa, Israel.

¹Center for Combinatorics, LPMC, Nankai University, 300071 Tianjin, P. R. China

²Department of Mathematics and Department of Computer Science, University of York, YO10 5DD York, United Kingdom

toufik@math.haifa.ac.il, ss54@york.ac.uk

ABSTRACT

A set partition is said to be (k, d) -noncrossing if it avoids the pattern $12 \cdots k12 \cdots d$. We find an explicit formula for the ordinary generating function of the number of (k, d) -noncrossing partitions of $\{1, 2, \dots, n\}$ when $d = 1, 2$.

Keywords: partitions, forbidden subsequences, kernel method.

2000 MATHEMATICS SUBJECT CLASSIFICATION: 05A05, 05A15

1. INTRODUCTION

A *partition* Π of the set $[n] = \{1, 2, \dots, n\}$ is a collection B_1, B_2, \dots, B_d of nonempty disjoint subsets of $[n]$. The elements of a partition are called *blocks*. We assume that B_1, B_2, \dots, B_d are listed in the increasing order of their minimum elements, that is $\min B_1 < \min B_2 < \dots < \min B_d$. The set of all partitions of $[n]$ with d blocks is denoted by $P(n, d)$. The cardinality of $P(n, d)$ is the well-known Stirling number of the second kind [8], which is usually denoted by $S(n, k)$. Any partition Π can be written in the *canonical sequential form* $\pi_1 \pi_2 \cdots \pi_n$, where $i \in B_{\pi_i}$ (see, e.g. [4]). From now on, we identify each partition with its canonical sequential form. For example, if $\Pi = \{1, 4\}, \{2, 5, 7\}, \{3\}, \{6\}$ is a partition of $[7]$, then its canonical sequential form is $\pi = 1231242$ and in such a case we write $\Pi = \pi$.

The *reduced form* of a word π on the alphabet $\{a_1, a_2, \dots, a_d\}$, where $a_1 < a_2 < \dots < a_d$, is a word π' obtained by renaming the letters of π . Specifically, the letter a_i is renamed i for all $i = 1, 2, \dots, d$. For example, the reduced form of the word 135351 is 123231. We say that a partition Π *avoids* $\tau = \tau_1 \cdots \tau_k$ if its canonical sequential form $\pi = \pi_1 \pi_2 \cdots \pi_n$ does not contain a subsequence $\pi' = \pi_{i_1} \cdots \pi_{i_k}$, such that the reduced form of π' equals the reduced form of τ . For example, the partition $\pi = 1231242$ avoids 12321.

In [4, 5], Klazar proved that the number of *noncrossing partitions* of $[n]$, that is partitions avoiding 1212, and the number of *nonnesting partitions* of $[n]$, that is partitions avoiding 1221, are given by the n -th Catalan number $\frac{1}{n+1} \binom{2n}{n}$. These results of Klazar have been extended in several recent directions. For instance, Chen *et al.* [2, 3] studied m -regular noncrossing, k -noncrossing and k -nonnesting partitions. Recall that a partition Π is called *m -regular* if $|x_1 - x_2| \geq m$, for any two distinct elements x_1, x_2 in the

same block of Π . A partition π is called k -noncrossing if it avoids $12 \cdots k12 \cdots k$ and k -nonnesting if it avoids $12 \cdots kk \cdots 21$. For further works on this subject, the reader is referred to Sagan [7] and the references therein.

In this paper, we generalize the concept of k -noncrossing partitions to (k, d) -noncrossing partitions. A partition is said to be (k, d) -noncrossing if it avoids $12 \cdots k12 \cdots d$. Let $\mathcal{N}_{k,d}(n)$ be the set of all (k, d) -noncrossing partitions of $[n]$. Note that $\mathcal{N}_{k,k}(n)$ is the set of k -noncrossing partitions of $[n]$ (see [3]).

For $d = 0$, it is easy to see from the definitions that the number of $(k, 0)$ -noncrossing partitions of $[n]$ is the same as the number of partitions of $[n]$ with at most $k - 1$ blocks. Thus,

$$\#\mathcal{N}_{k,0}(n) = \sum_{i=0}^{k-1} S(n, i),$$

where $S(n, i)$ is the Stirling number of the second kind. In this paper we give a complete enumeration for two cases of (n, d) -noncrossing partitions, in which d is either 1 or 2.

$k \setminus n$	0	1	2	3	4	5	6	7	8	9	10	11	12
2	1	1	2	5	14	42	132	429	1430	4862	16796	58786	208012
3	1	1	2	5	15	51	188	731	2950	12235	51822	223191	974427
4	1	1	2	5	15	52	202	856	3868	18313	89711	450825	2310453
5	1	1	2	5	15	52	203	876	4112	20679	109853	608996	3488806
6	1	1	2	5	15	52	203	877	4139	21111	115219	666388	4045991

TABLE 1. Number the $(k, 2)$ -noncrossing partitions of $[n]$ for $k = 2, 3, 4, 5, 6$ and $n = 0, 1, \dots, 12$.

Table 1 presents the number of $(k, 2)$ -noncrossing partitions in $\mathcal{P}_k(n) = \mathcal{N}_{k,2}(n)$, where $k = 2, 3, 4, 5, 6$. We will show that the ordinary generating function

$$\sum_{n \geq 0} \#\mathcal{P}_k(n)x^n$$

for the number of partitions in $\mathcal{P}_k(n)$ is rational in x and $\sqrt{(1 - kx)^2 - 4x^2}$. Namely, we prove the following result.

Theorem 1.1. *Let $k \geq 2$ and let*

$$y_k = \frac{1 - (k - 2)x - \sqrt{(1 - kx)^2 - 4x^2}}{2x(1 - (k - 2)x)}.$$

Then the ordinary generating function for the number of $(k, 2)$ -noncrossing partitions of $[n]$ is given by

$$\sum_{n \geq 0} \#\mathcal{P}_k(n)x^n = \frac{\frac{x^{k-1}y_k}{1-xy_k} + \sum_{j=0}^{k-2} \sum_{i=0}^j (-1)^{i+j} x^i \beta_{i,j}}{1 - \sum_{j=0}^{k-2} \sum_{i=0}^j (-1)^{i+j} ix \beta_{i,j}},$$

where $\beta_{j,j} = 1$ and

$$\beta_{i,j} = jx \prod_{s=i+1}^{j-1} (sx - 1)$$

for $i = 0, 1, \dots, j - 1$.

The proof of Theorem 1.1, see Section 2, is based on looking at $\mathcal{P}_k(n)$ as a disjoint union of subsets $\mathcal{P}_{k,\ell}(n)$, depending on some parameter ℓ . Then, we obtain linear recurrence relations with two indices, n and ℓ , for the number of partitions in these subsets. In order to solve the recurrence relations, we make use of the *kernel method technique* (see, e.g., [1]). The subsets $\mathcal{P}_{k,\ell}(n)$ can be defined as the collection of all ℓ -increasing partitions in $\mathcal{P}_k(n)$. Recall that $\pi = \pi_1 \cdots \pi_n$ is an ℓ -increasing partition of $[n]$ if $\pi_i = i$ for all $i = 1, 2, \dots, \ell$ and $\tau_{\ell+1} \neq \ell + 1$. Let us denote by $\mathcal{Q}_\ell(n)$ the set of all m -increasing partitions of $[n]$, with $m \leq \ell$. Directly from the proof of Theorem 1.1, we can obtain a formula for the number of partitions of $[n]$ in $\mathcal{Q}_\ell(n)$.

Corollary 1.2. *Let $\ell \geq 0$. The ordinary generating function for the number of partitions in $\mathcal{Q}_\ell(n)$ is*

$$I_\ell(x) = \sum_{n \geq 0} \#\mathcal{Q}_\ell(n)x^n = \frac{1 + \sum_{j=1}^{\ell} \sum_{i=0}^j (-1)^{i+j} x^i \beta_{i,j}}{1 - \sum_{j=1}^{\ell} \sum_{i=0}^j (-1)^{i+j} ix \beta_{i,j}},$$

where $\beta_{i,j}$ is defined in Theorem 1.1.

Theorem 1.1 gives two particular results, namely $k = 2$ and $k = 3$. For $k = 2$,

$$\sum_{n \geq 0} \#\mathcal{P}_2(n)x^n = 1 + \frac{xy_2}{1 - xy_2} = \frac{1}{1 - xy_2} = y_2,$$

where $y_2 = \frac{1 - \sqrt{1 - 4x}}{2x}$. Thus, the number of partitions in $\mathcal{P}_2(n)$ is given by the n -th Catalan number. For $k = 3$,

$$\sum_{n \geq 0} \#\mathcal{P}_3(n)x^n = \frac{1 + \frac{x^2 y_3}{1 - xy_3}}{1 - x} = \frac{3 - 3x - \sqrt{1 - 6x + 5x^2}}{2(1 - x)},$$

where $y_3 = \frac{1 - x - \sqrt{1 - 6x + 5x^2}}{2x(1 - x)}$. It follows that the number of partitions in $\mathcal{P}_3(n)$ is given by the n -th binomial transform of the Catalan number $\sum_{i=0}^n (-1)^i 3^{n-i} \binom{n}{i} \binom{i}{\lfloor i/2 \rfloor}$ (see [6, Sequence A007317]). For $k = 2$, there is a combinatorial proof that the number of partitions in $\mathcal{P}_2(n)$ is given by the n -th Catalan number. For $k = 3$, the formula above counts the number of Schröder paths with no peaks at even level of length n (see [6, Sequence A007317]). It would be interesting to find a bijective proof of this result.

Another bonus from the proof of Theorem 1.1 is the ordinary generating function for the number of $(k, 1)$ -noncrossing partition of $[n]$. Specifically, we prove the following result.

Theorem 1.3. *Let $k \geq 2$. Then the ordinary generating function for the number of $(k, 1)$ -noncrossing partitions of $[n]$ is given by*

$$\sum_{n \geq 0} \#\mathcal{N}_{k,1}(n)x^n = \frac{1 - x + (1 - x) \sum_{j=1}^{k-2} \sum_{i=0}^j (-1)^{i+j} x^i \beta_{i,j} + \sum_{i=0}^{k-1} (-1)^{i+k-1} x^i \beta_{i,k-1}}{1 - x - x(1 - x) \sum_{j=1}^{k-2} \sum_{i=0}^j (-1)^{i+j} i \beta_{i,j} - x \sum_{i=0}^{k-1} (-1)^{i+k-1} i \beta_{i,k-1}},$$

where $\beta_{i,j}$ is defined in Theorem 1.1.

For example, Theorem 1.3, for $k = 2, 3, 4, 5, 6$, gives the following ordinary generating functions for the number of $(k, 1)$ -noncrossing partitions of $[n]$:

$$\begin{aligned} \sum_{n \geq 0} \#\mathcal{N}_{2,1}(n)x^n &= \frac{1-x}{1-2x}, & \sum_{n \geq 0} \#\mathcal{N}_{3,1}(n)x^n &= \frac{1-3x+x^2}{(1-x)(1-3x)}, \\ \sum_{n \geq 0} \#\mathcal{N}_{4,1}(n)x^n &= \frac{1-6x+9x^2-3x^3}{(1-x)(1-2x)(1-4x)}, & \sum_{n \geq 0} \#\mathcal{N}_{5,1}(n)x^n &= \frac{1-10x+32x^2-37x^3+11x^4}{(1-x)(1-2x)(1-3x)(1-5x)}, \\ \sum_{n \geq 0} \#\mathcal{N}_{6,1}(n)x^n &= \frac{1-15x+81x^2-192x^3+189x^4-53x^5}{(1-x)(1-2x)(1-3x)(1-4x)(1-6x)}. \end{aligned}$$

The numbers of $(k, 1)$ -noncrossing partitions of $[n]$ with $k = 2, 3, 4, 5, 6$ are given by

$$\begin{aligned} \#\mathcal{N}_{2,1}(n) &= 2^{n-1}, \\ \#\mathcal{N}_{3,1}(n) &= \frac{1}{6}(3^n + 3), \\ \#\mathcal{N}_{4,1}(n) &= \frac{1}{24}(4^n + 6 \cdot 2^n + 8), \\ \#\mathcal{N}_{5,1}(n) &= \frac{1}{120}(5^n + 10 \cdot 3^n + 20 \cdot 2^n + 45), \\ \#\mathcal{N}_{6,1}(n) &= \frac{1}{720}(6^n + 15 \cdot 4^n + 40 \cdot 3^n + 135 \cdot 2^n + 264). \end{aligned}$$

2. PROOFS

Let us denote by $F_k(x)$ the generating function for the number of partitions in $\mathcal{P}_k(n)$:

$$F_k(x) = \sum_{n \geq 0} \#\mathcal{P}_k(n)x^n.$$

Here, instead of dealing with recurrence relations with two indices n and ℓ , as we mentioned in the introduction, we deal with recurrence relations in terms of ordinary generating functions with a single index ℓ . Let us denote by $F_{k,\ell}(x)$ the generating function for the number of partitions in $\mathcal{P}_{k,\ell}(n)$:

$$F_{k,\ell}(x) = \sum_{n \geq 0} \#\mathcal{P}_{k,\ell}(n)x^n$$

Here, for the case $\ell = 0$ we have $F_{k,\ell}(x) = 1$. Clearly, $F_k(x) = \sum_{i \geq 0} F_{k,i}(x)$. Our main result is based on the construction of linear recurrence relations with a single index for the ordinary generating function $F_{k,\ell}(x)$. As we will see later, since the recurrences contain the expression $\sum_{i \geq \ell} F_{k,i}(x)$, we define for clarity

$$G_{k,\ell}(x) = \sum_{i \geq \ell} F_{k,i}(x).$$

The expression $G_{k,\ell}(x)$ is the ordinary generating function for the number of j -increasing partitions of $[n]$, with $j \geq \ell$. It follows directly from the definitions that the generating function $G_{k,\ell}(x)$ is well-defined, since

$$G_{k,\ell}(x) = F_k(x) - \sum_{i=0}^{\ell-1} F_{k,i}(x).$$

In our first lemma, we find the recurrence relation for $F_{k,\ell}(x)$, where $1 \leq \ell \leq k-1$.

Lemma 2.1. *For all $1 \leq \ell \leq k-1$,*

$$F_{k,\ell}(x) = \ell x G_{k,\ell}(x) + x^\ell.$$

Proof. Let $\pi = \pi_1\pi_2\cdots\pi_n \in \mathcal{P}_{k,\ell}(n)$. If $n > \ell$ then $\pi_{\ell+1} \leq \ell$. Thus $\pi \in \mathcal{P}_k(n)$ if and only if the reduced form of $\pi' = \pi_1 \cdots \pi_\ell \pi_{\ell+2} \cdots \pi_n$ is a partition in $\cup_{j \geq \ell} \mathcal{P}_{k,j}(n-1)$. Therefore,

$$F_{k,\ell}(x) = \ell x(F_{k,\ell}(x) + F_{k,\ell+1}(x) + \cdots) + x^\ell = \ell x G_{k,\ell}(x) + x^\ell,$$

where x^ℓ counts the unique ℓ -increasing partition of $[\ell]$, namely $12 \cdots \ell$, as required by the statement. \square

The above observation together with the the definition of $G_{k,\ell}(x)$ gives the following system:

$$(2.1) \quad \begin{cases} F_{k,0}(x) & = 1 \\ xF_{k,0}(x) + F_{k,1}(x) & = xF_k(x) + x \\ 2xF_{k,0}(x) + 2xF_{k,1}(x) + F_{k,2}(x) & = 2xF_k(x) + x^2 \\ \vdots & \\ (k-1)xF_{k,0}(x) + \cdots + (k-1)xF_{k,k-2}(x) + F_{k,k-1}(x) & = (k-1)xF_k(x) + x^{k-1}. \end{cases}$$

Next, we find an explicit formula for $F_{k,\ell}(x)$ in terms of $F_k(x)$.

Lemma 2.2. *For all $1 \leq \ell \leq k-1$,*

$$F_{k,\ell}(x) = \sum_{i=0}^{\ell} (-1)^{i+\ell} (ixF_k(x) + x^i) \beta_{i,\ell},$$

where $\beta_{\ell,\ell} = 1$ and $\beta_{i,\ell} = \ell x \prod_{j=i+1}^{\ell-1} (jx-1)$ for $i = 0, 1, \dots, \ell-1$.

Proof. With the use of Cramer's Rule on (2.1), we obtain

$$F_{k,\ell}(x) = \sum_{i=0}^{\ell} (-1)^{i+\ell} (ixF_k(x) + x^i) \begin{vmatrix} (i+1)x & 1 & 0 & \cdots & 0 & 0 \\ (i+2)x & (i+2)x & 1 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ (\ell-1)x & (\ell-1)x & (\ell-1)x & \cdots & (\ell-1)x & 1 \\ \ell x & \ell x & \ell x & \cdots & \ell x & \ell x \end{vmatrix}.$$

By making use of the formula

$$\begin{vmatrix} ax & 1 & 0 & \cdots & 0 & 0 \\ (a+1)x & (a+1)x & 1 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ (b-1)x & (b-1)x & (b-1)x & \cdots & (b-1)x & 1 \\ bx & bx & bx & \cdots & bx & bx \end{vmatrix} = bx \prod_{j=a}^{b-1} (jx-1),$$

which holds by induction on $b \geq a$, we obtain

$$F_{k,\ell}(x) = \sum_{i=0}^{\ell} (-1)^{i+\ell} (ixF_k(x) + x^i) \beta_{i,\ell},$$

as claimed. \square

Now, before completing the proof of our main result, Theorem 1.1, let us present two applications of Lemma 2.2. The first one is the ordinary generating function for the number of partitions in $\mathcal{Q}_\ell(n)$; the second one is the ordinary generating function for the number of $(k, 1)$ -noncrossing partitions of $[n]$.

2.1. Enumerating partitions in $\mathcal{Q}_\ell(n)$. The formula of the ordinary generating function $I_\ell(x)$ for the number of partitions in $\mathcal{Q}_\ell(n)$ can be obtained as follows. From the definition of the set $\mathcal{Q}_\ell(n)$ and from the proof of Lemma 2.2 for $\ell < k$, we obtain that the ordinary generating function for the number of m -increasing partitions in $\mathcal{Q}_\ell(n)$ is given by

$$I_{\ell,m}(x) = \sum_{i=0}^m (-1)^{i+m} (ixI_\ell(x) + x^i) \beta_{i,m}.$$

On the other hand, $I_\ell(x) = \sum_{m=0}^{\ell} I_{\ell,m}(x)$. Combining these two equations, we obtain

$$I_\ell(x) = \sum_{j=0}^{\ell} I_{\ell,j}(x) = 1 + \sum_{j=1}^{\ell} \sum_{i=0}^j (-1)^{i+j} (ixI_\ell(x) + x^i) \beta_{i,j}.$$

The solution of this equation gives a formula for $I_\ell(x)$, as stated in Corollary 1.2.

2.2. Enumerating $(k, 1)$ -noncrossing partitions of $[n]$. Let $J_k(x)$ be the ordinary generating function for the number of $(k, 1)$ -noncrossing partitions of $[n]$, that is,

$$J_k(x) = \sum_{n \geq 0} \#\mathcal{N}_{k,1}(n) x^n.$$

More generally, let $J_{k,\ell}(x)$ be the ordinary generating function for the number of $(k, 1)$ -noncrossing ℓ -increasing partitions of $[n]$. Then, a similar argument as in the proof of Lemma 2.2 gives that

$$(2.2) \quad J_{k,\ell}(x) = \sum_{i=0}^{\ell} (-1)^{i+\ell} (ixJ_k(x) + x^i) \beta_{i,\ell}, \quad \ell = 1, 2, \dots, k-1,$$

with $J_{k,0}(x) = 1$. On the other hand,

$$(2.3) \quad J_{k,\ell}(x) = x^{\ell+1-k} J_{k,k-1}(x), \quad \ell = k, k+1, k+2, \dots$$

To prove this observation, let π be any $(k, 1)$ -noncrossing ℓ -increasing partition of $[n]$ with $\ell \geq k$. Then $\pi_1 \pi_2 \cdots \pi_\ell = 12 \cdots \ell$ and $\pi_{\ell+1} < \ell + 1$. Since π avoids $12 \cdots k1$, then $\pi_i \notin \{1, 2, \dots, \ell + 1 - k\}$ for all $i \geq \ell + 1$. Thus, the number of $(k, 1)$ -noncrossing ℓ -increasing partition of $[n]$ with $\ell \geq k$ is the same as the number of $(k, 1)$ -noncrossing $(k-1)$ -increasing partition of $[n - \ell - 1 + k]$. This is equivalent to $J_{k,\ell}(x) = x^{\ell+1-k} J_{k,k-1}(x)$, for all $\ell \geq k$. Therefore, using the fact that $J_k(x) = \sum_{\ell \geq 0} J_{k,\ell}(x)$, and the two equations (2.2) and (2.3), we can write

$$J_k(x) = 1 + \sum_{\ell=0}^{k-1} \sum_{i=0}^{\ell} (-1)^{i+\ell} (ixJ_k(x) + x^i) \beta_{i,\ell} + \sum_{\ell \geq k} x^{\ell+1-k} J_{k,k-1}(x).$$

Again, by (2.2), we have

$$J_k(x) = 1 + \sum_{\ell=0}^{k-1} \sum_{i=0}^{\ell} (-1)^{i+\ell} (ixJ_k(x) + x^i) \beta_{i,\ell} + \frac{x}{1-x} \sum_{i=0}^{k-1} (-1)^{i+k-1} (ixJ_k(x) + x^i) \beta_{i,k-1}.$$

The solution of this equation gives a formula for $J_k(x) = \sum_{n \geq 0} \#\mathcal{N}_{k,1}(n)x^n$ as stated in Theorem 1.3.

2.3. Proof of Theorem 1.1. We need some extra notation before completing the proof of Theorem 1.1. Let

$$H_k(x, y) = \sum_{\ell=0}^{k-2} F_{k,\ell}(x)y^\ell$$

and

$$F_k(x, y) = \sum_{\ell \geq 0} F_{k,\ell}(x)y^\ell.$$

From Lemma 2.2, we can observe that

$$H_k(x, y) = \sum_{\ell=0}^{k-2} y^\ell \left(\sum_{i=0}^{\ell} (-1)^{i+\ell} (ixF_k(x, 1) + x^i)\beta_{i,\ell} \right)$$

and that

$$F_k(x, 1) = F_k(x),$$

where $\beta_{\ell,\ell} = 1$ and $\beta_{i,\ell} = \ell x \prod_{j=i+1}^{\ell-1} (jx - 1)$, for $i = 0, 1, \dots, \ell - 1$. Now, let us focus on the generating functions $F_{k,\ell}(x)$, where $\ell \geq k - 1$.

Lemma 2.3. *For all $j \geq 0$,*

$$F_{k,k-1+j}(x) = x^{k-1+j} + \sum_{i=0}^{j-1} x^{j+1-i} G_{k,k-1+i}(x) + (k-1)xG_{k,k-1+j}(x).$$

Proof. The case $j = 0$ holds on the basis of Lemma 2.1. Let us assume that $j \geq 1$. Let $\pi = \pi_1\pi_2 \cdots \pi_n$ be any partition in $\mathcal{P}_{k,k-1+j}(n)$ such that $\pi_{k+j} = i \leq k - 1 + j$. We want the equation of the generating function for the number of partitions in $\mathcal{P}_{k,k-1+j}(n)$, namely $F_{k,k-1+j}(x)$. Let us consider the following two cases:

- If $1 \leq i \leq j$ then π is such that $\pi_p \notin \{i+1, i+2, \dots, j+1\}$, where $p \geq k+1+j$. Thus, the contribution of this case is

$$x^{j+1-i}(F_{k,k-1+i}(x) + F_{k,k+i}(x) + \cdots) = x^{j+1-i}G_{k,k-1+i}(x).$$

- If $j+1 \leq i \leq k-1+j$ then π satisfies the above conditions if and only if the reduced form of $\pi_1 \cdots \pi_{k-1+j}\pi_{k+1+j} \cdots \pi_n$ is a partition in $\cup_{i \geq 0} \mathcal{P}_{k,k-1+j+i}(n-1)$. Thus, the contribution of this case is

$$x(F_{k,k-1+j}(x) + F_{k,k+j}(x) + \cdots) = xG_{k,k-1+j}(x).$$

Putting together the above cases, $i = 1, 2, \dots, k-1+j$, we obtain that

$$F_{k,k-1+j}(x) = x^{k-1+j} + \sum_{i=0}^{j-1} x^{j+1-i} G_{k,k-1+i}(x) + (k-1)xG_{k,k-1+j}(x),$$

where x^{k-1+j} counts the unique $(k-1+j)$ -increasing partitions of $[k-1+j]$, namely $12 \cdots (k-1+j)$. \square

Now we have a formula for the generating function $F_k(x, y)$:

Proposition 2.4. *We have*

$$\begin{aligned} \left(1 + \frac{x^2 y^2}{(1-y)(1-xy)} + \frac{(k-1)xy}{1-y}\right) (F_k(x, y) - H_k(x, y)) \\ = \frac{(xy)^{k-1}}{1-xy} + \frac{y^{k-1}}{1-y} \left(\frac{x^2 y}{1-xy} + (k-1)x\right) (F_k(x, 1) - H_k(x, 1)). \end{aligned}$$

Proof. Lemma 2.1 together with Lemma 2.3 give

$$\begin{aligned} F_k(x, y) \\ = \frac{1}{1-xy} + \sum_{j=1}^{k-2} jxG_{k,j}(x)y^j + (k-1)x \sum_{j \geq k-1} G_{k,j}(x)y^j + \frac{x^2 y}{1-xy} \sum_{j \geq k-1} G_{k,j}(x)y^j \\ = \frac{1}{1-xy} + \sum_{j=1}^{k-2} (F_{k,j}(x) - x^j)y^j \\ + x \left(\frac{xy}{1-xy} + k-1\right) \left(\frac{y^{k-1}}{1-y}(F_k(x, 1) - H_k(x, 1)) - \frac{y}{1-y}(F_k(x, y) - H_k(x, y))\right), \end{aligned}$$

which is equivalent to

$$\begin{aligned} \left(1 + \frac{x^2 y^2}{(1-y)(1-xy)} + \frac{(k-1)xy}{1-y}\right) (F_k(x, y) - H_k(x, y)) \\ = \frac{(xy)^{k-1}}{1-xy} + \frac{y^{k-1}}{1-y} \left(\frac{x^2 y}{1-xy} + (k-1)x\right) (F_k(x, 1) - H_k(x, 1)), \end{aligned}$$

as claimed. \square

The functional equation in the statement of Proposition 2.4 can be solved systematically using the *kernel method technique* (see [1]). Let

$$y = y_k = \frac{1 - (k-2)x - \sqrt{(1-kx)^2 - 4x^2}}{2x(1 - (k-2)x)}$$

be one of the roots of the equation $1 + \frac{x^2 y^2}{(1-y)(1-xy)} + \frac{(k-1)xy}{1-y}$. Then Proposition 2.4 gives

$$F_k(x, 1) - H_k(x, 1) = \frac{x^{k-1} y_k}{1 - xy_k}.$$

Therefore, Lemma 2.2 gives

$$F_k(x, 1) - \sum_{j=0}^{k-2} \sum_{i=0}^j (-1)^{i+j} (ixF_k(x, 1) + x^i) \beta_{i,j} = \frac{x^{k-1} y_k}{1 - xy_k},$$

which implies our main result, Theorem 1.1.

Acknowledgment. Part of this work has been carried out while the second author was visiting the Center for Combinatorics of Nankai University. The financial support of the Center is gratefully acknowledged. Also, the authors would like to thank the anonymous referees for a number of valuable comments, which helped to improve the presentation of this paper.

REFERENCES

- [1] C. BANDERIER, M. BOUSQUET-MÉLOU, A. DENISE, P. FLAJOLET, D. GARDY, AND D. GOUYOU-BEAUCHAMPS, Generating functions for generating trees, Formal Power Series and Algebraic Combinatorics (Barcelona, 1999), *Discr. Math.* **246**:1-3 (2002) 29–55.
- [2] WILLIAM Y. C. CHEN, EVA Y.P. DENG, AND ROSENA R.X. DU, Reduction of m -regular noncrossing Partitions, *Europ. J. Combin.* **26**:2 (2005) 237–243.
- [3] WILLIAM Y. C. CHEN, EVA Y.P. DENG, ROSENA R.X. DU, R. P. STANLEY, AND CATHERINE H. YAN, Crossings and Nestings of Matchings and Partitions, *Trans. Amer. Math. Soc.*, to appear. arXiv: math.CO/0501230.
- [4] M. KLAZAR, On abab-free and abba-free set partitions, *Europ. J. Combin.* **17** (1996) 53–68.
- [5] M. KLAZAR, On trees and noncrossing partitions, *Discr. Appl. Math.* **82** (1998) 263–269.
- [6] N.J.A. SLOANE, The Online Encyclopedia of Integer Sequences, www.research.att.com/~njas/sequences/.
- [7] B. E. SAGAN, Pattern avoidance in set partitions, preprint.
- [8] R. P. STANLEY, Enumerative Combinatorics, Vol. 1, Cambridge University Press, Cambridge, UK, 1996.
- [9] M. WACHS AND D. WHITE, p, q -Stirling numbers and set partition statistics, *J. Combin. Theory, Series A*, **56**:1 (1991) 27–46.