# BALANCED HERMITIAN METRICS FROM SU(2)-STRUCTURES

M. FERNANDEZ, A. TOMASSINI, L. UGARTE AND R. VILLACAMPA ´

ABSTRACT. We study the intrinsic geometrical structure of hypersurfaces in 6-manifolds carrying a balanced Hermitian SU(3)-structure, which we call balanced  $SU(2)$ -structures. We provide conditions which imply that such a 5manifold can be isometrically embedded as a hypersurface in a manifold with a balanced SU(3)-structure. We show that any 5-dimensional compact nilmanifold has an invariant balanced SU(2)-structure as well as new examples of balanced Hermitian SU(3)-metrics constructed from balanced SU(2)-structures. Moreover, for  $n = 3, 4$ , we present examples of compact manifolds, endowed with a balanced  $SU(n)$ -structure, such that the corresponding Bismut connection has holonomy equal to  $SU(n)$ .

# 1. INTRODUCTION

Let  $(J, q)$  be a Hermitian structure on a manifold M, with Kähler form F and Lie form  $\theta$ . The 3-form  $JdF$  is the torsion of the Bismut connection of  $(J, g)$ , that is, the unique Hermitian connection with totally skew-symmetric torsion [\[2\]](#page-18-0). If  $JdF$  is closed and non-zero (so  $(M, J, g)$ ) is not a Kähler manifold) the Hermitian structure is called *strong Kähler with torsion*.

Hermitian structures for which the Lie form  $\theta$  vanishes identically or, equivalently,  $F^2 = F \wedge F$  is closed, are called *balanced*. If in addition, there is an SU(3)structure with Kähler form F and a complex volume (3,0)-form  $\Psi = \Psi_+ + i\Psi_-$  on M such that  $dF^2 = d\Psi_+ = d\Psi_- = 0$ , we say that  $(F, \Psi_+, \Psi_-)$  is a balanced SU(3)structure. Such structures are a generalization of integrable SU(3)-structures, which are defined by a triplet  $(F, \Psi_+, \Psi_-)$  satifying  $dF = d\Psi_+ = d\Psi_- = 0$ . Balanced SU(3)-structures have been very useful in physics for the construction of explicit compact supersymmetric valid solutions to the heterotic equations of motion in dimension six [\[7\]](#page-18-1).

Recently, Conti and Salamon [\[5\]](#page-18-2) introduced the notion of hypo structures on 5-manifolds as those structures corresponding to the restriction of an integrable  $SU(3)$ -structure on a 6-manifold M to a hypersurface N of M. They are a generalization in dimension 5 of Sasakian-Einstein metrics. In fact, Sasakian-Einstein metrics correspond to Killing spinors and hypo structures are induced by generalized Killing spinors. In terms of differential forms, a *hypo structure* on a 5-manifold N is determined by a quadruplet  $(\eta, \omega_i, 1 \leq i \leq 3)$  of differential forms, where  $\eta$  is a nowhere vanishing 1-form and  $\omega_i$  are 2-forms on N satisfying certain relations (see [\(5\)](#page-2-0) in Section [2\)](#page-1-0). If the forms  $\eta$  and  $\omega_i$  satisfy

(1) 
$$
d\eta = -2\omega_3, \quad d\omega_1 = 3\eta \wedge \omega_2, \quad d\omega_2 = -3\eta \wedge \omega_1,
$$

Date: November 14, 2018.

<sup>2000</sup> Mathematics Subject Classification. 53C55, 53C25, 32C10.

Key words and phrases. SU(2)-structures, evolution equations, balanced metrics, Bismut connection, holonomy.

then N is a Sasakian-Einstein manifold, that is, a Riemannian manifold such that  $N \times \mathbb{R}$  with the cone metric is Kähler and Ricci flat [\[3\]](#page-18-3), so  $N \times \mathbb{R}$  has holonomy contained in  $SU(3)$  or, equivalently, it has an integrable  $SU(3)$ -structure.

In the general case of a hypo structure, in [\[5\]](#page-18-2) it is proved that a real analytic hypo structure on N can be lifted to an integrable SU(3)-structure on  $N \times I$ , for some open interval I, if  $(\eta, \omega_i, 1 \leq i \leq 3)$  belongs to a one-parameter family of hypo structures  $(\eta(t), \omega_i(t), 1 \leq i \leq 3)$  satisfying the evolution equations [\(10\)](#page-6-0) given in Section [3.](#page-5-0) Moreover, any oriented hypersurface of a 6-manifold with an integrable SU(3)-structure is naturally endowed with a hypo structure (see Section [2](#page-1-0) for details).

Our purpose in this paper is to study the geometrical structure of an oriented hypersurface in a 6-manifold equipped with a balanced SU(3)-structure, which we call balanced  $SU(2)$ -structure. These are  $SU(2)$ -structures defined by a quadruplet  $(\eta, \omega_i, 1 \leq i \leq 3)$  of differential forms satisfying the relations given by [\(4\)](#page-2-1), and they can be considered as a generalization to dimension five of the holomorphic symplectic structures in dimension four. We prove that any compact nilmanifold of dimension 5 has a balanced  $SU(2)$ -structure although three of the 5-nilmanifolds do not admit an invariant hypo structure [\[5\]](#page-18-2). Furthermore, we provide conditions under which a balanced  $SU(2)$ -structure on N can be lifted to a balanced  $SU(3)$ structure on  $N \times \mathbb{R}$ . This allows us to exhibit examples of (non-invariant) balanced Hermitian metrics and, in particular of complex structures, on nilmanifolds not admitting invariant balanced Hermitian metrics.

To this end, in Section [2](#page-1-0) we show that there exists a balanced  $SU(2)$ -structure on any oriented hypersurface  $f: N \longrightarrow M$  of a 6-manifold M with a balanced  $SU(3)$ -structure. Furthermore, we prove that any balanced  $SU(2)$ -structure on N can be lifted to a balanced Hermitian SU(3)-structures on  $N \times \mathbb{R}$  if and only if it satisfies the *evolution balanced*  $SU(2)$  *equations* [\(9\)](#page-6-1) established in Theorem [3.1](#page-5-1) of Section [3.](#page-5-0)

On the other hand, in Proposition [2.6](#page-4-0) we characterize the circle bundles  $S^1$  $N \to X$  with a balanced SU(2)-structure induced by a holomorphic symplectic structure on X. In particular, we get that the product  $X \times \mathbb{R}$  has a balanced SU(2)-structure. Then, in Section [3](#page-5-0) we solve the evolution balanced  $SU(2)$  equations [\(9\)](#page-6-1) for the compact 5-nilmanifolds not having a hypo structure as well as for the balanced SU(2)-structures on manifolds which are the total space of a circle bundle over Kodaira-Thurston manifold with a certain holomorphic symplectic structure. In this way, we get new examples of balanced Hermitian SU(3)-metrics.

Finally, in Section [4](#page-8-0) we describe in detail several examples of compact manifolds obtained as quotient of solvable Lie groups, and endowed with a balanced  $SU(n)$ structure, for  $n = 3, 4$ . In particular, we show that the holonomy of the Bismut connection is equal to  $SU(n)$ . Moreover, many of such examples are holomorphic parallelizable.

### 2. Balanced SU(2)-structures

<span id="page-1-0"></span>In this section we introduce a special type of SU(2)-structures on 5-manifolds, namely *balanced* SU(2)-structures, which permit construct new Hermitian balanced metrics by solving suitable evolution equations. First we recall some facts about SU(2)-structure on a 5-dimensional manifold. (For more details, we refer e.g. to [\[5\]](#page-18-2).) Let N be a 5-dimensional manifold and let  $L(N)$  be the principal bundle of linear frames on N. An  $SU(2)$ -structure on N is an  $SU(2)$ -reduction of  $L(N)$ . We have the following (see [\[5,](#page-18-2) Prop.1])

**Proposition 2.1.** SU(2)-structures on a 5-manifold N are in 1 : 1 correspondence with quadruplets  $(\eta, \omega_1, \omega_2, \omega_3)$ , where  $\eta$  is a 1-form and  $\omega_i$  are 2-forms on N satisfying

$$
\omega_i \wedge \omega_j = \delta_{ij} v, \quad v \wedge \eta \neq 0,
$$

for some 4-form v, and

$$
i_X \omega_3 = i_Y \omega_1 \Rightarrow \omega_2(X, Y) \ge 0,
$$

where  $i_X$  denotes the contraction by X.

Equivalently, an SU(2)-structure on N can be viewed as the datum of  $(\eta, \omega_3, \Phi)$ , where  $\eta$  is a 1-form,  $\omega_3$  is a 2-form and  $\Phi = \omega_1 + i\omega_2$  is a complex 2-form such that

$$
\eta \wedge \omega_3^2 \neq 0, \qquad \Phi^2 = 0
$$
  

$$
\omega_3 \wedge \Phi = 0, \qquad \Phi \wedge \overline{\Phi} = 2\omega_3^2
$$

and  $\Phi$  is of type  $(2,0)$  with respect to  $\omega_3$ .

As a corollary of the last Proposition, we obtain the useful local characterization of  $SU(2)$ - structures (see [\[5\]](#page-18-2)):

Corollary 2.2. If  $(\eta, \omega_1, \omega_2, \omega_3)$  is an SU(2)-structure on a 5-dimensional manifold N, then locally, there exists a basis of 1-forms  $\{e^1, \ldots, e^5\}$  such that

$$
\eta = e^1
$$
,  $\omega_1 = e^{24} + e^{53}$ ,  $\omega_2 = e^{25} + e^{34}$ ,  $\omega_3 = e^{23} + e^{45}$ 

As a consequence, SU(2)-structures naturally arise on hypersurfaces of 6 manifolds with an SU(3)-structure. Indeed, let  $f : N \longrightarrow M$  be an oriented hypersurface of a 6-manifold M endowed with an SU(3)-structure  $(F, \Psi_+, \Psi_-)$  and denote by  $\mathbb U$  the unit normal vector field. Then N inherits an SU(2)-structure  $(\eta, \omega_1, \omega_2, \omega_3)$  given by

<span id="page-2-2"></span>(2) 
$$
\eta = -i_{\mathbb{U}}F, \quad \omega_1 = i_{\mathbb{U}}\Psi_-, \quad \omega_2 = -i_{\mathbb{U}}\Psi_+, \quad \omega_3 = f^*F.
$$

Conversely, an SU(2)-structure  $(\eta, \omega_1, \omega_2, \omega_3)$  on N induces an SU(3)-structure  $(F, \Psi_+, \Psi_-)$  on  $N \times \mathbb{R}$  given by

(3) 
$$
F = \omega_3 + \eta \wedge dt, \qquad \Psi = \Psi_+ + i\Psi_- = (\omega_1 + i\omega_2) \wedge (\eta + idt),
$$

where  $t$  is a coordinate on  $\mathbb{R}$ .

**Definition 2.3.** An SU(2)-structure  $(\eta, \omega_1, \omega_2, \omega_3)$  on a 5-dimensional manifold N is called balanced if it satisfies the following equations

(4) 
$$
d(\omega_1 \wedge \eta) = 0, \qquad d(\omega_2 \wedge \eta) = 0, \qquad d(\omega_3 \wedge \omega_3) = 0.
$$

<span id="page-2-1"></span><span id="page-2-0"></span>In [\[5\]](#page-18-2), an  $SU(2)$ -structure is said to be *hypo* if

(5) 
$$
d(\omega_1 \wedge \eta) = d(\omega_2 \wedge \eta) = d\omega_3 = 0.
$$

Hence, it is obvious that any hypo structure is balanced. However, there are nilmanifolds admitting no invariant hypo structure, but having balanced SU(2) structures. In fact, if the Lie algebra underlying a 5-dimensional compact nilmanifold N is isomorphic to  $(0, 0, 0, 12, 14)$ ,  $(0, 0, 12, 13, 23)$  or  $(0, 0, 12, 13, 14+23)$ , then

.

there is no invariant hypo structure on  $N$  (see [\[5\]](#page-18-2)). It is easy to check that the  $SU(2)$ -structure given by

 $\eta = e^1$ ,  $\omega_1 = e^{24} + e^{53}$ ,  $\omega_2 = e^{25} + e^{34}$ ,  $\omega_3 = e^{23} + e^{45}$ ,

satisfies [\(4\)](#page-2-1) on each one of these three Lie algebras. Therefore, we get the following

# Proposition 2.4. Any 5-dimensional compact nilmanifold has an invariant balanced  $SU(2)$ -structure.

There exist also 5-dimensional solvable non-nilpotent Lie algebras with no invariant hypo structure, but having a balanced SU(2)-structure. For example, let us consider the Lie algebra  $\mathfrak g$  whose dual is spanned by  $(e^1, \ldots, e^5)$  such that

$$
de1 = 0
$$
,  $de2 = 0$ ,  $de3 = e13$ ,  $de4 = -e14$ ,  $de5 = e34$ .

Then g is a 5-dimensional solvable non-nilpotent Lie algebra. The simply-connected Lie group G associated with g has a uniform discrete subgroup  $\Gamma$ , so that  $N = \Gamma \backslash G$ is a compact solvmanifold. In fact, the manifold  $N$  is the topological product of the unit circle by the compact solvable 4-dimensional manifold, studied in [\[1\]](#page-18-4), which is a circle bundle over the compact solvmanifold  $Sol(3)$ . A straightforward calculation shows that the following forms

$$
\eta = e^1
$$
,  $\omega_1 = e^{24} + e^{53}$ ,  $\omega_2 = e^{25} + e^{34}$ ,  $\omega_3 = e^{23} + e^{45}$ .

satisfy

$$
d(\omega_1 \wedge \eta) = d(\omega_3 \wedge \eta) = d(\omega_2^2) = 0,
$$

and thus they define a balanced  $SU(2)$ -structure on N. However, N has not invariant hypo SU(2)-structures. First, using Hattori's theorem [\[11\]](#page-18-5), we have that the real cohomology groups of N of degree  $\leq 2$  are

$$
H^{0}(N) = \langle 1 \rangle, \quad H^{1}(N) = \langle [e^{1}], [e^{2}] \rangle, \quad H^{2}(N) = \langle [e^{12}] \rangle.
$$

Now, let us suppose that N has an invariant hypo  $SU(2)$ -structure  $(\eta, \omega_1, \omega_2, \omega_3)$ . Then

$$
\omega_3 = ae^{12} + be^{13} + ce^{14} + fe^{34},
$$

for some real numbers a, b, c and f. Therefore,  $\omega_3^2 = 2af e^{1234}$ , and so  $\eta = e^5 +$  $\sum_{i=1}^{4} \lambda_i e^i$  since  $\omega_3^2 \wedge \eta$  is a volume form. On the other hand,

$$
\omega_1 = \sum_{i,j=1}^4 a_{ij} e^{ij},
$$

and

$$
\omega_2 = \sum_{i,j=1}^4 b_{ij} e^{ij},
$$

for some real numbers  $a_{ij}$  and  $b_{ij}$ . Now, the conditions  $d(\omega_1 \wedge \eta) = d(\omega_2 \wedge \eta) = 0$ imply that

$$
\omega_1 = a_{13}e^{13} + a_{14}e^{14} + a_{34}e^{34}
$$

,

and

$$
\omega_2 = b_{13}e^{13} + b_{14}e^{14} + b_{34}e^{34},
$$

which implies that  $\omega_1^2 = \omega_2^2 = 0$ . This is not possible for an SU(2)-structure on N.

Balanced SU(2)-structures on 5-manifolds are related to Hermitian balanced structures in six dimensions as the next proposition shows. First, recall that a balanced SU(3)-structure on a 6-manifold M is an SU(3)-structure  $(F, \Psi = \Psi_+ +$  $i\Psi$ <sub>-</sub>) (where F is the Kähler form of an almost Hermitian structure and  $\Psi$  =  $\Psi_+ + i\Psi_-$  is a complex volume form) such that  $F^2$  and  $\Psi$  are closed. The latter condition implies that the underlying almost complex structure is integrable. Notice that any balanced  $SU(3)$ -structure is in particular half-flat [\[4,](#page-18-6) [12,](#page-19-0) [6\]](#page-18-7), that is, it satisfies  $dF^2 = d\Psi_+ = 0$ .

**Proposition 2.5.** Let  $f: N \longrightarrow M$  be an immersion of an oriented 5-manifold into a 6-manifold with an  $SU(3)$ -structure. If the  $SU(3)$ -structure is balanced then the SU(2)-structure on N given by  $(2)$  is balanced.

*Proof.* From [\(2\)](#page-2-2) it follows that  $\omega_1 \wedge \eta = f^* \Psi_+$ ,  $\omega_2 \wedge \eta = f^* \Psi_-$  and  $\omega_3 \wedge \omega_3 = f^* F^2$ . Now, if  $F^2$  and  $\Psi$  are closed then the induced structure is half-balanced.  $\square$ 

Let  $(X, J)$  be a complex surface. By definition, a *holomorphic symplectic struc*ture on X is the datum of a d-closed and non-degenerate  $(2,0)$ -form  $\omega$  on X. Let g be a J-Hermitian metric on X and  $\omega_3$  be the fundamental form of  $(q, J)$ . Then, up to a conformal change, we may assume that

$$
\omega_3^2 = \omega_1^2 = \omega_2^2.
$$

Then we have the following

<span id="page-4-0"></span>**Proposition 2.6.** Let  $(X, J)$  be a complex surface equipped with a holomorphic symplectic structure  $\omega = \omega_1 + i\omega_2$ , and let  $\omega_3$  be the Kähler form of a J-Hermitian metric. Then, for any integral closed 2-form  $\Omega$  on X annihilating  $\cos\theta \omega_1 + \sin\theta \omega_2$ and  $\sin \theta \omega_1 - \cos \theta \omega_2$  for some  $\theta$ , there is a principal circle bundle  $\pi \colon N \longrightarrow X$ with connection form  $\rho$  such that  $\Omega$  is the curvature of  $\rho$  and such that the SU(2)structure  $(\eta, \omega_1^{\theta}, \omega_2^{\theta}, \omega_3^{\theta})$  on N given by

$$
\eta = \rho,
$$
  
\n
$$
\omega_1^{\theta} = \pi^*(\cos \theta \omega_1 + \sin \theta \omega_2),
$$
  
\n
$$
\omega_2^{\theta} = \pi^*(-\sin \theta \omega_1 + \cos \theta \omega_2),
$$
  
\n
$$
\omega_3^{\theta} = \pi^*(\omega_3)
$$

is a balanced SU(2)-structure.

*Proof.* As previously remarked, we may assume that  $\omega_1^2 = \omega_2^2 = \omega_3^2$ . Since  $d\omega_1 =$  $d\omega_2 = 0$  and  $d\eta = \pi^*(\Omega)$ , a simple calculation shows that

$$
d(\omega_1^{\theta} \wedge \eta) = \omega_1^{\theta} \wedge d\eta = \pi^*((\cos \theta \omega_1 + \sin \theta \omega_2) \wedge \Omega) = 0,
$$

and

$$
d(\omega_2^{\theta} \wedge \eta) = \omega_2^{\theta} \wedge d\eta = \pi^* ((-\sin \theta \omega_1 + \cos \theta \omega_2) \wedge \Omega) = 0.
$$

The existence of a principal circle bundle in the conditions above follows from a well known result by Kobayashi [\[13\]](#page-19-1).

**Remark 2.7.** Notice that  $\Omega = 0$  satisfies the hypothesis in the previous proposition for each  $\theta$  and one gets the trivial circle bundle  $N = X \times \mathbb{R}$  with the balanced  $SU(2)$ -structure which is the natural extension to N of the holomorphic symplectic structure on X.

<span id="page-5-2"></span>Remark 2.8. Following [\[9,](#page-18-8) Def.1.1], a symplectic couple on an oriented 4-manifold X is a pair of symplectic forms  $(\omega_1, \omega_2)$  such that  $\omega_1 \wedge \omega_2 = 0$  and  $\omega_1^2, \omega_2^2$  are volume forms defining the positive orientation. A symplectic couple is called conformal if  $\omega_1^2 = \omega_2^2$ . By [\[9,](#page-18-8) Thm.1.3] it follows that X admits a conformal symplectic couple if and only if X is diffeomorphic to  $(a)$  a complex torus,  $(b)$  a K3 surface or (c) a primary Kodaira surface. In the hypothesis of prop. [2.6,](#page-4-0)  $(\omega_1, \omega_2)$  defines a conformal symplectic couple on the 4-manifold X and  $\omega_3$  is a non-degenerate 2-form on X such that  $\omega_i \wedge \omega_3 = 0$ ,  $i = 1, 2$  and  $\omega_1^2 = \omega_2^2 = \omega_3^2$ 

Next we illustrate the construction given in Proposition [2.6](#page-4-0) by showing principal circle bundles over holomorphic symplectic manifolds in the cases (a) and (c) of Remark [2.8.](#page-5-2) Let us consider the closed 4-manifold  $X = \Gamma \backslash G$ , where the Lie algebra  $\mathfrak g$  of G has the following structure equations

<span id="page-5-5"></span>(6) 
$$
de^1 = 0
$$
,  $de^2 = 0$ ,  $de^3 = 0$ ,  $de^4 = -\epsilon e^{23}$   $(\epsilon = 0, 1)$ .

Clearly X is the Kodaira-Thurston manifold for  $\epsilon = 1$  and a 4-torus for  $\epsilon = 0$ . Consider the complex structure  $J$  on  $X$  defined by the complex  $(1,0)$ -forms

$$
\varphi^1 = e^1 + ie^4
$$
,  $\varphi^2 = e^2 + ie^3$ ,

so that

$$
\omega = \varphi^1 \wedge \varphi^2 = (e^{12} + e^{34}) + i (e^{13} - e^{24}) = \omega_1 + i \omega_2
$$

is a holomorphic symplectic structure on X. The metric g on X given by  $g =$  $\sum_{i=1}^{4} e^{i} \otimes e^{i}$  is J-Hermitian and the fundamental form of  $(g, J)$  is precisely  $\omega_3 =$  $e^{14} + e^{23}$ , so we are in the conditions of Proposition [2.6.](#page-4-0) Observe that  $\omega_3$  is closed only when X is a 4-torus, i.e.  $\epsilon = 0$ .

Now let  $\Omega$  be a closed 2-form on X such that

(7) 
$$
\Omega \wedge (\cos \theta \omega_1 + \sin \theta \omega_2) = 0 = \Omega \wedge (\sin \theta \omega_1 - \cos \theta \omega_2),
$$

for some  $\theta$ . A direct calculations shows that

<span id="page-5-3"></span>
$$
\Omega = a(e^{12} - e^{34}) + b(e^{13} + e^{24}) + (\epsilon - 1)c_1 e^{14} + c_2 e^{23}
$$

satisfies [\(7\)](#page-5-3) for all  $\theta$ , where  $a, b, c_1, c_2$  are constant. Applying Proposition [2.6,](#page-4-0) we have a balanced SU(2)-structure on the total space N, which is non-hypo if  $\epsilon = 1$ . In this case it is easy to see that  $N$  is a compact 5-nilmanifold with underlying Lie algebra isomorphic to either  $(0, 0, 0, 0, 12)$  or  $(0, 0, 0, 12, 13 + 24)$ .

# 3. Evolution equations and Hermitian balanced metrics

<span id="page-5-0"></span>Next we establish evolution equations that allow the construction of new balanced Hermitian metrics in dimension six from balanced SU(2)-structures in dimension five.

<span id="page-5-1"></span>**Theorem 3.1.** Let  $(\eta(t), \omega_1(t), \omega_2(t), \omega_3(t))$  be a family of SU(2)-structures on a 5-manifold N, for  $t \in I = (a, b)$ . Then, the SU(3)-structure on  $M = N \times I$  given by

<span id="page-5-4"></span>(8) 
$$
F = \omega_3(t) + \eta(t) \wedge dt, \qquad \Psi = (\omega_1(t) + i\omega_2(t)) \wedge (\eta(t) + idt),
$$

is balanced Hermitian if and only if  $(\eta(t), \omega_1(t), \omega_2(t), \omega_3(t))$  is a balanced SU(2)structure for any  $t$  in the open interval  $I$ , and the following evolution equations

(9) 
$$
\begin{cases} \partial_t(\omega_1 \wedge \eta) = -d\omega_2, \\ \partial_t(\omega_2 \wedge \eta) = d\omega_1, \\ \partial_t(\omega_3 \wedge \omega_3) = -2 d(\omega_3 \wedge \eta), \end{cases}
$$

are satisfied.

*Proof.* A direct calculation shows that the  $SU(3)$ -structure given by [\(8\)](#page-5-4) satisfies

<span id="page-6-1"></span>
$$
dF^{2} = d(\omega_{3} \wedge \omega_{3}) + (\partial_{t}(\omega_{3} \wedge \omega_{3}) + 2 d(\omega_{3} \wedge \eta)) \wedge dt,
$$

and

$$
d\Psi = d(\omega_1 \wedge \eta) - (\partial_t(\omega_1 \wedge \eta) + d\omega_2) \wedge dt + i d(\omega_2 \wedge \eta) - i (\partial_t(\omega_2 \wedge \eta) - d\omega_1) \wedge dt.
$$

The forms  $F^2$  and  $\Psi$  are both closed if and only if  $(\eta(t), \omega_1(t), \omega_2(t), \omega_3(t))$  is a balanced SU(2)-structure for any  $t \in I$ , and satisfies the equations [\(9\)](#page-6-1).

Remark 3.2. Notice that in the special case when the balanced SU(2)-structures in the family are hypo, the last equation in [\(9\)](#page-6-1) reduces to

$$
\omega_3 \wedge (\partial_t \omega_3 + d\eta) = 0.
$$

Hence, any solution of the hypo evolution equations introduced in [\[5\]](#page-18-2), namely

<span id="page-6-0"></span>(10) 
$$
\partial_t(\omega_1 \wedge \eta) = -d\omega_2, \quad \partial_t(\omega_2 \wedge \eta) = d\omega_1, \quad \partial_t \omega_3 = -d\eta,
$$

is trivially a solution of  $(9)$ , and the resulting  $SU(3)$ -structure is integrable because  $F$  in [\(8\)](#page-5-4) is closed.

Next we give some explicit solutions to the balanced evolution equations [\(9\)](#page-6-1). Our first example is the 5-manifold  $N = X \times \mathbb{R}$ , where X is the Kodaira-Thurston manifold. Let us consider the non-hypo balanced  $SU(2)$ -structure given by  $(6)$  for  $\epsilon = 1$  and with zero curvature, namely

$$
\eta = e^5
$$
,  $\omega_1 = e^{12} + e^{34}$ ,  $\omega_2 = e^{13} - e^{24}$ ,  $\omega_3 = e^{14} + e^{23}$ 

The family of non-hypo balanced SU(2)-structures on N given by  $\eta(t) = e^5$  and

 $\omega_1(t) = e^{12} + e^3(e^4 - t e^5), \ \omega_2(t) = e^{13} + (e^4 - t e^5)e^2, \ \omega_3(t) = e^1(e^4 - t e^5) + e^{23},$ coincides with the previous one for  $t = 0$  and satisfies the evolution equations [\(9\)](#page-6-1) for any  $t \in \mathbb{R}$ .

Denote by G the simply-connected nilpotent Lie group with Lie algebra  $(0, 0, 0, 0, 12)$ , so that  $N = \Gamma \backslash G$  for some lattice  $\Gamma$  of maximal rank in G. It follows from Proposition [3.1](#page-5-1) that the SU(3)-structure on  $G \times \mathbb{R}$  given by

<span id="page-6-2"></span>(11) 
$$
F = e^{14} + e^{23} - t e^{15} + e^5 \wedge dt,
$$

$$
\Psi_+ = e^{125} + e^{345} - (e^{13} - e^{24} + t e^{25}) \wedge dt,
$$

$$
\Psi_- = e^{135} - e^{245} + (e^{12} + e^{34} - t e^{35}) \wedge dt,
$$

is balanced Hermitian.

Next we find explicit solutions of [\(9\)](#page-6-1) for the non-hypo nilpotent Lie algebras  $(0, 0, 0, 12, 14)$  and  $(0, 0, 12, 13, 23)$ .

.

Lie algebra  $(0,0,0,12,14)$ : It is straightforward to check that the family of balanced SU(2)-structures given by

$$
\eta(t) = \sqrt[3]{\frac{2-3t}{2}}e^1,
$$
  
\n
$$
\omega_1(t) = \frac{1}{2} \left( \sqrt[3]{\frac{2}{2-3t}} - \frac{2-3t}{2} \right) e^{23} + \sqrt[3]{\frac{2-3t}{2}} e^{24} - \sqrt[3]{\frac{2}{2-3t}} e^{35},
$$
  
\n
$$
\omega_2(t) = \sqrt[3]{\frac{2}{2-3t}} e^{25} + \sqrt[3]{\frac{2-3t}{2}} e^{34},
$$
  
\n
$$
\omega_3(t) = e^{23} - \frac{1}{2} \left( 1 - \frac{2-3t}{2} \sqrt[3]{\frac{2-3t}{2}} \right) e^{24} + e^{45},
$$

satisfies the evolution equations [\(9\)](#page-6-1) for  $t \in \mathbb{R} - \{2/3\}$ . Observe that the volume form of the associated Riemannian metric along the family is given by

$$
\omega_1(t) \wedge \omega_1(t) \wedge \eta(t) = 2\sqrt[3]{\frac{2-3t}{2}}e^{12345},
$$

so the orientation for  $t \in (-\infty, 2/3)$  is opposite to the orientation for  $t \in (2/3, \infty)$ .

Let  $I = (-\infty, 2/3)$  and denote by G the simply-connected nilpotent Lie group with Lie algebra  $(0, 0, 0, 12, 14)$ . The basis of 1-forms on the product manifold  $G \times I$ given by

$$
\alpha^1 = e^2, \qquad \alpha^2 = e^3, \quad \alpha^3 = \sqrt[3]{\frac{2-3t}{2}} e^4, \quad \alpha^4 = \frac{1}{2} \sqrt[3]{\frac{2}{2-3t}} (e^2 + 2e^5) - \frac{2-3t}{4} e^2,
$$
  

$$
\alpha^5 = \sqrt[3]{\frac{2-3t}{2}} e^1, \quad \alpha^6 = dt,
$$

is orthonormal with respect to the Riemannian metric associated to the balanced  $SU(3)$ -structure on  $G \times I$ . The Hermitian balanced structure on  $G \times I$  is given by

$$
F = e^{23} - \frac{1}{2}e^{24} + e^{45} + \frac{2-3t}{4}\sqrt[3]{\frac{2-3t}{2}}e^{24} + \sqrt[3]{\frac{2-3t}{2}}e^{1} \wedge dt,
$$
  
\n
$$
\Psi_{+} = \frac{1}{2}e^{123} - e^{135} - \frac{2-3t}{4}\sqrt[3]{\frac{2-3t}{2}}e^{123} + \sqrt[3]{\frac{(2-3t)^{2}}{4}}e^{124}
$$
  
\n(12)  
\n
$$
- \left(\sqrt[3]{\frac{2}{2-3t}}e^{25} + \sqrt[3]{\frac{2-3t}{2}}e^{34}\right) \wedge dt,
$$
  
\n
$$
\Psi_{-} = e^{125} + \sqrt[3]{\frac{(2-3t)^{2}}{4}}e^{134}
$$

<span id="page-7-0"></span>
$$
\Psi_{-} = e^{125} + \sqrt[3]{\frac{(2-3t)^2}{4}} e^{134} \n+ \left(\frac{1}{2}\sqrt[3]{\frac{2}{2-3t}} e^{23} - \frac{2-3t}{4}e^{23} + \sqrt[3]{\frac{2-3t}{2}} e^{24} - \sqrt[3]{\frac{2}{2-3t}} e^{35}\right) \wedge dt.
$$

Lie algebra  $(0,0,12,13,23)$ : A direct calculation shows that the family of balanced SU(2)-structures given by

$$
\eta(t) = \frac{2}{2-t}e^3,
$$
  
\n
$$
\omega_1(t) = \frac{2-t}{2}(e^{15} + e^{42}),
$$
  
\n
$$
\omega_2(t) = \frac{t(2-t)(t-4)}{4}e^{12} + \frac{2-t}{2}(e^{14} + e^{25}),
$$
  
\n
$$
\omega_3(t) = e^{12} - \frac{t(2-t)^2(t-4)}{8}e^{25} - \frac{(2-t)^2}{4}e^{45},
$$

satisfies the evolution equations [\(9\)](#page-6-1) for  $t \in \mathbb{R} - \{2\}$ . Observe that the volume form of the associated Riemannian metric along the family is given by

$$
\omega_1(t) \wedge \omega_1(t) \wedge \eta(t) = -2e^{12345},
$$

so it remains constant.

Let  $I = (-\infty, 2)$  and denote by G the simply-connected nilpotent Lie group with Lie algebra  $(0, 0, 12, 13, 23)$ . The basis of 1-forms on the product manifold  $G \times I$ given by

$$
\alpha^1 = e^1
$$
,  $\alpha^2 = e^2$ ,  $\alpha^3 = \frac{2-t}{2}e^5$ ,  $\alpha^4 = \frac{t(2-t)(t-4)}{4}e^2 + \frac{2-t}{2}e^4$ ,  
\n $\alpha^5 = \frac{2}{2-t}e^3$ ,  $\alpha^6 = dt$ ,

is orthonormal with respect to the Riemannian metric associated to the balanced  $SU(3)$ -structure on  $G \times I$ . The Hermitian balanced structure on  $G \times I$  is given by

<span id="page-8-1"></span>(13) 
$$
F = e^{12} - \frac{t(2-t)^2(t-4)}{8}e^{25} - \frac{(2-t)^2}{4}e^{45} + \frac{2}{2-t}e^{3} \wedge dt,
$$

$$
\Psi_{+} = -e^{135} + e^{234} - \frac{2-t}{2}(\frac{t(t-4)}{2}e^{12} + e^{14} + e^{25}) \wedge dt,
$$

$$
\Psi_{-} = -e^{134} - e^{235} + \frac{t(t-4)}{2}e^{123} + \frac{2-t}{2}(e^{15} - e^{24}) \wedge dt.
$$

As a consequence of our previous examples we conclude:

Corollary 3.3. The 6-dimensional simply-connected nilpotent Lie groups  $H_8$ ,  $H_{16}$  and  $H_{17}$  corresponding to the Lie algebras  $\mathfrak{h}_8 = (0, 0, 0, 0, 0, 12)$ ,  $\mathfrak{h}_{16} =$  $(0, 0, 0, 12, 14, 24)$  and  $\mathfrak{h}_{17} = (0, 0, 0, 0, 12, 15)$  have balanced Hermitian SU(3)structures.

It is worthy to remark that  $H_8$  and  $H_{16}$  have invariant complex structures, but none of them admit invariant compatible balanced metric [\[15\]](#page-19-2). On the other hand,  $H_{17}$  has no invariant complex structures.

Finally, recall that according to  $[12]$  if M is a 6-manifold with a family  $(F(s), \Psi_+(s), \Psi_-(s))$  of half-flat structures,  $s \in I = (a, b)$ , satisfying the evolution equations

$$
\partial_s \Psi_+ = dF, \qquad F \wedge \partial_s F = -d\Psi_-,
$$

then the product manifold  $M \times I$  has a Riemannian metric whose holonomy is contained in  $G_2$ ; in fact, the form  $\varphi = F(s) \wedge ds + \Psi_+(s)$  defining the  $G_2$ -structure is parallel. Therefore, the balanced Hermitian  $SU(3)$ -structures  $(11)$ ,  $(12)$  and  $(13)$ can be lifted to a metric with holonomy in  $G_2$ .

# <span id="page-8-0"></span>4. Holonomy of Bismut connection of balanced Hermitian metrics on solvmanifolds

In this section we provide some examples of compact Hermitian manifolds, endowed with an  $SU(n)$ -structure whose associated metric is balanced and such that the corresponding Bismut connection has holonomy equal to  $SU(n)$ , for  $n = 3$  and 4.

Let  $(M, J, g)$  be a Hermitian manifold and F be the Kähler form of the Hermitian structure  $(g, J)$ . Denote by  $\nabla^{LC}$  the Levi Civita connection of the metric g. Then the Bismut connection  $\nabla^B$  of  $(M, J, F, g)$  is characterized by the following formula

$$
g(\nabla_X^B Y, Z) = g(\nabla_X^{LC} Y, Z) + \frac{1}{2} T(X, Y, Z), \quad \forall X, Y, Z \in \Gamma(M, TM),
$$

where the torsion form  $T$  is given by

$$
T(X, Y, Z) = JdF(X, Y, Z).
$$

We will need to compute the curvature of  $\nabla^B$ . In order to do this, we will use the Cartan structure equations,

<span id="page-9-1"></span>(14)  
\n
$$
\begin{cases}\n de^{i} + \sum_{j=1}^{2n} \omega_{j}^{i} \wedge e^{j} = \tau^{i}, & i = 1, ..., 2n \\
\omega_{j}^{i} + \omega_{2n}^{j} = 0, & i, j = 1, ..., 2n,\n\end{cases}
$$
\n(15)  
\n
$$
\begin{cases}\n d\omega_{j}^{i} + \sum_{r=1}^{2n} \omega_{r}^{i} \wedge \omega_{j}^{r} = \Omega_{j}^{i}, & i, j = 1, ..., 2n\n\end{cases}
$$

$$
\left\{\n\begin{array}{l}\n\Omega^i_j + \Omega^j_{2n} = 0, & i, j = 1, \ldots, 2n,\n\end{array}\n\right.
$$

where  $\{e^1, \ldots, e^{2n}\}$  is an orthonormal coframe,  $\omega_j^i$  are the connection 1-forms,  $\tau^i$ are the torsion 2-forms and  $\Omega_j^i$  are the curvature 2-forms.

If 
$$
\{e_1, \ldots, e_{2n}\}
$$
 denotes the dual frame of  $\{e^1, \ldots, e^{2n}\}$ , then  $2n$ 

$$
\tau^i = \sum_{j < k=1}^{\infty} T_{ijk} e^j \wedge e^k \,, \quad i = 1, \ldots, 2n \,,
$$

where  $T_{ijk} = T(e_i, e_j, e_k)$ .

4.1. Dimension six. There are two 3-dimensional complex-parallelizable (non-Abelian) solvable Lie groups [\[14\]](#page-19-3). The complex structure equations are given by:

(I) 
$$
d\varphi^1 = d\varphi^2 = 0
$$
,  $d\varphi^3 = \varphi^{12}$ ;

(II) 
$$
d\varphi^1 = 0
$$
,  $d\varphi^2 = \varphi^{12}$ ,  $d\varphi^3 = -\varphi^{13}$ .

Here (I) is the Lie group underlying the Iwasawa manifold. In the following we show that they can be endowed with a balanced Hermitian  $SU(3)$ -structure and determine the holonomy of the corresponding Bismut connection.

<span id="page-9-0"></span>Theorem 4.1. Any 3-dimensional complex-parallelizable (non-Abelian) solvable Lie group has a Hermitian metric such that the holonomy of its Bismut connection is equal to SU(3).

*Proof.* First we show that for the standard Hermitian balanced structure  $(J, g)$  on the Lie group  $(I)$  the holonomy of its Bismut connection equal to  $SU(3)$ .

Let us consider the real basis  $(e^1, \ldots, e^6)$  given by

$$
\varphi^1 = e^1 + i e^2
$$
,  $\varphi^2 = e^3 + i e^4$ ,  $\varphi^3 = e^5 + i e^6$ .

In this case the structure equations are

$$
de^1 = de^2 = de^3 = de^4 = 0
$$
,  $de^5 = e^{13} - e^{24}$ ,  $de^6 = e^{14} + e^{23}$ .

Then, the complex structure  $J$  is given by

$$
Je1 = -e2, \quad Je2 = e1, \quad Je3 = -e4, \quad Je4 = e3, \quad Je5 = -e6, \quad Je6 = e5.
$$

The fundamental form F associated with the J-Hermitian metric  $g = \sum_{i=1}^{6} e^{i} \otimes e^{i}$ is given by

$$
F = e^{12} + e^{34} + e^{56}.
$$

Since  $dF = e^{136} - e^{145} - e^{235} - e^{246}$ , we get that g is balanced and the torsion  $T = JdF$  is given by

$$
T = -e^{135} - e^{146} - e^{236} + e^{245}.
$$

The non-zero connection 1-forms of the Bismut connection  $\nabla^B$  are

$$
\omega_5^1 = -e^3
$$
,  $\omega_6^1 = -e^4$ ,  $\omega_5^2 = e^4$ ,  $\omega_6^2 = -e^3$ ,  
\n $\omega_5^3 = e^1$ ,  $\omega_6^3 = e^2$ ,  $\omega_5^4 = -e^2$ ,  $\omega_6^4 = e^1$ .

The (linearly independent) curvature forms of the Bismut connection are

$$
\Omega_2^1 = 2e^{34}
$$
,  $\Omega_3^1 = -e^{13} - e^{24}$ ,  $\Omega_3^2 = e^{14} - e^{23}$ ,  $\Omega_4^3 = 2e^{12}$ ,

and the (linearly independent) curvature forms of the Bismut connection are

$$
\nabla_{E_1}^B(\Omega_2^1) = -2(e^{36} - e^{45}), \qquad \nabla_{E_2}^B(\Omega_2^1) = 2(e^{35} + e^{46}),
$$
  
\n
$$
\nabla_{E_3}^B(\Omega_4^3) = 2(e^{16} - e^{25}), \qquad \nabla_{E_4}^B(\Omega_4^3) = -2(e^{15} + e^{26}).
$$

Therefore,  $Hol(\nabla^B) = SU(3)$ .

For case (II) we consider the real basis  $(e^1, \ldots, e^6)$  given by

$$
\varphi^1 = e^1 + ie^2, \quad \varphi^2 = e^3 + ie^4, \quad \varphi^3 = e^5 + ie^6.
$$

In terms of this basis, the structure equations are

$$
de1 = de2 = 0,
$$
  $de3 = e13 - e24,$   $de4 = e14 + e23$   
 $de5 = -e15 + e26,$   $de6 = -e16 - e25.$ 

Then, the complex structure  $J$  is given by

$$
Je1 = -e2, \quad Je2 = e1, \quad Je3 = -e4, \quad Je4 = e3, \quad Je5 = -e6, \quad Je6 = e5.
$$

Then the fundamental form  $F$  associated with the  $J$ -Hermitian metric  $g = \sum_{i=1}^{6} e^{i} \otimes e^{i}$  is given by

$$
F = e^{12} + e^{34} + e^{56}.
$$

Since  $dF = 2(e^{134} - e^{156})$ , we get that g is balanced and the torsion  $T = JdF$  is given by

$$
T = -2(e^{234} - e^{256}).
$$

The non-zero connection 1-forms of the Bismut connection  $\nabla^B$  are

$$
\omega_5^1 = -e^3
$$
,  $\omega_6^1 = -e^4$ ,  $\omega_5^2 = e^4$ ,  $\omega_6^2 = -e^3$ ,  
\n $\omega_5^3 = e^1$ ,  $\omega_6^3 = e^2$ ,  $\omega_5^4 = -e^2$ ,  $\omega_6^4 = e^1$ .

The (linearly independent) curvature forms of the Bismut connection are

$$
\Omega_3^1 = -e^{13} - e^{24}, \quad \Omega_4^1 = -e^{14} + e^{23}, \quad \Omega_5^1 = -e^{15} - e^{26}, \quad \Omega_6^1 = -e^{16} + e^{25}
$$
  

$$
\Omega_4^3 = -2e^{34}, \qquad \Omega_5^3 = e^{35} + e^{46}, \qquad \Omega_6^3 = e^{36} - e^{45}, \qquad \Omega_6^5 = -2e^{56}.
$$

Therefore, the holonomy of  $\nabla^B$  is again equal to SU(3).

Next, we study the holonomy of the Bismut connection for Hermitian balanced metrics on 6-nilmanifolds. First of all, we recall [\[15\]](#page-19-2) that if a nilmanifold  $M = \Gamma \backslash G$ admits a Hermitian balanced metric (not necessarily invariant), then the Lie algebra **g** of G is isomorphic to  $\mathfrak{h}_1, \ldots, \mathfrak{h}_6$  or  $\mathfrak{h}_{19}^-$ , where  $\mathfrak{h}_1 = (0, 0, 0, 0, 0, 0)$  is the Abelian Lie algebra and

$\mathfrak{h}_2 = (0,0,0,0,12,34),$	$\mathfrak{h}_5 = (0,0,0,0,13+42,14+23),$
$\mathfrak{h}_3 = (0,0,0,0,0,12+34),$	$\mathfrak{h}_6 = (0,0,0,0,12,13),$
$\mathfrak{h}_4 = (0,0,0,0,12,14+23),$	$\mathfrak{h}_{19} = (0,0,0,12,23,14-35).$

,

where  $\mathfrak{h}_5$  is the Lie algebra underlying the Iwasawa manifold considered in Theo-rem [4.1.](#page-9-0) Therefore, the Lie algebra  $\mathfrak{h}_{19}^-$  is the unique 6-dimensional 3-step nilpotent Lie algebra admitting balanced structures,

On the other hand, in [\[8\]](#page-18-9)[Corollary 6.2, Example 6.3] an explicit balanced Hermitian metric on  $\mathfrak{h}_3$ ,  $\mathfrak{h}_4$ ,  $\mathfrak{h}_5$ ,  $\mathfrak{h}_6$  is given such that the holonomy of the corresponding Bismut connection is equal to SU(3) for  $\mathfrak{h}_4$ ,  $\mathfrak{h}_5$  and  $\mathfrak{h}_6$ . For  $\mathfrak{h}_3$  they show that there is a reduction of the holonomy group, whereas it remains to study the Lie algebra  $\mathfrak{h}_2$  because the particular coefficients given at the end of the proof of [\[8\]](#page-18-9)[Corollary 6.2 correspond to  $\mathfrak{h}_6$ .

Next we prove that  $\mathfrak{h}_2$  and  $\mathfrak{h}_{19}$ – also admit a balanced Hermitian SU(3)-structure such that the holonomy of its Bismut connection also equals  $SU(3)$ . That is, we prove the following

Theorem 4.2. Let M be a 6-dimensional nilmanifold admitting invariant balanced Hermitian structures  $(q, J)$ . If the first Betti number of M is  $\leq 4$ , then there is  $(g, J)$  such that the holonomy of its Bismut connection is equal to SU(3).

*Proof.* A 6-dimensional nilmanifold with  $b_1(M) \leq 4$  admitting invariant balanced Hermitian structures  $(g, J)$  has underlying Lie algebra isomorphic to  $\mathfrak{h}_2$ ,  $\mathfrak{h}_4$ ,  $\mathfrak{h}_5$ ,  $\mathfrak{h}_6$ or  $\mathfrak{h}_{19}^-$ .

In order to define a balanced Hermitian  $SU(3)$ -structure on  $\mathfrak{h}_2$  such that the holonomy of its Bismut connection is equal to SU(3), let us consider the structure equations

<span id="page-11-0"></span>(16) 
$$
de^1 = de^2 = de^3 = de^4 = 0
$$
,  $de^5 = e^{13} - e^{24}$ ,  $de^6 = -2e^{12} + e^{14} + e^{23} + 2e^{34}$ ,  
and the complex structure L given by

and the complex structure J given by

$$
Je1 = -e2, \quad Je2 = e1, \quad Je3 = -e4, \quad Je4 = e3, \quad Je5 = -e6, \quad Je6 = e5.
$$

Firstly, we notice that the structure equations [\(16\)](#page-11-0) correspond to the Lie algebra  $\n *h*<sub>2</sub>$ . To see this it is sufficient to express the equations with respect to the new basis

$$
f1 = -2 e2 + \sqrt{3} e3 + e4, \t f2 = e1 - \sqrt{3} e2 + 2 e3, \t f3 = 2 e2 + \sqrt{3} e3 - e4,f4 = e1 + \sqrt{3} e2 + 2 e3, \t f5 = -\sqrt{3} e5 - e6, \t f6 = -\sqrt{3} e5 + e6.
$$

 $\sum_{i=1}^{6} e^{i} \otimes e^{i}$  is given by  $F = e^{12} + e^{34} + e^{56}$ . Since Now, the fundamental form  $F$  associated to the J-Hermitian metric  $g =$ 

$$
dF = e^{136} - e^{246} + 2e^{125} - e^{145} - e^{235} - 2e^{345},
$$

we get that  $g$  is balanced and the torsion  $T$  is given by

$$
T = -2e^{126} - e^{135} - e^{146} - e^{236} + e^{245} + 2e^{346}.
$$

The (linearly independent) curvature forms for the Bismut connection are

$$
\Omega_2^1 = -2(2e^{12} - e^{14} - e^{23} - 3e^{34}), \qquad \Omega_3^1 = -(e^{13} + e^{24}),
$$
  
\n
$$
\Omega_4^1 = -(e^{14} - e^{23}), \qquad \Omega_5^1 = -2e^{46},
$$
  
\n
$$
\Omega_6^1 = 2e^{36}, \qquad \Omega_4^3 = 2(3e^{12} - e^{14} - e^{23} - 2e^{34}),
$$
  
\n
$$
\Omega_5^3 = -2e^{26}, \qquad \Omega_6^3 = 2e^{16}.
$$

Therefore,  $\text{Hol}(\nabla^B) = \text{SU}(3)$ .

We have given a balanced metric with Hol( $\nabla^B$ ) = SU(3) for each case.

To complete the proof, we show that there is a balanced Hemitian  $SU(3)$ structure  $(J, g)$  on  $\mathfrak{h}_{19}^-$  with holonomy of its Bismut connection equal to SU(3).

We know that the structure equations of  $\mathfrak{h}_{19}^-$  are

$$
de^1 = de^2 = de^3 = 0
$$
,  $de^4 = e^{12}$ ,  $de^5 = e^{23}$ ,  $de^6 = e^{14} - e^{35}$ .

We define the complex structure  $J$  given by

$$
Je1 = e3, \quad Je2 = 4e6, \quad Je3 = -e1, \quad Je4 = -e5, \quad Je5 = e4, \quad Je6 = -\frac{1}{4}e2.
$$

 $\sum_{i=1}^{6} e^{i} \otimes e^{i}$  is given by  $F = -e^{13} - e^{26} + e^{45}$  and Then, the fundamental form F associated to the J-Hermitian metric  $g =$ 

$$
dF = -e^{124} + e^{125} - e^{234} - e^{235}
$$

.

Therefore, we get that  $g$  is balanced and the torsion  $T$  is given by

$$
T = e^{146} - e^{156} - e^{346} - e^{356}.
$$

The non-zero connection 1-forms of the Bismut connection  $\nabla^B$  are

$$
\omega_2^1 = -\frac{1}{2}e^4, \qquad \omega_4^1 = -\frac{1}{2}e^2 - e^6, \qquad \omega_5^1 = \frac{1}{2}e^6, \qquad \omega_6^1 = -\frac{1}{2}e^5,
$$
  
\n
$$
\omega_3^2 = -\frac{1}{2}e^5, \qquad \omega_4^2 = \frac{1}{2}e^1, \qquad \omega_5^2 = -\frac{1}{2}e^3, \qquad \omega_4^3 = \frac{1}{2}e^6
$$
  
\n
$$
\omega_5^3 = \frac{1}{2}e^2 + e^6, \qquad \omega_6^3 = -\frac{1}{2}e^4, \qquad \omega_6^4 = \frac{1}{2}e^3, \qquad \omega_6^5 = \frac{1}{2}e^1.
$$

and the (linearly independent) curvature forms for the Bismut connection are

$$
\Omega_2^1 = -\frac{1}{4}(3e^{12} + 2e^{16} + e^{36}), \qquad \Omega_3^1 = \frac{1}{2}(e^{26} + e^{45}),
$$
  
\n
$$
\Omega_4^1 = -\frac{3}{4}(e^{14} - e^{35}), \qquad \Omega_5^1 = \frac{1}{4}(2e^{14} - e^{15} - e^{34} - 2e^{35}),
$$
  
\n
$$
\Omega_6^1 = -\frac{1}{4}(e^{16} + 3e^{23} + 2e^{36}), \qquad \Omega_4^2 = \frac{1}{4}(e^{24} - 2e^{46} - e^{56}),
$$
  
\n
$$
\Omega_5^2 = \frac{1}{4}(e^{25} + e^{46} - 2e^{56}), \qquad \Omega_6^2 = \frac{1}{2}(e^{13} - e^{45}).
$$

Therefore,  $Hol(\nabla^B) = SU(3)$ .

To finish this section, we show an example of a six-dimensional compact solvmanifold with a balanced metric such that the holonomy of its Bismut connection is equal to SU(3).

Let  $\mathfrak g$  be the solvable Lie algebra whose structure equations are

$$
de^1 = 0
$$
,  $de^2 = 0$ ,  $de^3 = e^1 \wedge e^3$ ,  $de^4 = -e^1 \wedge e^4$ ,  $de^5 = e^1 \wedge e^5$ ,  $de^6 = -e^1 \wedge e^6$ 

Let  $G$  be the simply-connected Lie group whose Lie algebra is  $\mathfrak{g}$ . It can be easily checked that, for any  $X \in \mathfrak{g}$ , ad<sub>X</sub> has real eigenvalues, i.e.  $\mathfrak{g}$  is completely solvable. Thus G has a uniform discrete subgroup Γ, such that  $M^6 = \Gamma \backslash G$  is a 6-dimensional compact solvmanifold.

Define an (integrable) complex structure  $J$  on  $M^6$  by setting

$$
Je1 = -e2, Je2 = e1, Je3 = -e5, Je4 = -e6, Je5 = e3, Je6 = e4,
$$

and a J-Hermitian structure g on M by  $g = \sum_{i=1}^{6} e^{i} \otimes e^{i}$ . Then  $F = \sum_{i=1}^{3} e^{2i-1} \wedge e^{2i}$ . Hence

$$
dF = 2\left(e^{135} - e^{146}\right)
$$

and consequently, the torsion form of the Bismut connection is given by

$$
T = 2\left(e^{246} - e^{235}\right)
$$

.

.

By the above expression, the only non-zero components of the torsion are

$$
T_{235} = -2 \,, \quad T_{246} = 2 \,.
$$

By solving [\(14\)](#page-9-1), we obtain that the non zero connection forms are given by

$$
\omega_3^1 = -e^3, \quad \omega_4^1 = e^4, \qquad \omega_5^1 = -e^5, \quad \omega_6^1 = e^6,
$$
  

$$
\omega_3^2 = e^5, \qquad \omega_4^2 = -e^6, \quad \omega_5^2 = -e^3, \quad \omega_6^2 = e^4,
$$
  

$$
\omega_4^3 = e^5, \qquad \omega_5^3 = e^2, \qquad \omega_5^4 = 0, \qquad \omega_6^4 = -e^2
$$

Therefore, we get

$$
\Omega_2^1 = 2\left(e^{35} + e^{46}\right), \quad \Omega_3^1 = -e^{13} - e^{25}, \quad \Omega_4^1 = -e^{14} - e^{26}, \quad \Omega_5^1 = -e^{15} + e^{23},
$$
  
\n
$$
\Omega_6^1 = -e^{16} + e^{24}, \qquad \Omega_3^2 = e^{15} - e^{23}, \qquad \Omega_4^2 = e^{16} - e^{24}, \qquad \Omega_5^2 = -e^{13} - e^{25},
$$
  
\n
$$
\Omega_6^2 = -e^{14} - e^{26}, \qquad \Omega_4^3 = e^{34} + e^{56}, \qquad \Omega_5^3 = -2e^{35}, \qquad \Omega_6^3 = e^{36} + e^{45}
$$
  
\n
$$
\Omega_5^4 = e^{36} + e^{45}, \qquad \Omega_6^4 = -2e^{46}, \qquad \Omega_6^5 = e^{34} + e^{56}.
$$

By the above expression for the curvature forms, we obtain that  $Hol(\nabla^B) = SU(3)$ . On the other hand, the manifold  $M$  has no Kähler structures. Indeed, in view of the Main Theorem in [\[10\]](#page-18-10), a compact solvmanifold of completely solvable type has a Kähler structure if and only if it is a complex torus. Finally, note that

$$
dF^2 = 2\left(e^{135} - e^{146}\right) \wedge \left(e^{12} + e^{35} + e^{46}\right) = 0,
$$

i.e. g is a balanced metric.

4.2. Balanced Hermitian metrics on compact 8-solvmanifolds. Here we present examples of compact balanced Hermitian 8-manifolds with an SU(4) structure such that the holonomy of its Bismut connection is all the Lie group  $SU(4)$ . Furthermore, according to [\[14\]](#page-19-3), all the examples are holomorphic parallelizable.

Example 4.3. Consider the Lie algebra defined by the complex structure equations

$$
d\varphi^{1} = d\varphi^{2} = 0
$$
,  $d\varphi^{3} = -\varphi^{12}$ ,  $d\varphi^{4} = -2\varphi^{13}$ .

Let  $e^1, \ldots, e^8$  be the basis given by  $e^{2j-1} + i e^{2j} = \varphi^j$ , for  $j = 1, \ldots, 4$ .

The corresponding real structure equations are

$$
de^1 = de^2 = de^3 = de^4 = 0
$$
,  $de^5 = -e^{13} + e^{24}$ ,  
\n $de^6 = -e^{14} - e^{23}$ ,  $de^7 = -2(e^{15} - e^{26})$ ,  $de^8 = -2(e^{16} + e^{25})$ .

Let  $J$  be the complex structure given by

$$
Je1 = -e2
$$
,  $Je2 = e1$ ,  $Je3 = -e4$ ,  $Je4 = e3$ ,  
\n $Je5 = -e6$ ,  $Je6 = e5$ ,  $Je7 = -e8$ ,  $Je8 = e7$ .

Then fundamental form F associated with the J-Hermitian metric  $g = \sum_{i=1}^{8} e^{i} \otimes e^{i}$  is given by  $F = \sum_{j=1}^{4} e^{2j-1} \wedge e^{2j}$ . Since

$$
dF = -e^{136} + e^{145} - 2e^{158} + 2e^{167} + e^{235} + e^{246} + 2e^{257} + 2e^{268},
$$

we get that  $q$  is balanced and the torsion  $T$  is given by

$$
T = JdF = e^{135} + e^{146} + 2e^{157} + 2e^{168} + e^{236} - e^{245} + 2e^{258} - 2e^{267}.
$$

The non-zero connection 1-forms of the Bismut connection  $\nabla^B$  are

$$
\omega_5^1 = e^3, \qquad \omega_6^1 = e^4, \qquad \omega_7^1 = 2e^5, \quad \omega_8^1 = 2e^6, \qquad \omega_5^2 = -e^4, \quad \omega_6^2 = e^3,
$$
  

$$
\omega_7^2 = -2e^6, \quad \omega_8^2 = 2e^5, \qquad \omega_5^3 = -e^1, \quad \omega_6^3 = -e^2, \qquad \omega_5^4 = e^2, \qquad \omega_6^4 = -e^1,
$$
  

$$
\omega_7^5 = -2e^1, \quad \omega_8^5 = -2e^2, \quad \omega_7^6 = 2e^2, \qquad \omega_8^6 = -2e^1.
$$

The following curvature forms for the Bismut connection are linearly independent:

$$
\Omega_3^1 = -e^{13} - e^{24}, \qquad \Omega_4^1 = -e^{14} + e^{23}, \qquad \Omega_5^1 = -4(e^{15} + e^{26}),
$$
  
\n
$$
\Omega_6^1 = -4(e^{16} - e^{25}), \qquad \Omega_4^3 = 2e^{12}, \qquad \Omega_6^5 = 2(3e^{12} - e^{34}),
$$
  
\n
$$
\Omega_7^5 = -2(e^{35} + e^{46}), \qquad \Omega_8^5 = -2(e^{36} - e^{45}), \qquad \Omega_8^7 = -8(e^{12} + e^{56}).
$$

This gives a 9-dimensional space. Moreover, the following 6 covariant derivatives of the curvature forms

$$
\nabla_{E_2}^B(\Omega_2^1) = -16(e^{57} + e^{68}), \nabla_{E_5}^B(\Omega_3^1) = -2(e^{37} + e^{48}), \nabla_{E_6}^B(\Omega_3^1) = -2(e^{38} - e^{47}),
$$
  
\n
$$
\nabla_{E_6}^B(\Omega_5^1) = -8(e^{58} - e^{67}), \nabla_{E_5}^B(\Omega_4^3) = -4(e^{18} - e^{27}), \nabla_{E_6}^B(\Omega_4^3) = 4(e^{17} + e^{28}),
$$

are linearly independent, therefore Hol( $\nabla^B$ )=SU(4).

Example 4.4. Consider the complex structure equations

$$
d\varphi^{1} = 0
$$
,  $d\varphi^{2} = \varphi^{12}$ ,  $d\varphi^{3} = c\varphi^{13}$ ,  $d\varphi^{4} = -(1+c)\varphi^{14}$ ,

where  $c(1+c) \neq 0$ . Let  $e^1, \ldots, e^8$  be the basis given by  $\varphi^j = e^{2j-1} + i e^{2j}$ , for  $j = 1, \ldots, 4.$ 

The corresponding real structure equations are

$$
de1 = de2 = 0,
$$
  $de3 = e13 - e24,$   $de4 = e14 + e23,$   
\n $de5 = c(e15 - e26)$   $de6 = c(e16 + e25), de7 = -(1 + c)(e17 - e28),$   
\n $de8 = -(1 + c)(e18 + e27).$ 

Let  $J$  be the complex structure given by

$$
Je1 = -e2
$$
,  $Je2 = e1$ ,  $Je3 = -e4$ ,  $Je4 = e3$ ,  
\n $Je5 = -e6$ ,  $Je6 = e5$ ,  $Je7 = -e8$ ,  $Je8 = e7$ .

Then  $g = \sum_{i=1}^{8} e^{i} \otimes e^{i}$  is a J-Hermitian metric and the fundamental form is given by  $F = \sum_{j=1}^{4} e^{2j-1} \wedge e^{2j}$ . Hence

$$
dF = 2\left(e^{134} + ce^{156} - (1+c)e^{178}\right)
$$

and consequently we get that  $g$  is balanced and the torsion  $T$  is expressed by

$$
T = JdF = -2\left(e^{234} + ce^{256} - (1+c)e^{278}\right).
$$

A direct computation shows that the non-zero connection 1-forms of the Bismut connection  $\nabla^B$  are

$$
\omega_3^1 = -e^3, \qquad \omega_4^1 = -e^4, \qquad \omega_5^1 = -ce^5, \quad \omega_6^1 = -ce^6, \quad \omega_7^1 = (1+c)e^7,
$$
  
\n
$$
\omega_8^1 = (1+c)e^8, \qquad \omega_3^2 = e^4, \qquad \omega_4^2 = -e^3, \qquad \omega_5^2 = ce^6, \qquad \omega_6^2 = -ce^5,
$$
  
\n
$$
\omega_7^2 = -(1+c)e^8, \quad \omega_8^2 = (1+c)e^7, \quad \omega_4^3 = 2e^2, \qquad \omega_6^5 = 2ce^2, \qquad \omega_8^7 = -2(1+c)e^2.
$$

The following curvature forms for the Bismut connection are linearly independent:

$$
\Omega_3^1 = -(\alpha^{13} + \alpha^{24}), \qquad \Omega_4^1 = -\alpha^{14} + \alpha^{23}, \qquad \Omega_5^1 = -c^2(\alpha^{15} + \alpha^{26}),
$$
  
\n
$$
\Omega_6^1 = -c^2(\alpha^{16} - \alpha^{25}), \qquad \Omega_7^1 = -(1+c)^2(\alpha^{17} + \alpha^{28}),
$$
  
\n
$$
\Omega_8^1 = -(1+c)^2(\alpha^{18} - \alpha^{27}),
$$
  
\n
$$
\Omega_4^3 = -2\alpha^{34}, \qquad \Omega_5^3 = -c(\alpha^{35} + \alpha^{46}), \qquad \Omega_6^3 = -c(\alpha^{36} - \alpha^{45}),
$$
  
\n
$$
\Omega_7^3 = (1+c)(\alpha^{37} + \alpha^{48}), \qquad \Omega_8^3 = (1+c)(\alpha^{38} - \alpha^{47}), \qquad \Omega_6^5 = -2c^2\alpha^{56},
$$
  
\n
$$
\Omega_7^5 = c(1+c)(\alpha^{57} + \alpha^{68}), \qquad \Omega_8^5 = c(1+c)(\alpha^{58} - \alpha^{67}), \qquad \Omega_8^7 = -2(1+c)^2\alpha^{78}.
$$

Therefore,  $Hol(\nabla^B)=SU(4)$  for any value of the parameter c.

Example 4.5. Consider the complex structure equations

$$
d\varphi^{1} = 0
$$
,  $d\varphi^{2} = \varphi^{12}$ ,  $d\varphi^{3} = -2\varphi^{13}$ ,  $d\varphi^{4} = \varphi^{14} - \varphi^{12}$ .

Let  $e^1, \ldots, e^8$  be the basis given by  $\varphi^j = e^{2j-1} + i e^{2j}$ , for  $j = 1, \ldots, 4$ .

The corresponding real structure equations are

$$
de1 = de2 = 0, de3 = e13 - e24, de4 = e14 + e23, de5 = -2(e15 - e26),de6 = -2(e16 + e25), de7 = -e13 + e17 + e24 - e28,de8 = -e14 + e18 - e23 + e27.
$$

Let  ${\cal J}$  be the complex structure given by

$$
Je1 = -e2, \t Je2 = e1, \t Je3 = -e4, \t Je4 = e3,\nJe5 = -e6, \t Je6 = e5, \t Je7 = -e8, \t Je8 = e7.
$$

The fundamental form F associated to the J-Hermitian metric  $g = \sum_{i=1}^{8} e^{i} \otimes e^{i}$  is given by  $F = \sum_{j=1}^{4} e^{2j-1} \wedge e^{2j}$ . In such a case,

$$
dF = 2e^{134} - e^{138} + e^{147} - 4e^{156} + 2e^{178} + e^{237} + e^{248}
$$

and the torsion  $T$  is given by

$$
T = JdF = e^{137} + e^{148} - 2e^{234} + e^{238} - e^{247} + 4e^{256} - 2e^{278}.
$$

Again, g is balanced. The non-zero connection 1-forms of the Bismut connection  $\nabla^B$  are

$$
\omega_3^1 = -e^3, \qquad \omega_4^1 = -e^4, \qquad \omega_5^1 = 2e^5, \quad \omega_6^1 = 2e^6, \qquad \omega_7^1 = e^3 - e^7,
$$
  
\n
$$
\omega_8^1 = e^4 - e^8, \qquad \omega_3^2 = e^4, \qquad \omega_4^2 = -e^3, \quad \omega_5^2 = -2e^6, \quad \omega_6^2 = 2e^5,
$$
  
\n
$$
\omega_7^2 = -e^4 + e^8, \quad \omega_8^2 = e^3 - e^7, \quad \omega_4^3 = 2e^2, \quad \omega_7^3 = -e^1,
$$
  
\n
$$
\omega_8^3 = -e^2, \qquad \omega_7^4 = e^2, \qquad \omega_8^4 = -e^1, \quad \omega_6^5 = -4e^2, \quad \omega_8^7 = 2e^2.
$$

The following 15 curvature forms for the Bismut connection are linearly independent:

$$
\Omega_3^1 = -2e^{13} + e^{17} - 2e^{24} + e^{28},
$$
\n
$$
\Omega_4^1 = -2e^{14} + e^{18} + 2e^{23} - e^{27},
$$
\n
$$
\Omega_5^1 = -4(e^{15} + e^{26}),
$$
\n
$$
\Omega_7^1 = e^{13} - e^{17} + e^{24} - e^{28},
$$
\n
$$
\Omega_8^1 = e^{14} - e^{18} - e^{23} + e^{27},
$$
\n
$$
\Omega_4^3 = 2(e^{12} - e^{34}),
$$
\n
$$
\Omega_5^3 = 2(e^{36} - e^{45}),
$$
\n
$$
\Omega_6^3 = 2e^{34} - e^{38} - 2e^{43} + e^{47},
$$
\n
$$
\Omega_7^5 = -8e^{56},
$$
\n
$$
\Omega_7^7 = 2(e^{35} + e^{46} + e^{57} + e^{68}),
$$
\n
$$
\Omega_8^7 = -2(e^{12} + e^{34} - e^{38} + e^{47} + e^{78}).
$$
\n
$$
\Omega_8^7 = -2(e^{12} + e^{34} - e^{38} + e^{47} + e^{78}).
$$

Therefore,  $Hol(\nabla^B)=SU(4)$ .

**Example 4.6.** Let  $\varphi^j = e^{2j-1} + i e^{2j}$ ,  $j = 1, ..., 4$  be complex (1,0)-forms satisfying

<span id="page-16-0"></span>(17) 
$$
d\varphi^1 = 0
$$
,  $d\varphi^2 = \varphi^{12}$ ,  $d\varphi^3 = -\varphi^{13}$ ,  $d\varphi^4 = -\varphi^{23}$ .

Hence

$$
\label{eq:2.12} \left\{ \begin{array}{l} d e^1 = 0, \;\; d e^2 = 0, \;\; d e^3 = e^{13} - e^{24}, \qquad d e^4 = e^{14} + e^{23}, \vspace{0.12cm} \\ d e^5 = - e^{15} + e^{26}, \;\; d e^6 = - e^{16} - e^{25}, \qquad d e^7 = - e^{35} + e^{46}, \;\; d e^8 = - e^{36} - e^{45} \,. \end{array} \right.
$$

Let  $\mathfrak g$  be the real 8-dimensional Lie algebra whose dual is spanned by  $\{e^1, \ldots, e^8\}$ , and let  $G$  be the simply-connected Lie group whose Lie algebra is  $\mathfrak{g}$ . It is immediate to see that g is a 3-step solvable but not completely solvable Lie algebra. In fact,  $\mathfrak{g}' = \langle e_3, e_4, e_5, e_6, e_7, e_8 \rangle$ ,  $\mathfrak{g}'' = \langle e_7, e_8 \rangle$ ,  $\mathfrak{g}''' = \{0\}$ . However, by [\[14\]](#page-19-3) it turns out that G has a uniform discrete subgroup  $\Gamma$ , such that  $M^8 = \Gamma \backslash G$  is a compact solvmanifold of dimension 8.

Define an (integrable) complex structure  $J$  on  $M^8$  by setting

$$
Je1 = -e2, Je2 = e1, Je3 = -e4, Je4 = -e3,Je5 = -e6, Je6 = e5, Je7 = -e8, Je8 = e7,
$$

and a J-Hermitian structure g on M by setting  $g = \sum_{i=1}^{8} e^{i} \otimes e^{i}$ . Then  $F = \sum_{j=1}^{4} e^{2j-1} \wedge e^{2j}$ . Hence

$$
dF = 2\left(e^{134} - e^{156}\right) - e^{358} + e^{468} + e^{367} + e^{457}
$$

and consequently

$$
T=2\left(e^{256}-e^{234}\right)-e^{467}+e^{357}+e^{458}+e^{368}\,.
$$

Therefore we obtain that the non zero connection and curvature forms of the Bismut connection are given respectively by

(18) 
$$
\omega_3^1 = -e^3
$$
,  $\omega_4^1 = -e^4$ ,  $\omega_5^1 = e^5$ ,  $\omega_6^1 = e^6$ ,  $\omega_3^2 = e^4$ ,  $\omega_4^2 = -e^3$   
\n(18)  $\omega_5^2 = -e^6$ ,  $\omega_6^2 = e^5$ ,  $\omega_4^3 = 2e^2$ ,  $\omega_7^3 = e^5$ ,  $\omega_8^3 = e^6$ ,  $\omega_7^4 = -e^6$   
\n $\omega_8^4 = e^5$ ,  $\omega_6^5 = -2e^2$ ,  $\omega_7^5 = -e^3$ ,  $\omega_8^5 = -e^4$ ,  $\omega_7^6 = e^4$ ,  $\omega_8^6 = -e^3$ 

<span id="page-17-0"></span>and by  
\n(19)  
\n
$$
\Omega_2^1 = 2(e^{34} + e^{56}),
$$
  $\Omega_3^1 = -e^{24} - e^{13},$   $\Omega_4^1 = e^{23} - e^{14},$   $\Omega_5^1 = -e^{26} - e^{15}$   
\n $\Omega_6^1 = e^{25} - e^{16},$   $\Omega_3^2 = -e^{23} + e^{14},$   $\Omega_4^2 = -e^{13} - e^{24},$   $\Omega_5^2 = e^{16} - e^{25}$   
\n $\Omega_6^2 = -e^{15} - e^{26},$   $\Omega_4^3 = 2(-e^{34} + e^{56}),$   $\Omega_7^3 = -e^{26} - e^{15},$   $\Omega_8^3 = e^{25} - e^{16}$   
\n $\Omega_7^4 = -e^{25} + e^{16},$   $\Omega_8^4 = -e^{26} - e^{15},$   $\Omega_6^5 = 2(-e^{56} + e^{34}),$   $\Omega_7^5 = -e^{24} - e^{13}$   
\n $\Omega_8^5 = e^{23} - e^{14},$   $\Omega_7^6 = -e^{23} + e^{14},$   $\Omega_8^6 = -e^{24} - e^{13},$   
\n $\Omega_8^7 = -2(e^{56} + e^{34}).$ 

Hence the six 2-forms

$$
\Omega_2^1 = 2(e^{34} + e^{56}), \quad \Omega_3^1 = -e^{24} - e^{13}, \quad \Omega_4^1 = e^{23} - e^{14},
$$
  

$$
\Omega_5^1 = -e^{26} - e^{15}, \quad \Omega_6^1 = e^{25} - e^{16}, \quad \Omega_4^3 = 2(-e^{34} + e^{56})
$$

are linearly independent. Therefore, dim(hol  $(\nabla^B)$ ) ≥ 6. We need to compute the covariant derivative of the curvature. By the above expression for the connection forms, we get:

$$
\nabla^{B}e^{1} = e^{3} \otimes e^{3} + e^{4} \otimes e^{4} - e^{5} \otimes e^{5} - e^{6} \otimes e^{6}
$$
  
\n
$$
\nabla^{B}e^{2} = -e^{4} \otimes e^{3} + e^{3} \otimes e^{4} + e^{6} \otimes e^{5} - e^{5} \otimes e^{6}
$$
  
\n
$$
\nabla^{B}e^{3} = -e^{3} \otimes e^{1} + e^{4} \otimes e^{2} - 2e^{2} \otimes e^{4} - e^{5} \otimes e^{7} - e^{6} \otimes e^{8}
$$
  
\n
$$
\nabla^{B}e^{4} = -e^{4} \otimes e^{1} - e^{3} \otimes e^{2} + 2e^{2} \otimes e^{3} + e^{6} \otimes e^{7} - e^{5} \otimes e^{8}
$$
  
\n
$$
\nabla^{B}e^{5} = e^{5} \otimes e^{1} - e^{6} \otimes e^{2} + 2e^{2} \otimes e^{6} + e^{3} \otimes e^{7} + e^{4} \otimes e^{8}
$$
  
\n
$$
\nabla^{B}e^{6} = e^{6} \otimes e^{1} + e^{5} \otimes e^{2} - 2e^{2} \otimes e^{5} - e^{4} \otimes e^{7} + e^{3} \otimes e^{8}
$$
  
\n
$$
\nabla^{B}e^{7} = e^{5} \otimes e^{3} - e^{6} \otimes e^{4} - e^{3} \otimes e^{5} + e^{4} \otimes e^{6}
$$
  
\n
$$
\nabla^{B}e^{8} = e^{6} \otimes e^{3} + e^{5} \otimes e^{4} - e^{4} \otimes e^{5} - e^{3} \otimes e^{6}
$$
.

By the above formulas, a straightforward computation, taking into account [\(19\)](#page-17-0), gives

$$
\nabla_{e_3}^B \Omega_2^1 = 2 \left( -e^{14} + e^{23} - e^{67} + e^{58} \right)
$$
  
\n
$$
\nabla_{e_4}^B \Omega_2^1 = 2 \left( e^{13} + e^{24} - e^{68} - e^{57} \right)
$$
  
\n
$$
\nabla_{e_5}^B \Omega_2^1 = 2 \left( e^{47} - e^{38} + e^{16} - e^{25} \right)
$$
  
\n
$$
\nabla_{e_6}^B \Omega_2^1 = 2 \left( e^{48} + e^{37} - e^{26} - e^{15} \right)
$$
  
\n
$$
\nabla_{e_6}^B \Omega_4^1 = -e^{35} - e^{28} - e^{46} - e^{17}
$$
  
\n
$$
\nabla_{e_4}^B \Omega_5^1 = -e^{18} + e^{27} + e^{36} - e^{45}
$$
  
\n
$$
\nabla_{e_3}^B \Omega_6^1 = e^{45} + e^{27} - e^{36} - e^{18}
$$
  
\n
$$
\nabla_{e_4}^B \Omega_6^1 = -e^{35} + e^{28} - e^{46} + e^{17}
$$
  
\n
$$
\nabla_{e_5}^B \Omega_6^1 = 2(e^{56} - e^{12}).
$$

Therefore, we have proved the following

**Proposition 4.7.** The holomorphic parallelizable solvmanifold  $M = \Gamma \backslash G$ , where  $\{\varphi^1,\ldots,\varphi^4\}$  are the complex  $(1,0)$ -forms satisfying  $(17)$ , it is endowed with an Hermitian metric with Kähler form

$$
F=\frac{i}{2}\sum_{j=1}^4\varphi^j\wedge\overline{\varphi}^j
$$

satisfying  $dF^3 = 0$  and the infinitesimal holonomy of the Bismut connection is  $\mathfrak{su}(4)$ .

Acknowledgments. This work has been partially supported through grant MCyT (Spain) MTM2005-08757-C04-02, by the Project MIUR "Geometric Properties of Real and Complex Manifolds" and by GNSAGA of INdAM.

#### **REFERENCES**

- <span id="page-18-4"></span>[1] L.C. de Andrés, L.A. Cordero, M. Fernández, J.J. Mencia, Examples of four-dimensional compact locally conformal Kähler manifolds, Geom. Dedicata 29 (1989), 227-232.
- <span id="page-18-3"></span><span id="page-18-0"></span>[2] Bismut J. M.: A local index theorem of non-Kähler manifolds,  $Math. Ann. 284 (1989), pp.$ 681–699.
- <span id="page-18-6"></span>[3] C.Boyer, K. Galicki, 3-Sasakian manifolds, Surveys Diff. Geom. 6 (1999), 123–184.
- [4] S. Chiossi, S. Salamon, The intrinsic torsion of SU(3) and  $G_2$  structures, In: Gil-Medrano, O., Miquel, V. (eds.), Differential geometry, Valencia, 2001, World Sci. Publishing, River Edge, NJ, 2002, 115–133.
- <span id="page-18-7"></span><span id="page-18-2"></span>[5] D. Conti, S. Salamon, Generalized Killing spinors in dimension 5, Trans. AMS 359 (2007), 5319–5343.
- <span id="page-18-1"></span>[6] D. Conti, A. Tomassini, Special symplectic six-manifolds, Q. J. Math. 58 (2007), 297–311.
- [7] M. Fernández, S. Ivanov, L. Ugarte, R. Villacampa, Non-Kähler Heterotic String Compactifications with non-zero fluxes and constant dilaton, [arXiv:0804.1648v](http://arxiv.org/abs/0804.1648)2 [math.DG].
- <span id="page-18-9"></span>[8] A. Fino, M. Parton, S. Salamon, Families of strong KT structures in six dimensions, Comment. Math. Helv. 79 (2004), 317–340.
- <span id="page-18-10"></span><span id="page-18-8"></span>[9] H. Geiges, Symplectic couples on 4-manifolds, Duke Math. J. 85 (1996), 701–711.
- [10] K. Hasegawa, A note on compact solvmanifolds with Kähler structures, Osaka J. Math. 43 (2006), 131-135.
- <span id="page-18-5"></span>[11] A. Hattori, Spectral sequence in the de Rham cohomology of fibre bundles, J. Fac. Sci. Univ. Tokyo 8 (1960), 298–331.
- <span id="page-19-0"></span>[12] N. J. Hitchin, Stable forms and special metrics. In: Fernández, M., Wolf J. (eds.), Global differential geometry: the mathematical legacy of Alfred Gray (Bilbao, 2000), Contemp. Math. 288, Amer. Math. Soc., providence, RI, 2001, 70–89.
- <span id="page-19-1"></span>[13] S. Kobayashi, Principal fibre bundles with the 1-dimensional toroidal group, Tôhoku Math. J. (56) 8 (1956), 29–45.
- <span id="page-19-3"></span>[14] I. Nakamura, Complex parallelisable manifolds and their small deformations, J. Diff. Geom. 10 (1975), 85–112.
- <span id="page-19-2"></span>[15] L. Ugarte, Hermitian structures on six dimensional nilmanifolds, Transform. Groups 12 (2007), 175–202.

M. Fernández: DEPARTAMENTO DE MATEMÁTICAS, FACULTAD DE CIENCIA TECNOLOGÍA, UNIversidad del País Vasco, Apartado 644, 48080 Bilbao, Spain  $E\text{-}mail\;address: \texttt{marisa.fernandez@ehu.es}$ 

A. Tomassini: Dipartimento di Matematica, Universita di Parma, Viale G.P. Usberti ` 53/A, 43100 Parma, Italy

E-mail address: adriano.tomassini@unipr.it

L. Ugarte and R. Villacampa: DEPARTAMENTO DE MATEMÁTICAS - I.U.M.A., UNIVERSIDAD DE Zaragoza, Campus Plaza San Francisco, 50009 Zaragoza, Spain

E-mail address: ugarte@unizar.es, raquelvg@unizar.es