Decoherence-free quantum information in Markovian systems

Manas K. Patra^{1,*} and Peter G. Brooke²

¹Department of Computer Science, University of York, Heslington, York, United Kingdom YO10 5DD

²Centre for Quantum Computer Technology and Department of Physics,

Macquarie University, Sydney, New South Wales 2109, Australia

Decoherence in Markovian systems can result indirectly from the action of a system Hamiltonian which is usually fixed and unavoidable. Here, we show that in general in Markovian systems, because of the system Hamiltonian, quantum information decoheres. We give conditions for the system Hamiltonian that must be satisfied if coherence is to be preserved. Finally, we show how to construct robust subspaces for quantum information processing.

Decoherence remains the most important obstacle to experimental realizations of quantum processors. One well-developed method of counteracting the effects of decoherence is to encode quantum information (QI) into decoherence-free subspaces and subsystems (DFSs) [1, 2, 3, 4, 5, 6, 7]. This form of passive error correction has been well-studied, and has led to (approximate) realizations of DFSs. For example, in ion traps Kielpinski et al. used a decoherence-free (DF) state of two trapped ions to enable encoded information to be stored longer than its unencoded counterpart [8], and in an optical system Kwiat et al. prepared a similar DF state using parametric down-conversion [9]. These proof-of-principle experiments have shown that encoding QI into DFSs improves storage lifetimes, and have partly justified the extensive theoretical investigations into DFSs.

The (strict) requirement for infinite-lifetime (DF) quantum information storage is that all qubits must be symmetrically coupled to the environment [1, 2, 3, 4, 5, 6, 7]. Most theoretical results regarding DFSs rely on this assumption, which is possible to obtain only for qubits that are colocated. As the number of qubits increases this approximation becomes less tenable. Furthermore, it has recently been shown that infinite-lifetime quantum information storage is not possible if the physical qubits of interest are not colocated [10]. So, at best we are left with the regime in which all physical qubits *approximately* experience the same Markovian environment—the qubits are very close together, but not colocated. A full characterization of decoherence in this regime is the purpose of this paper.

So, we examine the hitherto neglected case of decoherence in physical systems for which the exact symmetric coupling between the qubits is perturbatively broken. One physical example of this is given by closely spaced dipole-coupled qubits that approximately satisfy the requirements for Dicke superradiance [11]. For a full analysis of this experimentally accessible regime within the Lindblad master equation, the effects of both the Lamb-shift-type Hamiltonian and the decoherence operators must be included. The unitary evolution generated by the Lamb-shift-type Hamiltonian can cause DFQI to evolve into non-DF states, and so decay via the action of the decoherence operators. Another example is a Heisenberg-type interaction in a spin lattice. We make no assumptions with regards to the Lamb-shifttype Hamiltonian, and find that for Markovian systems stable quantum information is rare. We relax the requirement for infinite-lifetime information storage, and derive expressions for fidelities which depend on the relative strengths of the evolution operators. Finally, we show how to construct robust subspaces for quantum information processing.

For the system density-matrix ρ in the Hilbert space \mathcal{H} , the most general description of Markovian dynamics for initial decoupling between the system and the bath is given by the Lindblad master equation

$$\dot{\rho} = -i[\mathbf{H}, \rho] + L_D(\rho) \tag{1a}$$

$$L_D(\rho) = \frac{1}{2} \sum a_{ij} ([\mathbf{S}_i, \rho \mathbf{S}_j^{\dagger}] + [\mathbf{S}_i \rho, \mathbf{S}_j^{\dagger}]), \qquad (1b)$$

where (a_{ij}) is a time-independent Hermitian coefficient matrix, and L_D is a completely positive map. The presence of the decoherence operators \mathbf{S}_i in Eq. (1b) means that the evolution may not be unitary. But, if $L_D(\rho(t)) = 0$ then $\rho(t) = e^{-i\mathbf{H}t}\rho(0)e^{i\mathbf{H}t}$, so one defines DF dynamics as satisfying $L_D(\rho) = 0$. One can restrict to pure states $\rho = |\psi\rangle \langle \psi|$ [5] to give a sufficient condition for $L_D(\rho) = 0$ as $\mathbf{S}_i |\psi\rangle = c_i |\psi\rangle$. Thus, a DFS is defined as an invariant subspace $M \subset \mathcal{H}$ such that $\mathbf{S}_i |\psi\rangle = c_i |\psi\rangle \ \forall \ \psi \in M$. This condition is guaranteed if M is an eigenspace of \mathbf{S}_i with eigenvalue zero, which in many cases of interest is true for all generators. We note that in two important cases this is the only possibility: i) if the Lie algebra generated by \mathbf{S}_i is semisimple and ii) if all the generators satisfy $\mathbf{S}_{i}^{n} = 0$ for some n (are nilpotent). More precisely, a subspace $A \in \mathcal{H}$ is DF if $\mathbf{S}_i(A) = 0 \,\forall i$. Although, this condition seems somewhat stronger than the usual condition of a *common* invariant eigenspace with eigenvalue c_i (not necessarily zero) of \mathbf{S}_i the analysis below can be generalized by transforming the operators $\mathbf{S}_i \to \mathbf{S}_i - c_i I$. The evolution equation (1) has a unitary and nonunitary part, so even if $\rho(t_0)$ satisfies $L_D(\rho(t_0)) = 0$ it does not ensure that $L_D(\rho(t)) = 0$, for $t > t_0$. This motivates the following definition. A DFS A of \mathcal{H} is to be considered completely-decoherence-free (CDF) if for any state $|\alpha\rangle \in A$, $\rho(0) = |\alpha\rangle \langle \alpha|$ gives $L_D(\rho(t)) = 0$. This condition is weaker than that derived in Ref. [12], but stronger than that derived in Ref. [5].

We illustrate the regime of interest to this paper using the example of dipole-coupled qubits. For these qubits, there exists a regime for which $\mathbf{H} = \sum_{ij} \Delta_{ij} \mathbf{S}_{+}^{i} \mathbf{S}_{-}^{j}$. This occurs when the qubits are closely spaced [11]. It happens that for closely-spaced dipole-coupled qubits the spatially-dependent interatomic spontaneous emission described by the matrix (a_{ij}) no longer depends on index i, j, but the spatially-dependent interatomic coherent dipole-dipole interaction does. This means that there exists exact DF states that are acted upon by \mathbf{H} , causing transitions to non-DF states. This is the case for any separation greater than zero.

We begin with a criteria for CDF dynamics.

Proposition 1 Let V be a DFS in \mathcal{H} : $\mathbf{S}_i \cdot V = 0 \forall i$. A necessary and sufficient condition that V contain a CDFS W is $\mathbf{H} \cdot W \subset W$. In particular, \mathbf{H} can be diagonalized in W.

Proof Define $\rho'(t) = e^{i\mathbf{H}t}\rho(t)e^{-i\mathbf{H}t}$. The equation satisfied by ρ' is $\dot{\rho}' = \mathbf{L}'_D[\rho']$, where $\mathbf{L}'_D[\mathbf{S}'_i] = L_D[e^{i\mathbf{H}t}\mathbf{S}_ie^{-i\mathbf{H}t}]$. Hence, in this picture $\rho'(0)$ is DF iff $\mathbf{L}'_D[\rho'(0)] = 0$. The generic DFSs are spanned by vectors $|x\rangle$ such that $\mathbf{S}'_i |x\rangle = e^{i\mathbf{H}t}\mathbf{S}_ie^{-i\mathbf{H}t} |x\rangle = 0$. Let W be the subspace consisting of all such vectors. This must be satisfied for all t, so we have $\mathbf{H} \cdot W \subset W \forall i$. Conversely, if this is satisfied then $\mathbf{S}_i e^{-i\mathbf{H}t} |x\rangle = 0 \forall i$.

A useful consequence of the proposition is the following.

Corollary A subspace W is CDF iff $(ad(\mathbf{H}))^n(\mathbf{S}_i) \cdot W = 0$, $\forall n$ and *i*. This is equivalent to the condition $[\mathbf{H}^n, \mathbf{S}_i] \cdot W = 0$.

Proof The first condition follows from the proposition and the identity $e^{itad \mathbf{H}}(\mathbf{S}_i) = e^{it\mathbf{H}}\mathbf{S}_i e^{-it\mathbf{H}}$. The second condition is proved by induction [13].

The Pauli matrices $\{\sigma_x, \sigma_y, \sigma_z\}$ generate the Lie algebra su(2) and form a basis (with identity matrix I) for the space of observables for each qubit. Let $\mathbf{S}_a^i =$ $I \otimes \cdots \otimes \sigma_a \otimes I \cdots \otimes I, \ a \in \{x, y, z\}, \text{ with } \sigma_a \text{ only at}$ the i^{th} place. Let $\mathbf{S}^i_{\pm} = \mathbf{S}^i_x \pm i \mathbf{S}^i_y$, $\mathbf{S}_{\pm} = \sum_i \mathbf{S}^i_{\pm}$ and $\mathbf{S}_z = 2 \sum_i \mathbf{S}_z^i$. These operators define a representation of su(2) on the system Hilbert space $\mathcal{H} = \otimes^N \mathbb{C}^2 = \mathbb{C}^{2N}$, and satisfy $[\mathbf{S}_{\pm}, \mathbf{S}_{z}] = -2\mathbf{S}_{z}$ and $[\mathbf{S}_{+}, \mathbf{S}_{-}] = \mathbf{S}_{z}$. We denote a tensor basis for \mathbb{C}^{2N} as $|i_1\rangle \otimes \cdots |i_N\rangle$ where $i_k \in \{0,1\} = |i_1 \cdots i_N\rangle$, and we write $|j_1 \cdots j_k\rangle$ for a vector with ones at $j_1 \cdots j_k$ and zeros elsewhere. The Hamiltonian is written $\mathbf{H} = \sum_{ij} \Delta_{ij} \mathbf{S}^i_+ \mathbf{S}^j_-$, where (Δ_{ij}) is a Hermitian matrix. Without loss of generality, we take Δ , for $\Delta = (\Delta_{ij})$, to be real symmetric. We consider the case where $a_{ij} = a$, so there is only one Lindblad generator S_{-} . In this instance, Eq. (1b) takes the form $L_D(\rho) = \kappa (\mathbf{S}_- \rho \mathbf{S}_+ - \frac{1}{2} (\mathbf{S}_+ \mathbf{S}_- \rho + \rho \mathbf{S}_+ \mathbf{S}_-))$. Each irreducible representation (irrep) of \mathcal{H} is generated by a unique lowest weight vector with weight -(r-1) for irrep

dimension r that satisfies $\mathbf{S}_{-} |\alpha\rangle = 0$. Note that **H** leaves the weight spaces invariant, so we only consider DFSs of fixed weight.

The subspace V_1 is generated by the basis $B \equiv \{|i\rangle =$ $|0\cdots 010\cdots 0\rangle$ | for 1 in the *i*th place}. Here, we let **H** also stand for its restriction to V_1 . Then, $\mathbf{H} |i\rangle =$ $\sum_{k} \Delta_{ki} |k\rangle$. A state $|\mathbf{x}\rangle = \sum_{i} x_{i} |i\rangle$ is a lowest weight state iff $\sum_{i} a_i = 0$. There are N-1 such independent vectors which generate a DFS, called here D_1 . We wish to find out whether there are any CDFS $\neq 0$ inside D_1 . Note that we refer to the CDFS C of D_1 , which denotes the maximal subspace that is the sum of all CDFSs. The condition for CDF dynamics is $\mathbf{H} \cdot C \subset C$. So, for a nonzero subspace C to exist it is necessary and sufficient that **H** have an eigenvector in V_1 . In the fixed basis B, we represent an arbitrary $|\mathbf{x}\rangle = \sum_{i} x_i |i\rangle$ as a column vector $\mathbf{x}^T = (x_1, \cdots, x_N)$ in \mathbb{C}^N . It is clear that $|\mathbf{x}\rangle$ is an eigenvector of **H** iff **x** is an eigenvector of Δ with the same eigenvalue. Hence, there will be a nonzero CDFS iff Δ has an eigenvector **x** such that $Tr(\mathbf{x}) = 0$ where $\operatorname{Tr}(\mathbf{x}) = \sum_{i=1}^{N} x_{i}$. Suppose $\boldsymbol{\Delta}$ has a degenerate eigenvalue c. Then there are at least two independent vectors y and z. If we have Tr(y) = Tr(z) = 0, then there is a nonzero CDFS containing at least $|\mathbf{y}\rangle$ and $|\mathbf{z}\rangle$. Otherwise, suppose $\operatorname{Tr}(\mathbf{z}) = k \neq 0$. Then, $\mathbf{x} \equiv \mathbf{y} - \frac{\operatorname{Tr}(\mathbf{y})}{k} \mathbf{z}$ has trace zero and we have a nontrivial CDFS. This gives:

Proposition 2 A sufficient condition for the existence of nonzero CDFS is that the matrix Δ has a degenerate eigenvalue c. If c is m-fold degenerate, then the dimension of the CDFS $\geq m - 1$.

Although the main result of this paper concerns the rarity of CDFSs, there are two examples of physical systems that satisfy CDF dynamics. First, consider four qubits in a spatially symmetric configuration, e.g., the corners of a square lattice. Then, from symmetry considerations it is clear that the row (or column) sums of the matrix Δ is constant. This automatically guarantees the condition in the proposition. There is at least one CDFS containing $\sum_{i=1}^{4} |i\rangle$. Second, consider two dipole-coupled two-level atoms. In the Dicke limit, the single-excitation antisymmetric state is CDF because for the special case of two atoms, the dipole-dipole interaction does not cause information to leak from the DF state to the non-DF state. Consider now the general case for a single-excitation. Assume all the eigenvalues of Δ are non-degenerate. We seek a condition on Δ that will ensure the existence of an eigenvector in D_1 . Suppose $\mathbf{x} = (x_1, \cdots, x_N)^T$ is such an eigenvector with eigenvalue c. Then the corresponding eigenspace is 1dimensional, and we have the following set of equations $\Delta_{11}x_1 + \dots + \Delta_{1N}x_N = cx_1; \Delta_{N1}x_1 + \dots + \Delta_{NN}x_N = cx_N$; $x_1 + \cdots + x_N = 0$. If we set one of the components, say $x_N = 1$, these equations have a unique solution. We rewrite the first N equations as $(\Delta_{11} - c)x_1 + \cdots +$ $\Delta_{1,N-1} x_{N-1} = -\Delta_{1N}; \ \Delta_{N-1,1} x_1 + \dots + (\Delta_{N-1,N-1} - \Delta_{N-1}) x_1 + \dots + (\Delta_{N-1,N-1}) x_1 + \dots +$

 $c)x_{N-1} = -\Delta_{N-1,N}; \Delta_{N1}x_1 + \dots + \Delta_{NN} - c = 0.$ Let $\Gamma(c)$ denote the $(N-1) \times (N-1)$ matrix such that $\Gamma_{ij} = \Delta_{ij} - c\delta_{ij}, \ 1 \le i, j \le N - 1$ (for δ_{ij} the Kronecker δ). The uniqueness of $\mathbf{x}' \equiv (x_1, \dots, x_{N-1})^T$ implies that Γ is invertible. Writing $\mathbf{d} = -(\Delta_{1N}, \ldots, \Delta_{N-1,N})^T$ we have $\mathbf{x}' = \Gamma^{-1}(c)\mathbf{d}$. The last equation $\Delta_{N1}x_1 + \cdots +$ $\Delta_{NN} - c = 0$ can be written as $\mathbf{d}^T \mathbf{x}' - c + \Delta_{NN}$. Hence, we have $\mathbf{d}^T \Gamma^{-1}(c) \mathbf{d} - c + \Delta_{NN} = 0$. That is, $f(c) = \mathbf{d}^T \operatorname{adj}(\Gamma(c))\mathbf{d} - \det(\Gamma(c))(c - \Delta_{NN}) = 0$, where $\operatorname{adj}(\Gamma)$ is such that $\operatorname{adj}(\Gamma)\Gamma = \det(\Gamma)I$. From the condition $\operatorname{Tr}(\mathbf{x}) = 0$ and $x_N = 1$ we obtain $\operatorname{Tr}(\Gamma^{-1}(c)\mathbf{d}) = -1$. Hence, $g(c) = \operatorname{Tr}(\operatorname{adj}(\Gamma(c))\mathbf{d}) + \det(\Gamma(c)) = 0$. So, we get two polynomial equations in c whose coefficients are functions of Δ_{ij} . For a solution to exist the resultant of the two polynomials must vanish [14], and we get a polynomial relation $R(\Delta_{ij})$ among the Δ_{ij} which does not vanish identically. There are N(N+1)/2 = M independent parameters characterizing any real symmetric matrix. Hence, the space of all such matrices may be identified with \mathbb{R}^M . We have just seen that for Δ to have a nontrivial CDFS, it must satisfy (at least) one polynomial equation. So, there are Hamiltonians with matrix Δ which do not have any CDFS in V_1 . We will see that this is the norm rather that the exception—Hamiltonians with CDFS are rare.

We illustrate the above analysis with an example. The first excited subspace V_1 is spanned by $\{|1\rangle = |001\rangle, |2\rangle =$ $|010\rangle, |3\rangle = |100\rangle$. Let the matrix Δ corresponding to **H** restricted to V_1 be given by $\Delta_{12} = \Delta_{21} = x_3, \Delta_{13} =$ $\Delta_{31} = x_2$ and $\Delta_{23} = \Delta_{32} = x_1$ with diagonal elements zero. The CDFS condition translates to $x_3 = x_1$ or x_2 corresponding to eigenvalues $-x_2$ and $-x_1$ respectively. The third possibility $x_1 = x_2$ gives eigenvalue $-x_3$. Fixing $\underline{x_3} = \underline{x_1} = a$ the eigenvalues of Δ are 1 and $x_2 \pm \sqrt{x_2^2 + 8a^2}$. The only possibility for a degenerate eigenvalue is when $x_1 = x_2 = x_3$. We conclude that **H** restricted to the first excited subspace V_1 will have an eigenvector in D_1 (DF subspace) iff at least two of its off-diagonal entries are equal, and it will have two eigenvectors in D_1 if all three are equal. The parameter space of **H** can be identified with \mathbb{R}^3 as x_1, x_2 and x_3 the range over the real numbers. Then, the only Hamiltonians with eigenvectors in D_1 are characterized by the parameters that lie in the planes $x_1 = x_2, x_1 = x_3$ or $x_2 = x_3$. So, we conclude that the Hamiltonians which leave some DFS state in V_1 invariant is a negligible fraction of all the possible Hamiltonians—in general, quantum information will decay.

We know that the DFS in the m^{th} excited subspace V_m is spanned by vectors $|\mathbf{x}_m\rangle = \sum x_{i_1,\dots,i_m} |i_1,\dots,i_m\rangle$ such that $M' = \binom{N}{m-1}$ equations $\sum_{i_r=1}^N x_{i_1,\dots,i_r,\dots,i_m} = 0$ (*)—with all indices except i_r fixed—are satisfied. The

action of \mathbf{H} on V_m is more complicated:

$$\mathbf{H} |\mathbf{x}_{m}\rangle = \sum_{i_{1},\dots,i_{m}} x_{i_{1},\dots,i_{m}} (\sum_{k} \Delta_{ki_{1}} |ki_{2}i_{3}\cdots i_{m}\rangle + \cdots + \sum_{k} \Delta_{ki_{m}} |i_{1}i_{2}i_{3}\cdots k\rangle). \quad (2)$$

We require that $|\mathbf{x}_m\rangle$ satisfy $\mathbf{H} |\mathbf{x}_m\rangle = \lambda |\mathbf{x}_m\rangle$. We write the matrix representing **H** restricted to the subspace V_m as $\Delta^{(m)}$. The action of **H** on V_m is equivalent to that of $\Delta^{(m)}$ on \mathbb{R}^M for $M = \binom{N}{m}$ whose coordinates are given by $x_{i_1,\ldots,i_r,\ldots,i_m}$. We write the eigenvalue equations as in the previous section. Since the eigenvector must satisfy M' equations (*) we use them to write the last M' components of such a vector in terms of the first M - M', and substitute in the eigenvalue equation of $\Delta^{(m)}$. The resulting system of equations in M - M' variables must have rank less than M - M' for a nontrivial solution to exist. Let $\Delta'^{(m)}$ be the square matrix of the coefficients of the first M - M' equations. It must have determinant zero. This gives a polynomial equation in λ , the eigenvalue. Write $f(\lambda) = \det(\Delta'^{(m)}(\lambda))$, which shows the explicit dependence on the eigenvalue λ . We also have $q(\lambda) = \det(\Delta^{(m)}(\lambda)) = 0$, the original characteristic equation. The coefficients in $f(\lambda)$ and $g(\lambda)$ are functions of the variables Δ_{ij} . The necessary and sufficient condition that \mathbf{H} has an eigenvector in a DFS is that $f(\lambda)$ and $q(\lambda)$ have a common root: that is, the resultant $R(\Delta) = 0$. It can be shown that the resultant does not vanish identically. All possible Hamiltonians, parametrized by the real numbers Δ_{ii} , constitute a manifold of dimension N(N+1)/2. The Hamiltonians which have an eigenvector in a DFS in V_m lie on a submanifold of dimension strictly less than N(N+1)/2. Hence, using Sard's theorem [15] and generalizing to Hermitian matrices we arrive at the following theorem.

Theorem 1 Let X be the N^2 -dimensional real manifold of the parameters characterizing the possible Hamiltonians in the Lindblad master equation. Let S be the set of values of the parameters which characterize Hamiltonians that have at least one DF state other than the ground state evolving into a DFS state at all times. Then S is of measure zero in X.

Proof Let S_m be those members S that correspond to Hamiltonians that have some DF states in a fixed weight space V_m , m > 0 evolving into DF states at all times. From Proposition 1, we see that this is equivalent to the condition that the Hamiltonians in S_m have an eigenvector in the DFS in V_m . From the preceding discussion, S_m has measure zero in X. Since $S = \bigcup_m S_m$, S too has measure zero.

Informally, we can say that for almost all Hamiltonians there is no DF state other than the ground state which evolves into other DF states at all times. Therefore, the best we can hope for is to seek states which remain DF up to some orders of perturbation.

Corollary to Proposition 1 implies that to get DF states we should look for states that are annihilated by operators $\mathbf{S}_{-}, [\mathbf{H}, \mathbf{S}_{-}], [\mathbf{H}[\mathbf{H}, \mathbf{S}_{-}]], \dots$ We have seen that in general DFSs are not invariant under **H**. So, we seek invariance up to certain orders. The condition for states $|\alpha\rangle$ that are invariant up to first order (in **H**) is that $\mathbf{S}_{-} |\alpha\rangle = [\mathbf{H}, \mathbf{S}_{-}] |\alpha\rangle = 0$. Similarly, the second order condition is $\mathbf{S}_{-} |\alpha\rangle = [\mathbf{H}, \mathbf{S}_{-}] |\alpha\rangle =$ $[\mathbf{H}, [\mathbf{H}, \mathbf{S}_{-}]] |\alpha\rangle = 0.$ We assume that the initial state $\rho(0) = |\psi\rangle \langle \psi|$ is a DF state. Then $\rho(t) =$ $e^{-it(\operatorname{ad} \mathbf{H} + iL_D)}\rho(0)$ and if the evolution is unitary, $\rho'(t) =$ $e^{-it\mathbf{H}} |\psi\rangle \langle\psi| e^{it\mathbf{H}}$. Hence, we take $F^2(\rho'(t), \rho(t)) = \langle\psi| e^{it\mathbf{H}} e^{-it(\operatorname{ad}\mathbf{H}+iL_D)}\rho(0)e^{-it\mathbf{H}} |\psi\rangle$ as a measure of deviation from unitary evolution [16]. Writing $\operatorname{ad} \mathbf{H} =$ X, we have $F^2 = \langle \psi | e^{itX} e^{-it(X+iL_D)} | \psi \rangle$. Using the Zassenhaus formula [17], the (exact) fidelity can be written $F^2 = \langle \psi | e^{L_D} e^{-it^2 [X, L_D]/2} e^{-t^3 Z_3(X, L_D)} \cdots \rho(0) | \psi \rangle.$ where $Z_3(X, L_D) = [X, [X, L_D]]/6 + i[L_D, [L_D, X]]/3)$ We consider the following three approximate cases.

Case 1 (weak unitary part). The Hamiltonian **H** is replaced by ϵ **H**, and treated as a perturbation. Then up to first order in ϵ , $F^2 = \langle \psi | e^{tL_D} e^{-it^2 [L_D, X]/2} \cdots e^{c_k t^k (\operatorname{ad} L_D)^{k-1}(X)} \rho(0) | \psi \rangle$ where $c_k = (-1)^{k-1} \epsilon(k-1)i/k!, k > 1$. The state $\rho(0) = |\psi\rangle \langle \psi |$ is DF, so $F^2 = \langle \psi | (1 + \sum c_k t^k (\operatorname{ad} L_D)^{k-1}(X))\rho(0) | \psi \rangle$. For DF-states $\langle \psi | (\operatorname{ad} L_D)^{k-1}(X)\rho(0) | \psi \rangle = 0$ and $F^2 = 1$ [5, 18].

Case 2 (strong unitary part). The Lindblad part is treated as a perturbation to give $\dot{\rho} =$ $(-iad \mathbf{H} + \epsilon L_D)\rho$. Then up to first order in ϵ , $F^2 = \langle \psi | e^{tL_D} e^{it^2[X,L_D]/2} \cdots e^{d_k t^k (ad X)^{k-1}(L)} \rho(0) | \psi \rangle$ where $d_k = (-i)^k \epsilon / k!$, k > 1. Computing the series expansions to first order, we get $F^2 = \langle \psi | (1 + \sum d_k t^k (ad X)^{k-1}(L_D))\rho(0) | \psi \rangle$. Unlike the previous case $L_D(\rho(0)) = 0$ does not guarantee $F^2 = 1$ for all times. There are two possible courses of action. First, if the time scales of **H** are much shorter than L_D , zeroth order may suffice. Thus, instead of talking the initial state $\rho(0)$ as an eigenstate of Lindblad generators, we use an eigenstate of **H**. This method is used in Ref. [19].

Case 3 (short-time expansions). Note that the Zassenhaus formula shows that DFSs are stable up to first order in t. If we require stability up to order t^k , $k \ge 1$, we can work in the smaller subspaces W satisfying ad $\mathbf{H}^m(\mathbf{S}_i) \cdot W = 0$, $m = 0, \ldots, k - 1 \forall i$. Since we are dealing with finite-dimensional Hilbert spaces, it suffices to require stability up to order n, the dimension of the system space. However, we have seen that this is not possible in general (Theorem 1), so we have to be satisfied with smaller k.

We consider the universality of subspaces for external Hamiltonians that implement quantum gates. Let \mathbf{H}_c denote the external Hamiltonian and $\mathbf{H}_d = \sum_{ij} \Delta_{ij} \mathbf{S}_i^i \mathbf{S}_-^j$. The equation of motion is $\dot{\rho} = (-i(\operatorname{ad} \mathbf{H}_c + \operatorname{ad} \mathbf{H}_d) + L_d)\rho \equiv (Y + X + L_D)\rho$. Since we aim to work in DFSs, they must be invariant under any external Hamiltonian. Thus, $[\mathbf{H}_c, \mathbf{H}_i] = 0$, and $[Y, L_D] = 0$. Assuming that the characteristic time-scale of \mathbf{H}_c is much smaller than other operators (essential for feasible computation), then using the short time expansion we can construct robust subspaces for computation. For example, if we wish to terminate at third order, then a subspace W satisfying $L_D \cdot W = [\mathbf{H}_d, L_d] \cdot W = [\mathbf{H}_d, [\mathbf{H}_d, L_D]] \cdot W = 0$ will suffice.

We have shown that in Markovian systems the Lambshift-type system Hamiltonian generally causes transitions from DF states to non-DF states. The results presented here emphasize the importance of accounting for *both* unitary and nonunitary evolution in passive quantum error correction. Note that the results presented here, as long as the Lindblad operators do not cause transitions between irreps, can be extended to the finite temperature case.

- * Electronic address: manas@cs.york.ac.uk
- G. M. Palma, K.-A. Suominen, and A. K. Ekert, Proc. Roy. Soc. London Ser. A 452, 567 (1996).
- [2] L. M. Duan and G. C. Guo, Phys. Rev. Lett. 79, 1953 (1997).
- [3] L. M. Duan and G. C. Guo, Phys. Rev. A 57, 737 (1998).
- [4] P. Zanardi and M. Rasetti, Phys. Rev. Lett. 79, 3306 (1997).
- [5] D. A. Lidar, I. L. Chuang, and K. B. Whaley, Phys. Rev. Lett. 81, 2594 (1998).
- [6] E. Knill, R. Laflamme, and L. Viola, Phys. Rev. Lett. 84, 2525 (2000).
- [7] D. A. Lidar and K. B. Whaley, in *Irreversible Quantum Dynamics*, edited by F. Benatti and R. Floreanini (Springer-Verlag, Berlin, 2003), vol. 622, pp. 83–120.
- [8] D. Kielpinski, V. Meyer, M. A. Rowe, C. A. Sackett, W. M. Itano, C. Munroe, and D. J. Wineland, Science 291, 1013 (2001).
- [9] P. G. Kwiat, A. J. Berglund, J. B. Altepeter, and A. G. White, Science **290**, 498 (2000).
- [10] R. Karasik, K.-P. Marzlin, B. C. Sanders, and B. K. Whaley, Phys. Rev. A 76, 012331 (2007).
- [11] P. G. Brooke, K.-P. Marzlin, J. D. Cresser, and B. C. Sanders, Phys. Rev. A 77, 033844 (2008).
- [12] A. Shabani and D. A. Lidar, Phys. Rev. A 72, 042303 (2005).
- [13] J. E. Humphreys, Introduction to Lie Algebras and Representation Theory (Springer-Verlag, Berlin, 1972).
- [14] B. L. van der Waerden, Modern Algebra (Ungar, New York, 1953).
- [15] J. W. Milnor, Topology from the Differentiable Viewpoint (Princeton University Press, Princeton, 1997).
- [16] R. Jozsa, J. Mod. Opt **41**, 2315 (1994).
- [17] R. M. Wilcox, J. Math. Phys. 8, 962 (1967).
- [18] D. Bacon, D. A. Lidar, and K. B. Whaley, Phys. Rev. A 60, 1944 (1999).
- [19] P. G. Brooke, J. D. Cresser, and M. K. Patra, Phys. Rev. A 77, 062313 (2008).