

# Electroweak Physics with ATLAS

Arif Akhundov\*

*Institute of Physics, Azerbaijan Academy of Sciences, 370143 Baku, AZERBAIJAN*

The precision measurements of electroweak parameters of the Standard Model with the ATLAS detector at LHC are reviewed. An emphasis is put on the bridge connecting the ATLAS measurements with the SM analysis at LEP/SLC and the Tevatron.

## 1. INTRODUCTION

The Standard Model (SM) has been impressively confirmed by successful collider experiments at the particle accelerators LEP, SLC and Tevatron [1, 2, 3] during the last twenty years.

In July 2008 the Large Hadron Collider (LHC) at CERN started-up to operate and will reach a total energy of 10 TeV with colliding proton beams by October 2008 [4]. The general purpose of the ATLAS experiment [5] is designed to meet the main physics goals of the LHC: the discovery of the Higgs boson and supersymmetric particles [6, 7, 8].

In addition to its discovery potential the LHC constitutes a powerful tool to make precision tests of the SM [9, 10, 11]. During the initial two years of low luminosity running from  $5 \times 10^{31} \text{ cm}^{-2}\text{s}^{-1}$  to  $10^{33} \text{ cm}^{-2}\text{s}^{-1}$  the integrated luminosity in 2008 will be  $20 \text{ pb}^{-1}$  and in 2009  $100 \text{ pb}^{-1}$  per LHC experiment. This will be an ideal period to continue precision measurements started at LEP/SLC and the Tevatron [12] thanks to the very high rate of the SM processes.

## 2. PRECISION FOR THE MEASUREMENTS OF THE W AND TOP QUARK MASSES

The masses  $M_W$  and  $m_t$  are the physical input parameters of the SM and we should know them with high precision. The results of the measurements at LEP and the Tevatron are shown in Figure 1.




Figure 1: Comparison of the direct measurements of  $M_W$  and  $m_t$  at LEP2 and the Tevatron with the indirect determination through electroweak radiative corrections at LEP1 and SLD [1, 2, 3].

\*Electronic address: Arif.Akhundov@cern.ch

The higher order terms, radiative corrections or quantum corrections, contain the self-coupling of the vector bosons as well their interactions with the Higgs field and the top quark. Their calculation provides the theoretical basis for the electroweak precision tests [12]. Theoretical prediction of the SM on the quantum level depends on the masses  $M_W$  and  $m_t$  and on the as yet experimentally unknown Higgs boson  $M_H$  through the virtual presence of these particles in the loops. The electroweak radiative corrections receive contributions from the square of the top mass [13] and the logarithm of the Higgs mass [14]. As a consequence, precision data from LEP, SLC and Tevatron [1, 2, 3] pin down the allowed range of the mass  $M_W$  and  $m_t$ . The Figure 2 (left) compares the information on  $M_W$  and  $m_t$  obtained at LEP1 and SLD with the direct measurements performed at LEP2 and the Tevatron.

The principal method for measuring the W mass at the LHC employs the leptonic decay channels. The cross section for  $pp \rightarrow W + X$  with  $W \rightarrow l\nu$  ( $l = e, \mu$ ) is 15 nb, corresponding to  $\sim 10^6$  events for the luminosity  $10 \text{ fb}^{-1}$  [15]. The W mass is extracted from the distribution of the W transverse mass,  $M_T^W$ , given by

$$M_T^W = \sqrt{2p_T^l p_T^\nu (1 - \cos \Delta\Phi)}, \quad (1)$$

where  $\Phi$  is the azimuthal angle between the 3-momentums of charged lepton and the system X.

The large sample of the W events available at the LHC means that the uncertainty on the W mass,  $\Delta M_W$ , arising from the statistics will be very small. The largest experimental uncertainty and the limiting factor governing the precision on the W mass is the knowledge of the 4-momentum of charged lepton which will drive a total uncertainty  $\Delta M_W$  lower than 20 MeV [9]. At LHC  $\sigma(pp \rightarrow t\bar{t}) \sim 830 \pm 100 \text{ pb}$  which is 100 times more than at the Tevatron and




Figure 2: Left: The SM relationship for the masses  $M_W$  and  $m_t$  as function of  $M_H$  [1, 2, 3]. Right: Reconstructed top mass and combinatorial background (shaded area) from semileptonic  $t\bar{t}$  events ( $10 \text{ fb}^{-1}$ ) [16].

a rate of  $\sim 10^4$   $t\bar{t}$  events for the luminosity  $10 \text{ fb}^{-1}$ , 90% of which are produced through  $gg \rightarrow t\bar{t}$ . The  $t\bar{t}$  events are categorized by the subsequent decay of the two W bosons into three types: full hadronic(65%), semileptonic (30%) and dileptonic (5%) events. The semileptonic channel represents the most favorable route to measuring the top quark mass [16]. The top mass measurement proceeds by reconstructing the  $jjb$ -system from the two jets which are not b-tagged and combining with one of the b-tagged jets. The invariant mass  $M_{jjb}$  distribution for the luminosity  $10 \text{ fb}^{-1}$  is plotted in Figure 2 (right) together with the combinatorial background [16]. The top quark mass is extracted from this plot by fitting to the peak using Monte Carlo samples generated with different values of  $m_t$ .

The large sample of  $t\bar{t}$  events available at the LHC means that the uncertainty on the top quark mass,  $\Delta m_t$ , arising from the statistics also will be very small. The largest uncertainty will arise from the contribution of the final state radiation of massive particles [17]. Nevertheless, ATLAS could attain a precision of  $\sim 1 \text{ GeV}$  on the top quark mass in two years of low luminosity running.

### 3. CONCLUSIONS

The precision measurements of the W and top quark masses with the ATLAS detector at LHC are reviewed. The anticipated uncertainty in the ATLAS measurements  $\Delta M_W \sim 15$  MeV and  $\Delta m_t \sim 1$  GeV would constrain the mass of the Higgs boson to within 40%.

### Acknowledgments

I would like to thank the organizers of the ICHEP08 for kind invitation to present this poster. I am grateful to A. Di Ciaccio for support of this material. I thank P. Pralavorio for important comments and A. Shiekh for remarks.

### References

- [1] The LEP Collaborations ALEPH, DELPHI, L3 and OPAL and the LEP Electroweak Working Group, arXiv:hep-ex/0612034; <http://www.cern.ch/LEPEWWG/>.
- [2] The ALEPH, DELPHI, L3, OPAL and SLD Collaborations, the LEP Electroweak Working Group and the SLD Electroweak and Heavy Flavour Groups, *Phys. Rept.* **427**, 257 (2006).
- [3] CDF and D0 Collaborations, arXiv:0803.1683 [hep-ex].
- [4] Juergen Thomas on behalf of the ATLAS Collaboration, Hadron Collider Physics Symposium (HCP2008), Galena, Illinois, USA, 2008; arXiv:0808.1031 [hep-ex].
- [5] ATLAS Collaboration, ATLAS Detector and Physics Performance Technical Design Report, *CERN/LHCC 99-14/15*, (1999).
- [6] ATLAS Collaboration, Anna Di Ciaccio *et al.*, Talk at 5th Workshop on Science with the New Generation High Energy Gamma-ray Experiments (SciNe-GHE07): The Light of the Dark: Solving the Mysteries of the Universe, Frascati, Rome, Italy, 18-20 Jun 2007; in arXiv:0712.1548 [astro-ph], pp. 111-120.
- [7] John Ellis, Plenary talk at UAE-CERN Workshop on High Energy Physics and Applications, Al-Ain, UAE, 26-28 Nov. 2007; *AIP Conf. Proc.* **1006**, 3–11 (2008).
- [8] Robert Harlander, *J. Phys. G: Nucl. Phys.* **35**, 033001 (2008).
- [9] ATLAS Collaboration, P. Pralavorio *et al.*, Talk at International Europhysics Conference on High Energy Physics (HEP-EPS 2005), Lisbon, Portugal, 21-27 Jul 2005; *PoS HEP2005* **294**, (2006).
- [10] Jens Erler, Talk at 3rd International Workshop on From parity Violation to Hadronic Structure and More (PAVI06), Milos, Greece, 16-20 May 2006; arXiv: hep-ph/0607323.
- [11] Jens Erler, Talk at 12th Mexican School on Particles and Fields and 6th Latin American Symposium on High Energy Physics (VI-Silafae/XII-MSPF), Puerto Vallarta, Mexico, 1-8 Nov 2006; *AIP Conf. Proc.* **917**, 244-251 (2007).
- [12] Arif Akhundov, Plenary talk at UAE-CERN Workshop on High Energy Physics and Applications, Al-Ain, UAE, 26-28 Nov. 2007; *AIP Conf. Proc.* **1006**, 43-48 (2008).
- [13] A. A. Akhundov, D. Yu. Bardin and T. Riemann, *Nucl. Phys.* **B276**, 1 (1986).
- [14] A. A. Akhundov, D. Yu. Bardin and T. Riemann, *Phys. Lett.* **B166**, 111 (1986).
- [15] ATLAS Collaboration, M. Wielers *et al.*, Talk at 33rd International Conference on High Energy Physics (ICHEP 06), Moscow, Russia, 26 Jul - 2 Aug 2006; In Moscow 2006, ICHEP. pp. 669-672.
- [16] I. Borjanovic *et al.*, *Eur. Phys. J.* **C39S2**, 63 (2005).
- [17] A. A. Akhundov, D. Yu. Bardin, O. M. Fedorenko and T. Riemann, *Sov. J. Nucl. Phys.* **42**, 762 (1985).