

First Cosmological Constraints on Dark Energy from the Radial Baryon Acoustic Scale

Enrique Gaztañaga*

*Institut de Ciències de l'Espai (IEEC-CSIC), Facultat de Ciències,
Torre C5, Campus UAB, E-08193 Bellaterra (Barcelona), Spain*

Ramon Miquel†

*Institució Catalana de Recerca i Estudis Avançats,
Institut de Física d'Altes Energies, Edifici Cn, Facultat de Ciències,
Campus UAB, E-08193 Bellaterra (Barcelona), Spain*

Eusebio Sánchez‡

*Centro de Investigaciones Energéticas, Medioambientales y Tecnológicas,
Avenida Complutense 22, E-28040 Madrid, Spain*

(Dated: August 4, 2021)

We present cosmological constraints arising from the first measurement of the radial (line-of-sight) baryon acoustic oscillations (BAO) scale in the large scale structure traced by the galaxy distribution. Here we use these radial BAO measurements at $z = 0.24$ and $z = 0.43$ to derive new constraints on dark energy and its equation of state for a flat universe, without any other assumptions on the cosmological model: $w = -1.14 \pm 0.39$ (assumed constant), $\Omega_m = 0.24_{-0.05}^{+0.06}$. If we drop the assumption of flatness and include previous cosmic microwave background and supernova data, we find $w = -0.974 \pm 0.058$, $\Omega_m = 0.271 \pm 0.015$, and $\Omega_k = -0.002 \pm 0.006$, in good agreement with a flat cold dark matter cosmology with a cosmological constant. To our knowledge, these are the most stringent constraints on these parameters to date under our stated assumptions.

PACS numbers: 95.36.+x, 98.65.Dx, 98.80.Es

Before recombination, the Universe was filled by a plasma of coupled photons and baryons. On horizon crossing, cosmological fluctuations produced sound waves in this plasma and when recombination occurred, 380 000 years after the Big Bang, the distance covered by the sound wave was about 150 comoving Mpc. This is the so called sound horizon r_s , also known as the BAO scale r_{BAO} . This signature can be found both in the cosmic microwave background (CMB) and large scale structure (LSS), so the baryon acoustic peak can be used as a standard ruler in the Universe. The value of r_{BAO} depends on a few physical quantities, mostly the time to recombination and the time of matter domination, both of which are well known from measurements of CMB temperature anisotropies, i.e. through parameters $\Omega_m h^2$ and $\Omega_B h^2$, also in good agreement with large scale structure and primordial abundance measurements. Thus the spectrum of CMB fluctuations provide an accurate estimate of $r_{BAO} = 153 \pm 1.9$ Mpc, independent of the value of H_0 , dark energy equation of state w or the curvature of the Universe [1]. At the same time the CMB also provides an angular measurement of the BAO scale, which together with r_{BAO} , can be used for geometrical test, such as the measurement of curvature.

A series of recent papers [2, 3, 4, 5] have presented the

clustering of Luminous Red Galaxies (LRG, see [6]) in the latest spectroscopic Sloan Digital Sky Survey (SDSS) data releases, DR6 & DR7, which include over 75 000 LRG galaxies and sample over $1 \text{ Gpc}^3/h^3$ to $z=0.47$. The last paper [5] focuses on the study of the 2-point correlation function $\xi(\sigma, \pi)$, separated in perpendicular σ and line-of-sight π directions to find a significant detection of a peak at $r \simeq 110 \text{ Mpc}/h$ (for $H_0 = 100 \text{ h Km/s}$), which shows as a circular ring in the σ - π plane. There is also a significant detection of the peak along the line-of-sight (radial) direction both in sub-samples at low, $z=0.15$ - 0.30 , and high redshifts, $z = 0.40 - 0.47$. The overall shape and location of the peak in the 2-point and 3-point function are consistent with its originating from the recombination-epoch baryon acoustic oscillations. This has been used to produce, for the first time, a direct measurement of the Hubble parameter $H(z)$ as a function of redshift. This is based on a calibration to the BAO scale measured in the CMB [1]. The values of $H(z)$ are then used in [5] to compare to other cosmological constraints that provide estimates for H_0 .

Here, instead of calibrating the BAO distance, we will use the direct dimensionless measurement as a redshift scale (shown in Table I) to provide cosmological constraints that are independent of H_0 . This is somehow similar to what is done in Percival et al. [7] for the spherically-averaged (i.e. monopole) BAO detection. Both the 2-Degree-Field Galaxy Redshift Survey 2dFGRS and SDSS spectroscopic redshift surveys have been used to constrain cosmological parameters at BAO dis-

*Electronic address: gazta@ice.cat

†Electronic address: rmiquel@ifae.es

‡Electronic address: Eusebio.Sanchez@ciemat.es

tances [8, 9, 10, 11, 12]. Photometric LRG catalogs have also been used to obtain cosmological constraints [13, 14].

None of these earlier papers have constrained the baryon feature in the line-of-sight (LOS) direction. We will argue here that the radial BAO (hereafter RBAO) detection provides in practice fully independent constraints from the monopole (or angular) BAO measurements. It is therefore important to check if such independent constraints agree with the flat concordance cosmological constant model (Λ CDM) and, if so, what are the new combined constraints when we add the new data to previous results. Also note that in previous BAO detections the constraints come from either a fit to the overall shape of the correlation function or to the normalized wiggles in the power spectrum $P(k)$, while in the LOS measurement of r_{BAO} in [5] they only used the BAO peak location (identified by the position of maximum signal to noise around 90 – 120 Mpc/h) and not the shape. There are potential advantages and disadvantages of both methods. Using a fit to the overall shape is a more stringent test of the model, but puts more weight on the intermediate scales, where the statistical errors are smaller but where the modeling is more uncertain. Fitting the peak location is more model independent but, precisely because of this, provides little evidence for the physical origin of such peak. In the case of the analysis in [5], the physical evidence comes from the agreement between model and data both on the radial direction (see Figs. 13–15 in [5]) and on the overall $\xi(\pi, \sigma)$ plane, which shows a clear BAO ring (with the right monopole contribution) that blends into the extended LOS peak (see Fig. 10 in [5]). A model with no radial BAO is disfavored at 3.2σ .

To work with fiducial comoving scales, [5] uses a reference flat Λ CDM Hubble rate:

$$H_{ref}(z) = H_0 \sqrt{\Omega_m(1+z)^3 + 1 - \Omega_m} \quad (1)$$

with $\Omega_m = 0.25$ to determine fiducial values for r_{BAO} at two redshifts. Here, $\Delta z_{BAO}(z)$ are obtained from $r_{BAO}(z)$ in [5] as $\Delta z_{BAO}(z) = r_{BAO}(z)H_{ref}(z)/c$. Note that the measurements of $\Delta z_{BAO}(z)$ do not depend on H_0 , since the dependences contained in $r_{BAO}(z)$ and $H_{ref}(z)/c$ cancel out.

Previous BAO (e.g. [7, 8]) analyses have only looked at the monopole, which is the average of $\xi(\sigma, \pi)$ over orientations:

$$\xi_0(r) = \int_0^1 \xi(\sigma, \pi) d\mu \quad (2)$$

where $r = \sqrt{\sigma^2 + \pi^2}$ and $\mu = \pi/r$. The BAO peak found by [5] was located at radial $\pi \simeq 110$ Mpc/h for $\sigma < 5$ Mpc/h, which corresponds to $\mu > 0.999$. The contribution of radial modes with $\mu > 0.999$ to the monopole at $r = 110$ in the above integral is less than 1% even if we take the amplitude of $\xi(\sigma, \pi)$ to be 10 times larger in the radial π direction than in the transverse σ direction. Moreover, the covariance between radial and monopole

correlations has been shown to be less than 10% on scales larger than 40 Mpc/h (see Fig. 2 of [15]). Thus, in practice, we can regard here the BAO measurements from the monopole as independent from the RBAO measurements.

Once the radial BAO scale has been measured [5], there are several possible approaches to extract the cosmological parameters solely from this measurement. Percival *et al.* [7] propose three possible ways. We have used two of them, that are described below. The third one uses the ratio of BAO scales at different redshifts. We have not used it, since we have the BAO scale at two different redshifts, so only one ratio can be constructed, and the degeneracy in the determination of cosmological parameters is larger using this approach.

Sample z range	z_m	r_{BAO} Mpc/h	σ_{st}	σ_{sys}	Δz_{BAO}	σ_{st}	σ_{sys}
0.15-0.30	0.24	110.3	2.9	1.8	0.0407	0.0011	0.0007
0.40-0.47	0.43	108.9	3.9	2.1	0.0442	0.0015	0.0009

TABLE I: The BAO fiducial scale r_{BAO} in the LOS direction calculated with a flat reference $H_{ref}(z)$ cosmology of $\Omega_m = 0.25$, for two redshift slices: z_m is the respective pair-weighted mean redshift, and σ_{st} and σ_{sys} are the statistical and systematic errors on r_{BAO} (from [5]). Here we use the direct Δz_{BAO} measurement, shown in the sixth column, which relates to the fiducial scale as $\Delta z_{BAO} = r_{BAO}H_{ref}(z)/c$ and is independent of the value chosen for $H_{ref}(z)$. The last two columns show the corresponding errors.

We will derive cosmological constraints from the measured values of Δz_{BAO} shown in Table I, which can be expressed as

$$\Delta z_{BAO}(z_i) = \frac{H(z_i)r_s}{c}, \quad (3)$$

where r_s is the sound horizon at recombination. It is important to remark that the two measurements of Δz_{BAO} are independent, due to the analysis strategy used in [5], which determines the BAO scale in two well separate intervals of redshift.

Throughout this letter, we will assume a standard Friedmann-Lemaître-Robertson-Walker cosmology having as free parameters the matter density Ω_m , curvature Ω_k , Hubble constant h (with $H_0 = 100 h$ km/s/Mpc), and the parameter w of the equation of state of dark energy, i.e. the ratio of its pressure to its density $w = p_{DE}/\rho_{DE}$. Unless otherwise specified, a constant w is assumed, and thus, the standard Λ CDM model is recovered for $w = -1$. We will start by assuming also a flat Universe, $\Omega_k = 0$.

In order to constrain the dark energy parameters Ω_Λ ($\equiv 1 - \Omega_m$) and w , we can take two approaches:

a) We can express r_s in (3) as [21]

$$r_s = \frac{c}{(3\Omega_m H_0^2)^{1/2}} \int_0^{\frac{1}{1+z_d}} \frac{da}{(a + a_{eq})^{1/2} (1 + a \frac{3\Omega_b h^2}{4\Omega_\gamma h^2})^{1/2}} \quad (4)$$

Data Set	Ω_m	$\sigma(\Omega_m)$	Ω_k	$\sigma(\Omega_k)$	w	$\sigma(w)$
RBAO a)	0.242	$^{+0.061}_{-0.053}$	0	—	-1.14	$^{+0.38}_{-0.40}$
RBAO b)	0.248	± 0.053	0	—	-1.16	$^{+0.42}_{-0.45}$
RBAO + WMAP5 [1]	0.274	$^{+0.035}_{-0.036}$	0	—	-0.92	$^{+0.16}_{-0.22}$
RBAO + WMAP5 + SNe [16]	0.268	$^{+0.015}_{-0.014}$	0	—	-0.961	$^{+0.056}_{-0.058}$
RBAO a)	0.249	$^{+0.049}_{-0.036}$	-0.08	$^{+0.25}_{-0.18}$	-1	—
RBAO b)	0.223	± 0.099	-0.028	± 0.058	-1	—
RBAO + WMAP5 [1]	0.264	$^{+0.017}_{-0.016}$	-0.0025	± 0.0059	-1	—
RBAO + WMAP5 + SNe [16]	0.271	$^{+0.015}_{-0.014}$	-0.0026	± 0.0060	-1	—
RBAO + WMAP5 [1]	0.244	$^{+0.062}_{-0.052}$	-0.0049	$^{+0.0121}_{-0.0061}$	-1.13	$^{+0.37}_{-0.39}$
RBAO + WMAP5 + SNe [16]	0.271	$^{+0.015}_{-0.014}$	-0.0021	± 0.0062	-0.974	$^{+0.057}_{-0.059}$

TABLE II: Results from the cosmology fits with the radial BAO (RBAO) data and additional data sets assuming a constant equation of state parameter w . In the data set a) the BAO radial data are used, and the values and errors of $\Omega_m h^2$ and $\Omega_b h^2$ from [1] are input into the computation of $H(z_i)r_s$. For the data set b) the BAO radial data is combined with the measurement of $l_A(z^*)$ from [1]. In all cases, the WMAP5 measurements within the appropriate cosmology model (flat, cosmological constant, etc.) are used, although the differences are minute. See text for details. The other rows combine the RBAO redshift scales presented in Table I with external data sets. All fits have values of χ^2/N_{dof} close to one. In particular, the fit with RBAO, WMAP5 and SNe data within a non-flat constant- w cosmology (last row) has $\chi^2/N_{dof} = 312/306$.

with

$$a_{eq} = \frac{\Omega_\gamma h^2 (1 + 0.2271 N_{eff})}{\Omega_m h^2} \quad (5)$$

with z_d , the so-called drag redshift, written in terms of $\Omega_b h^2$ and $\Omega_m h^2$ (Eq. (3) in [1]), and take $\Omega_b h^2 = 0.02273 \pm 0.0066$ and $\Omega_m h^2 = 0.1329 \pm 0.0064$ from the five-year results of the Wilkinson Microwave Anisotropy Probe (WMAP5) [1] as external constraints (we could have also used determinations from nucleosynthesis or LSS measurements), and fix $\Omega_\gamma h^2 = 2.449 \times 10^{-5}$ and $N_{eff} = 3.04$. Note that the expression for $\Delta z_{BAO}(z_i)$ in Eq. (3) is independent of H_0 . Once the one-sigma WMAP5 constraints on $\Omega_b h^2$ and $\Omega_m h^2$ are included in the fit, the only remaining unknowns are $\Omega_\Lambda (\equiv 1 - \Omega_m)$ and w . Since we have two independent $\Delta z_{BAO}(z_i)$, we will be able to determine them.




FIG. 1: 68% CL constraints on the plane $\Omega_m - w$ when a flat Universe is assumed. In yellow, from the measurements of the RBAO scale plus $\Omega_b h^2$ and $\Omega_m h^2$. In pink, from RBAO plus l_A and z^* from WMAP5 [1]. In blue, from the RBAO scale plus WMAP5 and in red from the RBAO scale, plus WMAP5 plus supernovae [16]. Results are consistent among them and compatible with a Λ CDM cosmology.

b) Alternatively, we can use as external constraints the measurements by WMAP5 of the ratio $l_A(z^*)$ between the distance to the last scattering surface and $r_s(z^*)$ and of z^* itself, $l_A(z^*) = 302.14 \pm 0.87$ and $z^* = 1090.5 \pm 1.0$ with a $\sim 40\%$ positive correlation, when assuming a flat CDM model with constant equation of state w . Again, H_0 cancels out and we are left with only Ω_m and w as unknowns, which we then proceed to determine.

In all cases, a frequentist χ^2 fit including correlations is applied to the data, using the publicly available Minit code [17] for minimization and contour calculation. Systematic errors in $\Delta z_{BAO}(z_i)$ have been neglected, being subdominant (see Table I) and relatively uncertain. Including them does not change the results significantly. The results obtained for Ω_m and w and their 1- σ errors, $\sigma(\Omega_m)$ and $\sigma(w)$, are shown in rows 1 and 2 of Table II. Both determinations, a) and b), are consistent between them and in agreement with Λ CDM. The two-dimensional constraints at 68% CL can be seen in Fig. 1.

Next, we have combined the results obtained from the radial BAO scale with the full CMB distance constraints, using the WMAP5 measurements of the shift parameter $R(z^*)$, the acoustic scale $l_A(z^*)$ and the redshift at decoupling z^* presented in Table 10 of [1], and the covariance matrix in Table 11 therein. We have also added the supernovae set compiled by Kowalski *et al.* [16], which can be found in [22]. We have taken their data covariance




FIG. 2: 68% CL constraints on the plane Ω_m - w when a non-flat constant- w cosmology is assumed. In blue, from the RBAO scale plus WMAP5 [1] and in red from the RBAO scale, plus WMAP5 plus supernovae [16]. Results are consistent with a cosmological constant Universe with a precision around 5%.

matrix without systematical errors. Adding systematics does not change the results qualitatively. The results of the combination assuming flatness can be seen in rows 3 and 4 of Table II. We determine the equation of state parameter w , assumed constant, to be consistent with a cosmological constant to about 5%.

If, instead, we assume the Λ CDM model and drop the assumption of flatness, we obtain the results in rows 5–8 of Table II, which show consistency with a flat Universe at the 0.5% level.

We can now keep the equation of state parameter w free and drop the assumption of flatness. The constraints, shown in the last two rows of Table II remain very robust when combining the three data sets. The two-dimensional Ω_m - w contour at 68% CL is shown in Fig. 2. All results are in good agreement with Λ CDM. Moreover, the combination allows us to simultaneously determine w with a precision around 5%, and set the flatness of the Universe with a precision of about half of a percent. As a by-product, we obtain a Hubble parameter $h = 0.703 \pm 0.020$, consistent with most recent

results in the literature. If instead of RBAO we use in the combination the BAO monopole determination in [8], we find compatible results with errors that are 20–40% larger (see Table 6 in [16]). On the other hand, if we combine RBAO and the monopole in [8], we find no improvement in the error band with respect to the case with RBAO alone.

Finally, we can drop the assumption of constant w and, instead, use the parametrization $w(z) = w_0 + w_a \cdot z/(1+z)$ [18, 19]. Assuming a flat Universe, and combining RBAO with WMAP5 and supernova data, we get $w_0 = -1.05^{+0.11}_{-0.10}$ and $w_a = 0.44^{+0.37}_{-0.51}$, with an 87% anticorrelation, consistent with Λ CDM ($w_0 = -1$, $w_a = 0$). We conclude that current data show no evidence for a time-dependent w , although the constraint on w_a is still very weak.

In summary, we have determined for the first time the cosmological parameters Ω_m and w using the radial BAO scale, to be $\Omega_m = 0.24^{+0.06}_{-0.05}$ and $w = -1.14 \pm 0.39$. These results are perfectly consistent with the expectations of the Λ CDM cosmology. Moreover, when these results are combined with the cosmological distance constraints coming from the CMB [1] and type-Ia supernovae [16], we measure $\Omega_m = 0.271 \pm 0.015$, $\Omega_k = -0.002 \pm 0.006$, and $w = -0.974 \pm 0.058$. The result is in good agreement with a flat Λ CDM cosmology, and shows that the determination of the RBAO scale is consistent with all the other cosmological measurements.

In the future, currently planned BAO surveys like Baryon Oscillations Sloan Survey (BOSS) [23] or Physics of the Accelerating Universe (PAU) [20] will measure radial BAOs with greater precision up to redshifts close to $z = 1$, producing more stringent constraints on the properties of dark energy.

We would like to thank the “Centro de Ciencias de Benasque Pedro Pascual” for the hospitality while most of this paper was written. RM wants to thank Eric Linder for useful discussions. We acknowledge funding from the Spanish Ministerio de Ciencia y Tecnología (MEC) through projects AYA2006-06341, AYA2006-13735-C02-01 and AYA2006-13735-C02-02, and Consolider-Ingenio CSD2007-00060, together with EC-FEDER funding, and the research project 2005SGR00728 from Generalitat de Catalunya.

[1] E. Komatsu, et al., *ApJS* **180**, 330 (2009)
 [2] A. Cabré and E. Gaztañaga, *MNRAS* **393**, 1183 (2009)
 [3] A. Cabré and E. Gaztañaga, *MNRAS* **396**, 1119 (2009)
 [4] E. Gaztañaga, A. Cabré, F. Castander, M. Crocce, and P. Fosalba, *MNRAS* in press (2009), 0807.2448.
 [5] E. Gaztanaga, A. Cabre, and L. Hui, *MNRAS* in press (2009), 0807.3551.
 [6] D. J. Eisenstein et al., *AJ* **122**, 2267 (2001)

[7] W. J. Percival et al., *MNRAS* **381**, 1053 (2007)
 [8] D. J. Eisenstein, et al., *ApJ* **633**, 560 (2005)
 [9] D. J. Eisenstein et al., *Bulletin of the American Astronomical Society* (2004), vol. 36 p. 1460
 [10] A. G. Sánchez et al., *MNRAS* **366**, 189 (2006)
 [11] G. Hütsi, *A&A* **449**, 891 (2006)
 [12] G. Hütsi, *A&A* **459**, 375 (2006)
 [13] N. Padmanabhan et al., *MNRAS* **378**, 852 (2007)

- [14] C. Blake et al., MNRAS **374**, 1527 (2007)
- [15] A. G. Sanchez et al., ArXiv e-prints (2009), 0901.2570.
- [16] M. Kowalski, et al., ApJ **686**, 749 (2008)
- [17] F. James and M. Roos, Computer Physics Communications **10**, 343 (1975).
- [18] M. Chevallier and D. Polarski, Int. J. Mod. Phys. D **10**, 213 (2001)
- [19] E. V. Linder, Phys. Rev. Lett. **90**, 091301 (2003)
- [20] N. Benítez, et al., ApJ **691**, 241 (2009)
- [21] We have checked that we can safely neglect the terms proportional Ω_k and Ω_{DE} in Eq. 4.
- [22] <http://supernova.lbl.gov/Union/>
- [23] <http://www.sdss.org/news/releases/20080110.sdss3.html>