arXiv:0808.2077v3 [quant-ph] 6 Oct 2011

Bounds of concurrence and their relation with fidelity and frontier states^{*}

Zhihao Ma,¹ Fu-Lin Zhang,² Dong-Ling Deng,² and Jing-Ling Chen^{2,†}

¹Department of Mathematics, Shanghai Jiaotong University, Shanghai, 200240, P.R.China

² Theoretical Physics Division, Chern Institute of Mathematics, Nankai University, Tianjin, 300071, P.R.China

PHONE: 011+8622-2350-9287, FAX: 011+8622-2350-1532

The bounds of concurrence in [F. Mintert and A. Buchleitner, Phys. Rev. Lett. 98 (2007) 140505] and [C. Zhang *et. al.*, Phys. Rev. A 78 (2008) 042308] are proved by using two properties of the fidelity. In two-qubit systems, for a given value of concurrence, the states achieving the maximal upper bound, the minimal lower bound or the maximal difference upper-lower bound are determined analytically.

PACS numbers: 03.67.-a, 03.65.Ta

I. INTRODUCTION

Entanglement, which depicts the nonclassical connection between two parts of a quantum system, has been recognized as crucial in various field of quantum information in recent years [1–7]. Several measures have been proposed to quantify the degree to which a state is entangled, such as entanglement of formation [8–10], entanglement of distillation [11], relative entropy of entanglement [12], negativity [13, 14], and so on. For two-qubit systems, the entanglement of formation is equivalent to a computable quantity, which is referred to as *concurrence* [9, 10]. The concurrence of a pure two-qubit state $|\psi\rangle$ is given by

$$C(|\psi\rangle) = \sqrt{2(1 - \text{Tr}\rho^{A^2})} = \sqrt{2(1 - \text{Tr}\rho^{B^2})},$$
 (1)

where $\rho^A = \text{Tr}_B |\psi\rangle \langle \psi|$ is the partial trace of $|\psi\rangle \langle \psi|$ over subsystem *B*, and ρ^B has a similar meaning. For a mixed state, the concurrence is defined as the average concurrence of the pure states of the decomposition, minimized over all decompositions of $\rho = \sum_j p_j |\psi_j\rangle \langle \psi_j|$,

$$C(\rho) = \min \sum_{j} p_j C(|\psi_j\rangle).$$
(2)

It has been proved in [9, 10] that Eq. (2) can be expressed explicitly as

$$C(\rho) = \max\{0, \sqrt{\lambda_1} - \sqrt{\lambda_2} - \sqrt{\lambda_3} - \sqrt{\lambda_4}\}, \qquad (3)$$

in which $\lambda_1, ..., \lambda_4$ are the eigenvalues of the operator $R = \rho(\sigma_y \otimes \sigma_y)\rho^*(\sigma_y \otimes \sigma_y)$ in decreasing order and σ_y is the second Pauli matrix.

Recently, many works about detecting entanglement experimentally have been reported [15–20]. In one of the existing schemes, the authors of [16, 19] presented observable lower and upper bounds of the squared concurrence

$$2\left[\operatorname{Tr}\rho^{2} - \operatorname{Tr}\rho^{A^{2}}\right] \leq C^{2}(\rho) \leq 2\left[1 - \operatorname{Tr}\rho^{A^{2}}\right], \qquad (4)$$

where $C(\rho)$ is the concurrence for arbitrary dimensional states, taking the definitions in Eq. (1) and (2). These bounds can be easily obtained by a few experimental measurements on a twofold copy $\rho \otimes \rho$ of the mixed states.

It is shown that the bounds provide an excellent estimation of concurrence for weakly mixed (called quasipure in [16]) states [19], and they have certain relations with the degree of mixing for a mixed state and some properties of the linear entropy. In this letter, we give closer analysis of the bounds. First, we find the inequality (4) can be distinctly understood in the viewpoint of fidelity, which is another important concept in quantum information [1, 21-23]. The details are given in Sec. II. To show the departure of the bounds for a given concurrence, in Sec. III, the two-qubit frontier states are determined analytically. Namely, we present the quantum states, which achieve the maximal upper bound, the minimal lower bound or the maximal difference upper-lower bound, for a given value of entanglement. In Sec. IV, we discuss the relations of the bounds and concurrence or the linear entropy, and give a brief conclusion.

II. PROOF BY FIDELITY

Fidelity is a measure of *closeness* of two quantum states, which is shown to be closely related to entanglement in many aspects [9, 10, 24, 25]. The fidelity between two states ρ_1 and ρ_2 of a quantum system R reads [1, 21–23]

$$F(\rho_1, \rho_2) = \left[\operatorname{Tr}\left(\sqrt{\sqrt{\rho_1}\rho_2\sqrt{\rho_1}}\right) \right]^2.$$
 (5)

When $\rho_1 = |\phi\rangle\langle\phi|$ and $\rho_2 = |\varphi\rangle\langle\varphi|$ are two pure states, $F(\rho_1, \rho_2) = \text{Tr}(|\phi\rangle\langle\phi|\varphi\rangle\langle\varphi|) = |\langle\phi|\varphi\rangle|^2$. Let Q be a copy of R, and $|\psi_1\rangle$ is a purification of ρ_1 and $|\psi_2\rangle$ of ρ_2 into RQ, then

$$F(\rho_1, \rho_2) \ge F(\sigma_1, \sigma_2) = |\langle \psi_1 | \psi_2 \rangle|^2, \tag{6}$$

^{*}Physics Letters A 373 (2009) 1616-1620

[†]Email:chenjl@nankai.edu.cn

where $\sigma_1 = |\psi_1\rangle\langle\psi_1|$, $\sigma_2 = |\psi_2\rangle\langle\psi_2|$, and $\text{Tr}_Q\sigma_1 = \rho_1$, $\text{Tr}_Q\sigma_2 = \rho_2$. It is proved in [26], that fidelity can be bounded above by the so called super-fidelity $G(\rho_1, \rho_2)$:

$$F(\rho_{1}, \rho_{2}) \leq G(\rho_{1}, \rho_{2})$$

= $\operatorname{Tr} \rho_{1} \rho_{2} + \sqrt{(1 - \operatorname{Tr} \rho_{1}^{2})(1 - \operatorname{Tr} \rho_{2}^{2})}$
 $\leq 1.$ (7)

When ρ_1 and ρ_2 are two pure states, $G(\rho_1, \rho_2) = \text{Tr}\rho_1\rho_2 = F(\rho_1, \rho_2)$. The following is the proof of the inequality (4) by using the properties of fidelity and superfidelity mentioned above.

Proof. Suppose $\{|\psi_i\rangle\}$ is an decomposition minimizing the average concurrence in Eq. (2), $\rho_i = |\psi_i\rangle\langle\psi_i|$, and $\rho = \sum_i t_i \rho_i$, where $\sum_i t_i = 1$. The reduced density operator $\rho^A = \sum_i t_i \rho_i^A$, where $\rho_i^A = \text{Tr}_B \rho_i$. It is straightforward to obtain

$$C^{2}(\rho) = 2 \sum_{i,j} t_{i} t_{j} \sqrt{\left[1 - \operatorname{Tr} \rho_{i}^{A^{2}}\right] \left[1 - \operatorname{Tr} \rho_{j}^{A^{2}}\right]},$$

$$\operatorname{Tr} \rho^{2} = \sum_{i,j} t_{i} t_{j} \operatorname{Tr}(\rho_{i} \rho_{j}),$$

$$\operatorname{Tr} \rho^{A^{2}} = \sum_{i,j} t_{i} t_{j} \operatorname{Tr}(\rho_{i}^{A} \rho_{j}^{A}).$$
(8)

Then, one can notice that

$$1 \geq \operatorname{Tr}(\rho_i^A \rho_j^A) + \sqrt{\left[1 - \operatorname{Tr}\rho_i^{A^2}\right] \left[1 - \operatorname{Tr}\rho_j^{A^2}\right]} = G(\rho_i^A, \rho_j^A) \geq F(\rho_i^A, \rho_j^A).$$
(9)

On the other hand, ρ_i and ρ_j are two purifications of ρ_i^A and ρ_i^A , we obtain

$$1 \ge G(\rho_i^A, \rho_j^A) \ge F(\rho_i^A, \rho_j^A) \ge F(\rho_i, \rho_j) = \operatorname{Tr}(\rho_i \rho_j).(10)$$

Hence,

$$2 \ge C^{2}(\rho) + 2\operatorname{Tr}\rho^{A^{2}} = 2\sum_{i,j} t_{i}t_{j}G(\rho_{i}^{A},\rho_{j}^{A})$$
$$\ge 2\sum_{i,j} t_{i}t_{j}\operatorname{Tr}(\rho_{i}\rho_{j}) = 2\operatorname{Tr}\rho^{2}.(11)$$

This ends the proof.

III. FRONTIER STATES

In the above section, we show the properties of the fidelity lead to the upper and lower bounds of squared concurrence in Eq. (4). For the sake of brevity, we adopt the *tangle* [27] to replace concurrence, $\tau = C^2$. It is

bounded by $\tau_L \leq \tau \leq \tau_U$, where

$$\tau_{L} = 2 \operatorname{Tr} \rho^{2} - 2 \min\{ \operatorname{Tr} \rho^{A^{2}}, \operatorname{Tr} \rho^{B^{2}} \}, \tau_{U} = 2 - 2 \max\{ \operatorname{Tr} \rho^{A^{2}}, \operatorname{Tr} \rho^{B^{2}} \}.$$
(12)

The aim of this section is to derive the quantum states achieving the maximal τ_U , the minimal τ_L or the maximal their difference $\Delta = \tau_U - \tau_L$, for a given value of τ . These frontier state are denoted as ρ_U , ρ_L and ρ_Δ respectively in the following paragraphes. We only consider the two-qubit case, for which the concurrence has the exact formula as Eq. (3).

A. Frontier state of τ_U , ρ_U

It is easy to obtain, when the purities of the two subsystem $\text{Tr}\rho^{A^2} = \text{Tr}\rho^{B^2} = \frac{1}{2}$, $\tau_U = 1$ achieves the maximum. A straightforward choice of the frontier of τ_U is the isotropic state or say the Werner state [28, 29]

$$\rho_U = \rho_W = \frac{1-x}{4} I_2 \otimes I_2 + x |\Phi_+\rangle \langle \Phi_+|, \qquad (13)$$

where I_2 is the 2 × 2 identity matrix and $|\Phi_+\rangle = [|00\rangle + |11\rangle]/\sqrt{2}$ is one of the Bell basis. Then $\tau(\rho_U) = [\max\{0, \frac{3}{2}(x-\frac{1}{3})\}]^2$ and $\tau_U(\rho_U) = 1$ for any value of tangle.

B. Frontier state of Δ , ρ_{Δ}

The difference between the upper and lower bounds can be written as

$$\Delta = 2\left[1 - \mathrm{Tr}\rho^2\right] - 2\left|\mathrm{Tr}\rho^{A^2} - \mathrm{Tr}\rho^{B^2}\right|.$$
 (14)

The first part of Eq. (14) is the linear entropy [30], a measure for the degree of of mixture of the quantum state. In [31], the authors present a class of *maximally entangled* mixed states:

$$\rho_{MEMS} = \begin{bmatrix} g(\gamma) & 0 & 0 & \gamma/2 \\ 0 & 1 - 2g(\gamma) & 0 & 0 \\ 0 & 0 & 0 & 0 \\ \gamma/2 & 0 & 0 & g(\gamma) \end{bmatrix},$$
(15)

with $g(\gamma) = 1/3$ for $0 \le \gamma \le 2/3$ and $g(\gamma) = \gamma/2$ for $2/3 \le \gamma \le 1$. For a given value of tangle, its linear entropy

$$2\left[1 - \mathrm{Tr}\rho_{MEMS}^{2}\right] = 4g(\gamma)\left[2 - 3g(\gamma)\right] - \gamma^{2}, \quad (16)$$

reaches the maximum over all the two-qubit states, with $\tau = \gamma^2$. Straightforward calculations indicate the two subsystems have the equivalent purities $\text{Tr}\rho_{MEMS}^A{}^2 = \text{Tr}\rho_{MEMS}^B{}^2 = 2g^2(\gamma) - 2g(\gamma) + 1$. Therefore, the state in Eq. (15) makes Δ maximal, $\rho_{\Delta} = \rho_{MEMS}$, and the



FIG. 1: Plot of (a) 30000 randomly generated states in the τ_{U^-} τ plane; (b) and (c) 30000 randomly generated states weighted in the τ_L - τ and Δ - τ planes. The curves in the planes are ρ_U (dashed), ρ_L (solid) and ρ_{Δ} (dot-dashed). The small triangles in the planes indicate an intermediate state between ρ_U and ρ_{Δ} .

corresponding maximum equals the linear entropy in Eq. (16).

C. Frontier state of τ_L , ρ_L

The region of the two-qubit states in the τ_L - τ plane is shown in Fig. 1(b). The state ρ_L , with the minimal τ_L for a given τ , achieves the maximal τ when the value of τ_L is fixed. To derive ρ_L , we postulate it take the ansatz form

$$\rho_{ansatz} = \begin{bmatrix} x + \gamma/2 & 0 & 0 & \gamma/2 \\ 0 & a & 0 & 0 \\ 0 & 0 & b & 0 \\ \gamma/2 & 0 & 0 & y + \gamma/2 \end{bmatrix},$$
(17)

with the non-negative real parameters, $x + y + a + b + \gamma =$ 1. It comprises a mixture of the Bell state $|\Phi_{+}\rangle$ and a mixed diagonal state. There are two reasons for choosing this ansatz state. (i) The frontier states ρ_U and ρ_{Δ} determined above are relating to this form of ρ_{ansatz} . As shown in Fig. 1(a), when the tangle takes a large value, for instance 0.8 or larger, τ_U of ρ_{Δ} is close to 1, moreover its Δ is the maximum. This indicates ρ_{Δ} is a candidate for the frontier ρ_L . (ii) For a pure state, the lower bound $\tau_L = \tau$. When the state is mixed, τ_L deviates from the tangle τ . This can be shown by a special type states in Fig. 3(d), which will be discussed in Sec. IV. Therefore, we can speculate the value of τ_L is a token of purity of a state. Otherwise, all the maximally entangled states for a given degree of mixing derived in [31, 32] take the form in Eq. (17), with various combinations of entanglement and mixedness measures.

The tangle of ρ_{ansatz} is given by

$$\tau = \left\{ \max\{\gamma - 2\sqrt{ab}, 0\} \right\}^2, \tag{18}$$

with the maximum $\tau = \gamma^2$ when b = 0, as shown in [31]. One can surmise directly $\text{Tr}\rho^{A^2} = \text{Tr}\rho^{B^2}$, when τ_L reached its minimum for a fixed τ , which requires $x = y = (1 - a - \gamma)/2$. Then, the lower bound is given by

$$\tau_L = 2a^2 - 2a + \gamma^2, \tag{19}$$

with $a \in [0, 1 - \gamma]$. The upper bound of a holds $x = y \ge 0$. Therefore, when $1 - \gamma \ge 1/2$, or say $\gamma \le 1/2$, the minimum points of τ_L are a = 1/2 and $x = (1 - 2\gamma)/4$. In the other region $\gamma \ge 1/2$, the optimal solution occurs when $a = 1 - \gamma$ and x = 0. Thus, the frontier state, achieving the minimal τ_L for a fixed τ , can be written in the same form as ρ_Δ

$$\rho_L = \begin{bmatrix} f(\gamma) & 0 & 0 & \gamma/2 \\ 0 & 1 - 2f(\gamma) & 0 & 0 \\ 0 & 0 & 0 & 0 \\ \gamma/2 & 0 & 0 & f(\gamma) \end{bmatrix},$$
(20)

with $f(\gamma) = 1/4$ for $0 \le \gamma \le 1/2$ and $f(\gamma) = \gamma/2$ for $1/2 \le \gamma \le 1$. For a given degree of entanglement $\tau = \gamma^2$, its τ_L is given by

$$\tau_L = \begin{cases} \gamma^2 - 1/2, & 0 \le \gamma \le 1/2; \\ 3\gamma^2 - 2\gamma, & 1/2 \le \gamma \le 1. \end{cases}$$
(21)

To verify the frontier state ρ_L , we randomly generate one million two-qubit matrices, which indicate the region of physically acceptable states in the $\tau_L - \tau$ plane is encircled perfectly by the curve of the state in Eq. (20). We plot the curve comparing with 30000 weighted random states, namely the mixtures of random states and ρ_L with random weights, in Fig. 1(b). The curve joins with the one of ρ_U at $(\tau, \tau_L) = (0, -1/2)$, where $\rho_L = [|00\rangle\langle 00| + |11\rangle\langle 11| + 2|01\rangle\langle 01|]/4$ and $\rho_U = I_2 \otimes I_2/4$. When $\gamma \geq 2/3$, $\rho_L = \rho_\Delta$; and when $\gamma < 2/3$, the values of τ_L for ρ_L and ρ_Δ are less than 0.

D. Frontier state of Δ_R , ρ_{Δ_R}

In the above discussion, we notice, when the degree of entanglement $\tau \leq 4/9$, the lower bound τ_L has the possibility of being less than 0. But the value of concurrence, as a good defined entanglement measure, should be nonnegative [33, 34]. This suggests a more reasonable lower bound of the squared concurrence should be corrected as

$$\tau_{L_R} = \max\{\tau_L, 0\},\tag{22}$$

whose difference with the upper bound is given by

$$\Delta_R = \tau_U - \tau_{L_R} = \begin{cases} \tau_U, & \tau_L < 0; \\ \tau_U - \tau_L, & \tau_L \ge 0. \end{cases}$$
(23)

Then, both the frontier states ρ_L and ρ_{Δ} correspond with the minimal τ_{L_R} for a given value of τ . A subsequent question arises: what is the frontier state ρ_{Δ_R} , which achieve the maximal Δ_R for a given τ ?

There are two regions of τ , where the frontier state ρ_{Δ_R} can be determined directly. The first one occurs for $\tau \leq (2-\sqrt{3})/2$, and the second for $\tau \geq 4/9$. In the former case, the τ_L of ρ_U or say the Werner state is less than 0, whose $\tau_U = 1$. Hence, $\rho_{\Delta_R} = \rho_U = \rho_W$ in this region. In the later case, $\rho_{\Delta_R} = \rho_\Delta = \rho_{MEMS}$, since $\tau_L \geq 0$ for all the two-qubit states with $\tau \geq 4/9$.

In the in-between region with $(2 - \sqrt{3})/2 < \tau < 4/9$, $\tau_L(\rho_W) = (-1 + 2\sqrt{\tau} + 2\tau)/3 > 0$ and $\tau_L(\rho_{MEMS}) = -4/9 + \tau < 0$. Their upper bounds of tangle are constants, $\tau_U(\rho_W) = 1$ and $\tau_U(\rho_{MEMS}) = 8/9$. One can image an intermediate state, as shown by the black triangles (\blacktriangle and \blacktriangledown) in Fig. 1, whose τ_U and τ_L are intermediate between the ones of ρ_W and ρ_{MEMS} . When it transforms from ρ_{MEMS} to ρ_W , as expressed by the directions of the triangles, τ_U and τ_L are enhanced, whereas Δ decreases. Considering the relations in Eq. (23), we can postulate the maximum of Δ_R occurs when $\tau_L = 0$. The Werner state and ρ_{MEMS} , in this region, can be uniformed by

$$\rho_{\blacktriangle} = \frac{1}{2} \begin{bmatrix} 1-S & 0 & 0 & \gamma + \sqrt{S^2 - M^2} \\ 0 & S+M & 0 & 0 \\ 0 & 0 & S-M & 0 \\ \gamma + \sqrt{S^2 - M^2} & 0 & 0 & 1-S \end{bmatrix}$$

with the parameters $0 \leq M \leq S \leq 1$, $(\sqrt{3}-1)/2 < \gamma < 2/3$, and the tangle $\tau = \gamma^2$. When S = M = 1/3, $\rho_{\blacktriangle} = \rho_{MEMS}$; and when $S = (1 - \gamma)/3$ and M = 0, ρ_{\blacktriangle} returns into the Werner state in term of the concurrence γ [32].

Taking the intermediate state ρ_{\blacktriangle} as a candidate for ρ_{Δ_R} , we will determine the values of S and M for a given γ in the following. The bounds of tangle for the state ρ_{\blacktriangle}



FIG. 2: Curve of ρ_{Δ_R} is plotted in the $\Delta_R - \tau$ plane, in company with 30000 random states weighted based on it. Upper right is the magnified region nearby the two critical points, $\tau = (2 - \sqrt{3})/2$ and $\tau = \gamma_c^2$

are given by

$$\tau_L = -2S(1-S) + (\gamma + \sqrt{S^2 - M^2})^2,$$

$$\tau_U = 1 - M^2.$$
(24)

The condition $\tau_L = 0$ leads to

$$M = \mathcal{M}(S) = \sqrt{S^2 - \left[\sqrt{2S(1-S)} - \gamma^2\right]^2},$$
 (25)

with $S \in [\mathcal{S}_{-}(\gamma), \mathcal{S}_{+}(\gamma)]$. Here, $\mathcal{S}_{\pm}(\gamma) = (1 \pm \sqrt{1-2\gamma^2})/2$ keep the value of $\sqrt{S^2 - M^2}$ to be non-negative. Under such a condition, the maximum of Δ_R occurs at

$$S = S(\gamma) = \begin{cases} S_0(\gamma), & (\sqrt{3} - 1)/2 < \gamma < \gamma_c \\ S_-(\gamma), & \gamma_c < \gamma < 2/3 \end{cases},$$
(26)

where $\gamma_c \approx 0.38218$, and $\mathcal{S}_0(\gamma)$ is the root of the equation $\partial_S \mathcal{M}(S) = 0$ in the region $[\mathcal{S}_-(\gamma), \mathcal{S}_+(\gamma)]$.

Comparing the solutions of ρ_{Δ_R} for the three regions of τ , one can find it can be expressed in the form of ρ_{\blacktriangle} with

$$(S,M) = \begin{cases} ((1-\gamma)/3,0), & 0 \le \gamma < (\sqrt{3}-1)/2; \\ (\mathcal{S}(\gamma),\mathcal{M}(S)), & (\sqrt{3}-1)/2 \le \gamma \le 2/3; (27) \\ (1-\gamma,S), & 2/3 \le \gamma \le 2/3. \end{cases}$$

This frontier state ρ_{Δ_R} has been verified by one million random two-qubit states. In Fig. 2, we show the curve of ρ_{Δ_R} in the $\Delta_R - \tau$ plane, in company with 30000 random states weighted based on it.

IV. DISCUSSION AND CONCLUSION

In this letter, we show the properties of the fidelity lead to the upper and lower bounds of the squared concurrence for a biparticle system. The bounds are closely related to the schemes to observe entanglement experimentally. The two important concepts in quantum infor-



FIG. 3: Plot of the errors of the bounds, $\tau_U - \tau$ and $\tau - \tau_L$, versus τ and Δ , for the generalized Werner states. The curves show the results of $\theta = \pi/4$ (solid), $\theta = \pi/6$ (dashed), $\theta = \pi/12$ (dot-dashed) and $\theta = 0$ (dotted).

mation, entanglement and fidelity, have been pointed out to be related to each other in many aspects [9, 10, 24, 25]. Recently, the relation between entanglement and Berry phase begins to attract the attention of researchers [35]. On the other hand, the Uhlmanns mixed state geometric phase has the inherent connection with fidelity [36]. These suggest the bounds investigated in this letter may induce some characters of the geometric phase of mixed state.

As a further research of bounds of the squared concurrence, limited to the two-qubit case, we derived the frontier states, which corresponding to the maximal upper bound, the minimal lower bound or the maximal difference between the bounds, for a given value of concur-

- M. A. Nielsen and I. L. Chuang, *Quantum Computation* and *Quantum Information* (Cambridge University Press, Cambridge, 2000).
- [2] C. H. Bennett, G. Brassard, C. Crepeau, and *et al*, Phys. Rev. Lett. **70**, 1895 (1993).
- [3] C. H. Bennett and S. J. Wiesner, Phys. Rev. Lett. 69, 2881 (1992).
- [4] P. W. Shor, SIAM J. Comput. 26, 1484 (1997).
- [5] L. K. Grover, Phys. Rev. Lett. 79, 325 (1997).
- [6] A. K. Ekert, Phys. Rev. Lett. 67, 661 (1991).
- [7] D. Deutsch, A. Ekert, and *et al*, Phys. Rev. Lett. **77**, 2818 (1996).
- [8] C. H. Bennett, D. P. DiVincenzo, J. A. Smolin, and W. K. Wootters, Phys. Rev. A 54 (1996).
- [9] S. Hill and W. K. Wootters, Phys. Rev. Lett. 78, 5022 (1997).
- [10] W. K. Wootters, Phys. Rev. Lett. 80, 2245 (1998).
- [11] C. H. Bennett, G. Brassard, S. Popescu, B. Schumacher, J. Smolin, and W. K. Wootters, Phys. Rev. Lett. 76, 722 (1996).
- [12] V. Vedral, M. B. Plenio, K. Jacobs, and P. L. Knight, Phys. Rev. A 56, 4452 (1997).
- [13] K. Życzkowski, P. Horodecki, A. Sanpera, and M. Lewenstein, Phys. Rev. A 58, 883 (1998).
- [14] G. Vidal and R. F. Werner, Phys. Rev. A 65, 032314

rence. We also determined the state achieving the maximal difference between the upper bound and the modified lower bound, which was defined as the maximum of 0 and the lower bound. All of these frontier states were relating to this form of ρ_{ansatz} in Eq. (17), a mixture of a Bell base and a mixed diagonal state.

We haven't discussed more general properties of the bounds in this letter. Instead, we exhibit some characteristics, in Fig. 3, by a family of generalized Werner states [37],

$$\rho_W(\theta) = \frac{1-x}{4} I_2 \otimes I_2 + x |\Phi(\theta)\rangle \langle \Phi(\theta)|, \qquad (28)$$

which are the mixtures of an entangled pure state $|\Phi(\theta)\rangle = \cos \theta |00\rangle + \sin \theta |11\rangle$ with the completely random state. One can notice, for a given value of entanglement, both the largest errors of the upper and lower bounds occurs when $\theta = \pi/4$, corresponding with the originally defined Werner state in Eq. (13). When the linear entropy (equalling to Δ here) increases, the errors are enhanced monotonously. These results provided the preparatory knowledge to derive the frontier states in Sec. III.

Acknowledgments

This work is supported by the New teacher Foundation of Ministry of Education of P.R.China (Grant No. 20070248087). J.L.C is supported in part by NSF of China (Grant No. 10605013), and Program for New Century Excellent Talents in University, and the Projectsponsored by SRF for ROCS, SEM.

(2002).

- [15] B. Terhal, Phys. Lett. A 271, 319 (2000).
- [16] F. Mintert and A. Buchleitner, Phys. Rev. Lett. 98, 140505 (2007).
- [17] F. Mintert, M. Kuś, and A. Buchleitner, Phys. Rev. Lett. 95, 260502 (2005).
- [18] J. Eisert, F. Brandão, and K. Audenaert, New J. Phys. 9, 46 (2007).
- [19] C.-J. Zhang, Y.-X. Gong, Y.-S. Zhang, and G.-C. Guo, Phys. Rev. A 78, 042308 (2008).
- [20] X.-L. Niu, Y.-X. Gong, J. Li, L. Peng, C.-J. Zhang, Y.-S. Zhang, Y.-F. Huang, and G.-C. Guo, eprint arXiv: 0808.0776.
- [21] D. Bures, Trans. Am. Math. Soc. **135**, 199 (1969).
- [22] A. Uhlmann, Rep. Math. Phys. 9, 273 (1976).
- [23] A. Uhlmann, Rep. Math. Phys. 24, 229 (1986).
- [24] A. Uhlmann, Phys. Rev. A 62, 032307 (2000).
- [25] J.-M. Cai, Z.-W. Zhou, and G.-C. Guo, Phys. Lett. A 363, 392 (2007).
- [26] J. A. Miszczak, Z. Puchała, P. Horodecki, A. Uhlmann, and K.Życzkowski, Quantum Information and Computation 9, 0103 (2009).
- [27] V. Coffman, J. Kundu, and W. K. Wootters, Phys. Rev. A 61, 052306 (2000).
- [28] B. M. Terhal and K. G. H. Vollbrecht, Phys. Rev. Lett.

85, 2627 (2000).

- [29] R. F. Werner, Phys. Rev. A 40, 4277 (1989).
- [30] S. Bose and V. Vedral, Phys. Rev. A 61, 040101(R) (2000).
- [31] W. J. Munro, D. F. V. James, A. G. White, and P. G. Kwiat, Phys. Rev. A 64, 030302(R) (2001).
- [32] T.-C. Wei, K. Nemoto, P. M. Goldbart, P. G. Kwiat, W. J. Munro, and F. Verstraete, Phys. Rev. A 67, 022110 (2003).
- [33] V. Vedral, M. Plenio, M. Rippin, and P. Knight, Phys. Rev. Lett. 78, 2275 (1997).
- [34] M. Horodecki, P. Horodecki, and R. Horodecki, Phys. Rev. Lett. 84, 2014 (2000).
- [35] B. Basu, Europhys. Lett. **73**, 833 (2006).
- [36] A. Uhlmann, Ann. Phys. (Leipzig) **501**, 63 (1989).
- [37] D.-L. Deng and J.-L. Chen, Annals of Physics 324, 408 (2009).