

Multidimensional Chebyshev Systems (Haar systems) - just a definition

Ognyan Kounchev
Institute of Mathematics and Informatics
Bulgarian Academy of Sciences
and
IZKS, University of Bonn

May 22, 2018

Abstract

The notion of Chebyshev system was coined by S. Bernstein although many results were proved already by A. Markov. Chebyshev systems play important role in the one-dimensional Moment problem and Approximation and Spline theory. We generalize the notion of Chebyshev system for several dimensions and define the Multidimensional Chebyshev Systems of order N . We prove that this definition is satisfied by the solutions of a wide class of elliptic equations of order $2N$. This definition generalizes a very large class of Extended Complete Chebyshev systems in the one-dimensional case. This is the first of a series of papers in this area, which solves the longstanding problem of finding a satisfactory multidimensional generalization of the classical Chebyshev systems introduced already by A. Markov more than hundred years ago.

1 History of Chebyshev systems

1.1 Developments in the Moment problem by A. Markov and M. Krein

It was namely in the Moment problem where the notion of Chebyshev systems appeared for the first time on the big stage, and provided a very natural and beautiful generalizations of the results of Gauss, Jacobi, Chebyshev, Stieltjes, Markov, and others.

The classical Moment problem is defined as follows: Find a non-negative measure $d\mu$ such that

$$\int_a^b t^j d\mu(t) = c_j \quad \text{for } j = 0, 1, \dots, N.$$

The solution of the problem includes conditions on the constants c_j providing solvability. In the case of $N = 2n - 1$ the problem has been solved by the famous *Gauss-Jacobi quadrature*; this solution is based on the orthogonal polynomials P_n (of degree n) which are orthogonal with respect to the inner product defined by

$$\langle t^j, t^k \rangle := c_{j+k}.$$

The history is well described in the book of M. Krein and A. Nudelman "The Markov Moment Problem" [5], actually based on a 1951 paper of M. Krein devoted to the ideas of Chebyshev. There it is said that A. Markov has realized that one may consider successfully the Moment problem of the type

$$\int_a^b u_j(t) d\mu(t) = c_j \quad \text{for } j = 0, 1, \dots, N,$$

where the system of **continuous** functions $\{u_j(t)\}_{j=0}^N$ represent a **Chebyshev system** in the interval $[a, b]$, i.e. any linear combination

$$u(t) = \sum_{j=0}^N \alpha_j u_j(t)$$

has no more than N zeros in $[a, b]$.

Further, by U_N we will denote the subspace of the space of continuous functions $C([a, b])$ generated by the Chebyshev system, i.e.

$$U_N := \left\{ u(t) : u(t) = \sum_{j=0}^N \alpha_j u_j(t) \right\}. \quad (1)$$

In general, in areas other than Approximation theory and Moment problem, people have tried to find those properties of the one-dimensional algebraic polynomials which make them so nice. Apparently, the Chebyshev property seems to be such.

1.2 Further developments in Approximation theory and Spline theory

Let us remind also the famous Chebyshev alternance theorem which has been proved for **Chebyshev systems**, and which one would like to see in a multivariate setting (cf. [5], chapter 9, Theorem 4.4) :

Theorem 1 (Chebyshev-Markov) *Let $f \in C([a, b])$. A necessary and sufficient condition for the element $u_0 \in U_N$ to solve problem*

$$\inf_{u \in U_N} \|f - u\|_C = \inf_{u \in U_N} \left(\max_{x \in [a, b]} |f(x) - u(x)| \right) =: \delta$$

is the existence of $N + 2$ points

$$t_1 < \dots < t_{N+2}$$

such that

$$\delta \varepsilon (-1)^j = f(t_j) - u_0(t_j) \quad \text{for } j = 1, \dots, N + 2$$

where $\varepsilon = 1$ or $\varepsilon = -1$.

What concerns other areas where Chebyshev systems have found numerous applications, one has to mention the book of [11] which contains an exhaustive consideration of spline theory where splines are piecewise functions belonging to a Chebyshev system.

2 Definitions

Let us provide some basic definitions.

Consider the system of functions $\{u_j(t)\}_{j=0}^N$ defined on some interval $[a, b]$ in \mathbb{R} and the linear space defined in (1).

Definition 2 We call the system of functions $\{u_j(t)\}_{j=0}^N$ Chebyshev (or T -system) iff for every set of constants $\{c_j\}_{j=0}^N$ and every choice of the points $t_j \in [a, b]$ with

$$t_0 < t_1 < \cdots < t_N$$

there is a unique solution $u \in U$ of the equations

$$u(t_j) = c_j \quad \text{for } j = 0, 1, \dots, N.$$

It is equivalent to say that

$$u(t_j) = 0 \quad \text{for } j = 0, 1, \dots, N$$

implies

$$u \equiv 0.$$

Proposition 3 Assume that the space $U \subset C[a, b]$ is given. Then if for some set of knots $t_j \in [a, b]$ with

$$t_0 < t_1 < \cdots < t_N,$$

and for arbitrary constants $\{c_j\}$ we have unique solution $u \in U$ of the equations

$$u(t_j) = c_j \quad \text{for } j = 0, 1, \dots, N$$

it follows that $\dim U = N + 1$.

One may formulate the above in an equivalent way:

Proposition 4 The following are equivalent

1. the system $\{u_j(t)\}_{j=0}^N$ is T -system
2. for every $u \in U_N$ the number of zeros in the interval $[a, b]$ is $\leq N$.
3. the following determinants satisfy

$$D(u; t_0, t_1, \dots, t_N) := \det \begin{bmatrix} u_0(t_0) & u_0(t_1) & \cdots & u_0(t_N) \\ u_1(t_0) & u_1(t_1) & \cdots & u_1(t_N) \\ \cdot & \cdot & \cdot & \cdot \\ u_N(t_0) & u_N(t_1) & \cdots & u_N(t_N) \end{bmatrix} \neq 0$$

Definition 5 If $D(u; t_0, t_1, \dots, t_N) \geq 0$ then $\{u_j(t)\}_{j=0}^N$ is called T_+ -system.

2.1 Examples

The classical polynomials, the trigonometric polynomials in smaller intervals $[0, 2\pi]$!

1. the system

$$\{u_j(t) = t^{\alpha_j}\}_{j=0}^N \quad \text{on subintervals of } [0, \infty]$$

2. the system

$$u_j(t) = \frac{1}{s_j + t} \quad \text{for } 0 < s_0 < s_1 < \dots < s_N \quad \text{on closed subint. of } (0, \infty).$$

3. the system

$$u_j(t) = e^{-(s_j-t)^2} \quad \text{for } 0 < s_0 < s_1 < \dots < s_N \quad \text{on } (-\infty, \infty).$$

4. if $G(s, t)$ is the Green function associated with the operator

$$Lf = -\frac{d}{dx} \left(p \frac{df}{dx} \right) + qf$$

and some boundary conditions on the interval $[a, b]$, then the system

$$u_j(t) = G(s_j, t) \quad \text{for } 0 < s_0 < s_1 < \dots < s_N \quad \text{on closed subint. of } [a, b]$$

For further examples, see the monograph of M. Krein and A. Nudel'man [5] and of S. Karlin and W. Studden, [6].

3 Extended systems

We usually work with differentiable systems of functions, and we count the multiplicities of the zeros.

Definition 6 Let $\{u_j(t)\}_{j=0}^N \in C^N[a, b]$ be a T -system. We call it *Extended Chebyshev system* (ET -system) if in U_N we may uniquely solve the **Hermite interpolation problem**:

$$u^{(k)}(t_j) = c_{j,k} \quad \text{for } k = 0, 1, \dots, d_j$$

with arbitrary numbers $\{c_{j,k}\}$ where

$$\sum (d_j + 1) = N + 1.$$

It is equivalent to say that if for some $u \in U_N$ holds

$$u^{(k)}(t_j) = 0 \quad \text{for } k = 0, 1, \dots, d_k$$

then

$$u \equiv 0.$$

There are equivalent formulations with determinants and zeros:

Proposition 7 *The following are equivalent:*

1. the system $\{u_j(t)\}_{j=0}^N \in C^N[a, b]$ is an *ET*-system
2. for every $u \in U$ the number of zeros counted with the multiplicities is $\leq N$.
3. the **modified** determinants (see the detailed explanation in [5], end of section 1, chapter 2, or [6]),

$$\det \begin{bmatrix} u_0(t_0) & u'_0(t_0) & \cdots & u_0^{(k_1-1)}(t_0) & u_0(t_1) & \cdots & u_0(t_N) \\ u_1(t_0) & u'_1(t_0) & \cdots & u_1^{(k_1-1)}(t_0) & u_1(t_1) & \cdots & u_1(t_N) \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ u_N(t_0) & u'_N(t_0) & \cdots & u_N^{(k_1-1)}(t_0) & u_N(t_1) & \cdots & u_N(t_N) \end{bmatrix} \neq 0$$

There is a nice **characterization** of *ET*-systems. A basic example of *ET*-system is the following: Let $w_i \in C^{N-i}[a, b]$ be positive functions on $[a, b]$ for $i = 0, 1, \dots, N$. Then the functions

$$u_0(t) = w_0(t) \tag{2}$$

$$u_1(t) = w_0(t) \int_a^t w_1(t_1) dt_1 \tag{3}$$

$$u_2(t) = w_0(t) \int_a^t w_1(t_1) \int_a^{t_1} w_2(t_2) dt_2 \cdot dt_1 \tag{4}$$

$$\dots \tag{5}$$

$$u_N(t) = w_0(t) \int_a^t w_1(t_1) \int_a^{t_1} w_2(t_2) \cdots \int_a^{t_{N-1}} w_N(t_N) dt_N \cdots dt_1 \tag{6}$$

form an *ET*-system.

Definition 8 An T -system $\{u_k\}_{k=0}^N$ is called M_+ -system if every subsystem $\{u_k\}_{k=0}^m$ for $m = 0, 1, \dots, N$ is a T_+ -system.

There is an important result which is basically due to S. Bernstein (cf. [5], Theorem 4.1 and Theorem 4.2, chapter 2, and the footnote at the end).

Theorem 9 Every T -system $\{u_k\}_{k=0}^N$ on the interval $[a, b]$ may be linearly transformed to an M_+ -system $\{v_k\}_{k=0}^n$ on the interval (a, b) .

A similar result may be proved in the differentiable case.

Corollary 10 Every ET -system in $[a, b]$ may be linearly transformed to an ECT -system in (a, b) .

The role of the ECT systems becomes clear from the following fundamental result.

Theorem 11 If the space U_N is generated by an ECT -system of order N then it has a basis $\{v_j\}_{j=0}^N$ which is representable in the above form (2)-(6). Hence, U_N is a set of solutions to the following equation

$$L_N u = 0 \quad \text{for } t \in (a, b)$$

and

$$L_N = \prod_{j=0}^N \frac{d}{dt} \frac{1}{w_j(t)}. \quad (7)$$

The proof is available in [6] (chapter 11, Theorem 1.1) and for M_+ -systems in [5], Theorems 4.1, 4.2 and 5.2, chapter 2.

4 The multivariate case – attempts

We have to note that all "brute force generalizations" of the Chebyshev systems fail. We will mention some of them.

4.1 Generalization by zero sets – theorem of Mairhuber

Apparently, the first attempt has been to generalize the Chebyshev systems by considering the set of zeros:

Definition 12 *Let K be a compact topological space. The system of functions $\{u_j\}_{j=0}^N$ is called Chebyshev of order N iff*

$$Z(u) \leq N$$

for every $u \in U$.

The following result shows that there are no non-trivial examples of multidimensional systems satisfying Definition 12 (cf. [5], chapter 2, section 1).

Theorem 13 *(Mairhuber, 1956) The only spaces K having a Chebyshev system satisfy $K \subset \mathbb{R}$ or $K \subset \mathbb{S}^1$.*

The result of Theorem 13 is intuitively clear since "general position" function in $C(K)$ has a zero set which is a subset of K of codimension 1. In particular, if $K = \mathbb{R}^2$ then the zero set is roughly speaking union of some curves, and it would be more reasonable to speculate about the number of these components then to consider Definition 12 above. Speculating in this direction in the multidimensional case, in view of the Polyharmonic Paradigm [7], one may try to replace the points on \mathbb{R} by closed surfaces, and the spaces U by solutions of Elliptic PDEs. Going further, one may obtain some interesting results if one uses spheres in the case of the polyharmonic operator Δ^N by considering the space

$$U_N = \{u : \Delta^N u = 0 \text{ in } D\}$$

for some bounded domain $D \subset \mathbb{R}^n$. In particular, one may prove that if a polyharmonic function of order N in a domain D (a function satisfying $\Delta^N u = 0$ in D) is zero on a set of n concentric spheres, then $u \equiv 0$. This result has been proved apparently a long time ago by means of the Almansi theorem, see e.g. [1]. However these hopes to try to generalize Definition 12 are only vain. They have been broken by an example which has been published apparently for a first time in 1982, by Atakhodzhaev, [1]. It shows a non-zero biharmonic function which is zero on two embedded ovals in \mathbb{R}^2 .

4.2 Generalization by Haar property

The following result belongs to A. Haar, [5]:

Theorem 14 *Let the space $U_N \subset C[a, b]$ be generated by a Chebyshev system $\{u_j\}_{j=0}^N$. Then for every $f \in C[a, b]$ the best approximation problem*

$$\min_{u \in U_N} \|f - u\|_C$$

has unique solution.

Extending this definition to the multivariate case seems to be very reasonable but the work with best approximations is very heavy and until now has not led to success.

5 Systems to be generalized

One needs a new point of view on the Chebyshev systems which would make them generalizable to several dimensions. We propose the point of view of boundary value problems: We consider a special class of *ET*-systems which are "generalizable".

Definition 15 *We say that the system $\{u_j\}_{j=0}^{2N-1} \in C^{2N-1}[a, b]$ is a **Dirichlet type Chebyshev system**, or *DT*-system, in the interval $[a, b]$ if for every two points α and β in $[a, b]$ and for every set of constants c_j and d_j we are able to **solve uniquely** the following **interpolation problem**, with $u \in U_N$,*

$$\begin{aligned} u^{(k)}(\alpha) &= c_k & \text{for } k = 0, 1, \dots, N-1 \\ u^{(k)}(\beta) &= d_k & \text{for } k = 0, 1, \dots, N-1 \end{aligned}$$

Remark 16 *Obviously, all *ET*-systems are *DT*-systems but not vice versa.*

Let us state an equivalent formulation which we are going to mimic in the multivariate case.

Proposition 17 *The system $\{u_j\}_{j=0}^{2N-1} \in C^\infty[a, b]$ is a **Dirichlet type Chebyshev system** in the interval $[a, b]$ iff for every two points α and β in $[a, b]$, for every set of constants c_j and d_j , and for every $\varepsilon > 0$, we*

are able to **solve** the following approximate interpolation problem, with $u \in U_{2N} = \text{span} \{u_j\}_{j=0}^{2N-1}$,

$$|u^{(k)}(\alpha) - c_k| < \varepsilon \quad \text{for } k = 0, 1, \dots, N-1 \quad (8)$$

$$|u^{(k)}(\beta) - d_k| < \varepsilon \quad \text{for } k = 0, 1, \dots, N-1 \quad (9)$$

The proof is evident since the space U_{2N} is finite-dimensional.

6 The multivariate case

In the present section we will provide a multivariate generalization to the *DT*-systems of Definition 15.

One might use the properties of the Dirichlet type Chebyshev system provided in Proposition 17 as possible way to make a multivariate generalization. However we would like to have also the properties exposed by Theorem 11 as well.

Remark 18 *It is expected that a further research would prove equivalence between the solvability of problem (8)-(9) and the representation of the space U_N as a set of solutions to an equation $L_N u = 0$.*

Whatever the definition of *Multidimensional Chebyshev systems*, we would like to retain the properties in Theorem 11 and Proposition 17. In general, it would need in the future to make a proper refinement of these properties which would make then into two equivalent sets of conditions. However at the present moment we will restrict ourselves with some special though sufficiently wide generalization.

We consider a subspace of functions U with $U \subset C^\infty(D)$ for some bounded domain $D \subset \mathbb{R}^n$ such that its boundary ∂D is infinitely smooth, and assume that D locally "lies on one side of the boundary". These are the usual conditions for the solvability of Elliptic Boundary Value problems, see e.g. [9]. For simplicity assume that D is connected and simply connected as well.

In the following definition we will mimic the properties of the Chebyshev systems provided in Theorem 11.

Definition 19 *We will say that the elliptic operator P_{2N} of order $2N$ defined in the domain D , is **factorizable** if there exist N uniformly strongly elliptic*

operators $Q_2^{(j)}$ of second order, defined in the domain D , and satisfying the following properties:

1. Every operator $Q_2^{(j)}$ satisfies the maximum principle in \overline{D} .
2. Every operator $Q_2^{(j)}$ satisfies condition $(U)_s$ for uniqueness in the Cauchy problem in the small.¹
3. The following equality holds

$$P_{2N} = \prod_{j=1}^N Q_2^{(j)} = Q_2^{(1)} Q_2^{(2)} \cdots Q_2^{(N)}.$$

However in the next definition we will mimic the interpolation properties of the Chebyshev systems exposed by Proposition 17.

Definition 20 We say that the space U satisfies the **Multivariate BVP Interpolation** of order N iff the following conditions hold:

1. The **approximate solvability of BVP** on subdomains holds in the following sense: Let the "boundary differential operators" $B_j(x; D_x)$, $j = 1, 2, \dots, N$, with smooth coefficients and of orders $\leq 2N$, defined in \overline{D} be given. Let D_1 be an arbitrary subdomain of D with $D_1 \subset D$, and such that D_1 satisfies the above conditions as D , and also $D \setminus D_1$ has only non-compact connected components. Let $c_j \in C^\infty(\partial D_1)$ for $j = 0, 1, \dots, N - 1$. Then for every $\varepsilon > 0$ there exists an element $u \in U$ such that

$$|B_j u - c_j(x)| \leq \varepsilon \quad \text{for all } x \in \partial D_1, \text{ and } j = 1, \dots, N. \quad (10)$$

In the case of the whole domain, i.e. $D_1 = D$, inequality (10) holds with $\varepsilon = 0$.

2. The **unique solvability of BVP** on subdomains holds: If for some $u \in U$ holds

$$B_j u(x) = 0 \quad \text{for all } x \in \partial D_1, \text{ for } j = 1, \dots, N, \quad (11)$$

then

$$u \equiv 0.$$

¹The differential operator P satisfies condition $(U)_s$ for uniqueness in the Cauchy problem in the small in G provided that if G_1 is a connected open subset of G and $u \in C^r(G_1)$ is a solution to $P^*u = 0$ and u is zero on a non-empty subset of G_1 then u is identically zero. Elliptic operators with analytic coefficients satisfy this property (cf. [?], part II, chapter 1.4; [3], p. 402).

Remark 21 *In the case of the domain D we have unique solvability of the Elliptic BVP since we may take $\varepsilon = 0$!*

Remark 22 *For Theorem 23 below it is important to note that if an operator B_j is non-characteristic in every direction then it is elliptic, see [9], (Definition 1.4 in chapter 2, section 1.4).*

We would like for a **Multidimensional Chebyshev space** U to satisfy analogs to both Theorem 11 and Proposition 17. In this respect we may prove the following theorem which shows that it is better to define the **Multidimensional Chebyshev spaces** by means of Definition 19 than by means of Definition 20.

Theorem 23 *Let the elliptic operator P_{2N} in the domain D be **factorizable** by Definition 19. Then the space*

$$U_N = \{u \in H^{2N}(D) : P_{2N}u = 0 \text{ in } D\}$$

*satisfies the **Multivariate BVP interpolation** of Definition 20.*

Proof. We choose the following boundary operators

$$B_j = \prod_{i=N-j+2}^N Q_2^{(i)} \quad \text{for } j \geq 1,$$

and in particular, $B_1 = id$.

The uniqueness (11) follows by induction in N , from item 1 in Definition 19.

Item 1 in Definition 20 follows inductively in N . Let us consider for simplicity the case $N = 2$. Let $u_0 \in H^4(D_1)$ satisfy $P_4u_0 = Q^{(1)}Q^{(2)}u_0 = 0$ in D_1 . Let us put

$$w_0 = Q^{(2)}u_0$$

By the $(U)_s$ property of the operator $Q^{(1)}$, it follows that a **Runge** type theorem holds, namely, for every $\varepsilon > 0$ there exists a solution $w_\varepsilon \in H^2(D)$ of the equation $Q^{(1)}w_\varepsilon = 0$ in D and

$$\|w_\varepsilon - w_0\| \leq \varepsilon,$$

(cf. [8], [10], [3]). Now we want to find a solution $u_\varepsilon \in H^4(D)$ such that $Q^{(2)}u_\varepsilon = w_\varepsilon$ in D and $\|u_\varepsilon - u_0\|_{D_1} \leq 2\varepsilon$. But here we use the $(U)_s$ property of

the operator $Q^{(2)}$ since we compare $Q^{(2)}u_0 = w_0$ and $Q^{(2)}u_\varepsilon = w_\varepsilon$ and we know that $\|w_\varepsilon - w_0\| \leq \varepsilon$. By the $(U)_s$ property as above we may find a solution u_ε to the non-homogeneous equation which satisfies $\|u_\varepsilon - u_0\|_{D_1} \leq 2\varepsilon$.

This ends the proof.

■

Remark 24 *An alternative reference for the Runge-Lax-Malgrange type theorem is [4] (Theorem 4.4.5), [12] (Section 2.3, Theorems 2.1, 2.2, and Theorem 2.4 for which it is mentioned there, that it was formulated by F. Browder with error).*

7 Examples

For some integer $N \geq 1$ let us consider the space

$$U := \{u : \Delta^N u(x) = 0 \quad \text{in } D\}.$$

Proposition 25 *The space U satisfies Definition 19.*

The proof follows directly after we define the operators

$$B_j = \Delta^{j-1} \quad \text{for } j = 1, 2, \dots, N.$$

7.1 The one-dimensional case

We still have to check that the DT -systems defined in Definition 15 satisfy the multivariate counterpart in Definition 20.

Proposition 26 *Let us assume that in Definition 20 the space dimension is $n = 1$. Then the set U coincides with a **Dirichlet type Chebyshev system** DT of order N from Definition 15.*

Proof. Indeed, let us take the set $D = [a, b]$ and apply the interpolation property to the case $D_1 = D$. Then we know that for all constants c_j and d_j we have unique solvability of the problem

$$\begin{aligned} u^{(k)}(a) &= c_k & \text{for } k = 0, 1, \dots, N-1 \\ u^{(k)}(b) &= d_k & \text{for } k = 0, 1, \dots, N-1. \end{aligned}$$

Hence, U is $2N$ dimensional; we may take the solution v_j to the problem

$$\begin{aligned} u^{(k)}(a) &= \delta_{j,k} & \text{for } k = 0, 1, \dots, N-1 \\ u^{(k)}(b) &= 0 & \text{for } k = 0, 1, \dots, N-1 \end{aligned}$$

and the solution w_j to the problem

$$\begin{aligned} u^{(k)}(a) &= 0 & \text{for } k = 0, 1, \dots, N-1 \\ u^{(k)}(b) &= \delta_{j,k} & \text{for } k = 0, 1, \dots, N-1 \end{aligned}$$

and we make a basis for U . The approximate solvability of the Dirichlet problem for $D_1 \subset [a, b]$ implies now the exact solvability since U is finite-dimensional. Indeed, we will take a sequence of solutions $u_\varepsilon(t)$ and the limit.

■

8 Remarks

1. In the case of Multidimensional Chebyshev Systems we need the approximate solvability of the Dirichlet problem since we have infinite-dimensional spaces, and there is no equivalence between the uniqueness and the existence but we have a substitute which is the Fredholm property of the regular Elliptic BVPs. The one-dimensional Proposition 17 traces the smooth path for the multivariate generalization.

Acknowledgement 27 *The present research has been partially sponsored by project DO – 02 – 275, 08.12.2008 with the Bulgarian NSF.*

References

- [1] M. Atakhodzaev, Ill-posed internal boundary value problems for the biharmonic equation, VSP, 2002.
- [2] L. Bers, F. John and M. Schechter, Partial Differential Equations, Interscience, New York, 1964.
- [3] Felix Browder, Approximation by Solutions of Partial Differential Equations, American Journal of Mathematics, vol. 84, no. 1, p. 134, 1962; Functional analysis and partial differential equations. II, Mathematische Annalen, vol. 145, no. 2, pp. 81-226, 1962.

- [4] Hormander, L., The Analysis of Linear Partial Differential Operators I, Springer-Verlag, 1990.
- [5] Krein, M. , Nudelman A., The Markov Moment problem and Extremal problems, AMS translation from the Russian edition of 1973.
- [6] S. Karlin, W. Studden, Tchebycheff Systems: with applications in analysis and statistics, Intersci. Publ., 1968.
- [7] Kounchev, O., Multivariate Polysplines, Academic Press, San Diego, 2001.
- [8] P. Lax, A stability theorem for solutions of abstract differential equations and its application to the study of local behavior of solutions of elliptic equations, Comm. Pure Appl. Math. 9 (1956) 747–766.
- [9] J.-L. Lions and E. Magenes, Non-homogeneous Boundary Value Problems and Applications, Springer, Berlin-Heidelberg, 1970.
- [10] B. Malgrange, Existence et approximation des solutions des equations aux derivees partielles et des equations de convolution, Ann. Inst. Fourier (Grenoble) 6 (1955–1956) 271–355.
- [11] L. Schumaker, *Spline Functions: basic theory*, Academic Press, NY, 1983.
- [12] Tarkhanov, N., Approximation on compact sets by solutions of systems with surjective symbol, Russian Mathematical Surveys , 48(5) (1993) : 103.