

State Transfer in Highly Connected Networks and a Quantum Babinet Principle

D. I. Tsomokos¹, M. B. Plenio^{2,3}, I. de Vega⁴ and S. F. Huelga¹

¹Quantum Physics Group, STRI, School of Physics, Astronomy & Mathematics, University of Hertfordshire, Hatfield AL10 9AB, United Kingdom

²Institute for Mathematical Sciences, Imperial College London, London SW7 2PG, UK

³QOLS, Blackett Laboratory, Imperial College London, London SW7 2BW, UK and

⁴Max-Planck-Institut für Quantenoptik, Hans-Kopfermann-Str. 1, Garching D-85748, Germany

(Dated: February 16, 2019)

The transfer of a quantum state between distant nodes of planar networks, is considered. The fidelity of state transfer is calculated as a function of the connectivity in networks that are described by regular graphs. It is shown that perfect state transfer is achieved in a network of size N , whose structure is that of a $\frac{N}{2}$ -cross polytope graph, if N is a multiple of 4. If N is even, but not a multiple of 4, then almost perfect state transfer is achieved for larger networks of this kind. The result is reminiscent of the Babinet principle of classical optics, for which a quantum analog is provided. The quantum Babinet principle allows for the identification of complementary graphs that lead to the same fidelity of state transfer, in analogy with complementary screens providing identical diffraction patterns, and can be efficiently exploited to assess the performance of a priori unrelated graphs.

The dynamics of quantum many-body systems offers a rich variety of features. Often this dynamics is considered in 1-D chains, where it is amenable to exact analytical treatment in some cases [1] and, in others, efficient numerical methods have been developed for its simulation [2]. For more complicated lattice structures, henceforth referred to as graphs, few analytical treatments are known. An interesting problem that arises in this context is the interplay between the dynamics of quantum many-body systems and the properties of the underlying graph, which determines the interaction structure of the complex many-body system. A variety of interesting phenomena, two examples of which are perfect state transfer [3, 4] and the possibility of deciding the graph isomorphism problem [5], have recently been explored in such systems.

In the case of graphs with uniform nearest neighbour coupling, *perfect* state transfer (PST) has so far been proven possible only with rings of $N = 4$ spins, chains of $N = 2$ or $N = 3$ spins and with Cartesian products of such graphs, the so-called one-link and two-link hypercubes [4]. For larger networks, it appears that increasing the number of spins and the connectivity (number of interactions) tends to compromise the transmission of quantum information [6, 7]. In the static case it has been shown that higher connectivity and associated monogamy constraints frustrate the system and affect its quantum correlations [8]. On the other hand, if natural interactions are abandoned in favor of particular coupling schemes, in which only nearest neighbors interact and the interaction strength depends on their position relative to a fixed point, then perfect state transfer is possible in spin chains with large N [3, 4, 9, 10].

We investigate here whether it is possible to transfer perfectly a quantum state between two distant vertices of a highly-connected spin network, in which the interactions between spins are both permanent and homoge-




FIG. 1: (Color online) A circulant spin network can be used to transfer quantum states from A to B. In a network of $N = 6$ spins there are three possible configurations with connectivity (a) $C = 1$, (b) $C = 2$, and (c) $C = 3$, as shown. Network (b) is a 3-cross polytope graph (CPG).

neous. We show that in contrast to what is expected, PST it is indeed feasible for a highly connected network with arbitrarily large N . This result is obtained by using a unique configuration, namely a two-dimensional graph of the $\frac{N}{2}$ -cross polytope [11], which is dual to the hypercube in $\frac{N}{2}$ dimensions and isomorphic to a type of circulant graph [12]. After demonstrating this possibility for two-dimensional graphs, we analyze the physical mechanisms behind PST and make an analogy between this dynamics and a well known result in classical optics.

We begin by considering N spins- $\frac{1}{2}$ situated along a circle, as shown in Fig. 1. It is understood that if two spins are interacting, a line is drawn between them. The result is a graph $\mathcal{G} = (V, E)$; the vertices $V(\mathcal{G})$ represent the spin sites and the edges $E(\mathcal{G})$ represent pairwise interactions. The necessary information about the graph \mathcal{G} is contained in its adjacency matrix, $A(\mathcal{G})$, whose elements are given by $A_{ij} = 1$ if $\{i, j\} \in E(\mathcal{G})$ and are zero otherwise. We consider Hamiltonians of the form ($\hbar = 1$)

$$\mathcal{H} = \sum_{k=1}^N \omega_k \sigma_k^+ \sigma_k^- + \sum_{k \neq l} J_{k,l} (\sigma_k^- \sigma_l^+ + \sigma_k^+ \sigma_l^-), \quad (1)$$

where σ_k^+ (σ_k^-) are the raising and lowering operators for site k , $\hbar\omega_k$ is the local site excitation energies and $J_{k,l}$ denotes the hopping rate of an excitation between the sites k and l . The dynamics in this system preserves the total excitation number, defined by $\mathcal{N} = \sum_{k=1}^N \sigma_k^+ \sigma_k^-$. During dynamical evolution the state of the network, $|\Psi(t)\rangle = \exp(-i\mathcal{H}t)|\Psi_0\rangle$, where $|\Psi_0\rangle$ is the initial state, always remains in the same excitation sector because $[\mathcal{H}, \mathcal{N}] = 0$. In what follows we restrict our attention to the single-excitation sector, for simplicity. In this subspace the Hamiltonian of the system is equal to the adjacency matrix of the underlying graph, $\mathcal{H} = A(\mathcal{G})$, provided that the spin-spin interactions are homogeneous. Deviations due to engineering errors in the interactions are also examined later on. The network is prepared in the state $|\Psi_0(\mathbf{j})\rangle := |0_1 0_2 \dots 1_j \dots 0_N\rangle$, where only spin j is excited. The propagation of an arbitrary state $\alpha|0_j\rangle + \beta|1_j\rangle$, where $|\alpha|^2 + |\beta|^2 = 1$, is equivalent to the propagation of the state $|1_j\rangle$ (since the $+1$ eigenstate of Z_j , $|0_j\rangle$, does not evolve under \mathcal{H}). The aim is to transfer the excitation from j to $N/2 + j$, that is, to the vertex that is diametrically opposite from j across the ring – hence we initially consider that N is even. The state transfer is quantified by the fidelity

$$F(t) := |\langle \Psi_0(\mathbf{N}/2 + \mathbf{j}) | \exp(-i\mathcal{H}t) | \Psi_0(\mathbf{j}) \rangle|. \quad (2)$$

Perfect state transfer is achieved at a certain time t_0 if and only if $F(t_0) = 1$.

We now ask, ‘How is the fidelity of quantum state transfer influenced by the connectivity of a network?’ The connectivity, $C(\mathcal{G})$, is defined here as the number of edges that are incident on a vertex, counting only within the half-disc defined by that vertex and the opposite one. The extreme cases are those of a ring ($C = 1$) and a fully-connected network ($C = N/2$), but in general we have $C = 1, 2, \dots, N/2$ (see Fig. 1). Before we analyze this question analytically we consider numerically the fidelity $F(t)$ of Eq. (2) for $t \in [0, \Delta t]$, given the number of spins N and the connectivity C . The maximum fidelity, $\max(F_{\Delta t})$, is then determined for the interval Δt . Under these conditions it is observed in Fig. 2(a) that the fidelity is a non-monotonic and rather complicated function of the connectivity. However, it displays remarkable behavior for $C = \frac{N}{2} - 1$, which corresponds to a $2k$ -cross polytope graph (CPG) with $N = 4k$ spins, where k is a positive integer [11]. In this case perfect state transfer is achieved at $t_0 = \frac{\pi}{2} + n\pi$, that is $F(\frac{\pi}{2} + n\pi) = 1$, where $n \geq 0$ is an integer. For $k = 1$ we recover the known result [13] for a ring with $N = 4$. In Ref. [12] it was shown that circulant graphs of odd order *do not allow* perfect state transfer (so our choice of even N is justified) and, moreover, it was left as an open question if there exist circulant graphs of even order with $N > 4$ that support perfect state transfer. Our results show that such graphs do indeed exist: the $2k$ -CPG is isomorphic to the circulant graph $\text{Ci}_{4k}(1, 2, \dots, 2k - 1)$. In these networks every

spin interacts with every other spin, except for the one that is diametrically opposite to it – see Fig. 1(b) for an example. The appropriate choice of Δt is made by comparing trial values with the occurrence time of the first peak in the evolution of the fidelity for a spin ring (this evolution is shown in Fig. 2(b)). In Fig. 2(c) we show the evolution of the fidelity for a network with connectivity $C = \frac{N}{2} - 1 = 99$ (i.e., for a 100-CPG). It is seen that the fidelity becomes equal to 1 at $t_0 = \pi/2$.

For $N = 4k + 2$, we obtain a $(2k + 1)$ -CPG network. Here we find that the transfer is not perfect but rapidly improves with increasing N and that the times at which it occurs are close to $\frac{(2n+1)\pi}{2}$. Hence, provided the networks are large, the transfer of a quantum state is practically perfect on a $\frac{N}{2}$ -CPG network for all even N . These results will now be supported by analytical arguments.

Analytical results.— We now discuss the underlying mechanisms that make possible the perfect (for $N = 4k$) or near perfect state transfer (for $N = 4k + 2$) in these configurations and draw an analogy with the well known Babinet principle in classical optics.

The Hamiltonian H_{CPG} of the cross polytope graphs discussed before are of the form eq. (1) with $\omega_k = 0$ and

$$J_{k,l} = (1 - \delta_{l, \frac{N}{2} + k})(1 - \delta_{k,l}) = J_{l,k}.$$

The numerical observations may be obtained by direct analytical diagonalization of the Hamiltonian but a more transparent approach, which points in the direction of a more general principle, is also presented.

Let us denote by H_f the Hamiltonian corresponding to a fully connected graph, i.e. one where $J_{k,l} = (1 - \delta_{k,l}) = J_{l,k}$, and by H_{pair} the Hamiltonian where all opposite




FIG. 2: (Color online) (a) Maximum fidelity in the interval $[0, \Delta t = 100]$ against connectivity for a network of size $N = 200$ and homogeneous interactions. (b) Fidelity against time for the simple ring network. (c) Fidelity against time for the 100-cross polytope graph network.

pairs are connected, i.e. $J_{k,l} = \delta_{l, \frac{N}{2}+k} = J_{l,k}$ for $k = 1, \dots, \frac{N}{2}$.

We start by noting that $H_{CPG} = H_f - H_{pair}$ and that, furthermore, $[H_f, H_{pair}] = 0$. Therefore we find

$$e^{-iH_{CPG}t} = e^{iH_{pair}t}e^{-iH_f t}$$

As $H_{pair} = \sum_{k=1}^{N/2} (|k\rangle\langle \frac{N}{2} + k| + |\frac{N}{2} + k\rangle\langle k|)$ only couples opposite sites its dynamics is very simple. We find

$$e^{iH_{pair}t} = \mathbb{1} \cos t + i \sum_{k=1}^{N/2} \left(|k\rangle\langle \frac{N}{2} + k| + |\frac{N}{2} + k\rangle\langle k| \right) \sin t.$$

In particular, after a time $t = \frac{\pi}{2} + n\pi$ a starting state $|k\rangle$ will have been transformed to $(-1)^n i |\frac{N}{2} + k\rangle$.

To determine the dynamics of H_{CPG} we need to consider H_f . Fortunately, again this is a very simple dynamics. As it may be written $H_f = N|+\rangle\langle +| - \mathbb{1}$ with $|+\rangle = \frac{1}{\sqrt{N}} \sum_{k=1}^N |k\rangle$ we find

$$e^{-iH_f t} = [|+\rangle\langle +| e^{-iNt} + (\mathbb{1} - |+\rangle\langle +|)] e^{it}$$

Therefore, an initial state $|k\rangle$ is mapped onto itself, up to a global phase, under $e^{-iH_f t}$ when $Nt = 2\pi k$ with $k \in \mathbb{N}$.

As a consequence, the dynamics due to H_{CPG} allows for perfect state transfer if both $Nt = 2\pi k$ and $t = \frac{\pi}{2}$ are satisfied for the same t . This implies the condition

$$N = 4k$$

and hence explains the perfect state transfer for $2k$ -cross polytopes. As stated earlier, this could of course have been obtained from the direct diagonalisation of $H_{N/2-1}$ but the above clearly reveals the origin of the condition for perfect transfer. For $N = 4k + 2$ the analysis above immediately applies and shows that we do not have perfect state transfer at $t = \frac{\pi}{2}$.

More generally we can easily find the fidelity of transfer for $t = \frac{\pi}{2}$. Starting with $|k\rangle$ and using $e^{iH_{pair}t}|+\rangle = e^{it}|+\rangle$ we find at $t = \frac{\pi}{2}$ the state

$$\begin{aligned} e^{iH_{pair}t} e^{-iH_f t} |k\rangle &= \\ e^{iH_{pair}t} \left[\frac{1}{\sqrt{N}} |+\rangle e^{-iN\pi/2} + (|k\rangle - \frac{1}{\sqrt{N}} |+\rangle) \right] e^{i\pi/2} \\ &= - \left[\frac{1}{\sqrt{N}} |+\rangle e^{-iN\pi/2} + (|\frac{N}{2} + k\rangle - \frac{1}{\sqrt{N}} |+\rangle) \right]. \end{aligned}$$

Then

$$\begin{aligned} F &= \left| \langle \frac{N}{2} + k | e^{iH_{pair}t} e^{-iH_f t} |k\rangle \right|^2 = \left| \frac{1}{N} (e^{-iN\pi/2} - 1) + 1 \right|^2 \\ &= 1 - \frac{2}{N} \left(1 - \frac{1}{N} \right) \left[1 - \cos \frac{N\pi}{2} \right]. \end{aligned}$$

For $N = 4k$ we recover $F = 1$, while for $N = 4k + 2$ we find that $F = (1 - 2/N)^2$. Therefore, as $N \rightarrow \infty$, the fidelity approaches 1 and we obtain almost PST.




FIG. 3: (Color online) Illustration of Babinet's principle in an optical set up with Fraunhofer conditions. (a) An unobstructed plane wave is focused by a lens L and produces a diffraction pattern of amplitude $A(\vec{r})$ at the screen S . Away from the central spot it is $A(\vec{r}) \sim 0$. (b) Diffraction patterns resulting from complementary screens s and \bar{s} . Away from the center, the diffracted amplitudes obey $A_s(\vec{r}) = -A_{\bar{s}}(\vec{r})$ and the diffracted intensities should therefore be the same. Complementary apertures play the role of CPG and $pair$ graphs in quantum state transfer. Increasing \vec{r} corresponds to increasing the number of nodes N .

A Quantum Babinet Principle.— These results provide a clear insight into the basic mechanisms that facilitate state transfer in these systems. The key issue is the realization that a fully connected network in which some couplings $J_{k,l}$ are removed, can behave similarly to an initially unconnected network which is supplemented with the very same $J_{k,l}$ links. This result is in fact reminiscent of the Babinet principle of classical optics [14], which is illustrated in Fig. 3. At any point downstream from two complementary screens s and \bar{s} , whose opaque and transparent areas are swapped, the sum of the two diffracted amplitudes $A_s(\vec{r}) + A_{\bar{s}}(\vec{r})$ equals the amplitude diffracted from the unobstructed plane wave $A(\vec{r})$. Away from the central spike, this amplitude is zero and therefore $A_s(\vec{r}) = -A_{\bar{s}}(\vec{r})$, which leads to Babinet's prediction of identical diffracted light fields for complementary apertures.

In our context of state transfer through connected networks, the situation is similar in the sense that $e^{-iH_{CPG}t} e^{-iH_{pair}t} = e^{-iH_f t}$ because H_{CPG} and H_{pair} commute and that $e^{-iH_f t}$ for specific times t equals the identity (in the optical setting this is the situation when all incident light emerges unaffected). Of course in the quantum setting we have the added problem that $e^{-iH_{CPG}t} e^{-iH_{pair}t} \neq e^{-i(H_{CPG}+H_{pair})t}$ in general. The analog of the Babinet theorem does hold however for much more general settings than just that of commuting H_{CPG} and H_{pair} . Indeed, as before, let us assume that $H_f = N\mathbb{P}$ where $\mathbb{P} = |+\rangle\langle +|$. For a sequence of H_N that satisfies $\|\mathbb{P}H_N(\mathbb{1} - \mathbb{P}) + (\mathbb{1} - \mathbb{P})H_N\mathbb{P}\| = O(\frac{1}{\sqrt{N}})$ we compare the dynamics of H_N and $H_f - H_N \equiv H_c$ in the limit of large N . The following argument is not fully rig-

orous in that it does not provide detailed error estimates but these may be provided in a more detailed analysis.

Let us consider the dynamics under H_c in an interaction picture with respect to H_f when this Hamiltonian becomes time-dependent

$$H_{c,I}(t) = e^{iH_f t}(H_c - H_f)e^{-iH_f t} = -e^{iH_f t}H_N e^{-iH_f t}.$$

The corresponding time-evolution operator from t_1 until t_2 in the interaction picture will be denoted by $U_I(t_1 \rightarrow t_2)$. Now we note that $H_{c,I}(t)$ may contain rapidly oscillating terms (those coupling subspace defined by \mathbb{P} to the subspace defined by $\mathbb{1} - \mathbb{P}$) thanks to the action of $e^{iH_f t}$. These rapidly oscillating terms may be neglected for large N leading to a correction of order $1/N$ in the dynamics. Hence, we find

$$\begin{aligned} H_{c,I}(t) &\cong -(\mathbb{1} - \mathbb{P})H_N(\mathbb{1} - \mathbb{P}) - \mathbb{P}H_N\mathbb{P} \\ &= -H + \mathbb{P}H_N(\mathbb{1} - \mathbb{P}) + (\mathbb{1} - \mathbb{P})H_N\mathbb{P} \end{aligned}$$

As we had assumed earlier that $|\mathbb{P}H_N(\mathbb{1} - \mathbb{P}) + (\mathbb{1} - \mathbb{P})H_N\mathbb{P}|$ is of order $1/\sqrt{N}$ we find that $H_{c,I}(t)$ is well approximated by $-H_N$ up to corrections that decrease with increasing N . Hence $U_I(0 \rightarrow t) \approx e^{iH_N t}$, and we find that

$$e^{-iH_c t} = e^{-iH_f t}U_I(0 \rightarrow t) \cong e^{-iH_f t}e^{iH_N t}.$$

Now we consider the transfer fidelity from state $|k\rangle$ to $|\frac{N}{2} + k\rangle$, as an example. We find

$$\begin{aligned} \langle \frac{N}{2} + k | e^{-iH_c t} | k \rangle &\cong \langle \frac{N}{2} + k | e^{-iH_f t} e^{iH_N t} | k \rangle \\ &= \langle \frac{N}{2} + k | e^{iH_N t} | k \rangle + \langle \frac{N}{2} + k | + \rangle \langle + | e^{iH_N t} | k \rangle (e^{iNt} - 1) \\ &\cong \langle \frac{N}{2} + k | e^{iH_N t} | k \rangle \end{aligned}$$

where the difference decreases with increasing N . Therefore, the transition amplitudes according to the dynamics under $-H_N$ and H_c are asymptotically (in N) equal. Note that for a real Hamiltonian H_N we have $(\langle \frac{N}{2} + k | e^{iH_N t} | k \rangle)^* = \langle \frac{N}{2} + k | e^{-iH_N t} | k \rangle$ so that

$$|\langle \frac{N}{2} + k | e^{-iH_c t} | k \rangle| \cong |\langle \frac{N}{2} + k | e^{-iH_N t} | k \rangle|$$

again with an error that decreases with increasing N . This is the quantum Babinet principle.

Influence of disorder.— After presenting a theoretical framework allowing for PST and explaining its underlying physical mechanisms, we conclude with an analysis of realistic engineering errors in the interactions of a $2k$ -CPG network in order to assess the robustness of a possible experimental implementation. We take into account two types of errors: (i) disorder in the interactions, and (ii) random breaking of interactions. In the first case we assume that if p and q are interacting (i.e., if $J_{pq} = 1$

when the system is homogeneous) then the interaction strength can take any value in the interval $[1 - \delta, 1 + \delta]$, with equal probability. The amount of disorder is thus quantified by $\delta \in [0, 1]$. In the second case we assume that some interactions are randomly broken, that is, J_{pq} vanishes for a fixed number of pairs (p, q) . The number of broken interactions is $B \in [0, 1]$, given as a ratio to the total number of interactions in the network. The main results of Monte-Carlo simulations on $\frac{N}{2}$ -CPG networks with $N = 40, 80, 120, 200, 400$ spins, are as follows. For type-(i) errors we find that disorder up to $\delta = 0.02$ allows for almost perfect state transfer in smaller networks ($N < 100$). In particular, the maximum fidelity is strictly greater than 0.99, on average, with a worst-case fidelity peak of over 0.98 in the case of $N = 40$; while for $N > 100$ the average maximum fidelity is over 0.95 for disorder that is less than 2%. For type-(ii) errors we find that the random breaking of very few bonds, so that $B < 0.001$, still allows for very high quality state transfer, where the maximum fidelity is larger than 0.95, on average. However, the value of the worst-case fidelity peak fluctuates considerably on individual cases, depending on the positions of the broken bonds.

In this connection, let us illustrate the usefulness of the quantum Babinet principle in the context of transport of excitations through noisy connected graphs, a setting that has recently been introduced independently in [15] and [16]. Initially all population resides in a given site and we evaluate how much population may be transferred asymptotically to a selected target site. To this end, we let the target site be attached to a sink to which the population is transferred irreversibly. We want to analyze whether the presence of local dephasing can assist the excitation transfer. If the sink is attached to site $\frac{N}{2} + 1$, then the Babinet principle implies that the time evolution is essentially that of a system where only the opposite two sites are coupled, and we recover a situation for which it was proven in [16] that no dephasing enhanced transport is possible [17].

Summary.— We have shown that perfect state transfer is achieved in a network of size N , whose structure is that of a $\frac{N}{2}$ -cross polytope graph, if N is a multiple of 4. If N is even, but not a multiple of 4, then almost perfect state transfer is achieved for larger networks of this kind, so that F approaches 1 for $N \rightarrow \infty$. These results can be interpreted in terms of a quantum Babinet principle, which establishes the conditions required for having *complementary* graphs leading to same fidelity of state transfer, in analogy with the classical situation of obtaining identical diffraction patterns from complementary screens. As shown in various examples, invoking Babinet's principle alone can simplify the analysis of the performance of connected networks and therefore become a useful tool in tackling a variety of problems in quantum information theory.

Acknowledgements.— This work was supported by the

EU via Integrated Projects QAP and SCALA, STREP actions CORNER and HIP and the EPSRC through the QIP-IRC. DIT holds an EPSRC Postdoctoral Fellowship in Theoretical Physics (EP/D065305/1). MBP holds a Royal Society Wolfson Research Merit Award. I.D.V. acknowledges support from Ministerio de Educacion y Ciencia, Spain. We thank Neil Oxtoby for carefully reading the manuscript.

-
- [1] M. Takahashi, *Thermodynamics of One-Dimensional Solvable Models*, Cambridge University Press, 2005.
- [2] S. R. White, Phys. Rev. Lett. **69**, 2863 (1992).
- [3] M. B. Plenio, J. Hartley and J. Eisert, New. J. Phys. **6**, 36 (2004).
- [4] M. Christandl, N. Datta, A. Ekert and A. J. Landahl, Phys. Rev. Lett. **92**, 187902 (2004); M. Christandl, N. Datta, T. C. Dorlas, A. Ekert, A. Kay, and A. J. Landahl, Phys. Rev. A **71**, 032312 (2005).
- [5] K. Audenaert, Ch. Godsil, G. Royle and T. Rudolph, Journal of Combinatorial Theory **97**, 74 (2007); E-print arXiv:math/0507251.
- [6] T. J. Osborne and N. Linden, Phys. Rev. A **69**, 052315 (2004).
- [7] M. Paternostro, G. M. Palma, M. S. Kim and G. Falci, Phys. Rev. A **71**, 042311 (2005).
- [8] A. Ferraro, A. García-Saenz and A. Acín, Phys. Rev. A **76**, 052321 (2007).
- [9] M. A. Jafarizadeh and R. Sufiani, Phys. Rev. A **77**, 022315 (2008).
- [10] C. Di Franco, M. Paternostro, D. I. Tsomokos, S. F. Huelga, Phys. Rev. A **77**, 062337 (2008).
- [11] H. S. M. Coxeter, *Regular Polytopes*, 3rd ed., New York: Dover Publications (1973).
- [12] N. Saxena, S. Severini, I. E. Shparlinski, Int. J. Quant. Inf. **5**, 417 (2007).
- [13] M-H. Yung and S. Bose, Phys. Rev. A **71**, 032310 (2005).
- [14] See M. Babinet, C. R. Acad. Sci. **4**, 638 (1837) for the original reference. This result is featured in most classical optics textbooks, for instance, see pp. 49-50 of G. Brooker *Modern Classical Optics (Oxford Master Series in Physics)*, Oxford University Press, 2003.
- [15] M. Mohseni, P. Rebentrost, S. Lloyd and A. Aspuru-Guzik, arXiv:0805.2741; P. Rebentrost, M. Mohseni, and A. Aspuru-Guzik, arXiv:0806.4725; P. Rebentrost, M. Mohseni, I. Kassal, S. Lloyd, A. Aspuru-Guzik, arXiv:0807.0929.
- [16] M. B. Plenio and S. F. Huelga, arXiv:0807.4902.
- [17] More general settings that allow for dephasing assisted transport will be presented elsewhere.