

Inverse resonance scattering for Jacobi operators

Evgeny Korotyaev *

November 3, 2018

Abstract

We consider the Jacobi operator $(Jf)_n = a_{n-1}f_{n-1} + a_n f_{n+1} + b_n f_n$ on \mathbb{Z} with a real compactly supported sequences $(a_n)_{n \in \mathbb{Z}}$ and $(b_n)_{n \in \mathbb{Z}}$. We give the solution of two inverse problems (including characterization): $(a, b) \rightarrow \{\text{zeros of the reflection coefficient}\}$ and $(a, b) \rightarrow \{\text{bound states and resonances}\}$. We describe the set of "iso-resonance operators J ", i.e., all operators J with the same resonances and bound states.

Keywords: Jacobi operator, resonances, inverse problem

AMS Subject classification: 81Q10 (34L40 47E05 47N50)

1 Introduction

We consider the Jacobi operator J acting on ℓ^2 and given by

$$(Jf)_n = a_{n-1}f_{n-1} + a_n f_{n+1} + b_n f_n, \quad n \in \mathbb{Z}, \quad f = (f_n)_{n \in \mathbb{Z}} \in \ell^2(\mathbb{Z}),$$

where a real compactly supported sequence $q = (q_n)_{n \in \mathbb{Z}}$, $q_{2n-1} = b_n$, $q_{2n} = 1 - a_n$ satisfies

$$q = (q_n)_{n \in \mathbb{Z}} \in \mathfrak{X}_\nu^\tau = \mathfrak{X}_\nu^\tau(p) = \ell_{1+\nu, 2p-\tau}, \quad \text{for some } \nu, \tau \in \{0, 1\}, \quad p \in \mathbb{N},$$

$$\ell_{m,k} = \left\{ (q_n)_{n \in \mathbb{Z}} \in \ell^2(\mathbb{Z}) : q_m \neq 0, q_k \neq 0, q_{2s} < 1, q_n = 0, \text{ all } s \in \mathbb{Z}, n \in \mathbb{Z} \setminus [m, k] \right\}, \quad (1.1)$$

for some $m, k \in \mathbb{Z}$. For fixed $p \geq 1$ the sequence $q \in \mathfrak{X}_0^0$ has the max support and $q \in \mathfrak{X}_1^1$ has the min support and $q \neq 0$. It is well known that the spectrum of J has the form

$$\sigma(J) = \sigma_{ac}(J) \cup \sigma_d(J), \quad \sigma_{ac}(J) = [-2, 2], \quad \sigma_d(J) \subset \mathbb{R} \setminus [-2, 2]. \quad (1.2)$$

Define the new variable $z \in \mathbb{D}_1 = \{z \in \mathbb{C} : |z| < 1\}$ by $\lambda = \lambda(z) = z + \frac{1}{z}$. Here $\lambda(z)$ is a conformal mapping from \mathbb{D}_1 onto $\mathbb{C} \setminus [-2, 2]$. Denote by $\psi^\pm = (\psi_n^\pm(z))_{n \in \mathbb{Z}}$ the fundamental solutions to

$$a_{n-1}\psi_{n-1}^\pm + a_n\psi_{n+1}^\pm + b_n\psi_n^\pm = (z + \frac{1}{z})\psi_n^\pm, \quad n \in \mathbb{Z}, \quad (1.3)$$

*School of Math., Cardiff Univ., Senghennydd Road, Cardiff, CF24 4AG, UK, e-mail: KorotyaevE@cf.ac.uk

$$\psi_n^+(z) = z^n, \quad n > p, \quad \text{and} \quad \psi_n^-(z) = z^{-n}, \quad n \leq 0, \quad |z| \leq 1. \quad (1.4)$$

Define the Wronskian $\{f, u\}_n = a_n(f_n u_{n+1} - u_n f_{n+1})$ for sequences $u = (u_n)_{-\infty}^{\infty}$, $f = (f_n)_{-\infty}^{\infty}$. If f, u are some solutions of (1.3), then $\{f, u\}_n$ does not depend on n . The following identities hold true:

$$\psi^+ = A\tilde{\psi}^- + B\psi^-, \quad \text{on} \quad \mathbb{S}_0^1 = \mathbb{S}^1 \setminus \{\pm 1\}, \quad \tilde{\psi}^\pm = \psi^\pm(z^{-1}), \quad (1.5)$$

where $\mathbb{S}^1 = \{z \in \mathbb{C} : |z| = 1\}$, and $\tilde{v} = v(z^{-1})$ for a function $v(z)$ and

$$A = \frac{w}{1 - z^2}, \quad w = z\{\psi^+, \psi^-\}_n, \quad B = \frac{z^2 s}{\eta}, \quad s = \frac{\{\psi^+, \tilde{\psi}^-\}_n}{z^2}, \quad \eta = z - \frac{1}{z}. \quad (1.6)$$

Note that if $q = 0$, then $w = 1 - z^2$, $A = 1$, $B = 0$. The S-matrix is given by

$$\mathcal{S}_M(z) = \begin{pmatrix} A(z)^{-1} & R_-(z) \\ R_+(z) & A(z)^{-1} \end{pmatrix}, \quad z \in \mathbb{S}^1, \quad R_- = \frac{B}{A} = -\frac{z^3 s}{w}, \quad R_+ = -\frac{\tilde{B}}{A} = \frac{\tilde{s}}{zw}, \quad (1.7)$$

where $\frac{1}{A}$ is the transmission coefficient and R_\pm is the reflection coefficient. For each $z \in \mathbb{S}_0^1$ the scattering matrix \mathcal{S}_M is unitary and satisfies

$$|A(z)|^2 = 1 + |B(z)|^2 \quad \Leftrightarrow \quad w(z)w(z^{-1}) + \eta^2(z) = s(z)s(z^{-1}), \quad (1.8)$$

$$\det \mathcal{S}_M(z) = \frac{\overline{A}(z)}{A(z)} = -z^2 \frac{\overline{w}(z)}{w(z)} = -z^2 \frac{w(z^{-1})}{w(z)}. \quad (1.9)$$

In the case of compactly supported q it is convenient to work with the polynomials w, s . Thus \mathcal{S}_M is meromorphic in the complex plane \mathbb{C} and the poles of \mathcal{S}_M are the zeros of w . Let $\lambda_j, j \in \mathbb{Z}_N = \{n_-, \dots, -1, 1, \dots, n_+\}$, $N = n_+ - n_- \geq 0$ be the bound states of J and $\lambda_j = z_j + z_j^{-1}$, where z_j are all zeros of w in \mathbb{D}_1 and the sequence $E_N = (z_j)_{j \in \mathbb{Z}_N}$ satisfies

$$-1 < z_{n_-} < \dots < z_{-1} < 0 < z_1 < \dots < z_{n_+} < 1 \quad \text{for} \quad \text{some} \quad \pm n_\pm \geq 0. \quad (1.10)$$

Recall that w has only these real zeros in \mathbb{D}_1 (otherwise J has a non-real eigenvalue).

The main goal of this paper is to prove the following results:

- i) the mapping $(a, b) \rightarrow \{\text{bound states, zeros of the reflection coefficient}\}$ is 1-to-1 and onto,*
- ii) the mapping $(a, b) \rightarrow \{\text{bound states, resonances and some sequence } \sigma = (\sigma_n)_0^m, \sigma_n = \pm 1, \}$ is 1-to-1 and onto,*
- iii) we characterize "iso-resonance Jacobi operators J ", i.e., all operators J with the same resonances and bound states.*

The inverse spectral problem consists of the following parts:

- i) Uniqueness. Prove that the spectral data uniquely determine the potential.
- ii) Characterization. Give conditions for some data to be the spectral data of some potential.
- iii) Reconstruction. Give an algorithm for recovering the potential from the spectral data.
- iv) A priori estimates. Obtain estimates of the potential in terms of the spectral data.

We define the class of scattering data (s, E_N) , where $s(z)$ is a polynomial associated with the zeros of the reflection coefficient and $E_N = (z_j)_{j \in \mathbb{Z}_N}$ is a sequence of all zeros of w in \mathbb{D}_1 .

Definition S. By $\mathcal{S}_\nu^\tau = \mathcal{S}_\nu^\tau(p), \nu, \tau \in \{0, 1\}, p \in \mathbb{N}$ we will denote the class of (f, K_f^N) , for some $N \geq 0$, where f is a real polynomial (i.e., a polynomial with real coefficients) given by

$$f = Cz^\nu \prod_1^m (z - \zeta_n), \quad \text{where } C \in \mathbb{R} \setminus \{0\}, \quad \zeta_n \in \mathbb{C} \setminus \{0\}, \quad m = 2p - 1 - \tau - \nu,$$

$K_f^N = (z_j)_{j \in \mathbb{Z}_N}$ is a sequence of zeros of the function $f(z)f(z^{-1}) - \eta^2(z)$, which satisfy

$$-1 < z_{n_-} < \dots < z_{-1} < 0 < z_1 < \dots < z_{n_+} < 1, \quad \pm n_\pm \geq 0, \quad N = n_+ - n_-, \quad (1.11)$$

$$(-1)^j z_j f(z_j^{\pm 1}) > 0, \quad j \in \mathbb{Z}_N, \quad \text{and} \quad (-1)^{n_\pm} f(\pm 1) \geq 0. \quad (1.12)$$

Remark. i) Below we will show that if $q \in \mathfrak{X}_\nu^\tau$, then $(s, E_N) \in \mathcal{S}_\nu^\tau$, where $E_N = K_s^N$ is a sequence of all zeros of w in \mathbb{D}_1 . Note that $K_s^0 = \emptyset$. ii) It is possible that the function $s(z)s(z^{-1}) - \eta^2(z)$ has $N_1 > N$ zeros on $(-1, 1)$, and it is important that we take some of them with the needed properties (1.12). In particular, we emphasize that E_N is not uniquely determined by s .

Theorem 1.1. Let $\tau, \nu \in \{0, 1\}$ and $m \geq 3$. The mapping $\mathfrak{J} : \mathfrak{X}_\nu^\tau \rightarrow \mathcal{S}_\nu^\tau$ given by $\mathfrak{J}(q) = (s, E_N)$ is one-to-one and onto.

Remark. 1) In the proof of Theorem 1.1 we present algorithm for recovering the potential q from the spectral data (s, E_N) . The potential q is uniquely determined by the Marchenko equations (2.12)-(2.16) (in terms of (s, E_N)). Here we use the results about the Marchenko equation from [Te]. Standard spectral data for the inverse problem for the Jacobi operator J are $(s, z_j, m_j^\pm, j \in \mathbb{Z}_N)$, where m_j^\pm is so-called norming constant given by

$$m_j^\pm = \sum_{n \in \mathbb{Z}} \psi_n^\pm(z_j)^2, \quad j \in \mathbb{Z}_N, \quad (1.13)$$

see [Te]. In Theorem 1.1 instead of the norming constants we need the condition (1.12).

2) In Section 3 we show simple examples of the scattering data for the case $p = 1, 2$.

3) We briefly indicate how to prove Theorem 1.1. Firstly, we show that if $q \in \mathfrak{X}_\nu^\tau$, then $(s, E_N) \in \mathcal{S}_\nu^\tau$. Here we check condition (1.12). Secondly, we consider the inverse mapping. Suppose $(s, K_s^N) \in \mathcal{S}_\nu^\tau$. Then in order to determine w we solve the functional equation (1.8) in some class of polynomials (see Theorem 3.2) and this gives the reflection coefficient R_\pm . Thirdly, (1.17), (1.18) yield the norming constants m_j^\pm . Then we check that $R_\pm, z_j, m_j^\pm, j \in \mathbb{Z}_N$ satisfy conditions from Theorem 2.3 of Teschl [Te], which we recall for the sake of the reader in Sect.3. Then we obtain a bijection of our mapping.

4) Assume that $s(\cdot) = 0$ for some $q \in \mathfrak{X}_\nu^\tau$. Then (1.8) yields $w(z)w(z^{-1}) = -\tau(z)^2, z \in \mathbb{C} \setminus \{0\}$. It is impossible since w is a polynomial and we have a contradiction. In fact we deduce that if $s(\cdot) = 0$ for some compactly supported "potential" q , then $q = 0$.

Let $\#(f, I)$ denote the number of zeros of a function f on the set I .

Definition W. By $\mathcal{W}_\nu^\tau = \mathcal{W}_\nu^\tau(p), \nu, \tau \in \{0, 1\}, p \in \mathbb{N}$ we will denote the class of polynomials

$$w = C_w \prod_1^m (z - \rho_n), \quad C_w \in \mathbb{R} \setminus \{0\}, \quad m = 2p - 1 - \tau - \nu, \quad \rho_n \in \mathbb{C}, \quad w(0) > 0, \quad (1.14)$$

i) w is real on \mathbb{R} and $|w(z)| \geq |\eta(z)|$ for any $|z| = 1$, where $\eta = z - \frac{1}{z}$,
ii) w has only real simple zeros $z_j, j \in \mathbb{Z}_N$ in $\overline{\mathbb{D}}_1$, and the sequence $E_N = (z_j)_{j \in \mathbb{Z}_N}$ and the function $F(z) = w(z)w(z^{-1}) + \eta^2(z)$ satisfies

$$\begin{aligned} -1 < z_{n_-} < \dots < z_{-1} < 0 < z_1 < \dots < z_{n_+} < 1, \quad \text{for some } \pm n_{\pm} \geq 0, \quad N = n_+ - n_- \geq 0, \\ \frac{1}{2} \#(F, (z_{n_{\pm}}, z_{n_{\pm}}^{-1})) &= \text{even} \geq 0, \quad \#(F, I_j) = \text{even} \geq 2, \\ I_0 &= (z_{-1}, z_1), \quad I_j = (z_j, z_{j+1}), \quad j \in \mathbb{Z}_N \setminus \{-1, n_{n_+}\} = \{n_-, \dots, -2, 1, 2, \dots, n_+ - 1\}. \end{aligned} \quad (1.15)$$

We describe the properties of s, w and the sequence of zeros $E_N = (z_j)_{j \in \mathbb{Z}_N} \subset (-1, 1)^N$.

Proposition 1.2. *Let $q \in \mathfrak{X}_{\nu}^{\tau}$ for some $\tau, \nu \in \{0, 1\}$. Then $(s, E_N) \in \mathcal{S}_{\nu}^{\tau}, w \in \mathcal{W}_{\nu}^{\tau}$ and for each $j \in \mathbb{Z}_N$ the following identities hold true*

$$\psi^+(z_j) = B(z_j)\psi^-(z_j), \quad B(z_j^{-1})B(z_j) = -1, \quad s(z_j^{-1})s(z_j) = \eta^2(z_j), \quad (1.16)$$

$$m_j^+ = -z_j A'(z_j)B(z_j) = \frac{z_j^2}{\eta^2(z_j)} w'(z_j)s(z_j), \quad (1.17)$$

$$m_j^+ = m_j^- B^2(z_j), \quad (1.18)$$

$$z_j(-1)^j s(z_j^{\pm 1}) > 0, \quad (-1)^j z_j w'(z_j) > 0, \quad (1.19)$$

$$(-1)^{n_{\pm}} w(\pm 1) \geq 0, \quad s(\pm 1) = \pm w(\pm 1). \quad (1.20)$$

Moreover, let $w(z) = \sum_1^{2p} \check{w}_n z^{n-1}$, $s(z) = \sum_1^{2p} \check{s}_n z^{n-1}$, $\check{w} = (\check{w}_n)_1^{2p}, \check{s} = (\check{s}_n)_1^{2p} \in \mathbb{R}^{2p}$ and let $Vh = (0, h_1, \dots, h_{2p-1}), h = (h_n)_1^{2p}$. Then

$$2 + (\check{s}, \check{s}) = (\check{w}, \check{w}), \quad (V^2 \check{s}, \check{s}) = 1 + (V^2 \check{w}, \check{w}) \quad (V^{2k-1} \check{s}, \check{s}) = (V^{2k-1} \check{w}, \check{w}), \quad k = 1, \dots \quad (1.21)$$

Below we will sometimes write $w(z, q), s(z, q), \dots$, instead of $w(z), s(z), \dots$, when several potentials are being dealt with. For $q \in \mathfrak{X}_{\nu}^{\tau}$ the iso-resonance set of potentials is given by

$$\text{Iso}(q) = \{r \in \mathfrak{X}_{\nu}^{\tau} : w(\cdot, q) = w(\cdot, r)\}. \quad (1.22)$$

We will describe $\text{Iso}(q)$. Assume that we know w and we need to recover the polynomial s . Due to Theorem 1.1 the function $F(z) = w(z)w(z^{-1}) + \eta^2(z) = s(z)s(z^{-1})$ has the zeros $t_n, t_{n+m} = t_n^{-1}, n = 1, \dots, m$ counted with multiplicity and given by

$$\begin{aligned} 0 < |t_1| \leq \dots \leq |t_m| \leq 1, \quad (t_n)_1^m \subset U = \mathbb{D}_1 \cup \overline{\mathbb{S}}_+^1, \quad \mathbb{S}_+^1 = \mathbb{C}_+ \cap \mathbb{S}^1, \\ \arg t_n \in [0, 2\pi), \quad \text{and} \quad \text{if } |t_n| = |t_k|, \quad \arg t_n \leq \arg t_k \Rightarrow n \leq k. \end{aligned} \quad (1.23)$$

Note that if $|t_n| = 1$, then $\text{Im } t_n \geq 0$. Hence $(t_n)_1^m$ is a uniquely defined sequence of all zeros $\neq 0$ of F in the set U . Thus $t_n, t_n^{-1}, n = 1, \dots, m$ are all zeros of F and t_n or t_n^{-1} is a zero of s .

If $q \in \mathfrak{X}_{\nu}^{\tau}$, then $s = C_s z^{\nu} \prod_1^m (z - \zeta_n)$, where each $\zeta_n \neq 0, n = 1, \dots, m$ and recall that $\nu \in \{0, 1\}$. The sequence $\sigma = (\sigma_n)_0^m$ is defined by

$$\sigma = (\sigma_n)_0^m \in \{\pm 1\}^{m+1}, \quad \sigma_0 = \text{sign } C_s, \quad \text{and} \quad \zeta_n = t_n^{\sigma_n}, \quad n = 1, \dots, m. \quad (1.24)$$

For each $w \in \mathcal{W}_\tau^\nu$ we define a set Ξ_w of all possible sequences $\sigma = (\sigma_n)_0^m$ by

$$\Xi_g = \left\{ \sigma = (\sigma_n)_0^m \subset \{-1, 1\}^{m+1} : (s, E_N) \in \mathcal{S}_\nu^\tau, \text{ where } s = Cz^\nu \prod_1^m (z - t_n^{\sigma_n}), \sigma_0 = \text{sign } C \right\}.$$

In particular we have:

- I) If $w(1) = w(-1) = 0 = N$, then $(\sigma_n)_0^m$ is any sequence from $\{-1, 1\}^{m+1}$, under the condition that f is real.
- II) If $w(1) \neq 0$ (or $w(-1) \neq 0$), then $(-1)^{n+s(1)} > 0$ (or $(-1)^{n-s(1)} < 0$) gives $\sigma_0 = \text{sign } C_s$.
- III) If $N \geq 1$, then condition $(-1)^j z_j s(z_j) > 0$ for some $j \in \mathbb{Z}_N$ gives $\sigma_0 = \text{sign } C_s$.
- IV) If $N \geq 2$, then the function s has an odd number ≥ 1 of zeros on each of the intervals $(z_{n-}, z_{n+1}), \dots, (z_{-2}, z_{-1}), (z_{-1}, z_1)$ and $(z_1, z_2), \dots, (z_{n+1-}, z_{n+})$.

Our goal is to show that the spectral data Ξ_w give the "proper" parametrization of the set $\text{Iso}(q)$. Our main Theorem 1.3 shows that $\sigma \in \Xi_w$ are **almost free parameters**. Namely, we prove that if the function $w(z, q)$ is fixed, then each σ_n can be changed in an almost arbitrary way.

Theorem 1.3. *Let $\tau, \nu \in \{0, 1\}$ and $m \geq 3$.*

- i) *The mapping $\mathfrak{J}_R : \mathfrak{X}_\nu^\tau \rightarrow \{(w, \Xi_w), w \in \mathcal{W}_\nu^\tau\}$ given by $q \rightarrow (w, (\sigma_n)_0^m)$ is one-to-one and onto.*
- ii) *Let $q \in \mathfrak{X}_\nu^\tau$. Then the mapping $\Psi : \text{Iso}(q) \rightarrow \Xi_w$, given by $r \rightarrow \sigma(r)$ (see (1.24)) is a bijection between the set of potentials $r \in \text{Iso}(q)$ and the set of sequences $\sigma(r) \in \Xi_w, w = w(\cdot, q)$.*

A great number of papers are devoted to the inverse problem for the Schrödinger operator, (see a book [M], papers [Fa], [DT], [Me] and ref. therein).

A lot of papers are devoted to the resonances for the 1D Schrödinger operator, see [F], [K1], [K2], [K3], [S],[Z], [Z1]. We recall that Zworski [Z] obtained the first results about the distribution of resonances for the Schrödinger operator with compactly supported potentials on the real line. Korotyaev obtained the characterization (plus uniqueness and recovering) of S -matrix for the Schrödinger operator with a compactly supported potential on the real line [K1] and on the half-line [K2]. In [K3] for the Schrödinger operator on the half line the stability result was given: if $\varkappa^0 = \{\varkappa^0\}_1^\infty$ is a sequence of zeros (eigenvalues and resonances) of the Jost function for some real compactly supported potential q_0 and $\varkappa - \varkappa^0 \in \ell_\varepsilon^2$ for some $\varepsilon > 1$, then \varkappa is the sequence of zeros of the Jost function for some unique real compactly supported potential.

There are a lot of papers and books devoted to the scattering for Jacobi operators, see [C1], [C2], [CC], [CK], [G1], [G2], [NMPZ], [Te], [T1], [T2]. In the case of Jacobi operators also there are papers about the inverse resonance problem, see [BNW], [MW], [DS1], [DS2]. In particular, some progress was made by Damanik and Simon in [DS1], [DS2], where they described the S -matrix for Jacobi operators on the half-lattice both for finite-support and exponentially decay perturbations.

2 Proof of main Theorems

We recall well-known facts from [Te].

Lemma 2.1. *Let $q \in \ell_{1,2p}^2$. Then each function $\psi_{p-n}^+(z), n = 1, 2, \dots, p$ is a real polynomial and satisfies*

$$\psi_p^+(z) = \frac{z^p}{a_p}, \quad \psi_{p-n}^+(z) = \frac{z^{p+n}}{\eta_n} \left(c_p - \frac{c_p \beta_{p-n+1} + b_p a_p^2}{z} + \frac{O(1)}{z^2} \right), \quad c_n = 1 - a_n^2, \quad (2.1)$$

$$\psi_2^+(z) = \frac{z^{2p-2}}{\eta_2} \left(c_p - \frac{c_p \beta_3 + b_p a_p^2}{z} + \frac{O(1)}{z^2} \right), \quad (2.2)$$

$$\psi_1^+(z) = \frac{z^{2p-1}}{\eta_1} \left(c_p - \frac{c_p \beta_2 + b_p a_p^2}{z} + \frac{O(1)}{z^2} \right) \quad (2.3)$$

as $z \rightarrow \infty$, where $\beta_n = b_n + b_{n+1} + \dots + b_p$ and $\eta_n = a_n a_{n+1} \cdots a_p$. Moreover,

$$\psi_{p-n}^+(z) = \frac{z^{p-n}}{\eta_{p-n}} (1 - z \beta_{p-n+1} + O(z^2)), \quad \psi_1^+(z) = z \frac{1 - z \beta_2 + O(z^2)}{\eta_1} \quad \text{as } z \rightarrow 0, \quad (2.4)$$

$$\psi_1^-(z) = z^{-1}, \quad \psi_2^-(z) = \frac{1 - z b_1}{a_1 z^2}. \quad (2.5)$$

Lemma 2.2. *Let $q \in \ell_{1,2p}^2$. Then*

$$w(z) = -(b_1 - z^{-1})\psi_1^+(z) - a_1 \psi_2^+(z), \quad z \neq 0, \quad (2.6)$$

$$s(z) = (1 - b_1 z^{-1})\psi_1^+(z) - z^{-1} a_1 \psi_2^+(z), \quad (2.7)$$

$$w(z) = \frac{z^{2p-1}}{\eta_1} \left(-b_1 c_p + \frac{c_p(c_1 + \beta_2 b_1) + b_1 b_p a_p^2}{z} + \frac{O(1)}{z^2} \right) \quad \text{as } z \rightarrow \infty$$

$$w(z) = \frac{1 - z \beta_1 + O(z^2)}{\eta_1} \quad \text{as } z \rightarrow 0, \quad (2.8)$$

$$s(z) = \frac{z^{2p-1}}{\eta_1} \left(c_p - \frac{c_p \beta_1 + b_p a_p^2}{z} + \frac{O(1)}{z^2} \right) \quad \text{as } z \rightarrow \infty$$

$$s(z) = \frac{1}{\eta_1} \left(-b_1 + z(c_1 + b_1 \beta_2) + O(z^2) \right) \quad \text{as } z \rightarrow 0. \quad (2.9)$$

Furthermore, if $b_1 = c_p = 0$, then

$$w(z) = -\frac{z^{2p-3}}{\eta_1} (c_1 b_p + O(1/z)) \quad \text{as } z \rightarrow \infty. \quad (2.10)$$

Proof. Using (1.6) and Lemma 2.1, we obtain (2.6), (2.7). Then asymptotics from Lemma 2.1 imply (2.8), (2.9). If $b_1 = c_p = 0$, then (2.2), (2.3) imply

$$w = -a_1\psi_2^+ + \frac{\psi_1^+}{z} = \frac{a_1^2 z^{2p-3}}{\eta_1} (b_p + O(z^{-1})) - \frac{z^{2p-3}}{\eta_1} (b_p + O(z^{-1})) = -\frac{z^{2p-3}}{\eta_1} (c_1 b_p + O(z^{-1})).$$

■

Proof Proposition 1.2. Identities (1.16) follow from (1.5), (1.8). Recall the following identity

$$A'(z_j) = -\frac{1}{z_j} \sum_{n \in \mathbb{Z}} \psi_n^+(z_j) \psi_n^-(z_j), \quad j \in \mathbb{Z}_N, \quad (2.11)$$

see (10.34) in [Te]. Then using (1.16) we obtain (1.17). Similar arguments give (1.18).

Using $w(0) > 0$ and (1.17), we obtain $w'(z_1) < 0$ and $w'(z_2) > 0, \dots$. Moreover, due to (1.16), (1.17) we have (1.19). Identity (2.6), (2.7) give (1.20).

Substituting $w = \sum_1^{2p} \tilde{w}_n z^{n-1}$ and $s = \sum_1^{2p} \tilde{s}_n z^{n-1}$ into (1.8) we obtain (1.21). ■

We need some results about the inverse problems from [Te] for $q \in \ell_1^1 = \{h = (h_n)_{n \in \mathbb{Z}} : \sum(1 + |n|)|h_n| < \infty\}$. Define the Marchenko operator $\mathfrak{F}_n : \ell^2(\mathbb{Z}_+) \rightarrow \ell^2(\mathbb{Z}_+)$ by

$$(\mathfrak{F}_n f)_k = \sum_{m \geq 0} F(2n + m + k) f_m, \quad f = (f_n)_0^\infty \in \ell^2(\mathbb{Z}_+), \quad \mathbb{Z}_+ = \mathbb{Z} \cap [0, \infty), \quad (2.12)$$

$n, k \geq 0$ where

$$F(n) = \widehat{R}_n^+ + \sum_{j=1}^N \frac{z_j^n}{m_j}, \quad \widehat{R}_n^\pm = \frac{1}{2\pi i} \int_{|z|=1} R_\pm(z) z^{n-1} dz, \quad n \in \mathbb{Z}, \quad (2.13)$$

and \widehat{R}_n^\pm are the Fourier coefficients of R_\pm . The Marchenko equation is given by

$$(I + \mathfrak{F}_n) \mathcal{K}_n(\cdot) = \eta_n^2 e_0, \quad \text{where } \mathcal{K}_n = (\mathcal{K}_n(m))_0^\infty, \quad e_n = (\delta_{n,k})_0^\infty \in \ell^2(\mathbb{Z}_+), \quad (2.14)$$

i.e.,

$$\mathcal{K}_n(k) + F(2n + k) + \sum_{m \geq 1} F(2n + k + m) \mathcal{K}_n(m) = \eta_n^2 \delta_{0,k}. \quad (2.15)$$

For each $n \in \mathbb{Z}$ these equations have unique "decreasing" solutions $(\mathcal{K}_n(k))_{k=0}^\infty$. The sequences $a_n, b_n, n \in \mathbb{Z}$ have the forms

$$a_n^2 = \frac{\Psi_n^0}{\Psi_{n+1}^0}, \quad b_n = \frac{\Psi_n^1}{\Psi_n^0} - \frac{\Psi_{n-1}^1}{\Psi_{n-1}^0}, \quad \Psi_n^k = \langle e_k, (I + \mathfrak{F}_n) e_0 \rangle, \quad k = 0, 1, \quad n \in \mathbb{Z}, \quad (2.16)$$

where $\langle \cdot, \cdot \rangle$ is the scalar product in $\ell^2(\mathbb{Z}_+)$. Recall the inverse spectral theorem from [Te].

Theorem 2.3. *The Faddeev mapping*

$$\ell_1^1(\mathbb{Z}) \oplus \ell_1^1(\mathbb{Z}) \rightarrow \mathfrak{S} = \{R_+, (z_j, m_j^+, j \in \mathbb{Z}_N) \in (-1, 1)^N \times \mathbb{R}_+^N, z_j \neq 0, N \geq 0\} \quad (2.17)$$

given by $(a_n - 1, b_n)_{n \in \mathbb{Z}} \rightarrow \{R_+(z), z \in \mathbb{S}^1, (z_j, m_j^+, j \in \mathbb{Z}_N)\}$ is one-to-one and onto, where the function R_+ , the eigenvalues z_j and the norming constants $m_j^+, j \in \mathbb{Z}_N$ satisfy the conditions

1) $R_+(z) = \overline{R_+(\overline{z})} = -\frac{s(\overline{z})}{zw(z)}, z \in \mathbb{S}_0 = \mathbb{S}^1 \setminus \{\pm 1\}$, and the function $R_+(z)$ is continuous in $z \in \mathbb{S}_0$ and satisfies

$$C|1 - z^2|^2 + |R_+(z)|^2 \leq 1, \quad \text{all } z \in \mathbb{S}_0, \quad \text{for some } C > 0. \quad (2.18)$$

2) The eigenvalues $z_j, j \in \mathbb{Z}_N$ are distinct and $m_j^- m_j^+ = w'(z_j)^2$.

3) The sequences $\widehat{R}^\pm = (\widehat{R}_n^\pm)_1^\infty$ defined in (2.13) with $R_-(z) = -R_+(\overline{z})\frac{A(\overline{z})}{A(z)}$ satisfy

$$\sum_{n \geq 1} n |\widehat{R}_n^\pm - \widehat{R}_{n+2}^\pm| < \infty. \quad (2.19)$$

We are ready to prove the first result.

Proof of Theorem 1.1. We consider the case $q \in \mathfrak{X}_0^0$ and $m = 2p - 1$, the proof of other cases is similar. If $q \in \mathfrak{X}_0^0$, then Lemma 1.2 gives that $(s, E_N) \in \mathcal{S}_0^0$, which yields a mapping $q \rightarrow (s, E_N)$ from \mathfrak{X}_0^0 into \mathcal{S}_0^0 .

We will show uniqueness. Let $q \in \mathfrak{X}_0^0$. Then Lemma 1.2 gives that $(s, E_N) \in \mathcal{S}_0^0$, and Theorem 3.2 gives a unique w . Moreover, Proposition 1.2 yields the norming constants $m_j^\pm, j \in \mathbb{Z}_N$. These data determine the compactly supported potential uniquely by Theorem 2.3. Then we deduce that the mapping $q \rightarrow (s, E_N)$ is an injection.

We will show surjection of the mapping $q \rightarrow (s, E_N)$. If $(s, E_N) \in \mathcal{S}_0^0$, then Theorem 3.2 gives unique $w \in \mathcal{W}_0^0$ and we have $R_+ = -\frac{s(z^{-1})}{zw(z)}$. If $n \geq 2p + 1$, then we have

$$\widehat{R}_n^+ = \frac{1}{2\pi i} \int_{|z|=1} R_+(z) z^{n-1} dz = -\frac{1}{2\pi i} \int_{|z|=1} \frac{z^{n-2} \widetilde{s}(z)}{w(z)} dz = -\sum_{j=1}^N \text{Res} \frac{z^{n-2} \widetilde{s}(z)}{w(z)} \Big|_{z_j} = -\sum_{j=1}^N \frac{z_j^n}{m_j},$$

since $n - 2 = (2p - 1) + (n - 2p - 1)$ and the function $z^{2p-1} \widetilde{s}$ is a polynomial. Thus

$$F(n) = 0 \quad \text{if} \quad n \geq 2p + 1, \quad (2.20)$$

and using (2.16) we obtain

$$\Psi_n^0 = 1 + F(2n), \quad \text{and} \quad a_n^2 = \frac{1 + F(2n)}{1 + F(2n + 2)} \quad (2.21)$$

thus $a_n^2 = 1$ if $n > p$. Similar arguments yield $b_n = 0$ if $n > p$. Then we deduce that

$$a_n = 1, \quad b_n = 0 \quad \text{if} \quad n \geq p + 1.$$

Moreover, similar arguments yield $b_n = 0, a_n = 1$ if $n \leq 0$. Then the asymptotics from Lemma 2.2 give that $q \in \mathfrak{X}_0^0$, which yields surjection. ■

Proof of Theorem 1.3. i) We consider the case $q \in \mathfrak{X}_0^0$, the proof of other cases is similar. Let $q \in \mathfrak{X}_0^0$. Then Proposition 1.2 yield $w(\cdot, q) \in \mathcal{W}_0^0$ and we have the mapping $q \rightarrow (w, \sigma)$, where the sequence $\sigma \in \Xi_w$ is given by (1.24), since $(s, E_N) \in \mathcal{S}_0^0$.

By Theorem 3.3, for each $(w, \sigma) \in (w, \Xi_w)$ there exists a unique $(s, E_N) \in \mathcal{S}_0^0$, then due to Theorem 1.1 the mapping $q \rightarrow (w, \sigma)$ is an injection.

We will prove that \mathfrak{J}_R is a surjection. Let $w \in \mathcal{W}_0^0$ have zeros $E_N = (z_j)_{j \in \mathbb{Z}_N}$ from $(-1, 1)$, $N \geq 0$ and let a sequence $\sigma = (\sigma_n)_0^m \in \Xi_w$ be defined by (1.24). By Theorem 3.2, there exists a unique $(s, E_N) \in \mathcal{S}_0^0$ such that (1.12) with E_N hold true. Then $(s, E_N) \in \mathcal{S}_0^0$ and, by Theorem 1.1, there exists a unique potential $q \in \mathfrak{X}_0^0$ with the scattering function $s(\cdot)$. Thus the mapping $\mathfrak{J}_{res} : \mathfrak{X}_\nu^\tau \rightarrow ((w, \sigma) \in \mathcal{W}_\nu^\tau \times \Xi_w)$ given by $q \rightarrow (w, (\sigma_n)_0^m)$ is one-to-one and onto, where $m = 2p - 1$.

ii) Furthermore, using similar arguments we deduce that if we fix $q \in \mathfrak{X}_\nu^\tau$, then the mapping $\Psi : r \rightarrow \sigma(s(r))$ is a bijection between the iso-resonance set of potentials $\text{Iso}(q)$ and the set of sequences $\Xi_w, w = w(\cdot, q)$. ■

3 Properties of polynomials s, w

In this section we will get the needed results about polynomials s, w . In order to solve the inverse problems $(a, b) \rightarrow$ (spectral data) we need the following results.

Lemma 3.1. *Let $g = C_g \prod_1^m (z - \rho_n)$ and $f = C_f \prod_1^m (z - \zeta_n)$ satisfy*

$$g(z)g(z^{-1}) + \eta^2 = f(z)f(z^{-1}), \quad \eta = z - z^{-1}, \quad \text{all } z \neq 0, \quad (3.1)$$

for some $m \geq 3$ and $g(0) \neq 0, f(0) \neq 0$. Then

$$\prod_1^m (\lambda - \mu_n) = \frac{\lambda^2 - 4}{C} + \prod_1^m (\lambda - \lambda_n), \quad \mu_n = \zeta_n + \frac{1}{\zeta_n}, \quad \lambda_n = \rho_n + \frac{1}{\rho_n}, \quad (3.2)$$

$$C = C_s^2 C_\zeta = C_w^2 C_\rho, \quad \frac{C_\rho}{C_\zeta} > 0, \quad C_\rho = \prod_1^m (-\rho_n), \quad C_\zeta = \prod_1^m (-\zeta_n), \quad (3.3)$$

$$\sum_1^m \mu_n = \sum_1^m \lambda_n, \quad \dots, \quad \prod_1^m (-\mu_n) + \frac{4}{C} = \prod_1^m (-\lambda_n). \quad (3.4)$$

Proof. Using $f = C_f \prod_1^m (z - \zeta_n)$ and $g = C_g \prod_1^m (z - \rho_n)$, we obtain

$$C_s^2 \prod_1^m (z - \zeta_n)(z^{-1} - \zeta_n) = \eta^2 + C_w^2 \prod_1^m (z - \rho_n)(z^{-1} - \rho_n).$$

Thus the identities $-\zeta_n(\lambda - \mu_n) = (z - \zeta_n)(z^{-1} - \zeta_n)$ and $-\rho_n(\lambda - \lambda_n) = (z - \rho_n)(z^{-1} - \rho_n)$, give

$$C_s^2 C_\zeta \prod_1^m (\lambda - \mu_n) = \lambda^2 - 4 + C_w^2 C_\rho \prod_1^m (\lambda - \lambda_n),$$

and $C_s^2 C_\zeta = C_w^2 C_\rho = C$, which yields $\frac{C_\zeta}{C_\rho} = \frac{C_w^2}{C_s^2} > 0$. ■

Assume that we know only the polynomial f in the equation (3.1) and we have to determine g . In order to do this we have to solve the equation (3.1) in some class of polynomials. The following Theorem will be used to determine the polynomial w if we know s .

Theorem 3.2. *i) Let functions f, g be analytic in $\mathbb{C} \setminus \{0\}$ and satisfy (3.1) and be real on $\mathbb{R} \setminus \{0\}$. Then $f^2(\pm 1) = g^2(\pm 1)$. Moreover, if $f(\pm 1) = 0$ then $(f')^2(\pm 1) = 8 + (g')^2(\pm 1) \geq 8$.
ii) Let $(f, K_f^N) \in \mathcal{S}_\nu^\tau$, for some $(\nu, \tau) \in \{0, 1\}$, $N \geq 0, m \geq 3$. Then there exists a unique polynomial $g \in \mathcal{W}_\nu^\tau$ satisfying (3.1).*

Remark. It is possible that the function $f(z)f(z^{-1}) - \eta^2(z)$ has more zeros on $(-1, 1) \setminus \{0\}$, and it is important that we make a special choice which, however, has to satisfy condition (1.15).

Proof. The statement i) is very simple and differentiating (3.1) we obtain $(f')^2(\pm 1) = 8 + (f'')^2(\pm 1)$ at $z = \pm 1$.

ii) Recall that $f = z^\nu C \prod_1^m (z - \zeta_n)$, where

$$C \in \mathbb{R} \setminus \{0\}, \quad 0 < |\zeta_1| \leq |\zeta_2| \leq \dots \leq |\zeta_m|, \quad m = 2p - 1 - \tau - \nu,$$

and $K_f^N = (z_j)_{j \in \mathbb{Z}_N}$ is some sequence of zeros of the function $f(z)f(z^{-1}) - \eta^2(z)$ such that

$$-1 < z_{n_-} < \dots < z_{-1} < 0 < z_1 < \dots < z_{n_+} < 1, \quad (-1)^{n_\pm} f(\pm 1) \geq 0, \quad z_j (-1)^j f(z_j^{\pm 1}) > 0, \quad (3.5)$$

$j \in \mathbb{Z}_N$. Then

$$f(z)f(z^{-1}) = C^2 \prod_1^m (z - \zeta_n)(z^{-1} - \zeta_n) = \frac{C_0}{z^m} \prod_1^m (z - \zeta_n)(z - \zeta_n^{-1}),$$

where $C_0 = C^2 C_\zeta$, $C_\zeta = \prod_1^m (-\zeta_n)$. Then $G(z) = f(z)f(z^{-1}) - \eta^2(z)$ satisfies

$$G(z) = \begin{cases} C_0 z^m (1 + O(z^{-1})) & \text{as } z \rightarrow \infty \\ C_0 z^{-m} (1 + O(z)) & \text{as } z \rightarrow 0 \end{cases}, \quad (3.6)$$

and thus

$$G(z) = -\eta^2(z) + \frac{C_0}{z^m} \prod_1^m (z - \zeta_n)(z - \zeta_n^{-1}) = \frac{C_0}{z^m} \prod_1^m (z - \rho_n)(z - \rho_n^{-1}), \quad (3.7)$$

where $\rho_n \neq 0, \rho_n^{-1}$ are the zeros of G counted with multiplicity and satisfying

$$0 < |\rho_1| \leq |\rho_2| \leq \dots \leq |\rho_N| < 1 \leq |\rho_{N+1}| \leq \dots \leq |\rho_m|, \\ \text{the set } \{\rho_1, \rho_2, \dots, \rho_N\} = \{z_j, j \in \mathbb{Z}_N\} \subset (-1, 1) \setminus \{0\}, \quad (3.8)$$

and Conditions i)-iv) in Definition W, since f satisfies Definition S and $G(z) = G(1/z)$ for all $z \neq 0$. Moreover, using $g_0(z) = \prod_1^m (z - \rho_n)$ we have

$$G(z) = C g_0(z) g_0(z^{-1}), \quad \text{where} \quad C = \frac{C_0}{g_0(0)}, \quad (3.9)$$

Note that Lemma 3.1 gives $C > 0$, then $g = C_* g_0$ and $g(0) > 0$, where C_* satisfies $C_* g_0(0) > 0, C_*^2 = C > 0$. By the construction of g , this function is unique. ■

In order to solve the inverse problems $\mathfrak{X}_\nu^\tau \rightarrow \mathcal{W}_\nu^\tau$ we need the following results. Assume that we have (w, σ) , where the function $w \in \mathcal{W}_\nu^\tau$ and the sequence $\sigma \in \Xi_w$, then we have to determine s uniquely. In order to do this we have to solve the equation (3.1) in class of polynomials $s \in \mathcal{S}_\nu^\tau$. The sequence σ will give uniqueness. The following Theorem will be used to determine the function s if we know (w, σ) .

Theorem 3.3. *Let $g \in \mathcal{W}_\nu^\tau$, for some $\nu, \tau \in \{0, 1\}$, $m \geq 3$. Then for each $\sigma \in \Xi_g$ defined in (1.24) there exists a unique $(f, K_f^N) \in \mathcal{S}_\nu^\tau$ satisfying (3.1).*

Proof. Recall that $g = C_g \prod_1^m (z - \rho_n)$, $m = 2p - 1 - \tau - \nu$ for some $C_g \in \mathbb{R} \setminus \{0\}$ and $\rho_n \in \mathbb{C} \setminus \{0\}$ such that:

- i) g is real on \mathbb{R} and if $+1$ and -1 are zeros, they are simple, and $g(0) > 0$,
- ii) $|g(z)| \geq |\eta(z)|$ for any $|z| = 1$, where $\eta = z - \frac{1}{z}$,
- iii) g has only simple zeros $z_{n_-}, \dots, z_{-1}, z_1, \dots, z_{n_+}$ in \mathbb{D}_1 for some $\pm n_\pm \geq 0$ such that $-1 < z_{n_-} < \dots < z_{-1} < 0 < z_1 < \dots < z_{n_+} < 1$.

Then we obtain

$$g(z)g(z^{-1}) = C_g^2 \prod_1^m (z - \rho_n)(z^{-1} - \rho_n) = \frac{C_0}{z^m} \prod_1^m (z - \rho_n)(z - \rho_n^{-1}),$$

where $C_0 = C_g^2 \prod_1^m (-\rho_n)$. Then $F(z) = \eta^2(z) + g(z)g(z^{-1})$ satisfies

$$F(z) = \begin{cases} C_0 z^m (1 + O(z^{-1})) & \text{as } z \rightarrow \infty \\ C_0 z^{-m} (1 + O(z)) & \text{as } z \rightarrow 0 \end{cases}, \quad (3.10)$$

and thus

$$F(z) = \frac{C_0}{z^m} \prod_1^{2m} (z - t_n) = \frac{C_0}{z^m} \prod_1^m (z - t_n)(z - t_n^{-1}), \quad (3.11)$$

where $t_n \neq 0$ are the zeros of F counted with multiplicity and satisfying

$$0 < |t_1| \leq |t_2| \leq \dots \leq |t_m|, \quad t_{n+m} = 1/t_n, \text{ all } n = 1, 2, \dots, m, \quad \arg t_n \in [0, 2\pi),$$

where if $|t_n| = |t_k|$, $\arg t_n \leq \arg t_k \Rightarrow n \leq k$ and if $|t_n| = 1$, then $\text{Im } t_n \geq 0$, since $F(z) = F(1/z)$ for all $z \neq 0$. Moreover, we have

$$F(z) = \frac{C_0}{z^m} \prod_1^m (z - \zeta_n)(z - \zeta_n^{-1}) = \frac{C_0}{f_0(0)} f_0(z) f_0(z^{-1}), \quad f_0(z) = \prod_1^m (z - \zeta_n), \quad (3.12)$$

where $\zeta_n = t_n^\sigma$, $(\sigma_n)_1^m \in \Xi_g$. Note that Lemma 3.1 gives $E = \frac{C_0}{f_0(0)} > 0$, then $f = C_* f_0$ and $g(0) > 0$, where $C_* = \sqrt{E}$. We need to chose the sign of C_* . By the construction of g , this function is unique. ■

Example p=1. In this case we have

$$w = \frac{1}{a_1} (1 - b_1 z - a_1^2 z^2), \quad s = \frac{1}{a_1} (-b_1 + c_1 z). \quad (3.13)$$

1) If we know $s = s_0 + 2s_1z$, then

$$a_1 = -s_1 + \sqrt{s_1^2 + 1}, \quad b_1 = -\frac{s_0}{s_1}. \quad (3.14)$$

2) If we know w , then

$$-a_1w = z^2 + \frac{b_1}{a_1^2}z - \frac{1}{a_1^2}, \quad \rho_1 = \frac{-b_1 - \sqrt{b_1^2 + 4a_1^2}}{4a_1^2} < 0, \quad \rho_2 = \frac{-b_1 + \sqrt{b_1^2 + 4a_1^2}}{4a_1^2} > 0, \quad (3.15)$$

where $\rho_1, \rho_2 \in \mathbb{R}$. Then

$$\rho_1\rho_2 = -\frac{1}{a_1^2}, \quad b_1 = \frac{\rho_1 + \rho_2}{\rho_1\rho_2}.$$

Example p=2. In this specific case we have

$$\psi_2^+ = \frac{z^2}{a_2}, \quad \psi_1^+ = \frac{c_2z^3 - b_2z^2 + z}{a_1a_2}, \quad (3.16)$$

and

$$w = \frac{1}{a_1a_2} \left(-z^3c_2b_1 + z^2(c_2 - a_1^2 + b_1b_2) - z(b_1 + b_2) + 1 \right), \quad (3.17)$$

$$s = \frac{1}{a_1a_2} \left(z^3c_2 - z^2(b_2 + b_1c_2) + z(c_1 + b_1b_2) - b_1 \right). \quad (3.18)$$

We obtain 4 cases:

1) The case $a_2 \neq 1, b_1 \neq 0$ have been considered since $m = 3$.

2) Let $a_2 = 1, b_2 \neq 1, b_1 = 0, a_1 \neq 1$. Then it is similar to the case $p = 1$ and we have:

$$w = \frac{-a_1^2z^2 - zb_2 + 1}{a_1}, \quad s = z \frac{-zb_2 + c_1}{a_1}. \quad (3.19)$$

3) If $a_2 = 1, b_2 \neq 0, b_1 \neq 0$, then

$$w = \frac{z^2(b_1b_2 - a_1^2) - z(b_1 + b_2) + 1}{a_1}, \quad s = \frac{-z^2b_2 + z(c_1 + b_1b_2) - b_1}{a_1}. \quad (3.20)$$

4) If $a_2 \neq 1, b_1 = 0, a_1 \neq 1$, then

$$w = \frac{z^2(c_2 - a_1^2) - zb_2 + 1}{a_1a_2}, \quad s = z \frac{z^2c_2 - zb_2 + c_1}{a_1a_2}. \quad (3.21)$$

Acknowledgments. The research was partially supported by EPSRC grant EP/D054621.

References

- [BNW] B. M. Brown; S. Naboko; R. Weikard, The inverse resonance problem for Jacobi operators Bull. London Math. Soc. 37(2005), 727–37.
- [C1] K. M. Case, The discrete inverse scattering problem in one dimension, J. Math. Phys. 15(1974), 143–146.
- [C2] K. M. Case, On discrete inverse scattering problems. II, J. Math. Phys. 14(1973), 916–920.
- [CC] K. M. Case; S. C. Chiu The discrete version of the Marchenko equations in the inverse scattering problem, J. Math. Phys. 14(1973), 1643–1647.
- [CK] K. M. Case and M. Kac, A discrete version of the inverse scattering problem, J. Math. Phys. 14(1973), 594–603.
- [DS1] D. Damanik; B. Simon, Jost functions and Jost solutions for Jacobi matrices, I. A necessary and sufficient condition for Szegő asymptotics, Invent. Math. 165 (2006), no. 1, 1-50.
- [DS2] D. Damanik; B. Simon, Jost functions and Jost solutions for Jacobi matrices. II. Decay and analyticity. Int. Math. Res. Not. 2006, Art. ID 19396, 32 pp.
- [DT] Deift, P., Trubowitz, E. Inverse scattering on the line, Commun. Pure and Applied Math., 32(1979), 121–251.
- [Fa] Faddeev L., Properties of the S-matrix of the one-dimensional Schrödinger equation, Trudy Mat. Inst. Steklov 73(1964), 314–333, English translation in AMST, 265, 139–166.
- [F] R. Froese, Asymptotic distribution of resonances in one dimension, Journal of Differential Equations 137 (1997), no. 2, 251–272.
- [G1] G. S. Guseinov, The determination of an infinite Jacobi matrix from the scattering data, Soviet Math. Dokl., 17(1976), 596–600.
- [G2] G. S. Guseinov, The inverse problem of scattering theory for a second-order difference equation on the whole axis, Soviet Math. Dokl., 17(1976), 1684–1688.
- [K1] E. Korotyaev, Inverse resonance scattering on the real line. Inverse Problems 21 (2005), no. 1, 325–341.
- [K2] E. Korotyaev, Inverse resonance scattering on the half line. Asymptot. Anal. 37 (2004), no. 3-4, 215–226.
- [K3] E. Korotyaev, Stability for inverse resonance problem. Int. Math. Res. Not. 2004, no. 73, 3927–3936.
- [MBO] F. G. Maksudov, Eh. M. Bajramov, R. U. Orudzheva, The inverse scattering problem for an infinite Jacobi matrix with operator elements, Russ. Acad. Sci., Dokl., Math. 45(1992), No.2, 366–370.
- [M] Marchenko V. Sturm-Liouville operator and applications. Basel: Birkhäuser 1986.
- [MW] M. Marletta; R. Weikard, Stability for the inverse resonance problem for a Jacobi operator with complex potential, Inverse Problems 23 (2007), 1677–1688.
- [Me] Melin A. Operator methods for inverse scattering on the real line. Comm. P.D.E. 10(1985), 677–786.
- [NMPZ] S. Novikov; S. Manakov; L. Pitaevski; V. Zakharov, Theory of solitons. The inverse scattering method. Consultants Bureau [Plenum], New York, 1984.
- [S] B. Simon, Resonances in one dimension and Fredholm determinants Journal of Functional Analysis 178 (2000), no. 2, 396–420.
- [Te] G. Teschl, Jacobi operators and completely integrable nonlinear lattices. Mathematical Surveys and Monographs, 72. AMS, Providence, RI, 2000.
- [T1] M. Toda, Theory of Nonlinear Lattices, 2nd enl. ed., Springer, Berlin, 1989.
- [T2] M. Toda, Theory of Nonlinear Waves and Solitons, Kluwer, Dordrecht, 1989.
- [Z] M. Zworski, Distribution of poles for scattering on the real line, Journal of Functional Analysis 73 (1987), no. 2, 277–296.
- [Z1] M. Zworski, A remark on isopolar potentials, SIAM Journal on Mathematical Analysis 32 (2001), no. 6, 1324–1326.