## Relation between Fresnel transform of input light field and Radon transform of Wigner function of the field

Hong-yi Fan and Li-yun Hu\*

Department of Physics, Shanghai Jiao Tong University, Shanghai, 200030, China \*Corresponding author: hlyun@sjtu.edu.cn; hlyun@sjtu.org

January 13, 2019

## Abstract

We prove a new theorem about the relationship between optical field Wigner function's Radon transform and optical Fresnel transform of the field, i.e., when an input field  $\psi(x')$  propagates through an optical [D(-B)(-C)A] system, the energy density of the output field is equal to the Radon transform of the Wigner function of the input field, where the Radon transform parameters are D, B. We prove this theorem in both spatial-domain and frequency-domain.

In optical communication theory every signal or image can be uniquely and indirectly described by a Wigner distribution function (WDF) [1, 2, 3]. The WDF (or named Wigner transform) of an optical signal field  $\psi(x')$  is defined as

$$W_{\psi}(\nu', x') = \int_{-\infty}^{+\infty} \frac{du}{2\pi} e^{i\nu' u} \psi^* \left(x' + \frac{u}{2}\right) \psi\left(x' - \frac{u}{2}\right), \tag{1}$$

 $W_{\psi}(\nu', x')$  involves both spatial distribution information and space-frequency distribution information of the signal.  $\nu$  is named space frequency.  $W_{\psi}(\nu', x')$  is said to be bilinear in the signal because the signal enters twice in its definition. The WDF undergoes certain variations if something happens to the signal. For examples, passage through a lens corresponds to a vertical shearing of the WDF, propagation in free space means a horizontal shearing of the WDF [4]. However, the WDF preserves space and space frequency marginal properties of any signal,

$$\int_{-\infty}^{+\infty} d\nu' W_{\psi}(\nu', x') = |\psi(x')|^2, \qquad (2)$$

$$\int_{-\infty}^{+\infty} dx' W_{\psi}(\nu', x') = |\tilde{\psi}(\nu')|^2, \qquad (3)$$

where  $\tilde{\psi}(\nu) = \int_{-\infty}^{+\infty} \frac{dx}{\sqrt{2\pi}} \psi(x) e^{ix\nu}$ . If one wants to reconstruct the Wigner function by using various probability distribution, obviously the position density  $|\psi(x')|^2$  and the space-frequency density  $|\tilde{\psi}(\nu')|^2$  were not enough, so the Radon transform [5, 6] of the Wigner function is introduced [7],

$$R(x) \equiv \iint_{-\infty}^{\infty} dx' d\nu' \delta\left(x - Dx' + B\nu'\right) W_{\psi}(\nu', x'), \tag{4}$$

R(x) is also a probability distribution along an infinitely thin phase space strip denoted by the real parameters D, B. The inverse relation of (4) is the foundation of optical tomographic imaging techniques (the techniques derive two-dimensional data from a three-dimensional object to obtain a slice image of the internal structure and thus have the ability to peer inside the object noninvasively.)

On the other hand, an optical system can be analyzed by using either diffraction theory or ray optics. It is worth setting up a connection between ray transfer matrix and diffraction theory, i.e., the diffraction integration written in terms of ray transfer matrix—named Collins' diffraction integral formula [8, 9, 10] which describes the propagation of a general beam  $\psi(x')$  through an (ABCD) optical paraxial system,

$$\phi(x) = \int_{-\infty}^{\infty} \mathcal{K}(x, x') \psi(x') \, dx', \tag{5}$$

where  $\phi(x)$  is the output field, AD - BC = 1, and  $\mathcal{K}(x, x')$  is the integral kernel

$$\mathcal{K}(x,x') = \frac{1}{\sqrt{2\pi i B}} \exp\left[\frac{i}{2B} \left(Ax'^2 - 2x'x + Dx^2\right)\right].$$
(6)

If the energy in the initial beam is normalized,  $\int_{-\infty}^{+\infty} |\psi(x')|^2 dx' = 1$ , then the output beam is normalized too,  $\int_{-\infty}^{+\infty} |\phi(x)|^2 dx = 1$ . Clearly, if the [ABCD] system is changed to [D(-B)(-C)A] system, then Eq. (5) should read

$$\phi(x) = \int_{-\infty}^{\infty} \vec{\mathcal{K}}(x, x') \psi(x') \, dx', \tag{7}$$

where the integral kernel  $\vec{\mathcal{K}}(x, x')$  is

$$\vec{\mathcal{K}}(x,x') = \frac{1}{\sqrt{-2\pi i B}} \exp\left[\frac{-i}{2B} \left(Dx'^2 - 2x'x + Ax^2\right)\right].$$
(8)

In this Letter, we shall reveal the following **theorem**:

When an input field  $\psi(x')$  propagates through an optical [D(-B)(-C)A] system, the energy density of the output field  $\phi(x)$  is equal to the Radon transform of the Wigner function of the input field, where the Radon transform parameters are D, B. The proof is demonstrated as follows.

With the use of Dirac  $\delta$ -function, we can re-express the Wigner function of input field  $\psi(x')$  in Eq. (1) as

$$W_{\psi}(\nu', x') = \iint_{-\infty}^{\infty} dx'' dx''' \int_{-\infty}^{+\infty} \frac{du}{2\pi} e^{i\nu' u} \psi^* (x'') \psi (x''') \delta \left(x'' - x' - \frac{u}{2}\right) \delta \left(x' - \frac{u}{2} - x'''\right)$$
$$= \frac{1}{\pi} \iint_{-\infty}^{\infty} dx'' dx''' \psi^* (x'') \psi (x''') e^{i2\nu' \left(x'' - x'\right)} \delta \left(2x' - x'' - x'''\right). \tag{9}$$

Substituting (9) into (4) we rewrite the Radon transform of  $W_{\psi}(\nu', x')$  as

$$R(x) = \frac{1}{\pi} \iint_{-\infty}^{\infty} dx' d\nu' \delta(x - Dx' + B\nu') \iint_{-\infty}^{\infty} dx'' dx''' \psi^*(x'') \psi(x''') e^{i2\nu'(x'' - x')} \delta(2x' - x'' - x''')$$
  
$$= \frac{1}{2\pi B} \iint_{-\infty}^{\infty} dx'' dx''' \psi^*(x'') \psi(x''') \exp\left\{\frac{i}{2B} [D(x''^2 - x'''^2) - 2x(x'' - x''')\right\}.$$
(10)

On the other hand, when the beam  $\psi(x')$  propagates through the [D(-B)(-C)A] optical system,

according to the Fresnel integration (7)-(8), we have

$$\begin{aligned} |\phi(x)|^{2} &= \int_{-\infty}^{\infty} dx' \vec{\mathcal{K}}(x,x') \psi(x') \int_{-\infty}^{\infty} dx'' \psi^{*}(x'') \vec{\mathcal{K}}^{*}(x,x'') \\ &= \frac{1}{2\pi B} \int_{-\infty}^{\infty} dx' \exp\left[\frac{-i}{2B} \left(Dx'^{2} - 2x'x + Ax^{2}\right)\right] \psi(x') \\ &\times \int_{-\infty}^{\infty} dx'' \exp\left[\frac{i}{2B} \left(Dx''^{2} - 2x''x + Ax^{2}\right)\right] \psi^{*}(x'') \\ &= \frac{1}{2\pi B} \iint_{-\infty}^{\infty} dx' dx'' \psi(x') \psi^{*}(x'') \exp\left\{\frac{i}{2B} \left[D\left(x''^{2} - x'^{2}\right) + 2x\left(x' - x''\right)\right]\right\}, \quad (11) \end{aligned}$$

which is the same as R(x) in (10). So combining (4), (10)-(11), (7) and (8) we reach the conclusion

$$\left|\frac{1}{\sqrt{-2\pi iB}}\int_{-\infty}^{\infty}\exp\left[\frac{-i}{2B}\left(Dx'^{2}-2x'x+Ax^{2}\right)\right]\psi\left(x'\right)dx'\right|^{2}=\iint_{-\infty}^{\infty}dx'd\nu'\delta\left(x-Dx'+B\nu'\right)W_{\psi}(\nu',x'),$$
(12)

where DA - BC = 1. The physical meaning of Eq. (12) is: when an input field propagates through an optical [D(-B)(-C)A] system, the energy density of the output field is equal to the Radon transform of the Wigner function of the input field. So far as our knowledge is concerned, this conclusion seems new.

Eq. (8) is the relationship between the input amplitude and output one in spatial-domain. Now we turn the above discussion to the case of space-frequency domain.

For a [D(-B)(-C)A] optical system, the Collins' diffraction integral formula in space-frequency (angle spectrum [11]) domain is [12]

$$\widetilde{\phi}(v) = \int_{-\infty}^{\infty} \widetilde{\mathcal{K}}(\nu, \nu') \,\widetilde{\psi}(\nu') \, d\nu', \tag{13}$$

where the kernel is

$$\widetilde{\mathcal{K}}(\nu,\nu') = \frac{1}{\sqrt{2\pi i C}} \exp\left[\frac{i}{2C} \left(D\nu^2 - 2\nu'\nu + A\nu'^2\right)\right].$$
(14)

On the other hand, the Wigner function expressed in terms of the space-frequency field is

$$W_{\tilde{\psi}}(\nu',x') = \int_{-\infty}^{+\infty} \frac{ds}{2\pi} e^{-ix's} \tilde{\psi}^* \left(\nu' + \frac{s}{2}\right) \tilde{\psi} \left(\nu' - \frac{s}{2}\right).$$
(15)

In space-frequency domain the Radon transform is along an infinitely thin phase space strip denoted by the real parameters A, C,

$$R(\nu) \equiv \iint_{-\infty}^{\infty} dx' dp' \delta\left(\nu - A\nu' + Cx'\right) W_{\tilde{\psi}}(\nu', x')$$
(16)

rather than the parameters D, B of the Radon transform (4) in spatial-domain. We want to examine if the above conclusion still holds in the space-frequency domain. For this propose, we rewrite (15) as

$$W_{\tilde{\psi}}(\nu', x') = \iint_{-\infty}^{\infty} d\nu'' d\nu''' \int_{-\infty}^{+\infty} \frac{ds}{2\pi} e^{-ix's} \tilde{\psi}^*(\nu'') \tilde{\psi}(\nu''') \delta\left(\nu'' - \nu' - \frac{s}{2}\right) \delta\left(\nu' - \frac{s}{2} - \nu'''\right)$$
$$= \iint_{-\infty}^{\infty} d\nu'' d\nu''' \tilde{\psi}^*(\nu'') \tilde{\psi}(\nu''') e^{-i2x'(\nu'' - \nu')} \delta\left(2\nu' - \nu'' - \nu'''\right)$$
(17)

Then we substitute (17) into (16), the result is

$$R(\nu) = \frac{1}{\pi} \iint_{-\infty}^{\infty} dx' d\nu' \delta\left(\nu - A\nu' + Cx'\right) \iint_{-\infty}^{\infty} d\nu'' d\nu''' \tilde{\psi}^*\left(\nu''\right) \tilde{\psi}\left(\nu'''\right) e^{-i2x'\left(\nu'' - \nu'\right)} \delta\left(2\nu' - \nu'' - \nu'''\right)$$
$$= \frac{1}{2\pi C} \iint_{-\infty}^{\infty} d\nu'' d\nu''' \tilde{\psi}^*\left(\nu''\right) \tilde{\psi}\left(\nu'''\right) \exp\left\{-\frac{i}{2C} \left[A\left(\nu''^2 - \nu'''^2\right) - 2\nu\left(\nu'' - \nu'''\right)\right]\right\}.$$
(18)

On the other hand, from (13) we calculate

$$\begin{split} |\tilde{\phi}(\nu)|^{2} &= \frac{1}{2\pi C} \int_{-\infty}^{+\infty} \tilde{\psi}(\nu') \exp\left[\frac{i}{2C} \left(D\nu^{2} - 2\nu'\nu + A\nu'^{2}\right)\right] d\nu' \\ &\times \int_{-\infty}^{+\infty} \tilde{\psi}^{*}(\nu'') \exp\left[\frac{-i}{2C} \left(D\nu^{2} - 2\nu''\nu + A\nu''^{2}\right)\right] d\nu'' \\ &= \frac{1}{2\pi C} \int \int_{-\infty}^{+\infty} d\nu' d\nu'' \tilde{\psi}(\nu') \tilde{\psi}^{*}(\nu'') \exp\left[-\frac{i}{2C} \left(A \left(\nu''^{2} - \nu'^{2}\right) - 2\nu \left(\nu'' - \nu'\right)\right)\right]. \end{split}$$
(19)

Comparing Eq. (19) with Eq. (18) and using (13)-(14) we see

$$\left|\frac{1}{\sqrt{2\pi iC}}\int \exp\left[\frac{i}{2C}\left(D\nu^2 - 2\nu'\nu + A\nu'^2\right)\right]\tilde{\psi}\left(\nu'\right)d\nu'\right|^2 = \iint_{-\infty}^{\infty} dx'dp'\delta\left(\nu - A\nu' + Cx'\right)W_{\tilde{\psi}}(\nu', x'),\tag{20}$$

this is the theorem expressed in space-frequency domain.

We now take an example to confirm the theorem. When the input field is described by a Gaussian function (Gaussian chirplet)

$$\psi_0\left(x'\right) \equiv \sqrt[4]{\frac{\epsilon}{\pi}} \exp\left\{-\left(\epsilon - i\beta\right)\frac{x'^2}{2}\right\}, \ \epsilon > 0,$$
(21)

which depicts a Gaussian windowed linear chirp signal [13]. Using Eq.(1) we obtain its Wigner function

$$W_{\psi_0}(\nu', x') = e^{-\epsilon x'^2} \sqrt{\frac{\epsilon}{\pi}} \int_{-\infty}^{+\infty} \frac{du}{2\pi} e^{-\frac{1}{4}\epsilon u^2 + i(\nu' - \beta x')u} \\ = \frac{1}{\pi} \exp\left\{-\left[\epsilon x'^2 + \frac{1}{\epsilon} (\nu' - \beta x')^2\right]\right\},$$
(22)

which shows that the energy of Gaussian chirplet is concentrated at  $\nu' = \beta x'$ . According to Eq. (4), the Radon transform of (22) with the parameters D, B is

$$\int_{-\infty}^{\infty} dx' d\nu' \delta \left(x - Dx' + B\nu'\right) W_{\psi_0}(\nu', x')$$

$$= \frac{1}{\pi D} \int_{-\infty}^{+\infty} d\nu' \exp\left\{-\frac{\epsilon}{D^2} \left(x + B\nu'\right)^2 - \frac{1}{\epsilon D^2} \left[\left(D - \beta B\right)\nu' - \beta x\right]^2\right\}$$

$$= \sqrt{\frac{\epsilon/\pi}{\left(D - B\beta\right)^2 + B^2\epsilon^2}} \exp\left[\frac{-\epsilon x^2}{\left(D - B\beta\right)^2 + B^2\epsilon^2}\right] \equiv R_0(x).$$
(23)

On the other hand, according to (7) and (8), the Fresnel transform of input field  $W_{\psi_0}(\nu', x')$  through a [D(-B)(-C)A] optical system is given by

$$\frac{1}{\sqrt{-2\pi iB}} \sqrt[4]{\frac{\epsilon}{\pi}} \int_{-\infty}^{\infty} dx' \exp\left[\frac{-i}{2B} \left(Dx'^2 - 2x'x + Ax^2\right) - (\epsilon - i\beta)\frac{x'^2}{2}\right]$$
$$= \sqrt[4]{\frac{\epsilon}{\pi}} \frac{1}{\sqrt{-i}} e^{-\frac{iA}{2B}x^2} \sqrt{\frac{1}{\epsilon B - i\left(\beta B - D\right)}} \exp\left[\frac{-x^2}{2B\left(B\epsilon + iD - i\beta B\right)}\right] \equiv \phi_0\left(x\right).$$
(24)

Comparing (24) with (23) and noticing

$$\frac{1}{2B\left(B\epsilon+iD-i\beta B\right)} + \frac{1}{2B\left(B\epsilon-iD+i\beta B\right)} = \frac{\epsilon}{\left(D-B\beta\right)^2 + B^2\epsilon^2}$$
(25)

we see

$$|\phi_0(x)|^2 = R_0(x), \qquad (26)$$

as expected.

In summary, we have derived a new theorem governing the connection between optical field's Fresnel transform and its Wigner function's Radon transformation, since both the Wigner function and the Fresnel transform are widely used in optical propagation, we hope this theorem would have new applications in the analysis of optical communication and optical tomography. For the application of Wigner function in deriving the quantum-mechanical photocount formula, we refer to [14].

This work was supported by the National Natural Science Foundation of China under grant 10775097.

## References

- [1] E. P. Wigner, "On the quantum correction for thermodynamic equilibrium", Phys. Rev. 40, 749 (1932).
- [2] M. j. Bastiaans, "The Wigner distribution function applied to optical signals and systems", Opt. Commun. 25, 26 (1978).
- [3] M. j. Bastiaans, "Wigner distribution function and its application to first-order optics", J. Opt. Soc. Am. 69, 1710 (1979).
- [4] A. W. Lohmann, "Image rotation, Wigner rotation, and the fractional Fourier transform," J. Opt. Soc. Am. A 10, 2181–2186 (1993).
- [5] J. Radon, "Uber die Bestimmung von Funktionen Durch Ihre Integralwerte Langs Gewisser Mannigfaltigkeiten", Ber. Verh. Saechs. Akad. Wiss. Leipzig Math.Phys.K1. 69, 262-267, (1917).
- [6] Y. Zhang, B. Gu, B. Dong, and G. Yang, "Optical implementations of the Radon–Wigner display for one-dimensional signals," Opt. Lett. 23, 1126-1128 (1998).
- [7] For quantum states' reconstruction, see e.g., Wolfgang P. Schleich, Quantum Optics in Phase Space, (Wiley-VCH, Birlin, 2001) and references therein
- [8] S. A. Collins, "Lens-system diffraction integral written in terms of matrix optic", J. Opt. Soc. Am. A 60, 1168-1177 (1970).
- [9] J. A. Arnaud, "Mode Coupling in First-Order Optics," J. Opt. Soc. Am. 61, 751-758 (1971).
- [10] H. Fan and H. Lu, "Collins diffraction formula studied in quantum optics," Opt. Lett. 31, 2622-2624 (2006).
- [11] J. W. Goodman, Introduction to Fourier Optics, (Mc Graw-Hill, 1968).
- [12] Z. Liu, X. Xiu and D. Fan, "Collins formula in frequency-domain and fractional Fourier transforms", Opt. Commun. 155, 7 (1998).
- [13] A. Papoulis, Signal Analysis, (McGraw-Hill, New York, 1977).
- [14] Hong-yi Fan and Li-yun Hu, "Two quantum-mechanical photocount formulas", Opt. Lett. 33, 443-445 (2008).