The Bi-Musquash Conjecture

by Jon Henry Sanders jon_sanders@partech.com

Bi-Musquash Conjecture

A construction (attributed to Conway) for 'doubling' a thrackled m-gon of odd length m to get a thrackled 2mgon (of even length) was given in [1]. Has anyone noticed that if this construction is applied to an odd length musquash we get a 'bi-musquash', i.e., a thrackled *n*-gon, n = 2m, m odd, with the property that row 1 of the intersection table generates the odd rows and row 2 generates the even rows? More precisely, for the definition of a 'bi-musquash' we require that¹ :

(d') If e_1 intersects edges in the order: $e_{k_1}, \ldots, e_{k_{n-3}}$, and e_2 intersects edges in the order $e_{j_1}, \ldots, e_{j_{n-3}}$ then for (odd) $i = 3, 5, \ldots, n-1$, edge e_i intersects edges in the order $e_{k_1+i-1}, \ldots, e_{k_{n-3}+i-1}$, and for (even) $i = 4, \ldots, n$, edge e_i intersects edges in the order : $e_{j_1+i-2}, \ldots, e_{j_{n-3}+i-2}$, where the edge subscripts are computed modulo n. Note that by this definition, an even musquash is automatically a bi-musquash. Remarks of [2,3] imply that bi-musquashes must have n/2 rotational symmetry and indeed the drawing of the bi-musquash of length 14 [Fig.1], as well the 'chordal representation' ² of it's Gauss word [Fig.2] exhibit this. I guess the conjecture to prove³ would be that bi- musquashes do not exist for n = 4m and that they are essentially⁴ unique for each n, with $n = 2 \pmod{4}$. The intersection table of bi-musquash 14 is shown below.

	3 7	12	5	10	3	13	6	11	4	9
	8	6	13	4	11	7	14	5	12	10
1) 9	14	7	12	5	1	8	13	6	11
1	L 10	8	1	6	13	9	2	7	14	12
1	2 11	2	9	14	7	3	10	1	8	13
1.	3 12	10	3	8	1	11	4	9	2	14
1	1 13	4	11	2	9	5	12	3	10	1
	L 14	12	5	10	3	13	6	11	4	2
	2 1	6	13	4	11	7	14	5	12	3
	3 2	14	7	12	5	1	8	13	6	4
	1 3	8	1	6	13	9	2	7	14	5
	5 4	2	9	14	7	3	10	1	8	6
	5 5	10	3	8	1	11	4	9	2	7
	7 6	4	11	2	9	5	12	3	10	8

Intersection Table of Bi-Musquash 14

The first two rows generate the table in the required way. In general, the conjecture is that the first two rows (generators) of a bi-musquash of length 2p, p odd, are either:

SET1

Row 1: p+1, p,2p-2,p-2,2p-4,p-4,...,p+3,3,2p-1,p-1,2p-3,p-3,...,p+4,4,p+2 **Row 2**: p+2,p+1, p-1,2p-1, p-3,2p-3,..., 4,p+4, p, 2p,p-2,2p-2,...,5,p+5,p+3

or

SET2 Row 1: p,2p-2,p-2,2p-4,p-4,...,p+3,3,2p-1,p-1,2p-3,p-3,...,p+4,4,p+2, p+1 Row 2: p+1, p-1,2p-1, p-3,2p-3,..., 4,p+4, p, 2p,p-2,2p-2,...,5,p+5,p+3,p+2,

each giving rise to a mirror image of the other.

¹ This is a modification of condition (d) of [2].

² i.e., the arrangement of the symbols of a Gauss word around a circle with a chord drawn between each pair of identical symbols

³ G. Cairns has pointed out to me that " there are two inequivalent musquash of the 6-gon; one being the Conway doubling of the 3-

gon". Thus the case n = 6 of the bi-musquash conjecture needs to be modified appropriately.

i.e., there exist precisely two and they are reflections of each other




Figure 1 Bi-Musquash 14




Figure 2 Chordal Representation of Bi-Musquash 14

References

1. D.R. Woodall, *Thrackles and deadlock*, Combinatorial Math. and Its Applications (D. Welsh, ed.), Academic Press, 1971, pp. 335–347.

- 2. G. Cairns and D.M. King, The Answer to Woodall's Musquash Problem, Discrete Math., 207 (1999), 25-32.
- 3. G. Cairns and D.M. King, All Odd Musquashes are Standard, Discrete Math., 226 (2001), 71-91.