On the Cayley semigroup of a finite aperiodic semigroup

Avi Mintz[∗]

October 30, 2018

Abstract

Let S be a finite semigroup. In this paper we introduce the functions $\varphi_s : S^* \to S^*$, first defined by Rhodes, given by $\varphi_s([a_1, a_2, \ldots, a_n]) =$ $[sa_1, sa_1a_2, \ldots, sa_1a_2 \cdots a_n]$. We show that if S is a finite aperiodic semigroup, then the semigroup generated by the functions $\{\varphi_s\}_{s\in S}$ is finite and aperiodic.

1 Introduction

Let S be a finite semigroup. Rhodes considered in 1965 [\[1\]](#page-23-0) a function defined on the free monoid S^* . He called this the machine of the semigroup. Formally, to each element s of the semigroup, he assigned a function $\varphi_s : S^* \to S^*$, defined by $\varphi_s([a_1, a_2, \ldots, a_n]) =$ $[sa_1, sa_1a_2, \ldots, sa_1a_2 \cdots a_n]$. Essentially, the φ_s arise from considering the Cayley graph to be a sequential machine and assigning a function to each state $s \in S$ considered as a start state. We call the semigroup generated by all such functions the Cayley semigroup of S , and denote it by $Cayley(S)$.

Rhodes showed that if S is finite, then S is aperiodic, that is, has no non-trivial subgroups, if and only if for every $s \in S$, there exists an $n \in \mathbb{N}$ such that $\varphi_s^n = \varphi_s^{n+1}$. This construction played a key role in the original Krohn-Rhodes Decomposition Theorem.

Grigorchuk and others have intensively studied semigroups and groups generated by finite automata and these have many important properties. Let \mathbb{Z}_2 denote the cyclic group of order 2, and denote its elements by $\{1, x\}$. The above gives two functions φ_1 and φ_x which are both invertible as functions $\mathbb{Z}_2^* \to \mathbb{Z}_2^*$. A natural question is to describe the group generated by them.

In [\[3\]](#page-23-1) and [\[4\]](#page-23-2) this group was proved to be $\mathbb{Z}_2wr\mathbb{Z}$, also known as the lamplighter group. Later on this result was generalized by Silva

[∗]This research was done as part of the requirement for a PhD in Bar Ilan University and was done under supervision of Stuart Margolis

and Steinberg to all abelian group in [\[6\]](#page-23-3) with some results on nonabelian groups. Recently, Grigorchuk and Zuk, used these techniques to calculate the spectrum of the lamplighter group [\[4\]](#page-23-2).

In this paper we study the Cayley semigroups of other finite semigroups. The first result is that if the semigroup S divides the semigroup T, then $Cayley(S)$ divides $Cayley(T)$. This motivates us to work with semigroups that are not divisible by groups, i.e. aperiodic semigroups. The main result, generalizing Rhodes $[1]$ is that if S is a finite aperiodic semigroup then $Cayley(S)$ is a finite and aperiodic semigroup.

2 Preliminaries

Let S be a semigroup. If S is not a monoid, we denote by S^1 the monoid $S \cup \{1\}$ with the obvious multiplication making 1 the identity element. If S happens to be a monoid already, S^1 will equal S. In a similar way we can add a zero to a semigroup S , if it doesn't have one. This semigroup is denoted by S^0 . For a set X we write X^* for the directed tree of all strings over X, and ϵ for the empty word. For all $s \in S$ define $\varphi_s : (S^1)^* \to S^*$ by $\varphi_s([a_1, a_2, \ldots, a_n]) = [sa_1, sa_1a_2, \ldots, sa_1a_2 \cdots a_n],$ and $\varphi_s(\epsilon) = \epsilon$. We note that unless S is a group, these functions are not invertible. We consider the semigroup generated by $\{\varphi_s\}_{s\in S}$, and denote it by Cayley(S). We will use φ_s, φ_t to denote generators of $Cayley(S)$ and f, g to denote elements of $Cayley(S)$.

2.1 Semigroup Actions

A semigroup S acts on the left of a set Y if there is a correspondence $S \times Y \to Y$ $(s, y) \mapsto sy$ that satisfies $(st)y = s(ty)$.

A special case of this is when Y has the structure of a rooted tree. In our case we mean that every $y \in Y$ has a depth, denoted by |y| and every two elements $y_1, y_2 \in Y$ have a least upper bound (lub) denoted by $y_1 \vee y_2$, which is the longest common path on the geodesics from the root to y_1 and y_2 .

We assume the reader is familiar with these notions, but mention that in the case of strings X^* over an alphabet X there is a natural structure of a tree.

For example $|abbab| = 5$ and $abbaab \vee abb = abb$.

For two strings w_1, w_2 , we call w_1 a prefix of w_2 if for some w_3 we have $w_1w_3 = w_2$. Thus $u \vee v$ is the longest common prefix of u and v.

Definition 1. Let $f: X^* \to X^*$. We say that f is a tree endomor*phism if for every* $w_1, w_2 \in W^*$ *we have*

- *1.* $|w_1| = |f(w_1)|$,
- *2.* $f(w_1 \vee w_2)$ *is a prefix of* $f(w_1) \vee f(w_2)$ *.*

2.2 Mealy Automata

We recall that a Mealy automata is a 5-tuple $\langle A, Q, i, \delta, \lambda \rangle$, with A, Q finite sets, $i \in Q$, $\delta: Q \times A \to Q$ and $\lambda: Q \times A \to A$. A Mealy

automata has a representation as a labeled directed graph. The vertices of the graph are the elements of Q , and for every q in Q and a in A there is an edge from q to $\delta(q, a)$ labeled with $a/\lambda(q, a)$. For every word $w \in A^*$, we have a walk on the graph, by starting at the vertex i and each time we see a letter a of w we walk from the state q to the state $\delta(q, a)$. Since this is a labeled graph, we can read the labels as we read the word $w = [a_1, a_2, \ldots, a_n]$ and obtain the output word $f_i(w) = [\lambda(i, a_1), \lambda(\delta(i, a_1), a_2), \ldots, \lambda(\delta(\cdots(\delta(i, a_1), \cdots, a_{n-1}), a_n)].$ Thus a Mealy automata defines a function $f_i: A^* \to A^*$.

We can also ignore the state i , and thus the Mealy automaton defines a collection of functions $A^* \to A^*$, each one determined by its initial state i.

Let S be a semigroup. The (right) Cayley graph of S is the graph described by a set of vertices S^1 , and for every s_1, s_2 in S there is an edge from s_1 to s_1s_2 .

We can consider the Cayley graph of a finite semigroup as a Mealy automaton by labeling the edge from s_1 to s_1s_2 with s_2/s_1s_2 . Using the formalities of Mealy automata this becomes $Q = S, A = S¹$ and $\delta(q, a) = \lambda(q, a) = qa.$

2.3 Examples

Let's consider the five semigroups of order 2, and their Cayley automata.

1. $S_1 = \langle a, b | a^2 = ab = a, b^2 = ba = b \rangle$, the left zero semigroup of order 2.

a $a,b,1/a$ $\overline{}$ b $a,b,1/b$ j

 $\varphi_a([x_1, x_2, \ldots, x_n]) = [ax_1, ax_1x_2, \ldots, ax_1x_2 \cdots x_n] = [a, a, \ldots, a].$ We get $Cayley(S_1) \cong S_1$.

2. $S_2 = \langle a, b | a^2 = ba = a, b^2 = ab = b \rangle$, the right zero semigroup of order 2.

$$
\bigcap^{a,1/a}_{a}\bigcap_{a} \bigoplus_{a/a}^{b/b} \bigcap^{b,1/b}
$$

 $\varphi_a([x_1, x_2, \ldots, x_n]) = [ax_1, ax_1x_2, \ldots, ax_1x_2 \cdots x_n].$ If we write this as $\varphi_a([x_1, x_2, ..., x_n]) = [y_1, y_2, ..., y_n]$, then we have $y_i = x_j$ where j is the largest number less than i such that $x_j \neq 1$. If no such j exists, then $y_i = a$ (this a comes from φ_a).

We verify that $\varphi_x \varphi_y = \varphi_y$ for $x, y \in \{a, b\}$ so that $Cayley(S_2) \cong \emptyset$ $S₂$.

3. $S_3 = (\{0,1\}, \cdot)$, the 2-element semilattice.

$$
\bigcap_{0 <_{0/0} 1}^{1/0} \bigcap
$$

 $0,$

 S_3 is a monoid so we don't need to add a 1 to the graph.

 $\varphi_1([x_1, x_2, \ldots, x_n]) = [1x_1, 1x_1x_2, \ldots, 1x_1x_2 \cdots x_n]$ $\varphi_0([x_1, x_2, \ldots, x_n]) = [0, 0, 0, 0, 0]$

It is clear that φ_0 inputs a string of length n and outputs a string of 0's of length n .

Let k be the smallest such that $x_k = 0$. Then $\varphi_1([x_1, x_2, \ldots, x_n]) =$ $[1, 1, \ldots, 1, 0, 0, \ldots, 0]$ where the first 0 is at the k place. We get $Cayley(S_3) \cong S_3$.

4. $S_4 = \langle x | x^2 = 0 \rangle$, the 2 element nil semigroup.

 $\boldsymbol{0}$ $0, x, 1/0$ $\xleftarrow{x,0/0} x$ $1/x$ l $\varphi_x([x_1, x_2, \ldots, x_n]) = [xx_1, xx_1x_2, \ldots, xx_1x_2 \cdots x_n] = [0, 0, \ldots, 0]$ $\varphi_x([1]) = [x]$ $\varphi_0([x_1, x_2, \ldots, x_n]) = [0, 0, \ldots, 0]$ We get $Cayley(S_4) \cong S_4$.

5. $S_5 = \mathbb{Z}_2 = \langle x | x^2 = 1 \rangle$, the cyclic group of order 2.

$$
\bigcirc^{0/0} \bigcirc \bigcirc \bigcirc \frac{1/1}{1/0} \bigcirc \bigcirc^{0/1}
$$

It is proved in [\[6\]](#page-23-3) that $Cayley(S) \cong \{a, b\}^+$, the free semigroup on two letters. In this case, φ_x and φ_1 are invertible and the group generated by $\{\varphi_x, \varphi_1\}$ is isomorphic to the lamplighter group $\mathbb{Z}_2wr\mathbb{Z}$.

Another interesting example is the following. Let S be any monoid and 1 the identity element. In general, $\varphi_1([a_1, a_2, \ldots, a_n])$ $[a_1, a_1 a_2, \ldots, a_1 a_2 \cdots a_n] \neq [a_1, a_2, \ldots, a_n]$, so the identity element of a semigroup is not the identity element of $Cayley(S)$. Actually, except for trivial cases, $Cayley(S)$ does not contain an identity element.

2.4 Finite Semigroups

We assume standard terminology, facts and notation of finite semigroup theory and refer the reader to [\[1\]](#page-23-0) and [\[2\]](#page-23-4) for more details.

Let A and B denote finite sets. Let G be a finite group. Let $C : B \times A \to G \cup \{0\}$. We usually think of C as a $B \times A$ matrix. Furthermore, assume C has the property that for every $a \in A$ ($b \in B$) there is a $b \in B$ $(a \in A)$ such that $C(b, a) \neq 0$. A semigroup can be constructed, as follows, from A, B, G, C , denoted by $\mathcal{M}^0(A, B, G; C)$ and called a Rees matrix semigroup.

Let $\mathcal{M}^0(A, B, G; C) = (A \times G \times B) \cup \{0\}$ as sets. 0 is a zero element. It remains to describe the product of elements in $A \times G \times B$.

 $(a_1, g_1, b_1) \cdot (a_2, g_2, b_2) = (a_1, g_1 C(b_1, a_2) g_2, b_2)$ if $C(b_1, a_2) \neq 0$ and $(a_1, g_1, b_1) \cdot (a_2, g_2, b_2) = 0$ otherwise.

We state without proof the following theorem.

Theorem 2 (Rees-Sushkevitch). *A finite semigroup* S *is* 0*-simple if and only if it is isomorphic to a Rees matrix semigroup.*

Corollary 3.

- *1. A simple aperiodic semigroup is of the form* A × B*, with multiplication* $(a_1, b_1) \cdot (a_2, b_2) = (a_1, b_2)$ *, and in particular, is an idempotent semigroup.*
- *2.* A 0*-simple aperiodic semigroup is of the form* $(A \times B) \cup \{0\}$, *with multiplication defined by* $(a_1, b_1) \cdot (a_2, b_2) = (a_1, b_2)C(b_1, a_2)$, *where* C *is a* $B \times A$ *matrix over* $\{0, 1\}$ *.*
- *3. A non regular* J *class of a finite aperiodic semigroup is of the form* A × B *with null multiplication.*

Proof.

- 1. If S is a simple aperiodic semigroup, then S^0 is a 0-simple semigroup, and thus S^0 is isomorphic to $M = \mathcal{M}^0(A, B, G; C)$. Since M has a subgroup isomorphic to G (this can be shown by choosing a, b such that $C(b, a) \neq 0$ and considering $\{(a, g, b)\}\mathfrak{g} \in G$, and M is aperiodic, we see that $G = \{1\}$. Since S is a subsemigroup of S^0 , the matrix C does not contain any zeros.
- 2. The proof of the second point is similar.
- 3. This follows from the basic theorems and definitions of the Schützenberger group. See [\[2\]](#page-23-4).

Another important corollary is the following.

Corollary 4. In a 0-simple semigroup, $s_1 s_2 \cdots s_n = 0$ if and only if $s_i s_{i+1} = 0$ *for some* $1 \le i \le n-1$ *.*

We would like to extend the last two corollaries to a semigroup acting on a 0-minimal ideal.

Lemma 5. *Let* S *be an aperiodic semigroup, and* $I \cong A \times B \cup \{0\}$ *a* 0-minimal regular ideal. Let $s \in S$ and $(a, b) \in I \setminus \{0\}$.

- 1. $s(a,b) = 0$ *if and only if* $s(a,b') = 0$ *for every* (a,b') *in the* \mathcal{R} *class of* (a, b)*.*
- 2. If $s(a, b) \neq 0$ then $s(a, b) = (a', b)$ for some $a' \in A$ which depends *only on* a *and* s*.*

We will write this a' as sa.

Proof. Both statements are a result of the fact that the \mathcal{R} relation is a left congruence (aRb) implies $xaRxb$. For the second point, choose an $a_b \in A$ such that $C(b, a_b) = 1$ and apply the associative law to $s(a, b)(a_b, b).$ \Box

Notation 6. This justifies an extended notation for the structure matrix of I. $C : S \times A \to \{0, 1\}$. $C(s, a) = 0$ if $s(a, b) = 0$ and $C(s, a) = 1$ if $s(a, b) \in A \times B$.

 \Box

Notation 7. Furthermore this shows that S has a left action on the set $A \cup \{0\}$, where sa is the a' in the formula $s(a, b) = (a', b)$. We have shown that this does not depend on choice of b.

With every $\mathcal J$ class of a semigroup S we can associate a 0-simple or null semigroup, which we will now define.

2.5 Trace of a $\mathcal J$ class

Let S be a semigroup, and J a $\mathcal J$ class of S. Let θ be an element disjoint from S and consider the following multiplication on J. If $j_1 \cdot j_2$ isn't in J we write their product as θ . The set $J \cup {\theta}$ with θ as a zero element is a semigroup called the trace of J , and denoted by J^{tr} . It is known (and easy to show) that $J^{tr} \prec S$. Another possible construction of J^{tr} is by defining T to be the ideal $S^1 J S^1 \setminus J$ and then $J^{tr} \cong S^1 J S^1 / T$.

2.6 Semidirect product of semigroups

Let S and T be finite semigroups. Suppose that T acts on S on the left. That is, for every $s \in S$ and $t \in T$ there is an element of S denoted by ^ts such that ^{t₁(t₂s) = t₁t₂ s and ^t(s₁s₂) = t^t_s^t₁s₂. In this case we can} define the (left) semidirect product of S and T whose underlying set is $S \times T$ with multiplication given by $(s_1, t_1)(s_2, t_2) = (s_1^{t_1} s_2, t_1 t_2)$. This semigroup is denoted by $S \rtimes T$. There is also the dual notion of a right semidirect product.

3 Fractalness Property of $Cayley(S)$

3.1 The Pascal Array

In [\[1\]](#page-23-0) Rhodes introduced the notion of a Pascal array. This gives some intuition to the action of $Cayley(S)$. This construction motivates many of the following proofs.

Let $s_1, s_2, \ldots, s_n \in S$ and $a_1, a_2, \ldots, a_k \in S^1$. We describe the $(n+1)\times (k+1)$ table (t_{ij}) . t_{00} is left empty, $t_{0i} = a_i$ and $t_{j0} = s_j$. The table is completed by the formula $t_{mn} = t_{m(n-1)} \cdot t_{(m-1)n}$.

The table describes the action of the function $\varphi_{s_n} \cdots \varphi_{s_3} \varphi_{s_2} \varphi_{s_1}$ on the word $[a_1, a_2, \ldots, a_k]$. The bottom row (excluding the first column) is the output.

3.2 Fractalness

Let $f = \varphi_{s_n} \dots \varphi_{s_2} \varphi_{s_1} \in Cayley(S)$ and let $v \in (S^1)^{\star}$. Since $Cayley(S)$ is a semigroup of tree endomorphisms, the function $g : (S^1)^* \to S^*$ defined by $f(vw) = f(v) \circ g(w) \forall w \in (S^1)^*$ is well-defined. We denote this function g by f_v .

The following theorem will show that f_v is in $Cayley(S)$ whenever $f \in Cayley(S)$ and $v \in (S^1)^*$. Let $p : S^* \setminus {\{\epsilon\}} \to S$ be the function defined by $p([v_1, v_2, \ldots, v_n]) = v_n$. Observe that for every $f \in Cayley(S)$ and every $w \in (S^1)^*$ $w \in (S^1)^*$ $w \in (S^1)^*$ we have $pf(w) \in S$ (and not¹ in S^*). This shows that writing $\varphi_{pf(w)}$ is meaningful.

Theorem 8. Let $f \in Cayley(S)$. Let $v \in (S^1)^*$, and let $f_v : (S^1)^* \to$ S^* *be defined by* $f(vw) = f(v) \circ f_v(w)$ *. If we write* f *as* $f = \varphi_{s_n} \dots \varphi_{s_2} \varphi_{s_1}$ then $f_v = \varphi_{(p\varphi_{s_n}\cdots \varphi_{s_1}(v))}\cdots \varphi_{(p\varphi_{s_2}\varphi_{s_1}(v))}\varphi_{(p\varphi_{s_1}(v))}$. We also have $f_{\epsilon} =$ *f.* Thus if $f \in Cayley(S)$, so is f_v for all $v \in (S^1)^*$.

Proof. This can be seen by observing the Pascal array. When we apply $\varphi_{s_n} \cdots \varphi_{s_1}$ to vw, we can look at the column of the last letter of v. The elements there are acting on w, describing the action of f_v on w. For a full proof we induct on n. If $n = 1$ then let $w = [w_1, w_2, \ldots, w_k]$ and $v = [v_1, v_2, \dots, v_l]$ and write $\overline{v} = v_1 v_2 \dots v_l$. Notice that $p\varphi_s(v) = s\overline{v}$ for every s. We need to show that $\varphi_s(vw) = \varphi_s(v) \circ \varphi_{p(\varphi_s(v))}(w)$.

Calculating gives us

$$
\varphi_s(vw) = \varphi_s([v_1, v_2, \dots, v_{l-1}, v_l, w_1, \dots, w_k])
$$

= $[sv_1, sv_1v_2, \dots, sv_1v_2v_{l-1}, s\bar{v}, s\bar{v}w_1, \dots, s\bar{v}w_1 \dots w_k]$
= $[sv_1, sv_1v_2, \dots, sv_1v_2v_{l-1}, s\bar{v}] \circ [s\bar{v}w_1, \dots, s\bar{v}w_1 \dots w_k]$
= $\varphi_s(v) \circ \varphi_{s\bar{v}}(w)$
= $\varphi_s(v) \circ \varphi_{p(\varphi_s(v))}(w)$

Let's assume this is correct for $n-1$. Let v and w be as before. Using the induction assumption (at the first $=$ sign) and the $n = 1$ case (at the second), we get:

 $\varphi_{s_n} \varphi_{s_{n-1}} \dots \varphi_{s_1}(vw) =$ $\varphi_{s_n}(\varphi_{s_{n-1}}\dots\varphi_{s_1}(v)\circ\varphi_{(p\varphi_{s_{n-1}}\cdots\varphi_{s_1}(v))}\cdots\varphi_{(p\varphi_{s_2}\varphi_{s_1}(v))}\varphi_{(p\varphi_{s_1}(v))}(w))=$ $\varphi_{s_n}(\varphi_{s_{n-1}}\dots\varphi_{s_1}(v))\circ\varphi_{(p\varphi_{s_n}(\varphi_{s_{n-1}}\cdots\varphi_{s_1}(v)))}\dots\varphi_{(p\varphi_{s_2}\varphi_{s_1}(v))}\varphi_{(p\varphi_{s_1}(v))}(w))=$ $\varphi_{s_n} \varphi_{s_{n-1}} \dots \varphi_{s_1}(v) \circ (\varphi_{(p \varphi_{s_n} \cdots \varphi_{s_1}(v))} \dots \varphi_{(p \varphi_{s_2} \varphi_{s_1}(v))} \varphi_{(p \varphi_{s_1}(v))})(w)$

 \Box

4 Preliminary Results

4.1 Cayley as a Functor

Lemma 9. *1.* If T is a quotient semigroup of S then $Cayley(T)$ is *a quotient of* Cayley(S)*.*

¹We do not identify strings of length 1 over S with S

2. If T is a subsemigroup of S then $Cayley(T) \prec Cayley(S)$.

- *Proof.* 1. Let $F : S \to T$ be a surjective morphism. We extend F to $F: S^1 \to T^1$ and verify that this is a semigroup morphism. Write \bar{s} for $F(s)$. Suppose we have in $Cayley(S)$ the following equation $\varphi_s([a_1, a_2, \ldots, a_n]) = [sa_1, sa_1a_2, \ldots, sa_1a_2 \cdots a_n].$ Then, applying $t \mapsto \overline{t}$ gives us $\varphi_{\overline{s}}([\overline{a_1}, \overline{a_2}, \ldots, \overline{a_n}]) = [\overline{sa_1}, \overline{sa_1a_2}, \ldots, \overline{sa_1a_2 \cdots a_n}]$ This shows that the mapping $s \mapsto \bar{s}$ can be extended to $\varphi_s \mapsto \varphi_{\bar{s}}$. We now extend this mapping to $Cayley(S) \rightarrow Cayley(T)$. It remains to show that this is well defined, i.e. if $\varphi_{s_n} \cdots \varphi_{s_2} \varphi_{s_1} =$ $\varphi_{r_m} \cdots \varphi_{r_2} \varphi_{r_1}$ in $Cayley(S)$ then $\varphi_{\overline{s_n}} \cdots \varphi_{\overline{s_2}} \varphi_{\overline{s_1}} = \varphi_{\overline{r_m}} \cdots \varphi_{\overline{r_2}} \varphi_{\overline{r_1}}$ in $Cayley(T)$. This follows from the previous calculation
	- 2. Denote by C the subsemigroup of Cayley(S) generated by $\{\varphi_t | t \in$ T}. An element of C acts on the tree $(S^1)^*$.

Assume first that $T = T^1$. In this case $(T^1)^*$ is a subtree of $(S^1)^*$ and is therefore invariant under the action of C , the map sending an element of C to its restriction on $(T^1)^*$ is an onto morphism $C \rightarrow Cayley(T)$.

We now consider the case $T \neq T^1$ and write 1_{T^1} for the identity of T^1 . We can identify 1_{T^1} with 1_{S^1} , and think of T^1 as a subsemigroup of S^1 , so again $(T^1)^*$ is a subtree of $(S^1)^*$, so $Cauleu(T)$ is a quotient of C.

 \Box

The above proof actually shows that *Cayley* is a functor, i.e. if $\psi : S \to T$ is a semigroup morphism, then ψ induces a $Cayley(\psi)$: $Cayley(S) \rightarrow Cayley(T)$.

Corollary 10. *Let* S and T be semigroups. If $S \prec T$ then $Cayley(S) \prec T$ $Cayley(T)$.

Remark 11. Notice that if S contains a zero element 0, then φ_0 is the zero element of $Cayley(S)$.

Remark 12. If $s \neq t$ then $\varphi_s \neq \varphi_t$.

Proof. This is immediate by considering the action of both elements on the word [1]. \Box

4.2 Idempotent Semigroups

Every $f \in Cayley(S)$ is a tree endomorphism, and therefore f is determined by its action on each node of the tree $(S^1)^*$ (where the action is some function $S^1 \to S$). By the portrait of f we mean the tree $(S^1)^*$, and for every node, a function $S^1 \to S$. We mention an obvious but important fact, that two elements of $Cayley(S)$ are equal if and only if the are decorated with the same portrait. See [\[3\]](#page-23-1) for more on this notation, and examples of how useful this is for proofs.

Notation 13. Denote by l_s the function $l_s : S^1 \to S$ given by $l_s(t) =$ st.

Every element of $Cauley(S)$ is determined by its action on the top level of the tree and by its restrictions to the subtrees. Thus, every $f \in Cayley(S)$ can be represented by a pair $((f_{[s]})_{s \in S}, l_s)$, where we recall the definition of $f_{[s]}$ as given in the fractalness section, and the fact that $f_{[s]} \in Cayley(S)$. If $f \in Cayley(S)$ is associated with the pair $((f_{[s]})_{s\in S}, l_s),$ then $f([a_1, a_2, \ldots, a_n]) = [l_s(a_1)] \circ f_{[sa_1]}([a_2, \ldots, a_n]).$ It is easy to see that for generators, $\varphi_s \mapsto ((\varphi_{sx})_{x \in S}, l_s)$, and that $(\varphi_t\varphi_s) \mapsto ((\varphi_{tsx}\varphi_{sx})_{x\in S}, l_{ts}).$

We can then extend this to any product of generators, $\varphi_{s_n} \dots \varphi_{s_2} \varphi_{s_1} \mapsto$ $((\varphi_{s_n\cdots s_2 s_1 x} \cdots \varphi_{s_2 s_1 x} \varphi_{s_1 x})_{x \in S}, l_{s_n \cdots s_2 s_1}).$

Lemma 14. *Let* S *be an idempotent semigroup. Then* $Cauleu(S) \cong S$.

Proof. Let $s, t \in S$. Consider the word $w = [a_1, a_2, \ldots, a_n]$. We will show that $\varphi_t \varphi_s(w) = \varphi_{ts}(w)$. Let $b_i = a_1 a_2 \cdots a_i$, and observe that since S is an idempotent semigroup we have $sb_ib_j = sb_j$ when $i \leq j$ for every $s \in S$. We now calculate $\varphi_t \varphi_s(w) = \varphi_t([sa_1, sa_1a_2, \ldots, sa_1a_2 \cdots a_n]) =$ $\varphi_t([sb_1, sb_2, \ldots, sb_n]) = [tsb_1, tsb_1b_2, \ldots, tsb_1b_2 \cdots b_n] = [tsb_1, tsb_2, \ldots, tsb_n] =$ $\varphi_{ts}([a_1, a_2, \ldots, a_n]) = \varphi_{ts}(w).$

Since the action of the generators is faithful by Remark [12,](#page-7-0) the mapping $s \mapsto \varphi_s$ is 1-1 and unto $Cayley(S)$. This completes the proof. \Box

Corollary 15. *Let* S *be a finite simple aperiodic semigroup. Then* $Cayley(S) \cong S$.

Proof. As seen before, a simple aperiodic semigroup is of the form $A \times B$, with A a left zero semigroup, and B a right zero semigroup. (i.e. $(a_1, b_1)(a_2, b_2) = (a_1, b_2)$), and in particular is an idempotent semigroup. By the above it immediately follows that $Cauleu(S) \cong$ S. \Box

4.3 Nilpotent Semigroups

We recall here the definition of a nilpotent semigroup. We say that a semigroup with zero is nilpotent of index n if for every $m \geq n$ we have $s_m \cdots s_2 s_1 = 0$ for every $s_m, \ldots, s_2, s_1 \in S$, and n is the smallest with this property. The following observation shows that a finitely generated nilpotent semigroup is finite. If S is such a semigroup, X a generating set and *n* the nilpotency index, then $|S| \leq \frac{|\overline{X}|^n - 1}{|X| - 1} + 1$ (unless $|X| = 1$ in which case the semigroup has *n* elements).

Here we will see another example where S is similar to $Cayley(S)$.

Lemma 16. If S is a finite nilpotent semigroup of index n then $Cayley(S)$ *is nilpotent with nilpotency index at most* n*.*

Proof. We use the wreath product notation. Let S be a nilpotent semigroup of index *n*. Let $\Pi_{i=1}^n \varphi_{s_i} = ((\Pi_{i=1}^n \varphi_{b_i})_{x \in S}, l_0)$ be a product of *n* generators, for some b_i who depend on *x*. Since every b_i is a product of at least one element of S, the portrait is decorated by an l_0 in each node.

Since S is finite, $Cayley(S)$ is a finitely generated nilpotent semigroup, and therefore is finite \Box

Recall that a monogenic semigroup is a semigroup which is generated by one element.

Corollary 17. *If* S *is a monogenic aperiodic semigroup, then* Cayley(S) *is nilpotent.*

A monogenic finite aperiodic semigroup is of the form $\{x, x^2, \ldots, x^n\}$ satisfying the relation $x^n = x^{n+1}$ for some $n \in \mathbb{N}$. Obviously, a monogenic semigroup is commutative.

In general (i.e. $|S|$ not too small) if S is monogenic, then $Cayley(S)$ is not monogenic and not even commutative. For example let $S = \leq$ $x|x^5 = x^6$. In Cayley(S) we have the following computation. $\varphi_x \varphi_{x^2}([1,1]) = \varphi_x([x^2, x^2]) = [x^3, x^5] \text{ and } \varphi_{x^2} \varphi_x([1,1]) = \varphi_{x^2}([x,x]) =$ $[x^3, x^4]$.

4.4 Basic Structure

We end this section with some structure facts about $Cayley(S)$. Recall that the definition of a monoid acting on a set does not require the identity of the monoid to have a trivial action.

Theorem 18. *Let* S *be an aperiodic monoid, and* 1 *the identity of* S*. The following are equivalent.*

- *1.* Cayley(S) *is a monoid, and the identity of* Cayley(S) *acts trivially on* S ∗ *.*
- 2. φ_1 has a trivial action on S^* .
- *3.* $S = \{1\}.$

Proof. (3) implies (2) implies (1) is obvious. Assume that $Cayley(S)$ has an identity which acts trivially on S^* . Let $f = \varphi_{s_n} \cdots \varphi_{s_2} \varphi_{s_1}$ be the identity. We have $[1] = f([1]) = [s_n \cdots s_2 s_1]$. This shows that $s_i \mathcal{J}$ 1, and in a finite aperiodic semigroup, this implies $s_i = 1$. We now have $f = \varphi_1^n$. Consider the word $[x, 1]$. $[x, 1] = \varphi_1^n([x, 1]) = [x, x^n]$. Therefore, $x^n = 1$, and by aperiodicity $x = 1$. This shows that $S =$ {1}. \Box

Theorem 19. *Let* S *be an aperiodic monoid, and* 1 *the identity of* S*. The following are equivalent.*

- *1.* φ_1 *is an idempotent.*
- 2. φ_1 *is regular.*
- *3.* S *is an idempotent monoid.*

Proof. We saw that if S is idempotent, then $S \cong Cayley(S)$. This shows that (3) implies (1) and (1) implies (2). Now suppose φ_1 is regular. For some $f \in Cayley(S), \varphi_1 f \varphi_1 = \varphi_1$. Considering the word [1], we have $\varphi_1 f \varphi_1([1]) = \varphi_1([1]) = [1]$. If we write $f = \varphi_{s_n} \cdots \varphi_{s_2} \varphi_{s_1}$, then $\varphi_1 f \varphi_1([1]) = [s_n \cdots s_2 s_1],$ thus $s_n \cdots s_2 s_1 = 1$. In an aperiodic semigroup, this implies $s_n = \cdots = s_2 = s_1 = 1$, and so $f = \varphi_1^n$ for some *n*. This gives us $\varphi_1 = \varphi_1^{n+2}$. Applying both functions on [x, 1] gives $[x, x] = [x, x^{n+2}]$. This gives $x = x^{n+2}$ and by aperiodicity, $x = x^2$ for every $x \in S$. \Box

Lemma 20. $Cayley(S \times T)$ *is a subsemigroup of* $Cayley(S) \times Cayley(T)$ *.*

Proof. Φ : $Cayley(S \times T) \rightarrow Cayley(S) \times Cayley(T)$ defined by Φ : $\Pi_{i=1}^n \varphi_{(s_i,t_i)} \mapsto (\Pi_{i=1}^n \varphi_{s_i}, \Pi_{i=1}^n \varphi_{t_i})$ maps $Cayley(S \times T)$ onto the subsemigroup of $Cayley(S) \times Cayley(T)$ of pairs of functions which can be written as elements of the same length. To see that this is well defined let $f \in Cayley(S \times T)$ map the string w to $w' = f(w)$, where $w, w' \in ((S \times T)^{1})^{*}$. If we denote by w_{S} the string obtained from w by replacing every pair (s_i, t_i) with s_i , then we can see that $\Phi(f)$ maps w_S to w_S' , regardless of the presentation of f as a product of generators. The same argument works for T .

To see that this is a morphism, let $f = \prod_{i=1}^n \varphi_{(s_i,t_i)}$, and $g =$ $\Pi_{i=1}^m \varphi_{(s'_i,t'_i)}$. We have $\Phi(f) = (\Pi_{i=1}^n \varphi_{s_i}, \Pi_{i=1}^n \varphi_{t_i})$, and $\Phi(g) = (\Pi_{i=1}^m \varphi_{s'_i}, \Pi_{i=1}^m \varphi_{t'_i})$. Thus

$$
\Phi(f)\Phi(g) =
$$
\n
$$
(\Pi_{i=1}^{n} \varphi_{s_i}, \Pi_{i=1}^{n} \varphi_{t_i}) (\Pi_{i=1}^{m} \varphi_{s'_i}, \Pi_{i=1}^{m} \varphi_{t'_i}) =
$$
\n
$$
(\Pi_{i=1}^{n} \varphi_{s_i} \Pi_{i=1}^{m} \varphi_{s'_i}, \Pi_{i=1}^{n} \varphi_{t_i} \Pi_{i=1}^{m} \varphi_{t'_i}) =
$$
\n
$$
\Phi(fg).
$$

 \Box

5 The Action on Minimal Ideals

We assume henceforth that all semigroups have an identity and a zero. We do not lose any generality for the main results since $Cayley(S) \prec$ $Cayley(S^1)$ and $Cayley(S) \prec Cayley(S^0)$.

Let S be a finite aperiodic semigroup, and I a 0-minimal ideal. We will show that the restriction of $Cayley(S)$ to I is finite and aperiodic.

Notation 21. Cayley(S, I) will denote the restriction of Cayley(S) to *I*^{*}, i.e. *Cayley*(*S*,*I*) is the semigroup of functions $I^* \to I^*$, generated by the functions $\{\varphi_s\}_{s\in S}$.

More generally, $Cayley(S, T)$ will denote the restriction of $Cayley(S)$ to the tree T^* for some ideal T of S .

5.1 The Rhodes Expansion

Let S be a finite semigroup. The right Rhodes expansion \hat{S} was introduced by Rhodes in 1969 [5].

Consider the set $M(S)$ of all strings in S^* of the form $[s_1, s_1s_2, s_1s_2s_3, \ldots, s_1s_2\cdots s_n].$ Define a multiplication on $M(S)$ by

$$
[s_1,\ldots,s_1s_2\cdots s_n][t_1,\ldots,t_1t_2\cdots t_m]=[s_1,\ldots,s_1s_2\cdots s_n,s_1\cdots s_nt_1,\ldots,s_1\cdots s_nt_1\cdots t_m]
$$

Consider the case where $s_1 \cdots s_{i-1} s_i \mathcal{R} s_1 \cdots s_{i-1}$. Equivalently, for some $x \in S$, $s_1 \cdots s_{i-1} s_i x = s_1 \cdots s_{i-1}$. A *one step reduction* replaces $w = [s_1, \ldots, s_1 \cdots s_{i-1}, s_1 \cdots s_{i-1} s_i, \ldots, s_1 \cdots s_n]$ with

 $w' = [s_1, \ldots, s_1 \cdots s_{i-2}, s_1 \cdots s_{i-1} s_i, \ldots, s_1 \cdots s_n].$ If no such i exists, w is called reduced. It is easy to see that every element w of $M(S)$ has a unique reduced form $red(w)$ obtained from w by applying finitely many one step reductions.

Definition 22. *The right Rhodes expansion of a finite semigroup* S *is the collection of reduced words in* $M(S)$ *with multiplication* $w \cdot w' =$ $red(ww')$ *. We denote it by* \hat{S}^R *.*

The Rhodes expansion is closely related to the following construction.

5.2 The Memory Semigroup of S

Let S be a finite semigroup. Define a multiplication on the set $S \times P(S)$ by

$$
(s,\alpha) \cdot (t,\beta) = (st,\alpha t \cup \{t\} \cup \beta)
$$

where, if $\alpha \in P(S)$ and $t \in S$, then $\alpha t = \{xt | x \in \alpha\}$. We will first verify that this multiplication is associative.

 $[(s,\alpha)(t,\beta)](x,\gamma) = (st,\alpha t\cup\{t\}\cup\beta)(x,\gamma) = (stx,\alpha tx\cup\{tx\}\cup\beta x\cup\beta x)$ ${x} \cup \gamma$ $(s, \alpha)[(t, \beta)(x, \gamma)] = (s, \alpha)(tx, \beta x \cup \{x\} \cup \gamma) = (stx, \alpha tx \cup \{tx\} \cup \beta x \cup \gamma)$ $\{x\} \cup \gamma$).

Definition 23. *We call the above the (right) memory semigroup of* S*, and denote it by* mem(S)*.*

Note that $f : mem(S) \to S, f(s, \alpha) = s$ is a surjective morphism. Thus $S \prec mem(S)$.

The finiteness of $mem(S)$ is trivial. Furthermore, we observe the following.

Lemma 24. *Let* S *be a finite semigroup. Then* mem(S) *is aperiodic if and only* S *is.*

Proof. A simple induction shows that $(s, \alpha)^n = (s^n, \alpha \cup \alpha \{s, s^2, \dots, s^{n-1}\} \cup$ $\{s, s^2, \ldots, s^{n-1}\}\)$. Thus if $s^n = s^{n+1}$ for every s in S then $(s, \alpha)^{n+1} =$ $(s, \alpha)^{n+2}$ for every (s, α) in $mem(S)$. The converse follows since $S \prec$ $mem(S).$ \Box

We now proceed with the action of $Cayley(S)$ on I^* .

5.3 Some Lemmas

Lemma 25. *Let* S *be an aperiodic semigroup. Let* $I \cong (A \times B) \cup \{0\}$ *be a* 0*-minimal ideal,* $f \in Cayley(S, I)$ *, and* $w \in I^* \setminus \{\epsilon\}.$

1. $f(w)$ *is of the form* $[(a, b_1), (a, b_2), \ldots, (a, b_k), 0, \ldots, 0]$ *, with* $a \in$ $A, b_i \in B$ *and* $k \geq 0$ *.*

- 2. The first k elements of w are $[(a_1, b_1), (a_2, b_2), \ldots, (a_k, b_k)]$, for *some* $a_i \in A$ *.*
- 3. Assume $k > 0$. If we write $f = \varphi_{s_n} \cdots \varphi_{s_2} \varphi_{s_1}$ then a is given by $s_n \cdots s_2 s_1(a_1, b_1) = (a, b_1)$ *, that is,* $s_n \cdots s_2 s_1 a_1 = a$ *, assuming* f(w) *is not a string of zeros.*
- *Proof.* 1. It is enough to show this for the case when f is a generator of $Cayley(S, I)$. It is also clear that once we have a 0 in $f(w)$ all elements beyond it will be 0. Let $w = [(a_1, b_1), (a_2.b_2), \ldots, (a_k, b_k)]$, and φ_s a generator.
	- $\varphi_s(w) =$ $[s(a_1, b_1), s(a_1, b_1)(a_2, b_2), \ldots, s(a_1, b_1)\cdots (a_k, b_k)] =$ $[s(a_1, b_1), s(a_1, b_2), \ldots, s(a_1, b_k)]$
	- 2. We cannot have a 0 in the first k entries of w because then we would also have a 0 in the first k entries of $f(w)$. We see that applying φ_s to w doesn't change b_i for $i \leq k$.
	- 3. The last statement is immediate.

 \Box

The following lemma will be needed to cover the last cases for the main theorem of this section.

Lemma 26. Let $\varphi_s \in Cayley(S, I)$ be a generator and $w = [\ldots, (a_i, b_i), (a_{i+1}, b_{i+1}), \ldots] \in$ *I*^{*}*. Furthermore assume that the first* $i - 1$ *entries of* w are $\neq 0$ *.*

- *1.* Assume $\varphi_s(w)$ *is non zero in the i position. Then* $\varphi_s(w)$ *has a zero in the* $i + 1$ *position if and only* $C(b_i, a_{i+1}) = 0$ *where* C *is the structure matrix of* $I \setminus \{0\}$ *.*
- 2. Let $f = \varphi_{s_n} \cdots \varphi_{s_2} \varphi_{s_1}$ and let $w = [(a, b), \ldots].$ Then $f(w) =$ $[0, 0, \ldots, 0]$ *if and only if* $s_n \cdots s_2 s_1(a, b) = 0$ *.*

Proof. A direct calculation can verify both facts. We notice that for the second statement there are two possibilities for f when $s_n \cdots s_2 s_1(a, b) =$ 0. Either $s_n \cdots s_2 s_1 = 0$ in which case f is the zero function. If not, $(s_n \cdots s_2 s_1)(a, b) = 0$ in which case f is not the zero function, but f acts as the zero function on any string of the form $[(a, b'), \dots]$ for any $b' \in B$. \Box

Proposition 27. *Let* S, I *be as before. Then* $Cayley(S, I) \prec mem(S)$ *.*

The proof is motivated by the following idea: Let $v \in I^*$, and let $f = \varphi_{s_n} \cdots \varphi_{s_1} \in Cayley(S, I)$. The output $f(v)$, is of the form $[(a, b_1), (a, b_2), \ldots, (a, b_k), 0, 0, \ldots, 0].$ Applying f doesn't change the b_i 's of v. The action of f on v is determined by the following two questions: 1. what is a? 2. what is k? (i.e. from what point does $f(v)$) consist of zeros?).

If $v = [(a_1, b_1), \ldots]$ then we answer the first question by (a, b_1) $s_n \cdots s_2 s_1(a_1, b_1)$, i.e. the first question is answered by the first component of an element of $mem(S)$.

For the second we will observe that the appearance of a 0 will depend on the set-component of mem(S). Each time we apply a φ_{s_i} to v we get a string of the form $[s_i \cdots s_2 s_1(a_1, b_1), \ldots],$ so the setcomponent determines the set of a's we will see along the way. Part 2 of Lemma [26](#page-12-0) tells us that if $v' = \varphi_{s_i} \cdots \varphi_{s_1}(v) = [(a, b_1), \ldots],$ then a 0 will be created in v' when we find a b_i such that $C(b_i, a) = 0$.

Proof. Denote by S^+ the free semigroup over S, with S considered as a set. Denote by $\Gamma: S^+ \to Cayley(S, I)$ the canonical morphism.

$$
\Gamma: ([s_n, \ldots, s_2, s_1]) \mapsto \varphi_{s_n} \cdots \varphi_{s_2} \varphi_{s_1}
$$

Let $\Phi: S^+ \to mem(S)$ be defined by

$$
\Phi: ([s_1]) \mapsto (s_1, \emptyset)
$$

By induction on n it is easy to see that

$$
\Phi : ([s_n, \cdots, s_2, s_1]) \mapsto (s_n \cdots s_2 s_1, \{s_{n-1} \cdots s_2 s_1, \ldots, s_2 s_1, s_1\})
$$

We will denote by \sim_{Γ} and \sim_{Φ} the congruences on S^{+} corresponding to Γ and Φ .

We show that $v_1 \sim_{\Phi} v_2$ implies $v_1 \sim_{\Gamma} v_2$. Let $v_1, v_2 \in S^+$ and let $f = \Gamma(v_1), g = \Gamma(v_2)$. Let $v_1 = [s_n, \ldots, s_2, s_1]$, and $v_2 = [t_m, \ldots, t_2, t_1]$. We want to show that $\Phi(v_1) = \Phi(v_2)$ implies $\Gamma(v_1) = \Gamma(v_2)$. This is equivalent to $f = g$ in $Cayley(S, I)$ if $(s_n \cdots s_1, \{s_{n-1} \cdots s_1, \ldots, s_1\}) =$ $(t_m \cdots t_1, \{t_{m-1} \cdots t_1, \ldots, t_1\})$. If f is the zero function then $s_n \cdots s_1(a, b)$ 0 for every $(a, b) \in A \times B$. Thus, $\Phi(v_1) = (s_n \cdots s_1, \alpha)$ for some $\alpha \in P(S)$, and since $\Phi(v_1) = \Phi(v_2)$, $\Phi(v_2) = (s_n \cdots s_1, \alpha)$. In other words, $s_n \cdots s_2 s_1 = t_m \cdots t_2 t_1$ and we see that g is also the zero function on I^* .

Throughout the rest of this proof, we will assume $f \neq \varphi_0 \neq g$. Write $\Phi(v_1) = \Phi(v_2) = (\sigma, \alpha), \sigma \in S, \alpha \in P(S)$, with $\sigma(a, b) \neq 0$ for some (a, b) . Finally, let $w \in I^*$.

Let k_1 denote the position of the last nonzero entry in $f(w)$ (and $k_1 = 0$ if $f(w) = [0, 0, \ldots, 0]$ and k_2 the position of the last nonzero entry in $g(w)$. By Lemma [25,](#page-11-0) and since $\Phi(v_1) = \Phi(v_2)$ (in particular $s_n \cdots s_2 s_1 = t_m \cdots t_2 t_1$ it follows that $f(w)$ equals $g(w)$ in the first $min(k_1, k_2)$ entries.

It remains to show that $k_1 = k_2$.

First, assume $k_1 = 0$. Let $w = [(a_1, b_1), \ldots]$. This means that $f(w) = [0, 0, \ldots, 0]$. Since $f \neq \varphi_0$ this means that $C(s_n \cdots s_2 s_1, a_1) =$ 0 i.e. $C(\sigma, a_1) = 0$ and so $g(w) = [0, 0, \ldots, 0]$ and $k_2 = 0$, so $k_1 = k_2$.

We may assume now that $k_1, k_2 > 0$. Without loss of generality $k_1 \leq k_2$.

If $k_1 = |w|$, then all entries of $f(w)$ are nonzero and since $k_1 \leq k_2$ we are done in this case. So we may assume that $k_1 < |w|$.

If w has a zero in the $k_1 + 1$ place then it is immediate that $g(w)$ has a zero in the k_1+1 place, so in this case $k_2 \leq k_1$. We may therefore assume that the first $k_1 + 1$ entries in w are not zero, and $w = [(a_1, b_1), (a_2, b_2), \ldots, (a_{k_1}, b_{k_1}), (a_{k_1+1}, b_{k_1+1}), \ldots].$

We know that $f(w)$ has a zero in the k_1+1 place, and $f = \varphi_{s_n} \cdots \varphi_{s_2} \varphi_{s_1}$. Let r be such that $\varphi_{s_r} \cdots \varphi_{s_2} \varphi_{s_1}(w)$ has a zero in the $k_1 + 1$ place but $\varphi_{s_{r-1}}\cdots\varphi_{s_2}\varphi_{s_1}(w)$ does not.

If $r = 1$ (i.e. if $\varphi_{s_1}(w)$ has a zero in the $k_1 + 1$ position) then part 1 of Lemma [26](#page-12-0) shows that $\varphi_{t_1}(w)$ has a zero in the $k_1 + 1$ position, and so $g(w)$ has a zero in the $k_1 + 1$ position. Thus $k_2 \leq k_1$ and the claim is proved.

Assume $r > 1$. Let $w' = \varphi_{s_{r-1}} \cdots \varphi_{s_1}(w) = [(c, b_1), \ldots, (c, b_{k_1}), (c, b_{k_1+1}), \ldots],$ where c is determined by $s_{r-1} \cdots s_2 s_1(a_1, b_1) = (s_{r-1} \cdots s_2 s_1 a_1, b_1) =$ (c, b_1) . Applying φ_{s_r} to w' creates a zero in the $k_1 + 1$ place and leaves the k_1 place nonzero. We calculate $\varphi_{s_r}(w') = [(s_r c, b_1), \ldots, (s_r c, b_{k_1}), (s_r c, b_{k_1}) (c, b_{k_1+1}), \ldots].$ Since $(s_r c, b_{k_1}) \neq 0$ and $(s_r c, b_{k_1})(c, b_{k_1+1}) = 0$ we see that the structure matrix of *I* has $C(b_{k_1}, c) = 0$. Thus, $C(b_{k_1}, s_{r-1} \cdots s_2 s_1) = 0$.

We observe that for $\Phi(v_2) = (\sigma, \alpha)$, α describes the set of elements of S that w is multiplied by, as we apply φ_{s_i} , $1 \leq i \leq n$. Since $\Phi(v_1)$ = $\Phi(v_2)$, and since $s_{r-1} \cdots s_2 s_1 \in \alpha$, we see that $s_{r-1} \cdots s_1$ belongs to the set-component of $\Phi(v_2)$. This implies that g can also be decomposed as $g = g_2 \varphi_t g_1$ for some $t \in S$ with $g_1 \in Cayley(S, I), g_2 \in Cayley(S, I) \cup$ $\{id\}$, and $g_1([a_1, b_1)]) = [s_{r-1} \cdots s_2 s_1(a_1, b_1)].$

We write $g_1(w) = [(d, b_1), \ldots, (d, b_{k_1}), (d, b_{k_1+1}), \ldots]$ (We assume that the k_1 and $k_1 + 1$ places are not zero or else we will have $k_2 \leq k_1$ and the proof is over). Since $[(d, b_1)] = g_1[(a_1, b_1)] = [(s_{r-1} \cdots s_2 s_1 a_1, b_1)] =$ $f_r[(a_1, b_1)] = [(c, b_1)]$ we see that $c = d$. We rewrite, $g_1(w) = [(c, b_1), \ldots, (c, b_{k_1}), (c, b_{k_1+1}), \ldots]$.

 \Box

Calculating $\varphi_t g_1(w) = [t(c, b_1), \dots, t(c, b_1) \dots (c, b_{k_1})(c, b_{k_1+1})],$ we see that the $k_1 + 1$ position has a right factor $(c, b_{k_1}) \cdot (c, b_{k_1+1})$. We noticed before that the structure matrix has $C(b_{k_1}, c) = 0$, so $\varphi_t g_1(w)$ has a zero in the $k_1 + 1$ position, and so does $g(w)$. We have showed that $k_2 \leq k_1$.

We now have $k_1 = k_2$. Thus we have showed that $f(w) = g(w)$ and thus $f = g$. Thus, $\Phi(v_1) = \Phi(v_2)$ implies $\Gamma(v_1) = f = g = \Gamma(v_2)$.

Since we have proved that $\sim_{\Phi} C \sim_{\Gamma}$, there is an induced surjective morphism from S^+/\sim_{Φ} onto $Cayley(S,I)\S^+/\sim_{\Gamma}$. Therefore, Cayley(S, I) divides $S^+/\sim \Phi$ which is a subsemigroup of $mem(S)$.

Theorem 28. *Let* S *be a finite aperiodic semigroup, and* I *a* 0*-minimal ideal. Then* Cayley(S, I) *is finite and aperiodic.*

This was proved for the regular case. For a nonregular 0-minimal ideal, the theorem is trivial. This is because such a $\mathcal J$ class is null, so for any $\varphi_s \in Cayley(S, I)$ and $w = [w_1, w_2, \ldots] \in I^*$ we have $\varphi_s(w) = [sw_1, sw_1w_2 = 0, 0, \ldots, 0],$ so the action of $\varphi_{s_n} \cdots \varphi_{s_1}$ on w is determined by $s_n \cdots s_1$, with $\varphi_{s_n} \cdots \varphi_{s_1}(w) = [s_n \cdots s_2 s_1 w_1, 0, \ldots, 0]$ and $Cayley(S, I)$ is the quotient of S obtained by identifying two elements if they act the same on the left of I.

For future purposes we extend the previous result.

Definition 29. *Let* S *be a finite aperiodic semigroup, and let* I *be a* 0-minimal ideal of *S*. Let $f \in Cayley(S, I)$, and $w = [v_1, \ldots, v_l] \in I^*$. *We say that a zero* is created *in the* i*th position of* w *by* f*, if there exists a decomposition of* f , $f = \varphi_{s_n} \cdots \varphi_{s_2} \varphi_{s_1}$ *such that*

- 1. $\varphi_{s_{k-1}} \cdots \varphi_{s_2} \varphi_{s_1}(w)$ *does not have a zero in the i position.*
- 2. $\varphi_{s_k} \cdots \varphi_{s_2} \varphi_{s_1}(w)$ *has a zero in the i position but is nonzero in the* $i - 1$ *position.*

If we need to be more specific we will say that the zero is created by the k*th generator of the above decomposition of* f*.*

Definition 30. *Let* S *be a finite aperiodic semigroup, and* I *a* 0 *minimal ideal of* S, such that $I \cong (A \times B) \cup \{0\}$. Let $f = \varphi_{s_n} \cdots \varphi_{s_2} \varphi_{s_1} \in$ $Cayley(S, I)$ and $a \in A$, such that $s_n \cdots s_2 s_1(a, b) \neq 0$ for any (and *therefore all*) $b \in B$.

- *1.* For $1 \le i \le n+1$ let $\pi_a(i) = s_{i-1} \cdots s_2 s_1 a$ (and $\pi_a(1) = a$). We *recall the notation* sa *from Notation [7.](#page-5-0)*
- 2. $new_a(s_n, \dots, s_1) = \{1 \leq i \leq n | \pi_a(i) \neq \pi_a(j) \forall j < i\}.$

Notice that $|new_a(s_n, \ldots, s_1)| \leq |A|$, for any $s_n, \ldots, s_1 \in S$.

Lemma 31. Let $f = \varphi_{s_n} \cdots \varphi_{s_2} \varphi_{s_1}$. Let $w \in Cayley(S)(I^*)$ (i.e. w *is the output of* $Cayley(S, I)$ *acting on some string in* I^* *) such that* w *is not a string of* 0*s.* Let $w = [(a_1, b_1), \ldots]$ *and let* $a = a_1$ *. If the kth generator of* f *creates a new zero, then* $k \in new_a(s_n, \ldots, s_2, s_1)$ *.*

Proof. Suppose the kth generator creates a zero in the *i*th entry. Without loss of generality $f = \varphi_{s_k} \cdots \varphi_{s_2} \varphi_{s_1}$. Let $f' = \varphi_{s_{k-1}} \cdots \varphi_{s_2} \varphi_{s_1}$. We then have $f'(w) = [(a', b_1), \ldots, (a', b_{i-1}), (a', b_i), \ldots],$ where $a' =$ $s_{k-1} \cdots s_2 s_1 a = \pi_a(k)$. This is true even if $k = 1$ and f' is the identity function, since we assume $w \in Cayley(S)(I^*)$. Since the zero is created in the ith entry we have

 $f(w) = \varphi_{s_k}(f'(w)) = [(s_ka', b_1), \ldots, (s_ka', b_{i-1}), (s_ka', b_{i-1})(a', b_i)]$ 0,...]. This shows that $C(b_{i-1}, a') = 0$. If $k = 1$ then it is immediate that $k \in new_a(s_n, \ldots, s_1)$. Suppose that $1 \lt k \notin new_a(s_n, \ldots, s_1)$. Then for some $l < k$ we have $s_{l-1} \cdots s_2 s_1 a = a'$ and by a similar calculation the zero will be created by the lth generator and not the kth generator. \Box

Corollary 32. *Let* S *be an aperiodic semigroup, and let* I *be a* 0 *minimal ideal of S. There exists an* $N \leq |A|^2 + 1 \in \mathbb{N}$ *such that every* $f = \varphi_{s_n} \cdots \varphi_{s_2} \varphi_{s_1} \in Cayley(S, I)$ can be written as $f = g_{k+1} \varphi_{s_{i_k}} g_k \cdots g_2 \varphi_{s_{i_1}} g_1$ *for some* $k \leq N$ *, with* $g_l \in Cayley(S, I) \cup \{id\}$ *and such that all zeros are created by the* $\varphi_{s_{i_j}}$, $1 \leq j \leq k$ *for every* $w \in I^*$,

Proof. In order to use the previous lemma w needs to be in $Cayley(S)(I^*)$. Instead of $f = \varphi_{s_n} \cdots \varphi_{s_2} \varphi_{s_1}$ acting on w we may consider $f =$ $\varphi_{s_n} \cdots \varphi_{s_2}$ acting on $\varphi_{s_1}(w)$.

For every $a \in A$ consider the set $new_a(s_n, \ldots, s_1)$, and let $X = \bigcup_{a \in A} new_a(s_n, \ldots, s_1)$. By the previous lemma, all the zeros must be created by the rth generator for some $r \in X$. In fact, every $a \in A$ contributes at most |A| elements to X. Thus, for any word w , the zeroes will be created by an rth generator for some $r \in X$. Thus, in the product $f = \varphi_{s_n} \cdots \varphi_{s_2} \varphi_{s_1}$ we can single out $|X|$ generators which are the only ones that can create a zero. We also have $|X| \leq |A|^2 + 1$. The +1 in this formula comes from the beginning of the proof. \Box

6 Finiteness

.

Before the next lemma, recall the definition of $p : S^+ \to S$ given by $p([a_1, a_2, \ldots, a_l]) = a_l$ and in particular, $p([a]) = a$.

Lemma 33. *Let* S *be a finite aperiodic semigroup, and* A×B *a regular* $\mathcal J$ *class of* S *. Let* $[v_1, v_2, \ldots, v_k] = v \in (A \times B)^+$ *. Let* $f \in Cayley(S)$ *. Write* $f = \varphi_{s_n} \cdots \varphi_{s_1}$ and $v_i = (a_i, b_i)$ *. Furthermore assume that* $f(v) \in (A \times B)^+$ *. Then*

$$
f_v = \varphi_{(s_n\cdots s_2s_1a_1,b_k)}\cdots\varphi_{(s_2s_1a_1,b_k)}\varphi_{(s_1a_1,b_k)}
$$

Proof. Since $f(v) \in (A \times B)^*$ we have $v_1v_2 \cdots v_k \in A \times B$, so $v_1v_2 \cdots v_i =$ (a_1, b_i) for $1 \leq i \leq k$. $\varphi_{s_1}(v) = [s_1v_1, s_1v_1v_2, \ldots, s_1v_1\cdots v_k] = [(s_1a_1, b_1), \ldots, (s_1a_1, b_k)].$ Proceeding from here (by inducting on i) we have

$$
\varphi_{s_i} \cdots \varphi_{s_1}(v) = [(s_i \cdots s_2 s_1 a_1, b_1), \ldots, (s_i \cdots s_2 s_1 a_1, b_k)]
$$

We now recall the formula for f_v from Theorem [8.](#page-6-1) We have $f_v =$ $\varphi_{(p\varphi_{s_n}\cdots\varphi_{s_1}(v))}\ldots\varphi_{(p\varphi_{s_2}\varphi_{s_1}(v))}\varphi_{(p\varphi_{s_1}(v))}.$ Since $p\varphi_{s_i}\ldots\rho_{s_1}(v)=(s_i\cdots s_2s_1a_1,b_k),$ we have our result.

Remark 34. With the previous definitions, assume that $f(v) \in (A \times$ B ⁺. Then $\varphi_1(v) \in (A \times B)^+$.

Proof. We write $f = \varphi_{s_n} \cdots \varphi_{s_2} \varphi_{s_1}$. We observe that every entry in the word $\varphi_1([v_1, v_2, \ldots, v_k]) = [v_1, v_1v_2, \ldots, v_1v_2 \cdots v_k],$ is $\mathcal L$ above the corresponding entry of $f(v)$. This follows from the fact that $f(v)$ can be obtained from $\varphi_1(v)$ by multiplying each element on the left by s_1 , and then by applying $\varphi_{s_n} \cdots \varphi_{s_2}$ to the resulting word, which again is just left multiplication. Since $v \in (A \times B)^+$ and $f(v) \in (A \times B)^+$ we have that $\varphi_1(v) \in (A \times B)^+$. \Box

Corollary 35. Let S be a finite aperiodic semigroup, $A \times B$ a J class *of* S. Let $[v_1, v_2, \ldots, v_k] = v \in (A \times B)^+$ *and let* $f \in Cayley(S)$ *. Write* $f = \varphi_{s_n} \cdots \varphi_{s_1}$ *and* $v_i = (a_i, b_i)$ *. Assume that* $f(v) \in (A \times B)^+$ *. Furthermore, let* $j = p\varphi_1(v) = (a_1, b_k)$ *. Then* $f_v = f_{[j]}$ *.*

Proof. We first show that $f([j]) \in (A \times B)^+$. We calculate $f([j]) =$ $[s_n \cdots s_2 s_1(a_1, b_k)]$. Since $f(v) \in (A \times B)^+$, and since the first letter of $f(v)$ is $s_n \cdots s_2 s_1(a_1, b_1)$ it follows that $s_n \cdots s_2 s_1(a_1, b_k) \in A \times B$, so $f([j]) \in (A \times B)^+$.

Now, $f_v = \varphi_{(s_n \cdots s_2 s_1(a_1,b_k))} \cdots \varphi_{(s_2 s_1(a_1,b_k))} \varphi_{(s_1(a_1,b_k))}.$ We also have $f_{[j]} = \varphi_{(s_n \cdots s_2 s_1(a_1,b_k))} \cdots \varphi_{(s_2 s_1(a_1,b_k))} \varphi_{(s_1(a_1,b_k))}.$

 \Box

Definition 36. *Let* S *be a finite semigroup. Let* T *be an ideal of* S *and* J *a* J *class of* S *not in* T *. We say that* J *is* directly above T *if* $J \cup T$ *is an ideal of S. Equivalently, J is a* 0*-minimal* J *class of* S/T *.*

Let S be a semigroup with an ideal T and a $\mathcal J$ class J directly above T. Let $w \in (T \cup J)^*$. Since every letter in w is in J or in T, w can be uniquely decomposed as $w = w_J$ or $w = w_J w'_T w'$ such that $w_J \in J^*, w'_T \in T$ and $w' \in (J \cup T)^*$. In the case that w is of the form $w = f(w_0)$ for some $f \in Cayley(S, T \cup J)$ and $w_0 \in (T \cup J)^*$ we can decompose $w = w_J w_T$ with $w_J \in J^*$ and $w_T \in T^*$. In both cases, we call w_J the prefix of w in J.

Definition 37. *Let* S *be a finite aperiodic semigroup. Let* T *be an ideal of* S, and let J be a J class directly above T. For an $f \in Cayley(S, T \cup$ J), the set of f-J-stable words is the set of words $st(f, J) \subset (T \cup J)^*$ *such that the prefix of* w *in* J has the same length as the prefix of $f(w)$ *in* J*.*

We observe that in general, the length of the prefix of w in J is larger or equal to the length of the prefix of $f(w)$ in J.

Consider the following relation on Cayley(S, T∪J). We write $f \sim q$ if and only if $st(f, J) = st(g, J)$ and the restrictions of f and g to this set is equal.

Lemma 38. *Let* $S, T, J, ∼$ *be as before.* $∼$ *is a congruence on* $Cayley(S, T \cup J)$ *.*

Proof. It is clear that \sim is an equivalence relation. Suppose that $f \sim f'$ and $g \sim g'$. Let $w \in st(fg, J)$. In particular this means that $w \in$ $st(g, J)$ and that $g(w) \in st(f, J)$. This implies $g(w) = g'(w)$ and that $f(g(w)) = f'(g(w)) = f'(g'(w))$. Thus $w \in st(f'g', J)$ and $fg(w) =$ $f'(g(w)) = f'g'(w)$. The dual argument completes the proof. \Box

Notation 39. We write $Cayley_{st-J}(S, T \cup J)$ for the quotient of $Cayley(S, T \cup J)$ by \sim .

We point out, that for $[f]_{\sim} \in Cayley_{st-J}(S, T \cup J)$ and $w \in st(f, J)$, $[f]_{\sim}(w)$ is always well defined, i.e. every $g \sim f$ can act on w and all these g's give the same g(w). We will refer to $[f]_{\sim}(w)$ as $\tilde{f}(w)$, whenever $[f]_{\sim}(w)$ is defined. In general we will write f for $[f]_{\sim}$.

Until now, the notation $Cayley(S,T)$ was used when T was an ideal of S. It can actually be used where T is any semigroup which S acts on from the left, by defining $\varphi_s([t_1, t_2, \ldots, t_k]) = [st_1, st_1t_2, \ldots, st_1t_2 \cdots t_k].$ However, in general, $Cayley(S, T)$ is no longer a divisor of $Cayley(S)$.

Let S be an aperiodic semigroup, T an ideal of S and J a $\mathcal J$ class directly above T. Let $\alpha: (T \cup J) \to J^{tr} \cong (T \cup J)/T$ be the ideal quotient. φ_s can act on strings in J^{tr} in the following way: Let $w = [j_1, \ldots, j_n] \in$ $(J^{tr})^*$. We have $\varphi_s([j_1, j_2, \ldots, j_n]) = [\overline{sj_1}, \overline{sj_1j_2}, \ldots, \overline{sj_1j_2 \ldots j_n}],$ with $\overline{s} = \alpha(s)$, i.e. $\overline{s} = s$ if $s \in J$ and $\overline{s} = 0$ otherwise.

Notation 40. We write $Cayley(S, J^{tr})$ for the semigroup of transformations $(J^{tr})^* \to (J^{tr})^*$ generated by $\{\varphi_s\}_{s \in S}$.

We define a morphism $Cayley(S, T \cup J) \rightarrow Cayley(S, J^{tr})$. Write f^{tr} for the image of f. f^{tr} 's action on $w \in (J^{tr})^*$ can be described as follows. We first replace any 0 in w with any element of T . Next, apply f to this new word. Finally, replace all elements of T with 0 (the zero of J^{tr}).

Let $f \in Cayley(S, J \cup T)$ and $w \in (J \cup T)^*$. Let f^{tr} be as defined above and let w^{tr} denote the word obtained from w be replacing elements of T with 0. We easily see that $w \in st(f, J)$ if and only if $w^{tr} \in st(f^{tr}, J)$ (*J* is a *J* class of J^{tr} , so $st(f^{tr}, J)$ makes sense).

We will use the above notations throughout the next proofs. We summarize.

Definition 41. *Let* S *be an aperiodic finite semigroup. Let* T *be an ideal and let* J *be a* J *class directly above* T *. We have defined:*

- 1. For $w \in (T \cup J)^*$, the word $w^{tr} \in (J^{tr})^*$ is obtained from w by *replacing elements of* T *with* 0*.*
- 2. Let $f \in Cayley(S, T \cup J)$. The function $f^{tr} \in Cayley(S, J^{tr})$ is given by $f^{tr}(w) = (f(w'))^{tr}$, where w' is any word in $(J \cup T)^*$ satisfying $(w')^{tr} = w$. It is obvious that this is well defined.
- *3.* Let f ∈ Cayley(S,T∪J). We define st(f^{tr} , J) to be the set of f^{tr} -*J*-stable words in $(J^{tr})^*$. For $w \in (J \cup T)^*$, we have $w \in st(f, J)$ *if and only if* $w^{tr} \in st(f^{tr}, J)$ *.*

We see that the mapping $f \mapsto f^{tr}$ is a morphism $Cayley(S, T \cup J) \rightarrow$ Cayley(S, J^{tr}), i.e. $f^{tr}g^{tr} = (fg)^{tr}$. We will now show that it induces a morphism $Cayley_{st-J}(S, T \cup J) \rightarrow Cayley_{st-J}(S, J^{tr})$.

Definition 42. Let $\tilde{f} \in Cayley_{st-J}(S, T \cup J)$ and let $w \in (J^{tr})^*$. *Write* $f^{tr}(w)$ *for the word* $(f(w))^{tr}$ *where f is an element of* Cayley(S, T∪ J) such that f maps to \tilde{f} , and the action of f on J^{tr} is understood *as in Notation [40.](#page-17-0) It is clear that this is well defined. Thus we have defined a mapping* $Cayley_{st-J}(S, T \cup J) \rightarrow Cayley_{st-J}(S, J^{tr}).$

Lemma 43. *Let* S *be an aperiodic semigroup,* T *an ideal of* S *and* J *a J class directly above T.* Let $\tilde{f}, \tilde{g} \in Cayley_{st-J}(S, T \cup J)$ *, and let* $f, g \in Cayley(S, T \cup J)$ *be preimages of* \tilde{f}, \tilde{g} *. Then* $(fg)^{tr} = f^{tr}g^{tr}$ *.*

Proof. This follows from the fact that the mapping $f \mapsto f^{tr}$ is a morphism, and that $\tilde{f} \mapsto f^{tr}$ is well defined. Furthermore, we see that this does not depend on the choice of f and g . \Box

We wish to extend all result we had previously obtained for $Cayley(S, I)$ where I was a 0-minimal ideal of S. When we consider $Cayley(S, J^{tr})$, J^{tr} is a 0-simple semigroup, however it's not an ideal of S.

Theorem 44. *Let* S *be an aperiodic semigroup, let* J *be a* J *class of* S, and let $T = S^1 J S^1 \setminus J$. Denote by $∼T$ the ideal congruence *associated with* T. Then $(J \cup T)/ \sim_T i s$ *a* 0*-minimal ideal of* S/\sim_T *and* $Cayley(S, J^{tr}) \cong Cayley(S/ \sim_T, (J \cup T)/ \sim_T)$ *. Furthermore, since* \sim_T *is* 1 − 1 *on J we have* $J/\sim_T = J$ *.*

Proof. It is obvious that $(J \cup T)/\sim_T$ is a 0-minimal ideal of S/\sim_T . To show the isomorphism, consider $\varphi_s \in Cayley(S, J^{tr})$. If $s \in T$, then the action of φ_s on $(J^{tr})^*$ is seen as sending any string to a string of zeros, so in this case φ_s corresponds to φ_0 where 0 is the 0 element of S/\sim_T and it sends any string in $((J\cup T)/\sim_T)^*$ to a string of zeros.

If $s \notin T$ then the action of φ_s on $(J^{tr})^*$ is the same as in $Cayley(S/\sim_T, (J\cup T)/\sim_T)$.

We can extend this from $\{\varphi_s\}$, the generators of $Cayley(S, J^{tr})$ to the whole semigroup, and obtain an isomorphism $Cayley(S, J^{tr}) \cong$
Cauleu(S/ ~π. (J++*T*)/ ~*π*). $Cayley(S/\sim_T, (J\cup T)/\sim_T$).

Let $Cayley(S,T)^{J^{tr}}$ denote the semigroup of all functions $J^{tr} \rightarrow$ $Cayley(S,T)$ with pointwise multiplication. We now define a right action of $Cayley_{st-J}(S, J^{tr})$ on $Cayley(S, T)^{J^{tr}}$. Let $\tilde{f} \in Cayley_{st-J}(S, J^{tr})$ and let $H: J^{tr} \to Cayley(S,T)$. Let $j \in J$. If $[j] \in st(\tilde{f}, J)$ then we let $H^{\tilde{f}}(j) = H(\tilde{f}(j))$. Otherwise we let $H^{\tilde{f}}(j) = \varphi_0$ (Recall that φ_0 , the function sending any string to a string of zeros is the zero element of $Cayley(S,T)$).

Notice the slight abuse of notation for $\hat{f}(j)$ where we identify elements of the semigroup S with strings of length 1. Formally, instead of $\tilde{f}(j)$ we should have written $p(\tilde{f}([j]))$, where p is the projection $p([v_1]) = v_1.$

To show that this is indeed an action, let $\tilde{f}, \tilde{g} \in Cayley_{st-J}(S, J^{tr}).$ We first need to show that $H^{\tilde{f}\tilde{g}}(j) = (H^{\tilde{f}})^{\tilde{g}}(j)$ for every $j \in J$. Consider the following cases:

- 1. If $j = 0$ then both sides are $H(0)$. [0] is a J-stable word for every function in $Cayley_{st-J}(S, T \cup J)$.
- 2. If $j \in J$ and $[j] \in st(\tilde{f}\tilde{g}, J)$ then we also have $[j] \in st(\tilde{g}, J)$, $j' = \tilde{g}(j)$ and $[\tilde{f}(j')] \in st(\tilde{g}, J)$. Now both sides of the equation equal $H(\tilde{f}(j'))$.
- 3. If $j \in J$ and $[j] \notin st(\tilde{f}\tilde{g}, J)$. This implies that either $[j] \notin st(\tilde{g}, J)$ or $[\tilde{g}(j)] \notin st(\tilde{f}, J)$. In both cases this gives φ_0 on both sides of the equation.

It is clear that $(FG)^{\tilde{h}} = F^{\tilde{h}}G^{\tilde{h}}$ for $F, G: J^{tr} \to Cayley(S, T)$, and $\tilde{h} \in Cayley_{st-J}(S, J^{tr}).$

We may now define the semidirect product $Cayley_{st-J}(S, J^{tr}) \rtimes$ $Cayley(S,T)^{J^{tr}}.$

Now for the main theorem of this section.

Theorem 45. *Let* S *be a finite semigroup. Let* T *be an ideal of* S*, and let J be a* $\mathcal J$ *class of* S *directly above* T *. Cayley*_{st−J}(S , $T \cup J$) *is isomorphic to a subsemigroup of* $Cayley_{st-J}(S, J^{tr}) \rtimes (Cayley(S, T))^{J^{tr}}$.

Before proceeding with the proof we would like to show the reasoning behind this theorem. The action of f on a stable word w can be thought of as two actions. The first is modifying the prefix of w in J to some other prefix in J. This can be thought of as $\tilde{f} \mapsto f^{tr}$. The second part is the action of \tilde{f} is on the remainder, the part in T^* . In general this action depends on the prefix of w in J . However, Corollary [35](#page-16-0) says that this action only depends on an element in J.

Proof. Let $\tilde{f} \in Cayley_{st-J}(S, T \cup J)$ and let f be any element in $Cayley(S, J \cup T)$ that maps onto \tilde{f} . Denote by \hat{f} the function \hat{f} : $J^{tr} \rightarrow Cayley(S,T)$ defined by

$$
(\hat{f}(j))(w) = \begin{cases} f(w) & j = 0\\ f_{[j]}(w) & [j] \in st(f, J), j \in J\\ \varphi_0(w) & [j] \notin st(f, j), j \in J \end{cases}
$$

restricted to T^* (i.e. $w \in T^*$). This is well defined. The first case because every word in T^* is J-stable. The second case, since if $f(j) \in$ J, then for every $w = [t_1, \ldots, t_k] \in T^*$ we can express $f_{[j]}(w)$ as the klength suffix of $f([j, t_1, t_2, \ldots, t_k])$. Furthermore this does not depend on the choice of f.

Let $\Pi : Cayley_{st-J}(S, T \cup J) \rightarrow Cayley_{st-J}(S, J^{tr}) \rtimes Cayley(S, T)^{J^{tr}}$ be given by $\Pi(\tilde{f}) \mapsto (f^{tr}, \hat{f}), \Pi$ is obviously well-defined. To show that it is a semigroup morphism, consider $\tilde{f}, \tilde{g} \in Cayley_{st-J}(S, T \cup J)$. We have $\Pi(\tilde{f})\Pi(\tilde{g}) = (f^{tr}, \hat{f})(g^{tr}, \hat{g})$. We have seen that $(fg)^{tr} = f^{tr}g^{tr}$. In order to show that Π is a morphism, it remains to show that for every $j \in J^{tr}$ the equation $(\widehat{fg}(j))(w) = (\widehat{f}^{(g^{tr})}(j)\widehat{g}(j))(w)$ holds for every $w \in T^*$.

If we choose j as the zero element of J^{tr} the equation is immediate. Choose a $j \in J$. If $[j] \notin st(g^{tr}, J)$ then we also have $[j] \notin st((fg)^{tr}, J)$, so $\widehat{fg}(j) = \varphi_0 = \widehat{f}^{(g^{tr})}(j)\varphi_0 = \widehat{f}^{(g^{tr})}(j)\widehat{g}(j)$. If $[j] \in st(g^{tr}, J)$, $g^{tr}([j]) = [j']$ and $[j'] \notin st(f^{tr}, J)$, then as before, we have $[j] \notin st((fg)^{tr}, J)$ so $\widehat{fg}(j) = \varphi_0 = \varphi_0 \widehat{g}(j) = \widehat{f}^{(g^{tr})}(j)\widehat{g}(j)$. Finally, we consider the case $[j] \in st((fg)^{tr}, J)$.

In this case $\widehat{fg}(j) = (fg)_{[j]}$, $\hat{g}(j) = g_{[j]}$, and $(\widehat{f})^{(g^{tr})}(j) = f_{[g(j)]}$, all restricted to T^* . Let $\varphi_{s_n} \cdots \varphi_{s_1}$ be any representative of f, and $\varphi_{t_m} \cdots \varphi_{t_1}$ a representative of g. Let $\bar{t} = t_m \cdots t_1$. By Lemma [33](#page-16-1) $fg(j) = (fg)_{[j]} = (\varphi_{s_n} \cdots \varphi_{s_1} \varphi_{t_m} \cdots \varphi_{t_1})_{[j]} =$

 $\varphi_{(s_n\cdots s_1t_m\cdots t_1j)}\cdots\varphi_{(s_1t_m\cdots t_1j)}\varphi_{(t_m\cdots t_1j)}\cdots\varphi_{(t_1j)}=f_{[\bar{t}j]}g_{[j]}=\hat{f}(\bar{t}j)\hat{g}(j).$ Since $[j] \in st(g^{tr}, J)$, we have $g^{tr}(j) = t_m \cdots t_1 j = \overline{t} j \in J$. Furthermore, since $[j] \in st((fg)^{tr}, J)$ we have $g^{tr}([j]) \in st(f^{tr}, J)$, i.e. $[\bar{t}j] \in st(f^{tr}, J)$. This completes the proof that Π is a morphism.

We now show that Π is $1 - 1$. Let $f, \tilde{g} \in Cayley_{st-J}(S, T \cup J)$ and assume that $\Pi(\tilde{f}) = \Pi(\tilde{g})$. As before we will write $\Pi(\tilde{f}) = (f^{tr}, \hat{f})$ and $\Pi(\tilde{g}) = (g^{tr}, \hat{g})$. Let $w \in st(f, J)$. We need to show that $\tilde{f}(w) = \tilde{g}(w)$ (this will show that $w \in st(q, J)$). Furthermore, since $f(w) = \tilde{f}(w)$ this is the same as showing that $f(w) = g(w)$. If $w \in T^*$ then this is clear since $f(w) = (\hat{f}(0))(w) = (\hat{g}(0))(w) = g(w)$. If $w \in J^*$, then (recall that w is in $st(f, J)$ so $f(w)$ is in J^*) $f(w) = f^{tr}(w)$ $g^{tr}(w)$. Since $g^{tr}(w)$ has no zeros (because $g^{tr}(w) = f^{tr}(w)$), we have that $g^{tr}(w) = g(w)$. We now assume that $f(w)$ has a prefix in J^* and a suffix in T^* . Write this as $f(w) = w' = w'_1 \circ w'_2$ and write a decomposition for w as $w = w_1 \circ w_2$ such that $|w_1| = |w'_1|$ and $|w_2| = |w'_2|$, $w \in st(f, J)$ implies that $w_1 \in J^*$ and $w_2 \in T(T \cup J)^*$ (the first letter of w_2 is in T). We need to show that $g(w) = w'$. By definition we have $g(w) = g(w_1) \circ g_{w_1}(w_2)$. Recall that $f(w_1) \in J^*$, which implies that $f(w_1) = f^{tr}(w_1) = g^{tr}(w_1) = g(w_1)$, It remains to show that $g_{w_1}(w_2) = w_2' = f_{w_1}(w_2)$. Let $w_1 = [(a_1, b_1), \cdots, (a_n, b_n)],$ and let $j = p\varphi_1(w_1) = (a_1, b_n)$. Corollary [35](#page-16-0) shows that $f_{w_1} = f_{[j]},$ and $g_{w_1} = g_{[j]}$. In particular the restriction of this equation to \tilde{T}^* remains valid. Summing up, we have $f_{w_1}(w_2) = f_{[j]}(w_2) = \hat{f}(j)(w_2) =$ $\hat{g}(j)(w_2) = g_{[j]}(w_2) = g_{w_1}(w_2)$ which completes the proof.

Corollary 46. *Let* S *be an aperiodic semigroup,* T *an ideal of* S*, and* J *a* J *class directly above* T. If $Cayley(S,T)$ *is finite then* $Cayley_{st-J}(S,T\cup$ J) *is finite.*

Proof. It is enough to show that $Cayley_{st-J}(S, J^{tr})$ is finite. This follows from Theorem [28](#page-14-0) and Theorem [44.](#page-18-0) When applying Theorem [44](#page-18-0) we recall that the quotient of a finite aperiodic semigroup is also aperiodic.

 \Box

Theorem 47. *Let* S *be an aperiodic semigroup,* T *an ideal of* S*, and* J *a J class directly above* T. If $Cayley(S,T)$ *is finite then* $Cayley(S,T\cup$ J) *is finite.*

Proof. We will do this by showing that for some constant N, every $f \in$ $Cayley(S, T \cup J)$ can be written as a product of N or less generators.

Let $f \in Cayley(S, T \cup J)$ be given, and let $f = \varphi_{s_n} \cdots \varphi_{s_2} \varphi_{s_1}$ be a presentation of f. Since the mapping $f \mapsto f^{tr}$ is a morphism $Cayley(S, T \cup J) \rightarrow Cayley(S, J^{tr}),$ we have $f^{tr} = \varphi_{s_n}^{tr} \cdots \varphi_{s_2}^{tr} \varphi_{s_1}^{tr}$, and element in $Cayley(S, J^{tr})$.

We observe here that the elements $\{\varphi_s^{tr}\}_{s\in S}$ are a generating set for $Cayley(S, J^{tr}).$

By Corollary [32](#page-15-0) there is a constant N' (depending only on J) such that we can write $f^{tr} = g_{k+1}^{tr} \varphi_{s_k}^{tr} g_k^{tr} \varphi_{s_{k-1}}^{tr} \cdots \varphi_{s_2}^{tr} g_2^{tr} \varphi_{s_1}^{tr} g_1^{tr}$ for some $k <$ N' , where the g_i^{tr} 's never create new zeros on any input.

It is obvious that for any word $w \in (T \cup J)^*$, w is g_i -J stable if and only if w^{tr} is g_i^{tr} -*J* stable. In other words, the g_i^{tr} never create new zeros if and only if the g_i never map a letter from J to a letter in T .

Since the g_i^{tr} never create zeros, we may say that the g_i^{tr} only act on g_i^{tr} -J stable words. This means that inside the above presentation of f, the g_i will only get to act on $J-g_i$ stable words, thus we can identify g_i with \tilde{g}_i , its image in $Cayley_{st-J}(S, T \cup J)$.

By Corollary [46](#page-21-0) $Cayley_{st-J}(S, T \cup J)$ is finite, so we may assume there is another constant l such that every g_i can be written as (or replaced by) a product of l or less generators, without changing the value of $f(w)$ for any word w.

This shows that there is a number N such that every $f \in Cayley(S, T\cup$ J) can be written as a product of N or less generators. \Box

Finally, we have the main theorem of this section.

Proposition 48. *If* S *is an aperiodic semigroup,* Cayley(S) *is finite.*

Proof. Choose an ordering of the $\mathcal J$ classes of $S \{J_i\}_{i=1}^n$, such that for every $1 \leq k \leq n$ we have $\bigcup_{i=1}^{k} J_i$ is an ideal of S. It is obvious that such an ordering exists and that $J_1 \cup J_2$ is a 0-minimal ideal. We have shown that $Cayley(S, J_1\cup J_2)$ is finite, and the last theorem shows that $Cayley(S, \bigcup_{i=1}^{k} J_i)$ is finite for every k. This completes the proof. 口

7 Aperiodicity

Lemma 49. *Let* S *be a finite semigroup. Let* T *be an ideal of* S *such that* $f^n = f^{n+1}$ *for every* $f \in Cayley(S,T)$ *for some* $n \in \mathbb{N}$ *. Let J* be a *J* class of *S* directly above *T*, such that $f^l = f^{l+1}$ for every $f \in Cayley(S, J^{tr})$ for some $l \in \mathbb{N}$. Then $f^{n+l} = f^{n+l+1}$ for all $f \in Cayley(S, T \cup J)$ *.*

Proof. Let $w \in (T \cup J)^*$. Let $f \in Cayley(S, T \cup J)$. Consider $w_0 =$ $(f)^{l}(w)$. w_0 has a prefix w_J in J^* and a suffix w_T in T^* . This implies that $(f^{tr})^l(w^{tr})$ has the same prefix w_J in J^* as $f^l(w)$, and a suffix of zeros. Furthermore, since $(f^{tr})^l(w^{tr}) = (f^{tr})^{l+1}(w^{tr})$, this gives $f^{tr}(w_J) = w_J.$

We now have

$$
f^{n+l}(w) =
$$

\n
$$
f^{n}(w_{0}) =
$$

\n
$$
f^{n}(w_{J} \circ w_{T}) =
$$

\n
$$
f^{n-1}(f(w_{J}) \circ f_{w_{J}}(w_{T})) =
$$

\n
$$
f^{n-1}(w_{J} \circ f_{w_{J}}(w_{T})) =
$$

\n... =
\n
$$
(w_{J} \circ f^{n}_{w_{J}}(w_{T})) =
$$

\n
$$
(w_{J} \circ f^{n+1}_{w_{J}}(w_{T})) =
$$

\n
$$
f^{n+1}(w_{J} \circ w_{T}) =
$$

\n
$$
f^{n+1}(w_{0}) =
$$

\n
$$
f^{n+l+1}(w).
$$

 \Box

Corollary 50. *Let* S *be a finite aperiodic semigroup. Then* $\exists n \in \mathbb{N}$ such that $f^n = f^{n+1}$ for all $f \in Cayley(S)$.

Proof. As before, let $\{J_i\}_{i=1}^k$ be an enumeration of the $\mathcal J$ classes of S , such that $\bigcup_{i=1}^k J_i$ is an ideal of S for all $1 \leq i \leq k$. We saw earlier that for some $N \in \mathbb{N}$, we have $f^N = f^{N+1}$ for every $f \in Cayley(S, J_1)$. Continuing by induction on the number of $\mathcal J$ classes in S , and by applying the previous lemma, the result follows. \Box

8 Summary

The results of this paper may be summed up as follows.

Proposition 51. *Let* S *be a finite semigroup. The following are equivalent.*

- *1.* S *is an aperiodic semigroup.*
- *2.* Cayley(S) *is finite.*
- *3.* Cayley(S) *is aperiodic.*

Proof. That (1) implies (2) and (1) implies (3), are the main results of this paper. Suppose that (1) does not hold. Then S contains a non trivial group G. By results of Silva and Steinberg [\[6\]](#page-23-3), $Cayley(G)$ is a free semigroup on |G| generators. Since $Cayley(G) \prec Cayley(S)$ we see that if (1) is not true, then (2) and (3) are not true. \Box

9 Acknowledgments

The author would like to thank the anonymous referee for his careful reading and helpful comments, and the Bar-Ilan President's Fellowship for funding this research.

References

- [1] M. Arbib, K. Krohn, and J. L. Rhodes. *Algebraic theory of machines, languages and semigroups*. Academic Press, New York, 1968.
- [2] A. H. Clifford and G. B. Preston. *The Algebraic Theory of Semigroups*. American Mathematical Society, 1961.
- [3] R. I. Grigorchuk, V. N. Nekrashevich and V. I. Sushchanskii. *Automata, Dynamical systems, and groups*, Proceedings of the Steklov Institute of Mathematics 231 (2000), 128-203.
- [4] R. I. Grigorchuk, A. \ddot{Z} uk. *The lamplighter group as a group generated by a 2-state automaton, and its spectrum*, Geom. Dedicata. 87 (2001), 209-244.
- [5] Rhodes J. *Infinite iteration of matrix semigroups, I. Structure theorems for torsion semigroups*. J. Algebra 98 (1986) 422-451.
- [6] P. V. Silva and B. Steinberg. *On a class of automata groups generalizing lamplighter groups*. International journal of algebra and computation 15 (2005), 1213-1235.