

Supersonic quantum communication

J. Eisert^{1,2} and D. Gross^{2,3}

¹*Institute of Physics and Astronomy, University of Potsdam, 14476 Potsdam, Germany*

²*Quantum Optics and Laser Science, Imperial College London, London SW7 2PE, UK*

³*Institut für Mathematische Physik, Technische Universität Braunschweig, 38106 Braunschweig, Germany*

When locally exciting a quantum lattice model, the excitation will propagate through the lattice. The effect is responsible for a wealth of non-equilibrium phenomena, and has been exploited to transmit quantum information through spin chains. It is a commonly expressed belief that for local Hamiltonians, any such propagation happens at a finite “speed of sound”. Indeed, the Lieb-Robinson theorem states that in spin models, all effects caused by a perturbation are limited to a causal cone defined by a constant speed (up to exponentially small corrections). In this work we show that for meaningful translationally invariant bosonic models with nearest-neighbor interactions, this belief is incorrect: We prove that one can encounter excitations which accelerate under the natural dynamics of the lattice and allow for reliable transmission of information faster than any finite speed of sound. The effect is only limited by the model’s range of validity (eventually by relativity). The result shows that non-equilibrium dynamics in bosonic models may involve far-away regions interacting with each other, even on short time scales and when the total energy in the system is bounded. It further suggests that chains of bosonic systems may serve as fast channels for quantum communication.

Quantum spin chains—or more generally, quantum spin models on a lattice—are ubiquitous in condensed matter physics and quantum optics. They share the fundamental feature that perturbations will propagate through the lattice at some characteristic “speed of sound” [1–4]. This effect plays an important role for a wealth of non-equilibrium phenomena in many-body systems, e.g., for the dynamics of relaxation processes towards equilibrium [5, 6]. In the context of quantum information science, it has been noted that excitations propagating through a spin chain may be used to transmit quantum information—thus turning a spin chain into a quantum channel. Here, the appealing feature is that the transport is not facilitated by engineered quantum gates, but rather by the natural time evolution of the lattice system [7–9].

Because in lattice models only neighboring systems interact with each other directly, it is intuitive to assume that the maximal propagation speed of excitations (i.e., the speed of sound) is finite and given by a value characteristic for each model. Indeed, an analogous statement is clearly true for relativistic systems, where a perturbation can have no influence outside its causal cone. Mathematical physics provides a rigorous justification for this observation in the form of Lieb-Robinson bounds [1]: in spin lattice systems, perturbations can spread only linearly in time, up to exponentially small corrections. Recently, an analogous result has been proven to hold for a class of bosonic systems [10].

Interestingly, familiar as the belief that propagation of excitations in local models happens with a finite velocity may be: it is not quite right. We demonstrate that certain well-defined local bosonic models allow excitations to accelerate to arbitrarily high velocities. The effect is only limited by the range of validity of the model (which must certainly break down with the onset of relativistic effects). It occurs even for single excitations with bounded energy, traveling along a one-dimensional chain of bosons with translationally invariant nearest-neighbor interactions. From the quantum information perspective, we show that the quantum channel associated with this chain has a strictly positive information capacity, even after a time sub-linear in the length of the chain. While




FIG. 1: In quantum spin chains, local excitation will travel no faster than some characteristic speed of sound. The phenomenon defines a causal cone, outside of which any influence is exponentially suppressed. For bosonic models, it is demonstrated that the causal region may be bent to the outside, covering distances exponential in time. Information can propagate from region A to B at arbitrary velocities, via the quantum channel defined by the free time evolution of the chain. Note that this contrasts the situation in disordered spin chains, where the causal region is curved to the inside [11].

the presented models are non-integrable, we derive the results rigorously, without resorting to numerical means. We do so by considering single excitation spaces and—in this context unusual—invoke ideas from convex optimization.

There are several conceptual consequences of these results. It is now clear that any analysis of non-equilibrium processes in bosonic models must incorporate the possibility of far-away regions exchanging information on short time scales. In particular, it seems likely that simulating short-term dynamics even of the low-energy sector of bosonic models is much harder than for spin chains (where Lieb-Robinson bounds are the basis for efficient algorithms [4, 12]). Further, the results highlight the non-triviality of Lieb-Robinson bounds for spin chains with finite-dimensional constituents.

More practically—while the models we present very strongly violate any bound on propagation speeds—they have reasonable physical properties. Related models with similar features could well be realized by tuning the parameters of suitable physical systems, such as *atoms in optical lattices*, or *coupled arrays of Josephson junctions*. This opens up the possibility of observing accelerating excitations experimentally and, potentially, of using bosonic chains as fast channels

for quantum communication (at least for a meaningful range of time and distances).

Local Hamiltonians and causality in spin chains. – A local Hamiltonian on n sites is of the form

$$H = \sum_{j=1}^n h_j, \quad (1)$$

where h_j acts non-trivially only on a finite number of adjacent sites. In what follows, we will restrict attention to the most relevant case of nearest-neighbor interactions. Quantum information transmission through spin chains with Hamiltonians as above has been extensively studied in the literature.

Before turning to bosonic models, let us first recall the precise situation for spin chains (d -level systems). The fact that there always exists a speed of sound—a maximal speed of information propagation—is the content of the following *Lieb-Robinson bound* [1, 2, 3]: If A and B are operators which act non-trivially only on some (distinct) regions of the chain, then there exist constants $\mu, C > 0$ and a velocity $v > 0$ such that

$$\| [A(t), B] \| \leq C \|A\| \|B\| e^{-\mu(\text{dist}(A,B) - v|t|)}, \quad (2)$$

for all times t . Here, $A(t) = e^{iHt} A e^{-iHt}$ is the time-evolved observable, $\|\cdot\|$ the operator norm, and $\text{dist}(A, B)$ denotes the number of sites between the supports of A and B (see Fig. 1). The above form may seem somewhat awkward at first sight. To get a more physical statement, one may verify that Eq. (2) implies that any effect a perturbation A can have on a distant observable B is exponentially suppressed outside the causal cone defined by $|t| \geq \text{dist}(A, B)/v$. In particular, any non-exponentially suppressed quantum communication using this spin chain can happen at most with velocity v [4]. Due to the intuitive nature of this statement the above bound is often taken for granted or even dismissed as being “trivial”.

Supersonic communication. – Roughly, we say that a model allows for “supersonic” communication, if its dynamics can carry information over distances m in time $t(m)$ which scales sub-linearly in m . We will make this concept precise below.

The setting is a chain of n bosonic systems with nearest-neighbor interactions and open boundary conditions. The interactions should be translationally invariant ($h_i = h_j$ in Eq. (1)), up to the obvious modifications at the boundary. We refer to the left sites $1, \dots, a$ as section A of the chain, whereas sites $a + m, \dots, n$ form part B . We assume that the system is initially in some factoring, translationally invariant pure state $|\psi\rangle = |\psi_0\rangle^{\otimes n}$. A party in control of region A may now try to communicate with a party at B by either creating some excitations in her end of the chain, or else leaving the system untouched. More precisely, in the first case party A would apply a unitary operator U_A to region A . At the receiving end, party B waits for some time t before probing whether a signal corresponding to some POVM element O_B is detected. The statistics are influenced by A 's decision and given by

$$P_1 = \text{tr}[O_B e^{-itH} U_A |\psi\rangle \langle \psi| U_A^\dagger e^{itH}]$$

in case A has excited the chain and

$$P_0 = \text{tr}[O_B e^{-itH} |\psi\rangle \langle \psi| e^{itH}]$$

in case A has not done so. The classical information capacity of the channel thus defined is a function of the *signal strength* $\delta = |P_0 - P_1|$. If δ scales as $1/\text{poly}(m)$, standard protocols involving polynomially many channels used in parallel may be employed to, say, “transmit radio signals through the quantum chain” with arbitrarily high fidelity. The last relevant quantity is the energy scale of the states involved, measured, e.g., by the variances

$$E_0^2 = \langle \psi | H^2 | \psi \rangle, \quad E_1^2 = \langle \psi | U_A^\dagger H^2 U_A | \psi \rangle.$$

Low values for E_0, E_1 imply that the states are largely contained in the low energy sector of H [13]. Set $\epsilon = \max\{E_0, E_1\}$.

Below, we define and discuss three increasingly strong ways in which bosonic models could potentially violate finite bounds on the maximum propagation speed for signals. We go on to establish the main result of this work: even the strongest scenario can be realized by reasonable Hamiltonians.

(i) *Models which allow for arbitrarily fast transmission of information, using polynomial resources.* More precisely, for every signal velocity m/t , there should be suitable encoding operations $U_A(m)$ and observables $O_B(m)$ such that the signal strength $\delta(m)$ is of order $1/\text{poly}(m)$. To obtain a reasonable protocol, the energy scale $\epsilon(m)$ should grow only polynomially in m . Note that the Lieb-Robinson bounds discussed earlier rule out the existence of such models for finite-dimensional spin chains. While the latter fact alone makes models of this type interesting objects of study, it may be argued that their existence would not be too surprising. Indeed, as energy and time take reciprocal roles in (quantum) mechanics, it is plausible that adding “more energy” to the system may lead to faster dynamics. This motivates the next, more stringent, situation.

(ii) *Models for which the signal velocity scales faster than the inverse energy.* In addition to the definitions above, we demand that $m\delta(m)/(t\epsilon(m)) \rightarrow \infty$ as $m \rightarrow \infty$. For such models, the phenomenon cannot just be explained by the fact that unbounded Hamiltonians allow for signals with higher energies and thus faster dynamics.

In scenario (i), (ii) above, information propagates at arbitrarily high velocities—yet the distance covered is still linear in time (so the causal regions are cones with arbitrarily wide opening angles). The final situation is more demanding, requiring that excitations “speed up” as they propagate.

(iii) *Models allowing for accelerating signals.* Here, we require that the signal strength δ , the energy scale ϵ and, in fact, the encoding operation U_A do not depend on the distance m , while the time t should scale sub-linearly in m .

In the next section, we construct a Hamiltonian which exhibits behavior of type (iii) (and hence also of type (i) and (ii)) in a quite radical fashion.

Models. – The type of models we subsequently allow for are governed by nearest-neighbor Hamiltonians of the form

$$H = \sum_{j=1}^{n-1} f_{j,j+1} + \sum_{j=1}^n g_j$$

with interaction term $f_{j,j+1}$ and on-site term g_j . For $f_{j,j+1} = a_j^\dagger a_{j+1} + h.c.$, and $g_j = a_j^\dagger a_j$ this is an instance of a *harmonic chain*. Harmonic versions of Lieb-Robinson bounds have recently been derived [10], and such chains have also been explored for quantum state transfer [9]. For an on-site interaction of the form $g_j = \mu a_j^\dagger a_j + U a_j^\dagger a_j (a_j^\dagger a_j - 1)$ this gives rise to the *Bose-Hubbard model*.

Specific “exchange interaction” model. – We will pay special attention to the following model, defined for bosons with spin 1 (so associated with the Hilbert space $\mathcal{H} = \mathcal{L}^2(\mathbb{R}) \otimes \mathbb{C}^3$). We define for site j the operators $A_{j;k,l} = |k, \uparrow\rangle\langle l, \downarrow|$ and $B_{j;k,l} = |k, \downarrow\rangle\langle l, \downarrow|$. The Hamiltonian is specified by setting

$$f_{j,j+1} = \sum_{k,l=0}^{\infty} (2l-1) (iA_{j;l,k}^\dagger B_{j+1;l,k} + h.c.),$$

$$g_j = 2 \sum_{k=0}^{\infty} (ik|k+1, \uparrow\rangle\langle k, \downarrow| + h.c.).$$

Note that $f_{j,j+1}$ may be looked at as a variant of the familiar exchange interaction. Clearly, H is a legitimate physical bosonic Hamiltonian with translationally invariant nearest-neighbor interactions. We will prove our claim three steps.

1. Mapping to an excitation Hamiltonian. – To start with, $|0, \downarrow; \dots, 0, \downarrow\rangle$ is an eigenstate of the Hamiltonian. If we now place a single excitation with spin \uparrow at the first site—so start with the initial state vector $|1, \uparrow; 0, \downarrow; \dots, 0, \downarrow\rangle$ —we see that time evolution will only couple this to state vectors of the form

$$||l\rangle\rangle = \begin{cases} |0, \downarrow; \dots, 0, \downarrow, \frac{l+1}{2}, \downarrow; 0, \downarrow; \dots, 0, \downarrow\rangle, & \text{if } l \text{ odd,} \\ |0, \downarrow; \dots, 0, \downarrow, \frac{l+2}{2}, \uparrow; 0, \downarrow; \dots, 0, \downarrow\rangle, & \text{if } l \text{ even,} \end{cases}$$

with particles at site $j = l/2 + 3/2$ and $j = l/2 + 1$, respectively (for $l = 0, \dots, 2n-1$). When considering only such excitations, we can hence pass to a new effective Hamiltonian E with Hilbert space $\mathcal{K} = \mathbb{C}^{2n}$ and initial condition $||0\rangle\rangle$,

$$E = \sum_{l=0}^{2n-1} i(l+1) (||l+1\rangle\rangle\langle\langle l| - ||l\rangle\rangle\langle\langle l+1|).$$

This Hamiltonian faithfully models the single excitation sector in the above sense. Note that $||l\rangle\rangle$ has now two roles: It both refers to a position in the original lattice, as well as the particle number in the original Hamiltonian. Also, to simplify notation later on, we will at this point pass to the half-open chain by setting $n = \infty$ [14].

2. Moments. – In this single excitation sector, time evolution corresponds to $\rho(t) = e^{-itE}\rho(0)e^{itE}$, for states on \mathcal{K} . It proves expedient to introduce the operators

$$X = \sum_{l=0}^{\infty} (l+1) ||l\rangle\rangle\langle\langle l|,$$

$$P = \sum_{l=1}^{\infty} (l+1) (||l+1\rangle\rangle\langle\langle l+| + |l\rangle\rangle\langle\langle l+1|).$$

Note that X corresponds to a discrete position operator, measuring twice the distance of the original model. There is a

lot of structure in this model: It is constructed such that the commutation relations between these operators form a closed algebra, and we find

$$i[E, X] = P, \quad i[E, P] = 4X - 2\mathbb{1}.$$

Using the familiar Baker-Hausdorff formula, the Heisenberg picture time evolution of X under E is given by

$$X(t) = e^{itE} X e^{-itE} = X + t[iE, X] + \frac{t^2}{2}[iE, [iE, X]] + \dots$$

One obtains a closed-form expression for $X(t)$ by exploiting the relations in the algebra $\mathcal{A} = \{E, X, P, \mathbb{1}\}$ to iteratively solve these nested commutators. Explicitly

$$X(t) = X + P \sum_{l=1}^{\infty} \frac{t^{2l-1} 4^{l-1}}{(2l-1)!} + (4X - 2\mathbb{1}) \sum_{l=1}^{\infty} \frac{t^{2l} 4^{l-1}}{(2l)!}$$

$$= X - P \frac{1}{2} \sinh(2t) + (X - \frac{1}{2} \mathbb{1}) (\cosh(2t) - 1),$$

where the final equality uses the respective Taylor series. For the time evolution of X starting from the single excitation $||0\rangle\rangle$, we hence find

$$\langle\langle 0|X(t)|0\rangle\rangle = \frac{1}{2}(1 + \cosh(2t)).$$

Thus, the expectation value of X is increasing exponentially in t . This fact alone, however, is not enough to show that we have signaling: It could be that the excitation develops a long asymptotic tail that leads to large first moments, but carries a small weight.

Thus, further information is needed. It turns out that knowledge of the second moments $\langle\langle 0|X(t)^2|0\rangle\rangle - \langle\langle 0|X(t)|0\rangle\rangle^2$ is sufficient to prove signaling using the convex optimization ideas below. An analogous—if more tedious—calculation arrives at

$$\langle\langle 0|X(t)^2|0\rangle\rangle = 1 + \frac{1}{4} \sinh^2(2t) + (\cosh(2t) - 1)$$

$$+ \frac{1}{4} (\cosh(2t) - 1)^2 = \cosh^2(t) \cosh(2t).$$

3. Hitting time from a convex optimization problem. – For some site $m > 1$ define the *hitting operator*

$$T = \sum_{l=2m-1}^{\infty} ||l\rangle\rangle\langle\langle l|.$$

So $P_1 = \text{tr}[\rho(t)T]$ is the signal a distant observer m sites away from the origin may receive, compared to $P_0 = 0$. Hence, the set $A = \{1\}$ is the single first site, whereas $B = \{m+1, \dots, \infty\}$ is the natural right part of the chain. Let us set $M = 2m-2$. We will now bound this expectation value by analytically solving a convex optimization problem:

$$\text{minimize} \quad \sum_{l=M+1}^{\infty} p_l,$$

$$\text{subject to} \quad \sum_{l=0}^{\infty} (l+1)p_l = a(t), \quad \sum_{l=0}^{\infty} (l+1)^2 p_l = b(t),$$

$\sum_{l=0}^{\infty} p_l = 1$, and $p_l \geq 0$ for all l , where $a(t) = (1 + \cosh(2t))/2$ and $b(t) = \cosh^2(t) \cosh(2t)$. So we minimize the signal, given first and second moments, as a worst case analysis. This is an (infinite-dimensional) linear program, and hence an instance of a convex optimization problem. We can readily get a bound to the optimal solution by identifying a suitable solution to the Lagrange dual problem [15]. This dual is found to be

$$\begin{aligned} & \text{maximize} && -d^T y, \\ & \text{subject to} && F^T y \geq -c, \end{aligned}$$

where $F_{1,j} = j$, $F_{2,j} = j^2$, $F_{3,j} = 1$ for $j = 1, 2, \dots$, and $d(t) = (a(t), b(t), 1)^T$. Also, $c = (0, \dots, 0, 1, 1, \dots)^T$, as a vector starting with M zeros. The latter is an optimization problem over y . Any solution we can identify of the dual will give a lower bound to the primal. It can be shown explicitly that the subsequent vector

$$y^* = (-2(1+M)/M^3, 1/M^4, (1+M)^2/M^2 - 1)^T.$$

is always a feasible solution of the dual problem. Taking the time $t^* = \log(M)$ gives

$$\begin{aligned} c^T x^* & \geq -d^T y^* = 2a(\log(M))(1+M)/M^3 \\ & - b(\log(M))/M^4 - (1+M)^2/M^2 + 1 =: g(M), \end{aligned}$$

For this function we have that $\lim_{M \rightarrow \infty} g(M) = 3/8$, and $g(M) > 1/5$ for $M \geq 9$. The solution of the dual will hence give rise to a lower bound of our primal problem. Hence, after a time t^* logarithmic in m , a signal of constant strength $\text{tr}[T\rho(t^*)] > 1/5$ will have reached party B ! This means, of course—within the validity of the model—that we can in principle signal at any speed over arbitrary distances: The signal will not even decay, and the Holevo- χ and the *classical information capacity* of the associated quantum channel are indeed constant. One can communicate with an exponentially accelerating signal of type (iii) in the above classification [16].

Bose-Hubbard-type models. – In a sketchier fashion, we

will now briefly address the question whether even Bose-Hubbard-type models

$$H = \sum_{j=1}^n \left((a_j^\dagger a_{j+1} + h.c.) + h(a_j^\dagger a_j) \right),$$

with some function $h : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ can display such a behavior. Indeed, it can be seen that if h increases in n sufficiently strongly, leading to highly interacting particles, then there exists for each chain length n an N (polynomial in n) such that the initial state $|\psi\rangle = |N+1\rangle|N\rangle^{\otimes(n-1)}$ —so an additional particle at site $A = \{1\}$ compared to $|N\rangle^{\otimes n}$ —will lead to a signal at site $B = \{n\}$ that is at most polynomially suppressed in n . The signal increases linearly in N . Hence, hopping models can also display violations of any finite bound on propagation speeds, at least in the sense (i) of the above classification.

Summary and Outlook. – In this work, we have shown that in systems of locally interacting bosons, excitations may accelerate and carry information faster than any finite speed of sound. This observation defies the intuition that information transfer happens at a well-defined speed, highlighting the significance of the Lieb-Robinson bound for spin chains. Looking ahead, the viability of such systems for the fast transport of quantum information could be explored. Here, numerical investigations of initially highly filled Bose-Hubbard systems would be relevant. In fact, such a setting does not appear to be out of the reach of current experiments: investigating propagation effects in non-equilibrium configurations of, say, Bose-Einstein condensates in optical lattices would be a highly interesting enterprise.

Acknowledgments. – We thank J. Preskill, T.J. Osborne, M.B. Hastings, A. Retzker, J. Kiukas, and F. Verstraete for fruitful discussions. We specifically thank T.J. Osborne for a key remark that was of use in the simplification of the calculation of moments. This work has been supported by the EU (QAP, COMPAS, CORNER), the EPSRC, QIP-IRC, Microsoft Research, and the EURYI.

-
- [1] E.H. Lieb and D.W. Robinson, *Commun. Math. Phys.* **28**, 251 (1972).
- [2] M.B. Hastings, *Phys. Rev. B* **69**, 104431 (2004); M.B. Hastings and T. Komatsu, *Commun. Math. Phys.* **265**, 781 (2006).
- [3] B. Nachtergaele and R. Sims, *Commun. Math. Phys.* **265**, 119 (2006).
- [4] J. Eisert and T.J. Osborne, *Phys. Rev. Lett.* **97**, 150404 (2006); S. Bravyi, M.B. Hastings, and F. Verstraete, *ibid.* **97**, 050401 (2006).
- [5] M. Cramer, C.M. Dawson, J. Eisert, and T.J. Osborne, *Phys. Rev. Lett.* **100** (2008); M. Cramer, A. Fleisch, I.A. McCulloch, U. Schollwöck, and J. Eisert, *ibid.* **101**, 063001 (2008).
- [6] P. Calabrese and J. Cardy, *Phys. Rev. Lett.* **96** (2006) 136801; A. Laeuchli and C. Kollath, *J. Stat. Mech.* (2008) P05018; N. Schuch, M.M. Wolf, K.G.H. Vollbrecht, J.I. Cirac, *New J. Phys.* **10**, 033032 (2008).
- [7] S. Bose, *Contemp. Phys.* **48**, 13 (2007).
- [8] M. Christandl, N. Datta, A. Ekert, and A.J. Landahl, *Phys. Rev. Lett.* **92**, 187902 (2004); T.S. Cubitt, F. Verstraete, and J.I. Cirac, *Phys. Rev. A* **71**, 052308 (2005); D. Burgarth, V. Giovannetti, and S. Bose, *Phys. Rev. A* **75**, 062327 (2007).
- [9] J. Eisert, M.B. Plenio, S. Bose, J. Hartley, *Phys. Rev. Lett.* **93**, 190402 (2004); M.B. Plenio, J. Hartley, and J. Eisert, *New J. Phys.* **6**, 36 (2004); M.J. Hartmann, M.E. Reuter, and M.B. Plenio, *ibid.* **8**, 94 (2006).
- [10] B. Nachtergaele, H. Raz, B. Schlein, and R. Sims, *arXiv:0712.3820*; M. Cramer, A. Serafini, and J. Eisert, *arXiv:0803.0890*; O. Buerschaper, M.M. Wolf, J.I. Cirac, in preparation.
- [11] C.K. Burrell and T.J. Osborne, *Phys. Rev. Lett.* **99**, 167201 (2007).
- [12] T.J. Osborne, *Phys. Rev. Lett.* **97**, 157202 (2006); M.B. Hastings, *Phys. Rev. B* **77**, 144302 (2008).
- [13] By Chebychev's inequality, the probability of observing an en-

ergy value exceeding tE_0 when performing a measurement on $|\psi\rangle$ is smaller than $1/t^2$.

- [14] With greater notational effort, the statements involving infinite chains can be made more precise, by e.g., passing to the Heisenberg picture and employing a C^* -algebraic language.
- [15] S. Boyd and L. Vandenberghe, *Convex optimization* (Cambridge University Press, Cambridge, 2004).
- [16] One could reasonably conjecture a ‘‘compatibility property’’: if two Hamiltonians V and W both satisfy a Lieb-Robinson bound, then also $V + W$. However, no such property can hold. A counter-example is obtained from H by including every second interaction term into V and setting $W = H - V$.

I. APPENDICES

For clarity of the argument, we present a few aspects of the above presentation in some more detail. Note that this material is identical with the above one.

Appendix A: The spin Hamiltonian

We consider a bosonic model with a spin-1 internal degree of freedom. The local Hilbert space is hence spanned by $\{|k, \uparrow\rangle, |k, \downarrow\rangle : k \in \mathbb{N}_0\}$. We will be able to keep track of local excitations in the following model. We take the Hamiltonian

$$H = \sum_{j=1}^n \left(\sum_{k,l=0}^{\infty} (2l-1) (i|k, \downarrow\rangle\langle l, \uparrow| \otimes |l, \downarrow\rangle\langle k, \downarrow| + h.c.) + g_j \right),$$

where the on-site interaction is taken to be

$$g_j = 2 \sum_{k=0}^{\infty} (ik|k+1, \uparrow\rangle\langle k, \downarrow| + h.c.).$$

The hopping term is a variant of an exchange interaction between neighboring sites (except from the spin, this would exactly be a standard exchange interaction). This Hamiltonian can also be written as

$$H = \sum_{j=1}^n \left(\sum_{k,l=0}^{\infty} (2l-1) (iA_{j;l,k}^\dagger B_{j+1;l,k} + h.c.) + g_j \right),$$

with the hopping operators at site j being defined as

$$A_{j;k,l} = |k, \uparrow\rangle\langle l, \downarrow|, \quad B_{j;k,l} = |k, \downarrow\rangle\langle l, \downarrow|,$$

The initial state vector is taken to be

$$|\psi\rangle = |0, \downarrow\rangle^{\otimes n}.$$

This is obviously an eigenstate of the Hamiltonian. Now we excite site 1, forming set $A = \{1\}$, by setting it to $|1, \uparrow\rangle$, while keeping the rest of the chain unaltered. We then see that the initial state vector after the excitation

$$|1, \uparrow\rangle|0, \downarrow\rangle^{\otimes(n-1)}$$

couples to

$$|0, \downarrow\rangle|1, \downarrow\rangle|0, \downarrow\rangle^{\otimes(n-2)},$$

and this to the former and to

$$|0, \downarrow\rangle|2, \uparrow\rangle|0, \downarrow\rangle^{\otimes(n-2)}.$$

Now the pattern is clear: All excitations are contained in

$$\text{span} \quad \{ |0, \downarrow\rangle^{\otimes k} |k, \downarrow\rangle |0, \downarrow\rangle^{\otimes(n-k-1)}, \\ |0, \downarrow\rangle^{\otimes k} |k+1, \uparrow\rangle |0, \downarrow\rangle^{\otimes(n-k-1)} \},$$

and each basis vector—except at the ends of the chain—couples to two further basis vectors. It is now a straightforward exercise to verify that in this subspace of excitations, one arrives at the above given effective Hamiltonian.

Appendix B: Convex optimization problem

The matrix F in the convex optimization problem defining our hitting time problem is given by

$$F = \begin{bmatrix} 1 & 2 & 3 & 4 & \dots \\ 1 & 4 & 9 & 16 & \dots \\ 1 & 1 & 1 & 1 & \dots \end{bmatrix}.$$

The first row captures the first moments, the second the second moments, whereas normalization of the probability distribution is enforced by the third one—together with the fact that all entries of x are positive. The vector $c = (0, \dots, 0, 1, \dots, 1)^T$, as a vector starting with M zeros. The primal problem can hence be written as

$$\begin{aligned} & \text{minimize} && c^T x, \\ & \text{subject to} && Fx = d, \\ & && x \geq 0, \end{aligned}$$

where $d(t) = (a(t), b(t), 1)^T$. This is a linear program in standard form, with a matrix inequality constraint, and positive elements of the objective vector x . In this form it is easiest to identify the dual problem. It is given by

$$\begin{aligned} & \text{maximize} && -d^T y, \\ & \text{subject to} && F^T y \geq -c, \end{aligned}$$

where now y is not constrained to be positive. The constraints on the vector y can equally be written as

$$jy_1 + j^2 y_2 + y_3 + \delta_{j>M} \geq 0 \quad (3)$$

for all j , where $\delta_{j>M}$ takes the value 1 if $j > M$ and is zero otherwise. The question is: Can we find for each M a choice of (y_1, y_2, y_3) and a time t such that we can meaningfully find a positive solution to the dual problem?

We can take any solution to the dual problem. We choose

$$\begin{aligned} y_1 &= -2(1+M)/M^3, \\ y_2 &= 1/M^4, \\ y_3 &= (1+M)^2/M^2 - 1, \end{aligned}$$

and will verify that it is indeed a solution with the appropriate properties. The intuition behind this construction is as follows: Consider the quadratic function $f : [0, \infty) \rightarrow \infty$ as

$$f(x) = xy_1 + x^2y_2 + y_3.$$

This is identical with the left hand side of Eq. (3), up to being defined on $[0, \infty)$. We require that f takes the value 0 for the first time exactly at $x = M$, and that $f(x) > -1$ for all x . Taking $y_2 = 1/M^4$ then gives rise to the above construction, satisfying Eq. (3) for all M and all j . We have $jy_1 + j^2y_2 + y_3 = 0$ exactly at $j = M$, which is a desirable feature to get a meaningful bound to the original problem at hand. This is hence always a feasible solution to the dual problem. We will therefore obtain a lower bound to the optimal solution of the primal problem $c^T x^*$ as

$$\begin{aligned} c^T x^* &\geq -d^T y \\ &= 2a(t)(1+M)/M^3 - b(t)/M^4 \\ &\quad - (1+M)^2/M^2 + 1. \end{aligned}$$

Choosing the time $t^* = \log(M)$, logarithmic in the distance, we find

$$\begin{aligned} c^T x^* &\geq 2a(\log(M))(1+M)/M^3 - b(\log(M))/M^4 \\ &\quad - (1+M)^2/M^2 + 1 =: g(M), \end{aligned}$$

Now it is not difficult to see that indeed, in the limit of large M we have

$$\lim_{M \rightarrow \infty} g(M) = 3/8,$$

and that $g(M) > 1/5$ for all $M \geq 9$. This means that for each large m —and hence each large M —we can arrive at a constant signal for an appropriate constant signal.

Appendix C: Bose-Hubbard-type models

In this final appendix, we briefly discuss Bose-Hubbard-type models of the form

$$H = \sum_{j=1}^n \left((a_j^\dagger a_{j+1} + h.c.) + h(a_j^\dagger a_j) \right),$$

with some function $h : \mathbb{R}^+ \rightarrow \mathbb{R}^+$, which will be specified later. As before, this Hamiltonian is modified in the obvious way at site n . That is, this is a particle-number preserving Hamiltonian with hopping and an on-site interaction that merely depends on particle number.

For each chain length n , we will consider a relevant n -dimensional subspace of the original Hilbert space

$$\mathcal{L}_N = \text{span}\{|j_1, \dots, j_n\rangle | \exists k : j_k = N+1, j_l = N \forall l \neq k\},$$

for a suitable integer N , and denote with $\mathcal{L}_N^\perp = \mathcal{H} \setminus \mathcal{L}_N$ the orthogonal complement of the infinite-dimensional Hilbert space. The initial state vector of the chain of length n is then taken to be

$$|\psi\rangle = |N+1\rangle|N\rangle^{\otimes(n-1)} \in \mathcal{L}_N.$$

This means that we fill up all number states with bosons at each site up to N , and add an additional particle at site $A = \{1\}$ as a local excitation. We will now see that we can then approximate the dynamics of the chain within a spin chain restricted to a single excitation \mathcal{L}_N arbitrarily well. Define for site j the operator $A_j = |N\rangle\langle N+1|$, then the above chain is approximated in its dynamics by the dynamics of

$$V = (N+1) \sum_{j=1}^n \left(A_j^\dagger A_{j+1} + h.c. \right).$$

This dynamics is specifically simple, and we can see that in a time t^* that is linear in N we will receive a signal at site $B = \{n\}$ that is at most polynomially suppressed in n . This clearly means that this signal will be received at an arbitrarily velocity, if N is sufficiently large. To see that this mapping can be done to arbitrary approximation, we need the subsequent observation:

Lemma 1 *For any Hermitian matrix M , partitioned as*

$$M = \begin{bmatrix} A & B \\ B^\dagger & C \end{bmatrix}.$$

we have that for all $t \in [0, \infty)$,

$$\lim_{x \rightarrow \infty} \sup_{K \in \mathcal{M}_x} \|O e^{itK} O^\dagger - e^{itA}\| = 0$$

where

$$\mathcal{M}_x = \left\{ M + \begin{bmatrix} 0 & 0 \\ 0 & P \end{bmatrix} : P \geq x \mathbb{1} \right\}, \quad O = [\mathbb{1} \ 0].$$

Here, $\|\cdot\|$ denotes the operator matrix or vector norm. To prove this, note that

$$\sum_{j=1}^d \lambda_j^\dagger \left(\begin{bmatrix} A & B \\ B^\dagger & C+P \end{bmatrix} \right) = \inf_Q \text{tr} \left[Q \begin{bmatrix} A & B \\ B^\dagger & C+P \end{bmatrix} \right],$$

where the infimum is taken over Q that are projectors of rank d , when A is a $d \times d$ -matrix. This infimum exists for any $M \in \mathcal{M}_x$, call it Q_M . Since $\|A\| = c$ is constant, we have that

$$\lim_{x \rightarrow \infty} \sup_{M \in \mathcal{M}_x} \left\| Q_M - \begin{bmatrix} \mathbb{1} & 0 \\ 0 & 0 \end{bmatrix} \right\| = 0.$$

Hence, since $|e^{it\lambda_j^\dagger(M)}| = 1$, the assertion follows.

Subsequently, the submatrix A will be identified with H restricted to \mathcal{L}_N , whereas C is the restriction to the orthogonal complement \mathcal{L}_N^\perp . Using this observation, we see that for each $\varepsilon > 0$ we can find a function $h : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ such that the following property is satisfied: For each chain length n there exists an N such that for the above hitting time t^*

$$\|e^{-it^*H}|\psi\rangle - e^{-it^*V}|\psi\rangle\| < \varepsilon$$

for all $t \in [0, t^*]$. But this means that we can signal with this chain, in the sense of (iii) in the above classification, as the class of restricted models has this property.