Fourier transform of function on locally compact Abelian groups taking value in Banach spaces [∗]

Yauhen Radyna, Anna Sidorik

Abstract

We consider Fourier transform of vector-valued functions on a locally compact group G, which take value in a Banach space X , and are square-integrable in Bochner sense. If G is a finite group then Fourier transform is a bounded operator. If G is an infinite group then Fourier transform $F: L_2(G, X) \to L_2(\widehat{G}, X)$ is a bounded operator if and only if Banach space X is isomorphic to a Hilbert one.

1 Fourier transform over groups R, Z, T

In the paper [\[1\]](#page-8-0) J. Peetre proved an extension of Hausdorf–Young's theorem describing image of $L_q(\mathbf{R})$ under Fourier transform. He considered vector-valued $x \in L_q(\mathbf{R}, X)$, $1 \le q \le 2$ on the real axis taking value in Banach space X , and integrable in Bochner sense, i.e. weakly measurable with finite norm

$$
||x||_{L_q(\mathbf{R},X)} = \left(\int_{\mathbf{R}} ||x(t)||_X^q dt\right)^{1/q}.
$$

J. Peetre noted, that for $q = 2$ in all known to him cases Fourier transform

$$
\mathcal{F}: L_2(\mathbf{R}, X) \to L_2(\mathbf{R}, X), \quad (\mathcal{F}x)(s) = \int_{\mathbf{R}} x(t)e^{-2\pi i s t} dt.
$$

was bounded only if X was isomorphic to a Hilbert space. In [\[3\]](#page-8-1) Polish mathematician S. Kwapien in fact proved the following

Theorem 1 Statements below are equivalent:

- 1) Banach space X is isomorphic to a Hilbert one.
- 2) There exists $C > 0$ such that for any positive integer n and $x_0, x_1, x_{-1}, \ldots, x_n, x_{-n} \in X$

$$
\int_0^1 \Big\| \sum_{k=-n}^n e^{2\pi ikt} \cdot x_k \Big\|^2 dt \le C \sum_{k=-n}^n \|x_k\|^2.
$$

3) There exists $C > 0$ such that for any positive integer n and $x_0, x_1, x_{-1}, \ldots, x_n, x_{-n} \in X$

$$
C^{-1} \sum_{k=-n}^{n} ||x_k||^2 \le \int_0^1 \left\| \sum_{k=-n}^n e^{2\pi ikt} \cdot x_k \right\|^2 dt.
$$

[∗]MSC2000: 46C15, 43A25; keywords: Hilbert space, Fourier transform, locally compact groups;

4) Fourier transform F initially defined on a dense subspace of simple functions $D_{\mathcal{F}} \subset L_2(\mathbf{R}, X)$,

$$
D_{\mathcal{F}} = \left\{ x(t) = \sum_{k=1}^{n} I_{A_k}(t) \cdot x_k \right\},\
$$

is a bounded operator. Here A_k are disjoint subset of finite measure in \mathbf{R} , I_{A_k} are indicators (i.e. functions equal to 1 on A_k and to 0 elsewhere), x_k are vectors in X.

Let us define Fourier transform of a vector-valued function over integers **Z** by

$$
\mathcal{F}_{\mathbf{Z}}: L_2(\mathbf{Z}, X) \to L_2(\mathbf{T}, X): (x_k) \mapsto \sum_{k \in \mathbf{Z}} e^{-2\pi i kt} \cdot x_k,
$$

Here **T** denotes a one-dimensional torus $T = R/Z$, which is isomorphic to [0, 1] with the length in the cathegory of spaces with a measure.

Statement 2) in Theorem [1](#page-0-0) means that $\mathcal{F}_{\mathbf{Z}}\mathcal{I}_{\mathbf{Z}}$ is a bounded operator (here $\mathcal{I}_{\mathbf{Z}}$ denotes the operator of changing variable \mathcal{I}_z : $(x_k) \mapsto (x_{-k})$, which is an isometry). That is why \mathcal{F}_z is bounded on a dense subspace of $L_2(\mathbf{Z}, X)$ consisting of compactly-supported functions, and can be continued to the whole $L_2(\mathbf{Z}, X)$.

Statement 3) in Theorem [1](#page-0-0) means that inverse Fourier transform

$$
\mathcal{F}_{\mathbf{Z}}^{-1}: L_2(\mathbf{T}, X) \to L_2(\mathbf{Z}, X)
$$

is a bounded operator.

2 Generalization and necessary facts

Fourier transform of Banach space-valued functions on a group different from $\mathbf{R},\mathbf{Z},\mathbf{T}$ (namely on additive group of p-adic field \mathbf{Q}_p) was considered in [\[4\]](#page-9-0).

Now it's natural to look at the general case of arbitrary locally compact group G. We consider functions on G taking value in Banach space X , that are square-integrable in Bochner sense, and Fourier transform

$$
\mathcal{F} \equiv \mathcal{F}_G : L_2(G, X) \to L_2(\widehat{G}, X), \quad (\mathcal{F}x)(\xi) = \int_G \langle \xi, t \rangle_G x(t) d\mu_G(t).
$$

Here \widehat{G} is Pontryagin dual to G (group of characters), $\langle \xi, t \rangle_G$ is the canonical pairing between \widehat{G} and G, μ_G is Haar measure.

First we recall some necessary results. We refer to [\[5\]](#page-9-1) for results in harmonic analysis, and to [\[6\]](#page-9-2) for structure theory of locally compact groups, Bruhat–Schwartz theory is exposed in [\[7\]](#page-9-3).

Fix a dual Haar measure $\mu_{\widehat{G}}$ on \widehat{G} such that scalar Steklov–Parseval's equality holds

$$
||\varphi||^2_{L_2(G)}=\int\limits_G|\varphi|^2d\mu_G=\int\limits_{\widehat{G}}|\mathcal{F}\varphi|^2d\mu_{\widehat{G}}=||\mathcal{F}\varphi||_{L_2(\widehat{G})}.
$$

Let $\mathcal{S}(G)$ denote Bruhat–Schwartz space of "smooth fastly decreasing" functions on G. It's useful to take a dense subspace

$$
D_{\mathcal{F}} = L_2(G) \otimes X \subset L_2(G, X),
$$

as initial domain of \mathcal{F} , where it acts by

$$
\mathcal{F}\Big(\sum_{k=1}^n \varphi_k(t) \cdot x_k\Big) = \sum_{k=1}^n \big((\mathcal{F}\varphi_k)(\xi) \cdot x_k\big).
$$

To show denseness of $D_{\mathcal{F}}$ and denseness of $\mathcal{S}(G) \otimes X \subset L_2(G) \otimes X$ consider indicator I_A of arbitrary measurable subset of finite measure (i.e. functions equal to 1 on A and to 0 elsewhere). Clearly, $I_A \in L_2(G)$, and it can be approximated by elements of $\mathcal{S}(G)$ using convolution with any δ -net consisting of Bruhat–Schwartz functions. By definition of $L_2(G, X)$ finite linear combinations

$$
\sum_{k=1}^{n} I_{A_k}(t) \cdot x_k \in L_2(G) \otimes X \equiv D_{\mathcal{F}}
$$

are dense in $L_2(G, X)$. Here A_k are measurable disjoint subsets of finite measure in G, I_{A_k} are indicators, and $x_k \in X$.

Due to the fact that scalar Fourier transform is a bijection from $L_2(G)$ into $L_2(G)$, and is a bijection from $\mathcal{S}(G)$ into $\mathcal{S}(\widehat{G})$, the restriction of vector-valued Fourier transform onto $L_2(G) \otimes X$ is a bijection into $L_2(\widehat{G}) \otimes X$, and its restriction onto $\mathcal{S}(G) \otimes X$ is a bijection to $\mathcal{S}(\widehat{G}) \otimes X$.

To handle the case of infinite group G we need a theorem describing structure of locally compact Abelian groups [\[6\]](#page-9-2).

Theorem 2 Let G be a locally compact Abelian group. Then G is a union of open compactly generated subgroups H. Topology of G is the topology of inductive limit. In turn, each compactlygenerated subgroup $H \subset G$ is a projective limit of elementary factor-groups H/K , where $K \subset H$ are compact. "Elementary" here means that H/K is isomorphic to cartesian product

$$
H/K \cong \mathbf{R}^{a_{H,K}} \times \mathbf{T}^{b_{H,K}} \times \mathbf{Z}^{c_{H,K}} \times \mathbf{F}_{H,K},
$$

where $a_{H,K} \geq 0$, $b_{H,K} \geq 0$, $c_{H,K} \geq 0$, and $\mathbf{F}_{H,K}$ is a finite group.

Definition 1 We say that G contains an **R**-component if for some H, K in Theorem [2](#page-2-0) number $a_{H,K}$ is positive in elementary factor-group $\mathbf{R}^{a_{H,K}} \times \mathbf{T}^{b_{H,K}} \times \mathbf{Z}^{c_{H,K}} \times \mathbf{F}_{H,K}$.

In the same way we use phrases "group G contains \mathbf{Z} -component", "group G contains \mathbf{T} -component".

Now recall some properties of Pontryagin duality.

Consider an exact sequence of homomorphisms (i.e. image of each homomorphism coincides the kernel of the following one)

$$
0 \to K \to G \to G/K \to 0, \quad 0 \to H \to G \to G/H \to 0,
$$

where K is compact, and H is open. Dual groups \widehat{G}/\widehat{K} , \widehat{G}/\widehat{H} can be identified with annihilators

$$
K_G^{\perp} = \{ \chi \in \widehat{G} : \ \forall g \in K, \ \langle \chi, g \rangle = 1 \}, \quad H_G^{\perp} = \{ \chi \in \widehat{G} : \ \forall g \in H, \ \langle \chi, g \rangle = 1 \}.
$$

Moreover, K_G^{\perp} is an open subgroup, and H_G^{\perp} is compact. One has dual oppositely-directed exact sequences

$$
0 \leftarrow \widehat{G}/K_G^{\perp} \leftarrow \widehat{G} \leftarrow K_G^{\perp} \leftarrow 0, \quad 0 \leftarrow \widehat{G}/H_G^{\perp} \leftarrow \widehat{G} \leftarrow H_G^{\perp} \leftarrow 0.
$$

If $K \subset H$, then $K_G^{\perp} \supset H_G^{\perp}$.

It's not hard to see that Fourier transform of a function on G , that is supported in open subgroup H and is constant on cosets of compact subgroup $K \subset H$, is a function on \tilde{G} supported in K_G^{\perp} and constant on cosets of H_G^{\perp} .

3 The case of an arbitrary local compact group

We will prove some necessary lemmas before formulating the main result.

Lemma 1 Assume that Banach space X is isomorphic to a Hilbert one, i.e. there exists inner product $(\cdot, \cdot)_X$ on X such that for some $C > 0$ the following inequality is true

$$
C^{-1}(x,x)_X^{1/2} \le ||x||_X \le C(x,x)_X^{1/2}.
$$

Then Fourier transform $\mathcal{F}: L_2(G,X) \to L_2(\widehat{G},X)$ is bounded.

Proof. Consider vector-valued Steklov–Parseval's equality

$$
(\mathcal{F}\varphi,\mathcal{F}\varphi)_{L_2(\widehat{G},X)}=\int_{\widehat{G}}(\mathcal{F}\varphi(\xi),\mathcal{F}\varphi(\xi))d\mu_{\widehat{G}}(\xi)=\int_G(\varphi(t),\varphi(t))d\mu_G(t)=(\varphi,\varphi)_{L_2(G,X)},
$$

which can be easily checked for $\varphi \in L_2(G) \otimes X$ by means of axioms for inner product, scalar Steklov– Parseval's equality and cross-norm's property

$$
(\varphi_1 \otimes x_1, \varphi_2 \otimes x_2)_{L_2(G,X)} = (\varphi_1, \varphi_2)_{L_2(G)} \cdot (x_1, x_2)_X.
$$

Then we have on a dense subspace

$$
||\mathcal{F}\varphi||_{L_2(\widehat{G},X)} \leq C(\mathcal{F}\varphi,\mathcal{F}\varphi)_{L_2(\widehat{G},X)} = C(\varphi,\varphi)_{L_2(G,X)} \leq C^2 ||\varphi||_{L_2(G,X)},
$$

so we can extend F by continuity onto the whole $L_2(G, X)$.

If G is a finite group, then space $L_2(G, X)$ is isomorphic to the finite Cartesian product X^G . Pontryagin's dual group \hat{G} is isomorphic to G itself. Self-dual Haar measure possesses the property $\mu_G(G) = \sqrt{|G|}$. Fourier transform, also known as discrete Fourier transform, becomes

$$
(\mathcal{F}\varphi)(\xi) = \frac{1}{\sqrt{|G|}} \sum_{t \in G} \langle \xi, t \rangle \varphi(t).
$$

Theorem 3 If G is a finite group, then Fourier transform $\mathcal{F}: L_2(G,X) \to L_2(\widehat{G},X)$ is bounded for any Banach space X.

Proof. It follows from inequality

$$
||\mathcal{F}\varphi||_{L_2(\widehat{G},X)}^2 = \sum_{\xi \in \widehat{G}} \left\| \frac{1}{\sqrt{|G|}} \sum_{t \in G} \langle \xi, t \rangle \varphi(t) \right\|_X^2 \le \frac{1}{|G|} \sum_{\xi \in \widehat{G}} \left(\sum_{t \in G} |\langle \xi, t \rangle| \cdot \|\varphi(t)\|_X \right)^2 =
$$

=
$$
\frac{1}{|G|} \sum_{\xi \in \widehat{G}} \left(\sum_{t \in G} \|\varphi(t)\|_X \right)^2 = \left(\sum_{t \in G} \|\varphi(t)\|_X \right)^2 \le |G| \sum_{t \in G} \|\varphi(t)\|_X^2 = |G| \cdot \|\varphi\|_{L_2(G,X)}^2.
$$

◭

Now pass to infinite groups.

Lemma 2 Let group G contain \mathbb{R} -component, \mathbb{T} -component or \mathbb{Z} -component. Then boundedness of Fourier transform

$$
\mathcal{F}: L_2(G, X) \to L_2(\widehat{G}, X)
$$

implies isomorphism of Banach space X to a Hilbert one.

Proof. Consider the case, when group G contains **R**-component. There are open compactly generated subgroup H in G and compact subgroup $K \subset H$ such that $H/K \cong \mathbb{R}^a \times \mathbb{T}^b \times \mathbb{Z}^c \times F$, where $a \geq 1$.

Let $\tau_1 : H \to H/K$ be a canonical projection, $\tau_2 : H/K \to \mathbf{R}$ be the projection on the first coordinate of \mathbf{R}^a , $\tau = \tau_2 \circ \tau_1$.

Consider helper functions $\psi_{\mathbf{R},i} \in L_2(\mathbf{R}), 2 \leq i \leq a, \ \psi_{\mathbf{T},j} \in L_2(T), 1 \leq j \leq b, \ \psi_{\mathbf{Z},k} \in L_2(\mathbf{Z}),$ $1 \leq k \leq c$, $\psi_F \in L_2(F)$, each of them having norm equal to 1 in corresponding space. Consider injection $J: L_2(\mathbf{R}, X) \to L_2(H, X)$,

$$
J: \varphi \mapsto \left((\varphi \circ \tau) \otimes \Big(\bigotimes_{i=2}^a \psi_{\mathbf{R},i}\Big) \otimes \Big(\bigotimes_{j=1}^b \psi_{\mathbf{T},j}\Big) \otimes \Big(\bigotimes_{k=1}^c \psi_{\mathbf{Z},k}\Big) \otimes \psi_F\right).
$$

It is easy to see that the injection J is isometric. There exists a unique injection $\hat{J}: L_2(\mathbf{R}, X) \to$ $L_2(\widehat{H}, X)$, which is also an isometry, and for which the following diagram is commutative

$$
L_2(\mathbf{R}, X) \xrightarrow{\mathcal{F}_{\mathbf{R}}} L_2(\mathbf{R}, X)
$$

$$
J \downarrow \qquad \downarrow \widehat{J}
$$

$$
L_2(H, X) \xrightarrow{\mathcal{F}_H} L_2(\widehat{H}, X).
$$

Space $L_2(H, X)$ can be identified with a closed subspace of $L_2(G, X)$ consisting of functions, that are 0 almost everywhere outside H. Space $L_2(\hat{H}, X)$ can be identified with a closed subspace of $L_2(\hat{G}, X)$ consisting of functions, which are constant on cosets of H^{\perp} (recall, that $\hat{H} \cong \hat{G}/H_G^{\perp}$).

Fourier transform \mathcal{F}_H is the restriction of \mathcal{F}_G and thus, bounded. Fourier transform $\mathcal{F}_{\mathbf{R}} =$ $(\tilde{J})^{-1}\mathcal{F}_HJ$ is continuous as a composition of continuous maps. By Theorem [1](#page-0-0) statement 4) space X is isomorphic to a Hilbert one.

Cases when group G contains **T**-component or **Z**-component are considered similarly.

Consider the case, when group G does not contain \mathbb{R}_2 . \mathbb{Z}_2 or T-elements. Then all compactly generated subgroups $H \subset G$ are projective limits of finite subgroups with discrete topology, i.e. they are profinite groups.

Profinite groups are characterized by the following lemma [\[9\]](#page-9-4).

Lemma 3 Topological group H is a profinite one if and only if it

a) possesses Hausdorff 's property;

b) is compact;

c) is totally disconnected, i.e. for any two points $x, y \in H$ there exists subset $U \subset H$ that is both open and closed, such that $x \in U$ and $y \notin U$.

Any profinite group H is either discrete (and finite by virtue of compactness) or non-discrete (and therefore infinite).

Consider non-discrete profinite group H. We normalize Haar measure on H with $\mu_H(H) = 1$.

Group H is a Lebesgue space, i.e. it is isomorphic as a space with measure to segment $[0,1]$ with length [\[5,](#page-9-1) [8\]](#page-9-5). This fact can be proved selecting sequence of compact subgroups $K_n \subset H$ such that $K_1 \subset K_2 \subset \dots$ and cardinality of quotient groups $M_n := |H/K_n|$ tends to $+\infty$. If one numbers cosets of K_n properly, he becomes able to identify them with the intervals of length $1/M_n$ in [0, 1] up to a subsets of zero measure. By τ denote this isomorphism of spaces with measure.

A system of functions $(r_i)_{i=1,2,...}$, similar to the Rademacher's system on [0, 1] can be constructed on group H. This is an orthogonal system of functions taking values $\{+1, -1\}$ on subsets of measure $\frac{1}{2}$. Saying in probability-theoretical language functions r_i are realizations of independent random variables taking values $\{+1, -1\}$ with probability $\frac{1}{2}$. One can simply assume $r_i = r_i^{\infty} \circ \tau$, where $r_i^{\infty}(t) = \sin 2^i \pi t$ on [0, 1] is the usual Rademacher's functions.

We need a criterion, which is proved in [\[3\]](#page-8-1).

Theorem 4 The following statements are equivalent:

1) Banach space X is isomorphic to a Hilbert one.

2) There exists constant $C > 0$ such that for any finite collection of vectors $x_1, x_2, ..., x_n \in X$ two-sided Khinchin's type inequality holds

$$
C^{-1} \sum_{i=1}^n \|x_i\|^2 \le \mathcal{E} \Big\| \sum_{i=1}^n r_i x_i \Big\|^2 = \int_H \Big\| \sum_{i=1}^n r_i(t) x_i \Big\|^2 dt \le C \sum_{i=1}^n \|x_i\|^2,
$$

where r_i are independent random variables taking values $\{+1, -1\}$ with probability $\frac{1}{2}$, and symbol E denotes expectation.

As in [\[3\]](#page-8-1) we formulate a Lemma, which shows importance of the system (r_i) on H and allows us to consider arbitrary basis in $L_2(H)$ instead of (r_i) . By dt denote Haar measure on H.

Lemma 4 Let X be a Banach space, (f_i) be an orthonormal complete system in $L_2(H)$. Assume that for some $C > 0$ and for any collection $x_1, x_2, \ldots, x_n \in X$ there is inequality

$$
\int_H \Big\|\sum_{i=1}^n f_i(t)x_i\Big\|^2 dt \le C \sum_{i=1}^n \|x_i\|^2 \quad \left(resp., \ C^{-1}\sum_{i=1}^n \|x_i\|^2 \le \int_H \Big\|\sum_{i=1}^n f_i(t)x_i\Big\|^2 dt \right).
$$

Then for the same constant $C > 0$ and for any collection $x_1, x_2, \ldots, x_n \in X$ one also has

$$
\int_H \Big\| \sum_{i=1}^n r_i(t) x_i \Big\|^2 dt \le C \sum_{i=1}^n \|x_i\|^2 \quad \left(resp., \ C^{-1} \sum_{i=1}^n \|x_i\|^2 \le \int_H \Big\| \sum_{i=1}^n f_i(t) x_i \Big\|^2 dt \right).
$$

Proof. As Rademacher's system (r_i) is orthonormal and (f_k) are complete, we can find for a given $\varepsilon > 0$ an increasing sequences of indices (k_j) , (m_j) and orthonormal sequence (h_j) such that

$$
h_j = \sum_{k=k_j}^{k_{j+1}-1} (h_j, f_k) \cdot f_k, \quad \int_H |h_j(t) - r_{m_j}(t)|^2 dt < \frac{\varepsilon}{2^j}.
$$

For a fixed n and fixed $x_1, x_2, \ldots, x_n \in X$ we have

$$
\int_{H} \Big\| \sum_{i=1}^{n} r_i(t) \cdot x_i \Big\|^2 dt = \int_{H} \Big\| \sum_{i=1}^{n} r_{m_i}(t) \cdot x_i \Big\|^2 dt.
$$

By the triangle inequality

$$
\left(\int_H \left\|\sum_{j=1}^n r_{m_j}(t) \cdot x_j\right\|^2 dt\right)^{1/2} \le \left(\int_H \left\|\sum_{j=1}^n (r_{m_j}(t) - h_j(t)) \cdot x_j\right\|^2 dt\right)^{1/2} + \left(\int_H \left\|\sum_{j=1}^n h_j(t) \cdot x_j\right\|^2 dt\right)^{1/2} \le \sqrt{\varepsilon} \left(\sum_{j=1}^n \|x_j\|^2\right)^{1/2} + \left(\int_H \left\|\sum_{j=1}^n h_j(t) \cdot x_j\right\|^2 dt\right)^{1/2}.
$$

As
$$
1 = ||h_j||^2 = \sum_{k=k_j}^{k_{j+1}-1} |(h_j, f_k)|^2
$$
, we get
\n
$$
\int_H \Big\|\sum_{j=1}^n h_j(t) \cdot x_j\Big\|^2 dt = \int_H \Big\|\sum_{j=1}^n \Big(\sum_{k=k_j}^{k_{j+1}-1} (h_j, f_k) \cdot f_k\Big) x_j\Big\|^2 dt \le
$$
\n
$$
\le C \sum_{j=1}^n \sum_{k=k_j}^{k_{j+1}-1} |(h_j, f_k)|^2 \|x_j\|^2 = C \sum_{j=1}^n \|x_j\|^2.
$$

Thus,

$$
\int_{H} \Big\| \sum_{i=1}^{n} r_i(t) \cdot x_i \Big\|^2 dt \le (\sqrt{\varepsilon} + \sqrt{C})^2 \sum_{i=1}^{n} \|x_i\|^2.
$$

As ε is arbitrary, we obtain the desired inequality. Proof in the case of reverse type inequality is analogous. \blacktriangleleft

Corollary 4.1 Let X be a Banach space, and (f_i) be a complete orthonormal system in $L_2(H)$. Space X is linearly isomorphic to a Hilbert one if and only if there exists constant $C > 0$ such that for any set of vectors $x_1, x_2, \ldots, x_n \in X$ one has

$$
C^{-1} \sum_{i=1}^{n} ||x_i||^2 \le \int_H \Big\|\sum_{i=1}^{n} f_i(t) \cdot x_i\Big\|^2 dt \le C \sum_{i=1}^{n} ||x_i||^2.
$$

In Corollary [4.1](#page-6-0) isomorphism of X to a Hilbert space follows from *two-sided* inequality. Knowledge of profinite groups' structure allows us to switch from lower estumate to upper one and vice versa as shown below.

First recall that Bruhat–Schwartz space on a profinite group H and on dual discrete \hat{H} consists of locally constant functions with compact support. Spaces $\mathcal{S}(H)$ and $\mathcal{S}(H)$ are inductive limit of finite-dimensional spaces and carry the strongest locally convex topology [\[7\]](#page-9-3).

Now we are going to study the case of vector-valued Fourier transform on a profinite non-discrete group H. As H is a compact infinite group, its dual \hat{H} is a discrete infinite group.

Lemma 5 Let X be a Banach space, and H be a profinite non-discrete group. The following statements are equivalent:

1) X is linearly isomorphic to a Hilbert space.

2) There exists constant $C > 0$ such that for any set of vectors $x_1, ..., x_n \in X$ and characters $\xi_1, ..., \xi_n \in \widehat{H}$ one has

$$
\int_{H} \Big\| \sum_{k=1}^{n} \langle \xi_k, t \rangle x_k \Big\|^2 dt \le C \sum_{k=1}^{n} \|x_k\|^2.
$$

2)' Fourier transform $\mathcal{F}_{\widehat{H}} : L_2(\widehat{H}, X) \to L_2(H, X)$ and inverse Fourier transform $F_H^{-1} = \mathcal{I}_H \mathcal{F}_{\widehat{H}}$ are bounded. Here \mathcal{I}_H is an isometrical operator of changing variable $x(t) \mapsto x(-t)$.

3) There exists constat $C > 0$ such that for any set of vectors $x_1, ..., x_n \in X$ and characters $\xi_1, ..., \xi_n \in \widehat{H}$ one has

$$
C^{-1} \sum_{k=1}^{n} \|x_k\|^2 \le \int_H \Bigl\| \sum_{k=1}^{n} \langle \xi_k, t \rangle x_k \Bigr\|^2 dt.
$$

3)' Inverse Fourier transform $\mathcal{F}_{\widehat{H}}^{-1}: L_2(H, X) \to L_2(\widehat{H}, X)$ and Fourier transform $\mathcal{F}_H = \mathcal{I}_H \mathcal{F}_{\widehat{H}}^{-1}$ are bounded.

Proof. Lemma [1](#page-3-0) yields implications $1) \Rightarrow 2)'$, $1) \Rightarrow 3)'$.

To prove equivalences $2' \Leftrightarrow 2$, $3' \Rightarrow 3$ it's enough to consider

$$
h = \sum_{k=1}^{n} I_{\{\xi_k\}} \cdot x_k \in \mathcal{S}(\widehat{H}) \otimes X,
$$

where $\xi_k \in \widehat{H}$, $x_k \in X$, $n \in \mathbb{N}$. By definition of Fourier transform $\mathcal{F} : L_2(\widehat{H}, X) \to L_2(H, X)$ and by chosen normalization of Haar measure on H we get

$$
||h||_{L_2(\widehat{H},X)}^2 = \sum_{k=1}^n ||x_k||^2, \quad ||\mathcal{F}h||_{L_2(H,X)}^2 = \int_H \Big|\Big|\sum_{k=1}^n \langle \xi_k, t \rangle x_k\Big|\Big|^2 dt.
$$

Equivalence follows from the density of $\mathcal{S}(\widehat{H}) \otimes X$ in $L_2(\widehat{H}, X)$, density of $\mathcal{S}(H) \otimes X$ in $L_2(H, X)$, and bijectivity of $\mathcal{F}_{\hat{H}}$, \mathcal{F}_{H} in the corresponding spaces.

By Corollary [4.1](#page-6-0) one also has 2) $\& 3) \Rightarrow 1$.

Now assume boundedness condition 2)'. Let's look at Fourier transform \mathcal{F}_H on a subspace $\mathcal{S}(H) \otimes$ X. For this consider arbitrary compact subgroup $K \subset H$, for which $|H/K| < +\infty$. We identify functions, which are constant on the cosets of K, with elements of $\mathcal{S}(H/K) \otimes X$.

By finiteness of H/K there exist an isomorphism $\alpha : \widehat{H/K} \to H/K$. Adjoint isomorphism $\alpha^* : \widehat{H/K} \to H/K$ is defined by

$$
\langle \xi_1, \alpha(\xi_2) \rangle_H = \langle \alpha^*(\xi_1), \xi_2 \rangle_{\widehat{H}}.
$$

Consider operator

$$
R_{\alpha}: \mathcal{S}(H/K) \otimes X \to \mathcal{S}(\widehat{H/K}) \otimes X : (R_{\alpha}\psi)(\xi') = \psi(\alpha(\xi')) \cdot |H/K|^{-\frac{1}{2}}.
$$

It's an isometry, because

$$
||R_{\alpha}\psi||^{2} = \sum_{\xi' \in \widehat{H/K}} ||R_{\alpha}\psi(\xi)||^{2} \mu_{\widehat{H/K}}(\xi) = \sum_{\xi' \in \widehat{H/K}} ||\psi(\alpha(\xi'))||^{2} \cdot |H/K|^{-\frac{1}{2}\cdot 2} =
$$

=
$$
[t := \alpha(\xi')] = \sum_{t \in H/K} ||\psi(t)||^{2} \cdot |H/K|^{-1} = \sum_{t \in H/K} ||\psi(t)||^{2} \mu_{H/K}(t) = ||\psi||^{2}.
$$

Now one has

$$
(\mathcal{F}_{\hat{H}}R_{\alpha}\psi)(t') = \sum_{\xi' \in \widehat{H/K}} \langle t', \xi' \rangle_{\widehat{H/K}} (\psi(\alpha(\xi'))|H/K|^{-\frac{1}{2}}),
$$

$$
(R_{(\alpha^*)}\mathcal{F}_{\hat{H}}R_{\alpha}\psi)(\xi) = \Big(\sum_{\xi' \in \widehat{H/K}} \langle \alpha^*(\xi), \xi' \rangle_{\widehat{H/K}} \psi(\alpha(\xi'))|H/K|^{-\frac{1}{2}}\Big)|H/K|^{-\frac{1}{2}} = [t := \alpha(\xi')] =
$$

$$
= \Big(\sum_{t \in H/K} \langle \alpha^*(\xi), \alpha^{-1}(t) \rangle_{\widehat{H/K}} \psi(t)\Big)|H/K|^{-1} = \sum_{t \in H/K} \langle \alpha^*(\xi), \alpha(\alpha^{-1}(t)) \rangle_{H/K} \psi(t)|H/K|^{-1} =
$$

$$
= \sum_{t \in H/K} \langle \xi, t \rangle_{H/K} \psi(t)\mu_{H/K}(t) = (\mathcal{F}_H \psi)(\xi).
$$

Thus, restriction of \mathcal{F}_H onto $\mathcal{S}(H/K)\otimes X$ has the same norm as $\mathcal{F}_{\widehat{H}}$ does. As K is arbitrary, it implies continuity of \mathcal{F}_H on $L_2(H, X)$, and implication $2 \Rightarrow 3$) is true. Implication $3 \Rightarrow 2$) can be proved in the same way.

Now one has enough implications to see the equivalence of all statements.

Let's pass to the general case of Fourier transform on a locally compact Abelian group G. This is our main result.

Theorem 5 Let X be a Banach space, let G be a locally compact Abelian group. Space X is linearly isomorphic to a Hilbert one if and only if Fourier transform

$$
\mathcal{F}: L_2(G,X) \to L_2(G,X)
$$

is bounded.

Proof. Existence of isomorphism is a sufficient condition for boundedness of $\mathcal F$ as shown in Lemma [1.](#page-3-0)

Now assume that Fourier transform is bounded.

If G contains \mathbb{R} -, \mathbb{T} - or Z-component, then isomorphism of X to a Hilbert space follows from Lemma [2,](#page-3-1) and we are done.

Otherwise, group G does not contain \mathbb{R} , \mathbb{T} or \mathbb{Z} -components. In this case all compactly generated open subgroups $H \subset G$ are profinite.

If some of these H is non-discrete, then we consider space $L_2(H, X)$ as a closed subspace in $L_2(G, X)$ (one can simply assume that function from $L_2(H, X)$ are zero outside H). We identify $L_2(H, X)$ with a subspace of $L_2(G, X)$ consisting of functions constant on cosets of annihilator $H_G^{\perp} \subset$ G. Fourier transform on $L_2(H, X)$ is the restriction of Fourier transform from $L_2(G, X)$ and is bounded. Isomorphism of X to a Hilbert space follows from Lemma [5,](#page-6-1) statement 3)'. And we are done.

If all subgroups $H \subset G$ considered are discrete (their compactness implies finiteness), then by Theorem [2](#page-2-0) group G is an *inductive* limit of discrete subgroups. By properties of Pontryagin duality (in the language of Cathegory theory one can say that passing to a dual group is an exact functor) dual group \widehat{G} is a projective limit of \widehat{H} . Groups \widehat{H} are dual to finite discrete H. Thus, \widehat{H} are isomorphic H, and are finite discrete themselves. Group \widehat{G} is profinite. As G is infinite, \widehat{G} is non-discrete.

Boundedness of Fourier transform $\mathcal{F}: L_2(G, X) \to L_2(\widehat{G}, X)$ is equivalent to boundedness of inverse Fourier transform $\mathcal{F}_{\widehat{G}}^{-1}$ on profinite non-discrete \widehat{G} . Isomorphism of X to a Hilbert space follows from Lemma [5](#page-6-1) statement 2 . \mathcal{A} , which is a set of \mathcal{A} , we have a set of \mathcal{A}

References

- $[1]$ Peetre J. Sur la transformation de Fourier des fonctions à valeurs vectorielles // Rend. Sem. Mat. Univ. Padova. – 1969, vol.42. – p.15–26.
- [2] Mikusinski J. The Bochner integral. – Acad. Press, 1978.
- [3] Kwapien S. Isomorphic characterizations of inner product spaces by orthogonal series with vectorvalued coefficients // Studia mathematica. – 1972, vol.XLIV. – p.583–595.
- [4] Radyna Ya.M., Radyno Ya.V., Sidorik A.G. Characterizing Hilbert spaces using Fourier transform over the field of p-adic numbers $//$ Dokl. Nats. Akad. Nauk Belarusi. – 2007, vol.51, No.5. – p. 17–22 (in Russian), [arXiv:0803.3646v](http://arxiv.org/abs/0803.3646)1 [math.FA].
- [5] Hewitt E., Ross K.A., Abstract Harmonic Analysis. – Vol.1. – Springer-Verlag, 1979.
- [6] Morris S. Pontryagin duality and the structure of locally compact abelian groups. – Cambridge University Press, 1977.
- [7] Bruchat F. Distributions sur un groupe localement compact et applications `a l'´etude des representations des groupes p-adiques // Bull. Soc. Math. France. – 1961, vol.89. – p.43–75.
- [8] Leonov N.N. Mathematical sociology. Sructural approximation approach. – Minsk, 2002 (in Russian).
- [9] Lenstra H. Profinite groups //<http://websites.math.leidenuniv.nl/algebra/Lenstra-Profinite.pdf>