

Fidelity approach to the disordered quantum XY model

Silvano Garnerone,* N. Tobias Jacobson, Stephan Haas, and Paolo Zanardi†

Department of Physics and Astronomy, University of Southern California, Los Angeles, CA 90089

(Dated: February 6, 2020)

We study the random XY spin chain in a transverse field by analyzing the susceptibility of the ground state fidelity, numerically evaluated through a standard mapping of the model onto quasi-free fermions. It is found that the fidelity susceptibility and its scaling properties provide useful information about the phase diagram. In particular it is possible to determine the Ising critical line and the Griffiths phase regions, in agreement with previous analytical and numerical results.

Introduction.— In the last few years concepts borrowed from quantum information theory have proven useful in characterizing the critical behavior of quantum many-body systems [1]. In particular, a geometric approach to the study of quantum phase transitions (QPTs), i.e. the fidelity analysis, has been shown to be an effective way of characterizing distinct phases of quantum systems [2, 3, 4, 5, 6, 7, 8]. Previously, the fidelity approach has been applied to a variety of homogeneous systems. In this work we extend these studies to disordered quantum systems. Specifically, we investigate the behavior of the fidelity susceptibility of the disordered XY model in a transverse field. It is well known that the presence of quenched disorder can have drastic effects on the critical properties of a quantum system. The appearance of new universality classes and novel states of matter such as the Griffiths phase are two important examples [9, 10, 11, 12]. The aim of the present work is to show what can be inferred about the physics of the disordered quantum system from the properties of the fidelity susceptibility.

The Hamiltonian of the disordered XY chain is given by

$$H = - \sum_{i=0}^{L-1} \left(\frac{1 + \gamma_i}{2} \sigma_i^x \sigma_{i+1}^x + \frac{1 - \gamma_i}{2} \sigma_i^y \sigma_{i+1}^y + \lambda_i \sigma_i^z \right), \quad (1)$$

where $\sigma_i^{\{x,y,z\}}$ are Pauli spin matrices, and γ_i and λ_i are sets of independent random coupling and field variables with distributions $\pi(\gamma_i)$ and $\rho(\lambda_i)$. Note that due to gauge symmetry the Hamiltonian (1) can be chosen to have only positive couplings and fields. This model can be mapped onto a system of quasi-free fermions with periodic boundary conditions, and an exact expression for the fidelity susceptibility is obtained which depends explicitly on the random parameters characterizing the ground state of the system. In this work, we investigate the statistical properties of the fidelity susceptibility for relevant regions of parameter space.

Gaussian distributions are used for the random vari-

ables,

$$\pi(x_i) = \rho(x_i) = \frac{1}{\sigma\sqrt{2\pi}} \exp \left\{ -\frac{1}{2} \left(\frac{x_i - x}{\sigma} \right)^2 \right\}, \quad (2)$$

where x_i is either the field or the coupling at position i on the lattice, x is the respective average value and σ is the variance.

Previous Results.— The pure XY chain has been analytically solved in [13]. In the absence of disorder two different quantum phase transitions are present. Following the standard notation, we refer to the QPT driven by the transverse magnetic field λ as the *Ising transition*, and the QPT driven by the coupling parameter γ as the *anisotropy transition*. The Ising transition separates a ferromagnetic ordered phase from a paramagnetic quantum-disordered phase, whereas the anisotropy critical line is the boundary between a ferromagnet ordered along the x direction and a ferromagnet ordered along the y direction.

A major improvement in the understanding of the effect of disorder on the physics of quantum magnets has been achieved with the use of the strong-disorder renormalization group technique (SDRG) by Dasgupta and Ma [14], and further developed by Fisher [10, 11]. The correctness of this method has been corroborated both by numerics [15, 16] and analytic exact studies [17, 18]. In the work of McKenzie and Bunder [17, 18] the critical behavior of the disordered XY chain in a transverse field has been studied using a mapping to random-mass Dirac equations. The properties of the solutions of these equations imply the disappearance of the anisotropy transition in the presence of disorder. Furthermore, Griffiths phases are predicted to appear both around the Ising critical line and the anisotropy $\gamma = 0$ line.

For $\gamma = 1$ the XY random chain reduces to the random transverse-field Ising chain (RTFIC), which is a prototypical model for disordered quantum systems. Since it is representative of the universality class of Ising transitions for all values of γ , let us briefly review what is known for this model. The Hamiltonian of the RTFIC is $H = - \sum_{i=0}^{L-1} [J_i \sigma_i^x \sigma_{i+1}^x + h_i \sigma_i^z]$, where J_i and h_i are random couplings and fields respectively. The system is at criticality when the average value of the field equals the average value of the coupling. Using the SDRG one obtains that, at the quantum critical point, the time scale τ and the length scale L are related by $\ln \tau \sim L^{1/2}$.

*Electronic address: garneron@usc.edu

†Also at Institute for Scientific Interchange, Viale Settimio Severo 65, I-10133 Torino, Italy

This results in an infinite value for the dynamical exponent z at criticality. The distribution of the logarithm of the energy gap, $\ln \epsilon$, at criticality broadens with increasing system size, in accordance with the scaling relation $\ln \epsilon \sim -L^{1/2}$ [16]. In the vicinity of the critical point the distribution of relaxation times is broad due to Griffiths singularities. This region of the parameter space, the Griffiths phase, is characterized by a dynamical exponent z which depends on the distance from the critical point. This dependence is one of the hallmarks of the Griffiths phase. Another important consequence of the presence of disorder is that the decay of the correlation functions is characterized by two kinds of correlation lengths, *typical* and *average*, with two different scaling exponents [10, 11].

Method.— The main idea of the fidelity approach is to detect QPTs through enhanced orthogonalization rates between ground states $|\Psi(x)\rangle$ infinitesimally close in parameter space. The orthogonalization is signaled by a drop in the fidelity, $F(x, x + \delta x) \equiv |\langle \Psi(x) | \Psi(x + \delta x) \rangle|$, at the critical point. The fidelity susceptibility is a related quantity with a more transparent physical meaning [4, 5], and whose behavior is more suitable for numerical analysis. It is defined as

$$\chi(x) = \lim_{\delta x \rightarrow 0} \frac{-2 \ln F(x, x + \delta x)}{\delta x^2}. \quad (3)$$

In [5] it was shown that χ is related to the dynamic structure factor of the relevant operator associated with the transition. A generalization of this result, valid for the so-called *geometric tensor*, has been given in [4].

Previous works have characterized the pure XY spin chain using the fidelity approach [2, 3, 19, 20] and the quantum Chernoff bound [21]. The mapping of the spin model onto the quasi-free fermion Hamiltonian,

$$H = \sum_{i,j=1}^L c_i^\dagger A_{ij} c_j + \frac{1}{2} \sum_{i,j=1}^L \left(c_i^\dagger B_{ij} c_j^\dagger + H.c. \right), \quad (4)$$

yields an explicit BCS-like form for the ground state $|\Psi\rangle = \mathcal{N} \exp\left(\frac{1}{2} \sum_{j,k=1}^L c_j^\dagger G_{jk} c_k^\dagger\right) |0\rangle$, where \mathcal{N} is a normalization factor.

The fidelity of the ground states evaluated at slightly different parameter values (coupling or magnetic field) x and $x + \delta x$ has a simple analytical expression. Defining the matrix $Z(x) \equiv A(x) - B(x)$ and the unitary part of the polar decompositions of $Z(x)$ and $\tilde{Z} \equiv Z(x + \delta x)$ as $T(x)$ and $\tilde{T} \equiv T(x + \delta x)$, respectively, the fidelity can be written as

$$F(Z, \tilde{Z}) = \sqrt{\left| \det \frac{T + \tilde{T}}{2} \right|}. \quad (5)$$

Note that the matrix G defining the ground state is simply the Cayley transform of T [19].

In the following, we will use an alternative expression for the fidelity susceptibility,

$$\chi(x) = \frac{1}{8} \|\partial_x T\|_F^2, \quad (6)$$

with $\|\cdot\|_F$ the Frobenius norm. Eq. (6) is obtained from (5) via standard algebra. We have numerically evaluated the fidelity susceptibility using (6) for relevant regions of parameter space of the disordered XY model. The numerical analysis has been performed on two sets of system sizes, i.e. $\{128, 256, 512\}$, and $\{400, 410, \dots, 500\}$ in steps of 10. We have taken 50,000 disorder realizations for all sizes except for those larger than 400, in which case we used 10,000 realizations.

Results.— We consider the Hamiltonian (1), where the couplings γ_i and the transverse fields λ_i are independent random variables with Gaussian distributions centered around $\lambda \equiv [\lambda_i]_{\text{ave}}$ and $\gamma \equiv [\gamma_i]_{\text{ave}}$, both with variance $\sigma = 0.1$. $[\cdot]_{\text{ave}}$ denotes the arithmetic mean over the disorder realizations.

A scaling analysis has been performed using arguments first developed in [4]. Following that reference, we can express the fidelity susceptibility as an integral in imaginary time $\chi = \int_{-\infty}^{\infty} d\tau \tau G(\tau)$, where $G(\tau) = \theta(\tau) \ll \partial_x H(\tau) \partial_x H(0) \gg$ is the connected correlation function of the conjugate operator in the Hamiltonian associated with the driving parameter in the transition, and θ is the Heaviside step function. For example, in the case of the Ising transition we have $G(\tau) = \theta(\tau) \ll \sum_{i,j} \sigma_i^z(\tau) \sigma_j^z(0) \gg$. The average fidelity susceptibility can then be written as $[\chi]_{\text{ave}} = \int_{-\infty}^{\infty} d\tau \tau [G(\tau)]_{\text{ave}} \sim L^{\Delta_\chi}$, where the finite-size scaling exponent Δ_χ of $[\chi]_{\text{ave}}$ is given by $\Delta_\chi = 2z + 2 - 2\Delta_O$ [4]. Δ_O is the scaling exponent of the QPT conjugate operator ($\sum_i \sigma_i^z$ in the case of the transition driven by λ), and in general Δ_O depends on the parameters γ and λ .

For the XY chain without disorder, in the quantum critical region χ scales as $\chi \sim L^2$, whereas away from the critical region $\chi \sim L$. Since for finite system sizes the quantum critical region has a finite width, Δ_χ is 1 for all but a narrow range of λ , having a maximum of 2 at $\lambda = 1$.

With this disorder-free behavior in mind, we now study $[\chi]_{\text{ave}}$ about the Ising transition, driven by the coupling λ . In our numerical studies we have focused on the case of the RTFIC, where $\gamma = 1$. Qualitatively all of our results on the critical behavior of the fidelity susceptibility hold true for other values of γ , since the universality class of the model does not change in the range $\gamma \in (0, 1]$.

Fig. 1(a) shows $[\chi]_{\text{ave}}$ as a function of λ . The averaged fidelity susceptibility displays a local maximum at the Ising critical point, $\lambda = 1$. Fig. 1(b) shows Δ_χ , the finite-size scaling exponent of $[\chi]_{\text{ave}}$, for different values of the transverse field λ . Note that far from the Ising critical point $[\chi]_{\text{ave}}$ scales strictly extensively, while in the vicinity of the critical point the scaling becomes superextensive and approximately quadratic in L at $\lambda = 1$. This scaling behavior is quite similar to that of the pure case, save for a slight broadening in the peak of Δ_χ . The presence of the Ising critical point and its associated quantum critical region makes it difficult to identify the effects on the finite-size scaling of χ due to the Griffiths phase.




FIG. 1: Ising transition at $\gamma = 1$. (a) Average fidelity susceptibility, $[\chi]_{\text{ave}}$, for $L = 512$ and 10^4 realizations, (b) the associated finite-size scaling exponent Δ_χ of $[\chi]_{\text{ave}}$ [22], (c) probability distribution of $\ln \chi$ at the Ising transition, (d) distribution of $\ln \chi$ away from the Ising transition.

However, in Fig. 1(c) and (d) we plot the distribution of the fidelity susceptibility over many realizations at the Ising critical point and away from it. We choose to plot the distribution of $\ln \chi$ instead of χ itself because, in analogy to other physical quantities, the presence of disorder greatly broadens the distribution. Looking at the scale on the x-axes, note that the probability density function of $\ln \chi$ away from criticality is much more sharply peaked than at criticality, and its shape is more symmetric away from the critical point. Here, the increased breadth of the fidelity susceptibility distribution may signal the Griffiths phase where the scaling fails to distinguish clearly the effect of disorder.

One characteristic of disordered quantum systems is the difference between typical and average quantities. We define the typical susceptibility as $\chi_{\text{typ}} \equiv 2^{\lfloor \log_2 \chi \rfloor_{\text{ave}}}$, which is equivalent to the geometric mean of χ over the different realizations of disorder. Qualitatively, the geometric mean weighs outlier values less than the arithmetic mean, providing a measure of the most representative values. For averages over positive quantities the arithmetic mean provides an upper bound for the geometric mean.

In Fig. 2(a) the difference between the average and typical fidelity susceptibilities, $([\chi]_{\text{ave}} - \chi_{\text{typ}})/\chi_{\text{typ}}$, near the Ising transition is shown, i.e. with $\chi(\lambda, \gamma = 1)$. Away from the critical point these two quantities cannot be distinguished within the system sizes and disorder strength we have considered. However, approaching the Ising critical point the more broadly spread fidelity susceptibility distribution results in a significant difference between the two. The irregular behavior of this quantity near the critical point is due to rare but large susceptibility realizations which have a stronger effect on the arithmetic mean than on the geometric mean.

Fig. 2(b) shows the difference between average and typical susceptibilities near the anisotropy line, with $\chi(\lambda = 0.2, \gamma)$. Notice that the location of the maximum of this difference coincides with the maximum of $[\chi]_{\text{ave}}$, shown in Fig. 3(a). Also, the local minimum at $\gamma = 0$ coincides with the local minimum of $[\chi]_{\text{ave}}$. Moreover, for values of γ near the anisotropy line the distribution of $\ln \chi$ broadens, as shown in Fig. 3(c).




FIG. 2: Difference between average and typical values, $\frac{[\chi]_{\text{ave}} - \chi_{\text{typ}}}{\chi_{\text{typ}}}$, of χ near the Ising transition (a), for $\gamma = 1$ and $L = 512$, and near the anisotropy line (b), for $\lambda = 0.2$ and $L = 500$.




FIG. 3: Elimination of anisotropy transition at $\gamma = 0$ due to disorder. (a) Average fidelity susceptibility $[\chi]_{\text{ave}}$, near $\gamma = 0$ for $L = 500$ and 10^4 realizations. (b) Finite-size scaling exponent of $[\chi]_{\text{ave}}$, (c) probability distribution of $\ln \chi$ at $\gamma = 0$ and at the value of γ corresponding to the maximum of $[\chi]_{\text{ave}}$, (d) distribution of $\ln \chi$ far away from the anisotropy line.

Although for the disordered XY model the line $\gamma = 0$ is not critical as it is in the pure case [17, 18], the presence of Griffiths singularities still has highly non-trivial effects on the fidelity susceptibility in the vicinity of $\gamma = 0$, as shown in Fig. 3(a). Specifically, in the presence of dis-

order the peak in $[\chi]_{\text{ave}}(\gamma)$ is shifted away from its original position in the pure limit, i.e. away from $\gamma = 0$. Note that $\gamma = 0$ is a special case for the XY chain, which can be mapped onto free fermions via the Jordan-Wigner transformation. Using renormalization group arguments, it was shown that the introduction of even infinitesimal disorder in such systems leads to Anderson localization in dimensions lower than $D=3$ [23]. Accordingly, the observed dip in $[\chi]_{\text{ave}}(\gamma)$ at $\gamma = 0$ indicates that the Griffiths regime in the vicinity of the critical anisotropy line of the pure XY chain is interrupted by an Anderson insulator which occurs at $\gamma = 0$. This is further corroborated by the non-monotonic dependence on γ of the associated scaling exponent shown in Fig. 3(b). At $\gamma = 0$, one finds $\Delta_\chi = 1$, whereas in the interval $0 < \gamma < 0.075$ the exponent $\Delta_\chi(\gamma)$ exhibits a non-universal dependence on the driving parameter γ , indicating the presence of a Griffiths regime. Note that the observed maximum is not to be seen as an indication of a QPT. Rather, it originates from the competition between the scaling properties of χ in the Griffiths phase and at the $\gamma = 0$ line.

To complete the discussion of the effects of disorder on the anisotropy transition, let us comment on the probability distributions of $\ln \chi$. In Figs. 3(c) and (d), we show $P(\ln \chi)$ at $\gamma = 0$, at the midpoint of the Griffiths regime where $\chi(\gamma)$ and $\Delta_\chi(\gamma)$ peak, and at a point far away from the anisotropy line. In analogy to the Ising transition, the probability distribution function in the Griffiths regime is broad and asymmetric, whereas far away from it

its shape is symmetric and its distribution is much more narrow. Moreover, at the Anderson localization line, it is more narrow than in the Griffiths regime, indicating that singularities due to rare events are less relevant at $\gamma = 0$.

Conclusions.— In this work we have applied the fidelity approach to the study of the disordered XY chain in an external magnetic field. We have found that the fidelity susceptibility may be able to provide a detailed phase diagram for this model. In the parameter region around the $\gamma = 0$ line the scaling analysis of the fidelity susceptibility shows the disappearance of the QPT and the emergence of a Griffiths phase, in accordance with previous analytical and numerical results. In the case of the Ising transition it is difficult from a numerical point of view to unambiguously distinguish the Griffiths phase from the extended critical region. We plan to further investigate the relevance of disorder on the fidelity susceptibility in future works. Other aspects that will be studied with more details are the extent of the Griffiths phase together with its dependence on disorder strength and the probability distribution of disorder.

We thank H. Saleur and L. Campos Venuti for helpful discussions. Computation for the work described in this paper was supported by the University of Southern California Center for High Performance Computing and Communications. We acknowledge financial support by the National Science Foundation under grant DMR-0804914.

-
- [1] L. Amico, R. Fazio, A. Osterloh and V. Vedral, Rev. Mod. Phys. **80**, 517 (2008)
 - [2] P. Zanardi and N. Paunkovic, Phys. Rev. E **74**, 031123 (2006)
 - [3] P. Zanardi, P. Giorda and M. Cozzini, Phys. Rev. Lett. **99**, 100603 (2007)
 - [4] L. Campos Venuti and P. Zanardi, Phys. Rev. Lett. **99**, 095701 (2007)
 - [5] W. L. You, Y. W. Li and S. J. Gu, Phys. Rev. E **76**, 022101 (2007)
 - [6] H. -Q. Zhou and J. P. Barjaktarevic, arXiv:cond-mat/0701608
 - [7] H. -Q. Zhou, J.-H. Zhao and B. Li, arXiv:0704.2940
 - [8] H. -Q. Zhou, arXiv:0704.2945
 - [9] S. Sachdev, Quantum Phase Transitions (Cambridge University Press, 1999)
 - [10] D. S. Fisher, Phys. Rev. Lett. **69**, 534 (1992)
 - [11] D. S. Fisher, Phys. Rev. B **51**, 6411 (1995)
 - [12] R. B. Griffiths, Phys. Rev. Lett **23**, 17 (1969)
 - [13] E. Lieb, T. Schultz and D. Mattis, Ann.Phys. **16**, 407 (1961)
 - [14] C. Dasgupta and S.K. Ma, Phys. Rev. B **22**, 1305 (1980)
 - [15] S. Haas, J. Riera and E. Dagotto, Phys. Rev. B **48**, 13174 (1993)
 - [16] A. P. Young and H. Rieger, Phys. Rev. B **53**, 8486 (1996)
 - [17] R. H. McKenzie, Phys. Rev. Lett. **77**, 4804 (1996)
 - [18] J. E. Bunder and R. H. McKenzie, Phys. Rev. B **60**, 344 (1999)
 - [19] M. Cozzini, P. Giorda and P. Zanardi, Phys. Rev. B **75**, 014439 (2007)
 - [20] P. Zanardi, M. Cozzini and P. Giorda, J. Stat. Mech. L02002 (2007)
 - [21] D. F. Abasto, N. T. Jacobson, and P. Zanardi, Phys. Rev. A **77**, 022327 (2008)
 - [22] This scaling analysis considered system sizes $L = 128, 256, \text{ and } 512$. Varying finite-size critical points were taken into account by translating the splines of $[\chi]_{\text{ave}}$ for each size along λ in order to give strictly extensive scaling far from criticality.
 - [23] E. Abrahams, P.W. Anderson, D.C. Licciardello, and T.V. Ramakrishnan, Phys. Rev. Lett. **42**, 673 (1979).