

Electromagnetic Light by Action of Polarized Atoms ^3He .

Minasyan V.N.

February 4, 2019

Abstract

First, it is predicted that polarized atoms ^3He increases a value of speed reflected electromagnetic waves. Thus, we prove that velocity of electromagnetic waves into gas consisting of polarized atoms ^3He is rather than one in vacuum.

PACS:01.55. + *b* General physics

1. INTRODUCTION.

The motivation for theoretical study of the Fermi gas of atoms ${}^3\text{He}$ is an attempt at a microscopic understanding of quantization of electromagnetic field. In this letter, we investigate an interaction between electromagnetic field represented as nonideal Bose-gas of elementary Bose-particles of electromagnetic field [1] with Fermi gas of polarized atoms ${}^3\text{He}$. In this context, we show that value of speed of reflected electromagnetic waves is increased by action polarized atoms ${}^3\text{He}$.

Theoretical description of quantization local electromagnetic field in vacuum within a model of Bose-gas of local electromagnetic waves, which are propagated by speed c in vacuum, was first proposed by Dirac [2]. In this context, the vector potential for local electromagnetic field $\vec{A}(\vec{r}, t)$ is determined by following form:

$$\begin{aligned}\vec{A}(\vec{r}, t) &= \int \left(\vec{A}_{\vec{k}} \exp^{i(\vec{k}\vec{r}+kct)} + \vec{A}_{\vec{k}}^{\dagger} \exp^{-i(\vec{k}\vec{r}+kct)} \right) d^3k = \\ &= \sum_{\vec{k}} \left(\vec{A}_{\vec{k}} \exp^{i(\vec{k}\vec{r}+kct)} + \vec{A}_{\vec{k}}^{\dagger} \exp^{-i(\vec{k}\vec{r}+kct)} \right)\end{aligned}\quad (1)$$

where $\vec{A}_{\vec{k}}^{\dagger}$ and $\vec{A}_{\vec{k}}$ are, respectively, the Fourier components of vector potentials electromagnetic field, which determine a solution of wave-equation

$$\nabla^2 \vec{A} - \frac{1}{c^2} \frac{d^2 \vec{A}}{dt^2} = 0 \quad (2)$$

with condition

$$\text{div} \vec{A} = 0 \quad (3)$$

Further, we shall pursue a fresh approach for quantization of classic electromagnetic field, which considers the Fourier components of vector potentials electromagnetic field $\vec{A}_{\vec{k},t}^{\dagger}$ and $\vec{A}_{\vec{k},t}$ via the Bose-operators "creation" and "annihilation" of Bose-particles with spin one. The law conservation for total number bosons of electromagnetic field provides the creation of photons of electromagnetic field.

11. ANALYSIS.

As it is known, the Hamiltonian of radiation \hat{H}_R is determined as:

$$\hat{H}_R = \frac{1}{8\pi V} \int \left(E^2 + H^2 \right) dV \quad (4)$$

The solution of Maxwell equations leads to following formulas

$$\vec{H} = \text{curl}\vec{A} \quad (5)$$

and

$$\vec{E} = -\frac{1}{c} \frac{d\vec{A}}{dt} \quad (6)$$

At finding of Eq.(4), Dirac used of the unnecessary expression:

$$\text{div}[\vec{A} \times \vec{H}] = H^2 + \vec{A}\nabla^2\vec{A} \quad (7)$$

and introduced by handling an unnecessary condition as $\int \text{div}[\vec{A} \times \vec{H}]dV = 0$ which implies a presence of term as $\int H^2dV = -\int \vec{A}\nabla^2\vec{A}dV$. However, it is easy to show that there is absence of contribution term $\int H^2dV$ in the Hamiltonian of radiation \hat{H}_R in Eq.(4). Indeed, to calculate of H^2 , we use of a supporting formulae from textbook [8]

$$\left[\vec{a} \times \vec{b}\right]^2 = a^2 \cdot b^2 - \left(\vec{a} \cdot \vec{b}\right)^2$$

which leads to

$$H^2 = \left(\text{curl}\vec{A}, \text{curl}\vec{A}\right) = \nabla^2 A^2 - \text{div}\vec{A}\text{div}\vec{A} = \nabla^2 A^2 \quad (8)$$

at application Eq.(3).

Therefore, at using of Eq.(1), we posses

$$\frac{1}{V} \int H^2 dV = \frac{1}{V} \int \nabla^2 A^2 dV = -\delta_{\vec{k}+\vec{k}_1} \left(\vec{k} + \vec{k}_1\right)^2 \left(\vec{A}_{\vec{k}} + \vec{A}_{-\vec{k}}^+\right) \left(\vec{A}_{\vec{k}_1} + \vec{A}_{-\vec{k}_1}^+\right) = 0$$

This reasoning is not allowing to introduce so-called an unnecessary condition as $\int \text{div}[\vec{A} \times \vec{H}]dV = 0$. Consequently, the Hamiltonian radiation in Dirac theory may take a following form:

$$\hat{H}_R = \frac{1}{8\pi V} \int E^2 dV = \frac{1}{8\pi V c^2} \int \left(\frac{d\vec{A}}{dt}\right)^2 dV \quad (9)$$

within suggestion $t = 0$, presented in approach of Dirac [2], with using of Eq.(1), we have

$$\frac{d\vec{A}}{dt} = ic \sum_{\vec{k}} k \left(\vec{A}_{\vec{k}} - \vec{A}_{\vec{k}}^+\right) e^{i\vec{k}\vec{r}} \quad (10)$$

Inserting value $\frac{d\vec{A}}{dt}$ from Eq.(10) into Eq.(9), we find the Hamiltonian of radiation \hat{H}_R by following form:

$$\begin{aligned}
\hat{H}_R &= \frac{1}{8\pi c^2 V} \int \left(\frac{d\vec{A}}{dt} \right)^2 dV = \\
&= -\frac{1}{8\pi} \sum_{\vec{k}, \vec{k}_1} \delta(k + k_1) |k| \cdot |k_1| \left(\vec{A}_{\vec{k}} - \vec{A}_{-\vec{k}}^+ \right) \left(\vec{A}_{\vec{k}_1} - \vec{A}_{-\vec{k}_1}^+ \right) = \\
&= -\frac{1}{8\pi} \sum_{\vec{k}} k^2 \left(\vec{A}_{\vec{k}} - \vec{A}_{-\vec{k}}^+ \right) \left(\vec{A}_{-\vec{k}} - \vec{A}_{\vec{k}}^+ \right) \quad (11)
\end{aligned}$$

On other hand, we may claim an important condition for conjunction of vectors electric and potential of electromagnetic fields $\vec{A} \cdot \vec{E} = 0$. This reasoning leads to the law conservation for square potential electromagnetic field $\vec{A}^2 = const$ which in turn introduces a contribution to the Hamiltonian of radiation. In this context, we determine the Hamiltonian of radiation \hat{H}_0 as

$$\hat{H}_0 = \hat{H}_R + \hat{H}_A \quad (12)$$

with

$$\hat{H}_A = -\frac{1}{8\pi V} \int \mu A^2 dV$$

where μ is the constant parameter which is similar to chemical potential for the Bose gas.

In this context, our calculation shows that

$$\begin{aligned}
\hat{H}_A &= -\frac{\mu}{8\pi V} \int A^2 dV = \\
&= -\frac{\mu}{8\pi} \sum_{\vec{k}, \vec{k}_1} \delta(k + k_1) \left(\vec{A}_{\vec{k}} + \vec{A}_{-\vec{k}}^+ \right) \left(\vec{A}_{\vec{k}_1} + \vec{A}_{-\vec{k}_1}^+ \right) = \\
&= -\frac{\mu}{8\pi} \sum_{\vec{k}} \left(\vec{A}_{\vec{k}} + \vec{A}_{-\vec{k}}^+ \right) \left(\vec{A}_{-\vec{k}} + \vec{A}_{\vec{k}}^+ \right) \quad (13)
\end{aligned}$$

Consequently, we obtain the Hamiltonian operator \hat{H}_0 in Eq.(12) by following way:

$$\hat{H}_0 = \sum_{\vec{k}} \left(\frac{k^2}{4\pi} - \frac{\mu}{4\pi} \right) \hat{A}_{\vec{k}}^+ \hat{A}_{\vec{k}} - \sum_{\vec{k}} \left(\frac{k^2}{8\pi} + \frac{\mu}{8\pi} \right) \left(\hat{A}_{\vec{k}}^+ \hat{A}_{-\vec{k}}^+ + \hat{A}_{-\vec{k}} \hat{A}_{\vec{k}} \right) \quad (14)$$

Obviously, we cannot arrive to diagonal form of operator Hamiltonian in Eq.(14) because as we have been saying the Fourier components of vector potentials electromagnetic field $\vec{A}_{\vec{k}}^+$ and $\vec{A}_{\vec{k}}$ are the Pseudo- Bose-operators "creation" and "annihilation" of Bose-waves with spin one. Therefore, we

express these components via Bose-operators of Bose-particles by following way, namely, we reformulate the vector potential electromagnetic field \vec{A} via superposition of second quantization wave functions of one boson with spin one in point coordinate \vec{r} at suggestion $t = 0$:

$$\vec{A}(\vec{r}) = \vec{A}_0 \sqrt{V} \left(\psi^+(\vec{r}) + \psi(\vec{r}) \right) \quad (15)$$

where \vec{A}_0 is the amplitude of the vector potential electromagnetic field; $\psi^+(\vec{r})$ and $\psi(\vec{r})$ are, respectively, second quantization wave functions of creation and annihilation of free Bose- particles for one boson in the space of coordinate \vec{r} [3]:

$$\psi(\vec{r}) = \frac{1}{\sqrt{V}} \sum_{\vec{k}} \hat{a}_{\vec{k}} \exp^{i\vec{k}\vec{r}} \quad (16)$$

$$\psi^+(\vec{r}) = \frac{1}{\sqrt{V}} \sum_{\vec{k}} \hat{a}_{\vec{k}}^+ \exp^{-i\vec{k}\vec{r}} \quad (17)$$

where the operators $\hat{a}_{\vec{k}}^+$ and $\hat{a}_{\vec{k}}$ are, respectively, Bose- operators of creation and annihilation for free boson with wave-vector \vec{k} , which in turn satisfy the Bose-commutation relations:

$$\left[\hat{a}_{\vec{k}}, \hat{a}_{\vec{k}'}^+ \right]_- = \delta_{\vec{k}, \vec{k}'}$$

$$\left[\hat{a}_{\vec{k}}, \hat{a}_{\vec{k}'} \right]_- = 0$$

$$\left[\hat{a}_{\vec{k}}^+, \hat{a}_{\vec{k}'}^+ \right]_- = 0$$

Hence we see the form of Eq.(15) coincides with one of Eq.(1) at suggestion $t = 0$, when $\hat{A}_{\vec{k}} = \vec{A}_0 \hat{a}_{\vec{k}}$ and $\hat{A}_{\vec{k}}^+ = \vec{A}_0 \hat{a}_{\vec{k}}^+$.

Consequently, within reformulation of the vector potential electromagnetic field \vec{A} in Eq.(15) as well as introducing of assumption that the kinetic energy of boson can be expressed via amplitude of intensity electromagnetic wave A_0^2 as $\frac{\hbar^2 k^2}{2m} = \frac{A_0^2 k^2}{4\pi}$, the Hamiltonian of radiation \hat{H}_0 in Eq.(14) represents as Hamiltonian of Gibbs, and in turn takes a following form in the Fock space:

$$\hat{H}_0 = \sum_{\vec{k}} \left(\frac{\hbar^2 k^2}{2m} - \frac{\hbar^2 \mu}{2m} \right) \hat{a}_{\vec{k}}^+ \hat{a}_{\vec{k}} - \sum_{\vec{k}} \left(\frac{\hbar^2 k^2}{2m} + \frac{\hbar^2 \mu}{2m} \right) \left(\hat{a}_{\vec{k}}^+ \hat{a}_{-\vec{k}}^+ + \hat{a}_{-\vec{k}} \hat{a}_{\vec{k}} \right) \quad (18)$$

where the effective mass of bosons equals to

$$m = \frac{2\pi\hbar^2}{A_0^2}$$

The law conservation for square potential electromagnetic field $\vec{A}^2 = \text{const}$ means conservation total number n of Electromagnetic bosons:

$$\int \psi^+(\vec{r})\psi(\vec{r})dV = n$$

or

$$\sum_{\vec{k}} \hat{a}_{\vec{k}}^+ \hat{a}_{\vec{k}} = n$$

The evaluation of energy levels of the operators \hat{H}_0 in Eq. (18) within diagonal form, we apply the Bogoliubov linear transformation [4]:

$$\hat{a}_{\vec{k}} = \frac{\hat{b}_{\vec{k}} + L_{\vec{k}} \hat{b}_{-\vec{k}}^+}{\sqrt{1 - L_{\vec{k}}^2}} \quad (19)$$

where $L_{\vec{k}}$ is a symmetrical function of the wave vector \vec{k} .

Thus, the diagonal form of operator Hamiltonian \hat{H}_0 is presented as:

$$\hat{H}_0 = 2 \sum_{\vec{k}} \xi_{\vec{k}} \hat{b}_{\vec{k}}^+ \hat{b}_{\vec{k}} \quad (20)$$

hence we infer that the Bose-operators $\hat{b}_{\vec{k}}^+$ and $\hat{b}_{\vec{k}}$ are, respectively, the "creation" and "annihilation" operators of free photons of electromagnetic field with energy

$$\xi_{\vec{k}} = \sqrt{\left(\frac{\hbar^2 k^2}{4m} - \frac{\hbar^2 \mu}{4m}\right)^2 - \left(\frac{\hbar^2 k^2}{4m} + \frac{\hbar^2 \mu}{4m}\right)^2} = \frac{\hbar k \sqrt{-\mu}}{2m} = \hbar k c \quad (21)$$

where c is the velocity of photons which equals to $c = \frac{\hbar \sqrt{-\mu}}{2m}$ or $\mu = -\frac{4m^2 c^2}{\hbar^2}$ which in turn implies that the Hamiltonian of radiation \hat{H}_0 in the space of Fock takes a form:

$$\hat{H}_0 = \sum_{\vec{k}} \left(\frac{\hbar^2 k^2}{2m} + mc^2\right) \hat{a}_{\vec{k}}^+ \hat{a}_{\vec{k}} - \sum_{\vec{k}} \left(\frac{\hbar^2 k^2}{4m} - \frac{mc^2}{2}\right) \left(\hat{a}_{\vec{k}}^+ \hat{a}_{-\vec{k}}^+ + \hat{a}_{-\vec{k}} \hat{a}_{\vec{k}}\right) \quad (22)$$

111. INTERACTION BOSONS of ELECTROMAGNETIC FIELD WITH POLARISED ATOMS OF ^3He .

In this context, we try now present the Hamiltonian \hat{H}_Q of the interaction between electromagnetic waves with Fermi gas consisting N neutral atoms ${}^3\text{He}$ which represent as fermions.

By application of method of second quantization for system of N fermions [4], we may rewrite down the second quantization wave functions for one atom ${}^3\text{He}$ in point of coordinate \vec{r} in following form:

$$\psi(\vec{r}, \sigma) = \frac{1}{\sqrt{V}} \sum_{\vec{k}} \hat{c}_{\vec{k}, \sigma} e^{i\vec{k}\vec{r}} \quad (23)$$

$$\psi^+(\vec{r}, \sigma) = \frac{1}{\sqrt{V}} \sum_{\vec{k}} \hat{c}_{\vec{k}, \sigma}^+ e^{-i\vec{k}\vec{r}} \quad (24)$$

where the operators $\hat{c}_{\vec{k}, \sigma}^+$ and $\hat{c}_{\vec{k}, \sigma}$ are, respectively, of creation and annihilation Fermi operators for fermion with wave -vector \vec{k} , by the value of its spin z-component $\sigma = \pm \frac{1}{2}$.

These operators satisfy to the Fermi commutation relations $[\cdot \cdot \cdot]_+$ as:

$$\left[\hat{c}_{\vec{k}, \sigma}, \hat{c}_{\vec{k}', \sigma'}^+ \right]_+ = \delta_{\vec{k}, \vec{k}'} \cdot \delta_{\sigma, \sigma'}$$

$$[\hat{c}_{\vec{k}, \sigma}, \hat{c}_{\vec{k}', \sigma'}]_+ = 0$$

$$[\hat{c}_{\vec{k}, \sigma}^+, \hat{c}_{\vec{k}', \sigma'}^+]_+ = 0$$

and

$$\int \psi^+(\vec{r}, \sigma) \psi(\vec{r}, \sigma) dV = \sum_{\vec{k}, \sigma} \hat{c}_{\vec{k}, \sigma}^+ \hat{c}_{\vec{k}, \sigma} = \hat{N} \quad (25)$$

In beginning, we consider the Hamiltonian H_Q of interaction of radiation with N neutral atoms ${}^3\text{He}$ in volume V :

$$\hat{H}_Q = \int \sum_{\sigma} \psi^+(\vec{r}, \sigma) H_q \psi(\vec{r}, \sigma) dV \quad (26)$$

where H_q presents the Hamiltonian of interaction between one neutral atom with electromagnetic field which was proposed by Dirac in [2]:

$$H_q = \frac{1}{m_e} \left[\left(\vec{p} + \frac{e\vec{A}}{c} \right)^2 - \vec{p}^2 \right] \quad (27)$$

where $\vec{p} = -i\hbar\nabla$; m_e and e are, respectively, the mass of and charge of electron.

Inserting (27) into (26), and using of Eq.(3) and

$$\frac{1}{V} \int e^{i\vec{k}\vec{r}} dV = \delta_{\vec{k}}$$

we may rewrite down:

$$\begin{aligned} \hat{H}_Q &= \frac{e^2}{m_e c^2 V} \sum_{\vec{k}_1, \sigma} \sum_{\vec{k}_2, \sigma} \sum_{\vec{k}_3} \sum_{\vec{k}_4} \hat{c}_{\vec{k}_1, \sigma}^+ \hat{c}_{\vec{k}_2, \sigma} \left(\vec{A}_{\vec{k}_3} + \vec{A}_{-\vec{k}_3}^+ \right) \times \\ &\times \left(\vec{A}_{\vec{k}_4} + \vec{A}_{-\vec{k}_4}^+ \right) \delta_{\vec{k}_2 + \vec{k}_3 + \vec{k}_4 - \vec{k}_1} - \\ &- \frac{e}{m_e c} \sum_{\vec{k}_1, \sigma} \sum_{\vec{k}_2, \sigma} \sum_{\vec{k}_3} \vec{k}_2 \hat{c}_{\vec{k}_1, \sigma}^+ \hat{c}_{\vec{k}_2, \sigma} \left(\vec{A}_{\vec{k}_3} + \vec{A}_{-\vec{k}_3}^+ \right) \delta_{\vec{k}_2 + \vec{k}_3 - \vec{k}_1} \end{aligned} \quad (28)$$

However, introducing of the random - phase approximation [5], proposed by Bohm-Pines, which suggests that

$$\sum_{\vec{k}_1} \hat{c}_{\vec{k}_1, \sigma}^+ \hat{c}_{\vec{k}_1 - \vec{k}, \sigma} \approx \sum_{\vec{k}_1} \hat{c}_{\vec{k}_1, \sigma}^+ \hat{c}_{\vec{k}_1 - \vec{k}, \sigma} \delta_{\vec{k}, 0} = \sum_{\vec{k}} \hat{c}_{\vec{k}, \sigma}^+ \hat{c}_{\vec{k}, \sigma} = N$$

we obtain

$$\begin{aligned} \hat{H}_Q &= \frac{e^2 N}{m_e c^2 V} \sum_{\vec{k}} \left(\vec{A}_{\vec{k}} + \vec{A}_{-\vec{k}}^+ \right) \left(\vec{A}_{-\vec{k}} + \vec{A}_{\vec{k}}^+ \right) = \\ &= \frac{e^2 A_0^2 N}{m_e c^2 V} \sum_{\vec{k}} \left(\hat{a}_{\vec{k}} + \hat{a}_{-\vec{k}}^+ \right) \left(\hat{a}_{-\vec{k}} + \hat{a}_{\vec{k}}^+ \right) \end{aligned} \quad (29)$$

Inserting $m = \frac{2\pi\hbar^2}{A_0^2}$ in (30), we determine

$$\hat{H}_Q = \hat{H}_A = \frac{2\pi\hbar^2 e^2 N}{m m_e c^2 V} \sum_{\vec{k}} \left(\hat{a}_{\vec{k}} + \hat{a}_{-\vec{k}}^+ \right) \left(\hat{a}_{-\vec{k}} + \hat{a}_{\vec{k}}^+ \right) \quad (30)$$

Due to interaction between electromagnetic waves with Fermi gas consisting N neutral atoms ^3He , we possess a following form of Hamiltonian reflected radiation $\hat{H} = \hat{H}_0 + \hat{H}_Q$ which by support of (18), has a following form:

$$\begin{aligned} \hat{H} &= \sum_{\vec{k}} \left(\frac{\hbar^2 k^2}{2m} + mc^2 + \frac{2\pi\hbar^2 e^2 N}{m m_e c^2 V} \right) \hat{a}_{\vec{k}}^+ \hat{a}_{\vec{k}} - \\ &- \sum_{\vec{k}} \left(\frac{\hbar^2 k^2}{4m} - \frac{mc^2}{2} - \frac{\pi\hbar^2 e^2 N}{m m_e c^2 V} \right) \left(\hat{a}_{\vec{k}}^+ \hat{a}_{-\vec{k}}^+ + \hat{a}_{-\vec{k}} \hat{a}_{\vec{k}} \right) \end{aligned} \quad (31)$$

which at using of Bogoliubovs transformations, presents in diagonal form:

$$\hat{H}_0 = 2 \sum_{\vec{k}} E_{\vec{k}} \hat{d}_{\vec{k}}^+ \hat{d}_{\vec{k}} \quad (32)$$

hence we infer that the Bose-operators $\hat{d}_{\vec{k}}^+$ and $\hat{d}_{\vec{k}}$ are, respectively, the "creation" and "annihilation" operators of reflected photons of electromagnetic field with energy

$$E_{\vec{k}} = \hbar k v \quad (33)$$

where v is the velocity of reflected photons, which equals to

$$v = c \sqrt{1 + \frac{\pi \hbar^2 e^2 N}{m^2 m_e c^4 V}} \quad (34)$$

In the letter [6], it is found the mass of the boson of an electromagnetic field which represents as a new fundamental constant:

$$m = \frac{m_e e^4}{4 \hbar^2 c^2} = 1.2 \cdot 10^{-35} \text{ kg} \quad (35)$$

Inserting m from (35) into (34), we can find

$$v = c \sqrt{1 + \frac{16 \pi a^3 N}{V}} \quad (36)$$

where $a = \frac{\hbar^2}{m_e e^2}$ is the Bohr radius.

Introducing the parameter

$$r_s = \left(\frac{3V}{4\pi N} \right)^{\frac{1}{3}} \frac{m_e e^2}{\hbar^2}$$

which characters a density of atomic gas, and then, we obtain

$$v = c \sqrt{1 + \frac{12}{r_s^3}} \quad (37)$$

Obviously, at high density of atomic gas $r_s \rightarrow 0$, and indeed $v \rightarrow \infty$. This fact confirms that the relativistic theory of Einstein is wrong but in the case of considering a low density gas $r_s \gg 1$, which corresponds to a ${}^3\text{He}$ gas, the speed of light in (36) is approximated to a meaning

$$v = c \left(1 + \frac{6}{r_s^3} \right) \quad (38)$$

Thus, we proved that velocity of electromagnetic waves into gas consisting polarized atoms ${}^3\text{He}$ is rather than one in vacuum, due to an interaction between electrons of atom and electromagnetic field. This fact may have significant output for application in instruments and devices, which work on the photo effect.

References

1. V.N. Minasyan ., Quasi-bosons of Electromagnetic Field (2008) ; arXiv:0808.0567v9 [physics.gen-ph]
2. P.A.M.. Dirac , The Principles of Quantum Mechanics, Oxford at the Clarendon press (1958), Lectures on Quantum Mechanics. Yeshiva University New York (1964)
3. K. Huang , Statistical Mechanics, (Wiley 1963).
4. N.N. Bogoliubov , Jour. of Phys.(USSR), **11**, 23 (1947)
5. D. Bohm , D. Pines ., Phys.Rev.B. **82**, 625 (1951); D. Bohm , D. Pines , Phys.Rev.B. **92**, 609 (1953).
6. V.N. Minasyan ., Quantized Equations of Maxwell and New Fundamental Constant in The Nature <http://arxiv.org/abs/0902.3057>. 2009, [physics.gen-ph]