

STABILITY OF SUPERFLUID FERMI GASES IN OPTICAL LATTICES

YOSHIHIRO YUNOMAE*

Department of Physics, Waseda University, Okubo, Shinjuku-ku, Tokyo 169-8555, Japan

**E-mail: yunon@kh.phys.waseda.ac.jp*

IPPEI DANSHITA

Department of Physics, Faculty of Science, Tokyo University of Science, Kagurazaka, Shinjuku-ku, Tokyo 162-8601, Japan

DAISUKE YAMAMOTO

Department of Physics, Waseda University, Okubo, Shinjuku-ku, Tokyo 169-8555, Japan

NOBUHIKO YOKOSHI

*Nanotechnology Research Institute, AIST, Tsukuba 305-8568, Japan
CREST(JST), Saitama 332-0012, Japan*

SHUNJI TSUCHIYA

*Department of Physics, Keio University, Hiyoshi, Kohoku-ku, Yokohama 223-8522, Japan
CREST(JST), Saitama 332-0012, Japan*

Critical velocities of superfluid Fermi gases in optical lattices are theoretically investigated across the BCS-BEC crossover. We calculate the excitation spectra in the presence of a superfluid flow in one- and two-dimensional optical lattices. It is found that the spectrum of low-lying Anderson-Bogoliubov (AB) mode exhibits a roton-like structure in the short-wavelength region due to the strong charge density wave fluctuations, and with increasing the superfluid velocity one of the roton-like minima reaches zero before the single-particle spectrum does. This means that superfluid Fermi gases in optical lattices are destabilized due to spontaneous emission of the roton-like AB mode instead of due to Cooper pair breaking.

Keywords: superfluidity; critical velocity; Anderson-Bogoliubov mode.

1. Introduction

According to Landau criterion, superfluid flow of bosonic particles propagates without dissipation, unless its group velocity exceeds a certain critical value and spontaneous creation of excitations is induced.¹ The direct experimental verifications of the criterion have been attempted mainly in superfluid ^4He ² and dilute atomic Bose gases.³ These studies have inspired various discussions and contributed to understanding of the superfluid phenomenon itself.

The recent realization of the BCS-BEC crossover in ultracold Fermi gases⁴ has raised

a new question, i.e., how does the critical velocity for bosonic superfluids change in the BCS regime? Recently, Miller *et al.* investigated critical velocities of superfluid Fermi gases across the BCS-BEC crossover in a moving one-dimensional (1D) optical lattice.⁵ They determined the critical velocities at which the abrupt decrease of fermionic pairs in the condensate occurs, when the velocity of the lattice potential is varied.

Theoretically, critical velocities in the BCS-BEC crossover have been investigated in a uniform system.⁶ It was claimed that the instability of superfluid flow in a uniform

system is induced by Cooper pair breaking in the BCS regime.⁶ In this paper, motivated by the experiment done by Miller *et al.*,⁵ we investigate critical velocities of superfluid Fermi gases in the BCS-BEC crossover in optical lattices. We calculate excitation spectra and determine critical velocities in 1D and 2D optical lattices. We find that the excitation spectrum of the low-lying Anderson-Bogoliubov (AB) mode^{6,7,8} has a characteristic roton-like structure, and the instability of superfluid flow in the BCS regime is induced by a different mechanism from that in a uniform system: the spontaneous emission of AB mode destabilizes the superfluidity.

2. Formalism

We consider two-component atomic Fermi gases with equal populations. We assume that the lattice potential is sufficiently deep that the tight-binding approximation is valid. Thus, the system can be described by a single-band Hubbard model

$$H = -J \sum_{\langle i,j \rangle, \sigma} (c_{i\sigma}^\dagger c_{j\sigma} + h.c.) + U \sum_i c_{i\uparrow}^\dagger c_{i\downarrow}^\dagger c_{i\downarrow} c_{i\uparrow}, \quad (1)$$

where $c_{j\sigma}$ is an annihilation operator of an atom with mass m and pseudospin $\sigma = \uparrow, \downarrow$ on the j -th site, J is a hopping energy between nearest-neighbor sites, and $U (< 0)$ is an on-site attractive interaction. We set $\hbar = 1$ throughout the paper.

We assume that a superfluid flows with the velocity \mathbf{v} in the coordinate system fixed with respect to the lattice potential, i.e. Cooper pairs have the center of mass quasi-momentum $\mathbf{q} = 2m\mathbf{v}$. The corresponding order parameter is given by

$$\Delta_{\mathbf{v}} = -\frac{U}{M} \sum_{\mathbf{k}} \langle c_{-\mathbf{k}+m\mathbf{v}\downarrow} c_{\mathbf{k}+m\mathbf{v}\uparrow} \rangle, \quad (2)$$

where M is the number of lattice sites. By diagonalizing the Hamiltonian within the

BCS mean-field approximation, we obtain the single-particle excitation energy as

$$E_{\mathbf{v}}^{\pm}(\mathbf{k}) = \eta_{\mathbf{v}}(\mathbf{k}) \pm \sqrt{(\varepsilon_{\mathbf{v}}(\mathbf{k}) - \mu)^2 + |\Delta_{\mathbf{v}}|^2}, \quad (3)$$

where $\eta_{\mathbf{v}}(\mathbf{k}) = 2J\Sigma_{\nu} \sin(mv_{\nu}d) \sin(k_{\nu}d)$ and $\varepsilon_{\mathbf{v}}(\mathbf{k}) = 2J\Sigma_{\nu}(1 - \cos(mv_{\nu}d) \cos(k_{\nu}d))$ with d being the lattice constant. The energy gap $\Delta_{\mathbf{v}}$ and the chemical potential μ are determined by solving the gap equation and the number equation self-consistently. This scheme interpolates the BCS and BEC regimes at $T = 0$.⁹

To determine the critical velocities, it is necessary to calculate the excitation spectra in the presence of a superfluid flow. For neutral fermionic superfluids, there exists a low-lying AB mode^{6,7,8} in addition to the single-particle excitation. The spectrum of the AB mode can be obtained from the pole of the density response function $\chi(\mathbf{q}, \omega)$.¹⁰ We calculate $\chi(\mathbf{q}, \omega)$ by employing the generalized random phase approximation (GRPA) developed by Côté and Griffin.¹⁰

3. Results

Conditions for Cooper pair breaking are determined from the single-particle excitation energy Eq. (3). When the absolute value of the superfluid velocity \mathbf{v} exceeds a certain critical value v_{pb} , one finds that $E_{\mathbf{v}}^+(\mathbf{k})$ becomes negative. The negative energy of the single-particle excitation indicates the occurrence of spontaneous Cooper pair breaking. Thus, the critical velocity for Cooper pair breaking is given by v_{pb} .^{5,6,11}

In order to compare the instabilities induced by two kinds of excitations, i.e., the single-particle excitation and AB mode excitation, it is necessary to monitor both the excitation spectra in parallel. For that purpose, it is useful to plot the dynamical structure factor $S(\mathbf{q}, \omega) = -\text{Im}\chi(\mathbf{q}, \omega)/\pi$. First, we investigate the case of 1D optical lattices. In Fig. 1, $S(q, \omega)$ is shown for (a) current-free and (b) current-carrying cases. In Fig. 1,




Fig. 1. Dynamical structure factors $S(q, \omega)$ in 1D optical lattices are shown for (a) current-free ($v = 0$) and (b) current-carrying cases ($v = 0.095/md$), when $n = 0.25$ and $U = -1.0J$. The dark region indicates large $S(q, \omega)$. In each panel, the upper gray region corresponds to the particle-hole continuum, and the lower dashed curve is the spectrum of the AB mode.

the upper gray regions are the particle-hole continuums, and the lower boundaries of the particle-hole continuums correspond to $\omega = \min[E_v^+(\mathbf{k} + \mathbf{q}) - E_v^-(\mathbf{k})]$. The lower curves in Fig. 1 are the spectrum of AB mode, which becomes phonon-like when $qd \ll 1$.⁷ It should be noted that the spectrum of the AB mode has roton-like minima at $|q| \simeq 2k_F$ where k_F is the Fermi wavevector. Such a roton-like structure also appears in the spectrum of AB mode in uniform 1D Fermi gases.¹² The energies of the roton-like minima decrease as one approaches the half-filling when $v = 0$. They eventually become




Fig. 2. Critical velocities v_{pb} and v_{ab} as functions of the filling n when $U = -1.0J$. The solid line represents v_{pb} and the dashed line represents v_{ab} .

zero at $k = \pi/d$ at the half-filling. This suggests that exciting density fluctuations with $k = \pi/d$ costs zero energy and the superfluid ground state is in competition with the charge density wave (CDW) state. Thus, the roton-like minima reflect the strong CDW fluctuations.

In the presence of a superfluid flow with $v > 0$, the spectrum slants towards the left side (see Fig. 1(b)). The critical velocity due to pair breaking v_{pb} coincides with the velocity at which the particle-hole continuum touches the line $\omega = 0$. Since the AB mode spectrum locates below the particle-hole continuum, one of the roton-like minima reaches zero at the velocity v_{ab} which is smaller than v_{pb} . According to the Landau criterion,¹ this means that the spontaneous emission of the AB mode is induced before the Cooper pair breaking occurs. In Fig. 2, the critical velocities v_{pb} and v_{ab} are plotted as functions of the filling n . One clearly sees that v_{ab} is always smaller than v_{pb} , and that both v_{ab} and v_{pb} decrease monotonically with increasing filling. The difference between the two velocities diminishes near the half-filling ($n \simeq 1$).

We proceed to discuss Fermi gases in 2D

4




Fig. 3. Dynamical structure factor $S(\mathbf{q}, \omega)$ in 2D optical lattices for $q_x = q_y$. We set $U = -4.5J$, $n = 0.5$, and $\mathbf{v} = (v, v)/\sqrt{2}$ with $v = 0.467/md$. The dark region indicates large $S(\mathbf{q}, \omega)$. The upper gray region corresponds to the particle-hole continuum, and the lower curve is the spectrum of AB mode.

optical lattices. In Fig. 3, we plot the dynamical structure factor $S(\mathbf{q}, \omega)$ for $q_x = q_y$, when $\mathbf{v} = (v/\sqrt{2}, v/\sqrt{2})$. Similarly to the 1D case, the spectrum of AB mode has a roton-like minimum in Fig. 3. It reaches zero before the particle-hole continuum does when v is increased. Thus, v_{ab} is smaller than v_{pb} , and the instability induced by emission of the AB mode also dominates in 2D optical lattices. Furthermore, it is found that the direction of the superfluid velocity \mathbf{v} affects the spectrum of AB mode. The energy of the roton-like minimum decreases more rapidly for $\mathbf{v} = (v/\sqrt{2}, v/\sqrt{2})$ than for $\mathbf{v} = (v, 0)$. This implies that the CDW fluctuation is enhanced when the supercurrent is flowing in the direction of (π, π) .

4. Conclusion

In summary, we have studied critical velocities of superfluid Fermi gases in 1D and 2D optical lattices using the attractive Hubbard model. We calculated the excitation spectra of single-particle excitation and AB mode by employing the GRPA. We have shown that the spectrum of AB mode has a roton-like

structure at $|q| \simeq 2k_F$, and it always lies below the particle-hole continuum due to the strong CDW fluctuations in optical lattices. As the velocity of the superfluid flow is increased, the energy of a roton-like minimum decreases and the instability due to the emission of roton-like excitations was shown to occur before Cooper pair breaking occurs. We note that Burkov and Paramekanti have proposed another type of current-induced instability of superfluid Fermi gases in optical lattices.¹³

Acknowledgments

We would like to thank K. Kamide, S. Kurihara, and T. Nikuni for valuable comments and discussions. I.D. and D.Y. are supported by a Grant-in-Aid from JSPS.

References

1. L. D. Landau, *J. Phys. (USSR)* **5**, 71 (1941).
2. D. R. Tilley and J. Tilley, *Superfluidity and Superconductivity*, (Bristol, Hilger 1991).
3. R. Onofrio *et al.*, *Phys. Rev. Lett.* **85**, 2228 (2000).
4. C. A. Regal *et al.*, *Phys. Rev. Lett.* **92**, 040403 (2004); M. W. Zwierlein *et al.*, *ibid.* **92**, 120403 (2004).
5. D. E. Miller *et al.*, *Phys. Rev. Lett.* **99**, 070402 (2007).
6. R. Combescot *et al.*, *Phys. Rev. A* **74**, 042717 (2006).
7. P. W. Anderson, *Phys. Rev.* **112**, 1900 (1958).
8. L. P. Pitaevskii *et al.*, *Phys. Rev. A* **71**, 053602 (2006).
9. A. J. Leggett in *Modern Trends in the Theory of Condensed Matter*, edited by A. Pekalski and R. Przystawa (Springer-Verlag, Berlin, 1980).
10. R. Côté and A. Griffin, *Phys. Rev. B* **48**, 10404 (1993).
11. M. Rodriguez and P. Törmä, *Phys. Rev. A* **69**, 041602(R) (2004).
12. T. Alm and P. Schuck, *Phys. Rev. B* **54**, 2471 (1996).
13. A. A. Burkov and A. Paramekanti, *Phys. Rev. Lett.* **100**, 255301 (2008).