

Can Heisenberg Chains Mirror a State?

Marcin Wieśniak

Centre for Quantum Technologies, National University of Singapore, 3 Science Drive 2, Singapore 117543, Singapore

Faithful exchange of quantum information can in future become a key part of many computational algorithms. Some Authors suggest to use chains of mutually coupled spins as channels for quantum communication. One can divide these proposals into the groups of assisted protocols, which require some additional action from the users, and natural ones, based on the concept of state mirroring. We show that the natural spin transfer is fundamentally not possible with chains of spins- $\frac{1}{2}$ coupled by the Heisenberg interaction, but without local magnetic fields.

Transferring information is one of the most important parts of classical computing. It is also the simplest. Classical wires are just pieces of metal, in which the electromagnetic field can freely propagate.

Quantum Mechanics also offers interesting computation algorithms [1, 2, 3], which can be implemented in various physical systems. In measurement-based techniques, such as the cluster state computation [4], there is no need to exchange quantum information, but this is at the expense of maintaining a large entangled state. One may say that in this computational process the information was initially exchanged between all the qubits in the lattice, later to be only processed locally.

In unitary-operation based proposals, the exchange of quantum information is a challenge. One can use the teleportation [5], again dependent on entangled states. They are often considered an additional resource and could be not easy to produce. The exchange of photons between more distant sites of the lattice is not a good solution as there is no faithful and controllable atom-photon interface [6]. The other, often discussed idea is sending the state through a quantum wire, a chain (or more generally, a graph) of mutually coupled qubits.

Before we answer the title question, let us briefly discuss some state transfer proposals, together with their possible disadvantages. The original idea of quantum communication state through a system of interacting spins- $\frac{1}{2}$ was introduced by Bose [7]. He has considered any Heisenberg lattice described by the following Hamiltonian:

$$H_1 = \sum_{i,j=1}^N J_{ij} \vec{\sigma}^{[i]} \cdot \vec{\sigma}^{[j]} + \sum_{i=1}^N B_i \sigma_z^{[i]}. \quad (1)$$

$\vec{\sigma}^{[i]}$ denotes the usual Pauli matrix vector acting on a qubit labeled i , and $\sigma_z^{[i]}$ is the z component thereof. J_{ij} denotes a coupling constant between two spins, and the local magnetic field is B_i .

A qubit can be encoded as follows: for the logical $|0\rangle$ we take the fully magnetized state, $|00\dots 0\rangle$, while logical $|1\rangle$ is translated into the flip of the first spin (the one closest to the sender), $|10\dots 0\rangle$. Due to the free evolution, the excitation $|1\rangle$ wanders over the whole system. The idea

of the protocol is simply to wait until the time, at which the average fidelity of the receiver's qubit to the input state, $F = \int \langle \psi_{in} | \rho_{out} | \psi_{in} \rangle d\mu(|\psi_{in}\rangle)$ ($\in d\mu(|\psi_{in}\rangle) = 1$) has a peak.

Bose discusses an example of the uniform Heisenberg chain ($J_{ij} = \frac{J}{2} \delta_{j,i+1}$, $B_i = B$). For the number of spins $N \approx 80$ and the maximum time of waiting $t_{\max} = 4000/J$ he observes F to be hardly above the classical limit of $2/3$. This shows the main problem in quantum state transfer. If energies are not mutually rational, the excitation is irreversibly spread over the wire. At some instances, the eigenstates interfere constructively enough to partially extract encoded information from the last spin, but in general, there is always some part of it left in the chain and hence lost to the receiver.

In fact, the non-periodic evolution is a feature of all but few lattices. The Reader should keep in mind that a complex quantum protocol could involve a number of transfer routines. On top of that, the environment interacts with the system causing decoherence [8, 9, 10, 11, 12]. The cumulative effect of these losses could cancel the benefit of quantum computing.

Bose has noticed, however, that even with a low fidelity of the transfer, one can constitute entanglement between the sender's and the receiver's spins. This entanglement could be accumulated over many runs, distilled [13], and used to teleport the message.

Subsequent solutions [8, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23] were aimed to minimize the loss of information. An example is "the valve scheme", which was introduced in [21]. This time, apart from the wire, one has an additional qubit ("the bucket"), a gate between which and the last spin of the chain is occasionally switched on by the receiver. This gate is time-dependent partial SWAP operation. Similarly as in the first scheme, logical $|0\rangle$ and $|1\rangle$ are represented by the magnetized state, and an excitation in the chain, respectively. The partial SWAP is designed in such a way, that it transfers the excitation only from the chain to the bucket. One needs to perform this operation many time to gather the complete information (in the asymptotic limit).

The secret behind the action of the valve is that it is being adjusted to the free evolution. In the first iteration, it simply acts as a full SWAP gate, as the bucket was

initialized in $|0\rangle$. Then we let the chain to freely evolve for some time, until we again switch on the valve. This time it needs to take into account the present state of the last spin in the chain. To use this protocol, one needs to precisely simulate the free dynamics. The good news is, it is enough to do the simulation only once and devise a circuit, which would “replay” it for every transfer. Still, the protocol requires a huge amount of computational power, as for a simple process of transfer.

An example of a lossless protocol was proposed by Burgarth and Bose in [8]. Instead of using one chain of spins, their dual-rail protocol involved two independent, arbitrary, but identical such systems (the scheme was later generalized for two non-identical systems [20]). The input is encoded by injecting the excitation into one or the other chain. At the output, the magnetization of the two last spins, $\sigma_z^{[N,1]} + \sigma_z^{[N,2]}$, is measured at some time determined numerically. If the result is 0, the superposition of the two receiver’s qubits is the same, as created by the sender initially. This measurement can be repeated until success.

An important feature of the dual rail protocol is that whenever the transfer is conclusive, it is also faithful. This is, needless to say, in the idealized case. The superposition can be much deformed if the two chains are not identical and the measurement is conducted after rather a long time. This protocol can also turn out inefficient if one includes decoherence. Two chains are a much bigger system than a single one, which therefore has much more ways to interact with the environment.

The above schemes, among others (e.g. [22, 23]), are examples of assisted state transfer protocols. Their success comes at the expense of an additional action taken by the user. This is as opposed to natural schemes, in which the only conducted operations are encoding and decoding. These proposals rely on mirroring chains [24]. A mirroring chain is symmetric with respect to its middle. The parity (symmetry or antisymmetry) of its Hamiltonian eigenstates matches the parity of the respective energy. This is known as the spectrum parity-matching condition (SPMC) [18]. At half of the period, the relative phase between the odd and the even component of the state is changed by π . Whatever was initialized at one end of the chain, can be now found at the opposite. An example of a system with this property was found by Christandl *et al.* [14], and independently by Nikolopoulos, Petrosyan, and Lambropoulos [15, 16]. It is an xx chain, the Hamiltonian of which is given by

$$H_2 = \sum_{i=1}^{N-1} J_{i,i+1} (\sigma_x^{[i]} \sigma_x^{[i+1]} + \sigma_y^{[i]} \sigma_y^{[i+1]}). \quad (2)$$

For the choice of coupling constants $J_{i,i+1} = \sqrt{i(N-i)}$,

the fully magnetized state has energy 0, while the one excitation subspace is built of states of energies $-\frac{N-1}{2}, -\frac{N-3}{2}, \dots, \frac{N-3}{2}, \frac{N-1}{2}$.

Different physical systems realize different couplings. The xx is typical for opto-atomic implementations [25]. The Heisenberg interaction is specific for some solid state technologies, e.g. quantum dots [26, 27, 28]. The above Hamiltonian with these coupling constants fails to realize a natural quantum wire if we replace the xx interaction between qubits with the latter type. In what remains, we are going to answer the question if it is possible to realize a natural state transfer in a chain of spin- $\frac{1}{2}$ in a system with the isotropic interaction between the nearest neighbors.

The problem is trivial if one allows local magnetic fields, specifically chosen for each qubit. This would make it possible to cancel the diagonal terms of the Hamiltonian (projected onto the one excitation subspace) and bring it to the form of (2). Such a precise adjustment of the magnetic field could be, nevertheless, challenging for a system of physical length of a few micrometers. We are rather interested in the case of a uniform magnetic field acting over the whole wire. The magnitude of this field is here chosen to be 0.

As shown by Katsura [29], the Hamiltonian of a linear system with xx interaction between nearest neighbors can be expressed in terms of a finite field of non-interacting fermions. This allows the chain presented in [14, 15, 16] to mirror all states, not only those in particular subspaces, provided a proper choice of the magnitude of the uniform magnetic field. $\sigma_z^{[i]} \sigma_z^{[i+1]}$ represents the interaction between two fermionic modes. After the Jordan-Wigner transformation, which constitutes the canonical anti-commutation relations,

$$a^{[k]} = \frac{1}{2} \left(\prod_{j<k} \sigma_z^{[j]} \right) (\sigma_x^{[k]} + i\sigma_y^{[k]}), \quad (3)$$

$$a^{\dagger[k]} = \frac{1}{2} \left(\prod_{j<k} \sigma_z^{[j]} \right) (\sigma_x^{[k]} - i\sigma_y^{[k]}), \quad (4)$$

the zz term takes the form of

$$\sigma_z^{[i]} \sigma_z^{[i+1]} = (2a^{\dagger[i]} a^{[i]} - 1)(2a^{\dagger[i+1]} a^{[i+1]} - 1). \quad (5)$$

Hence, the Heisenberg chain could possess the property of mirroring not in general, but only in certain subspaces—including the fully magnetized state (the fermionic vacuum) and the one excitation space. For the sake of our analysis, we can drop the first term, as we aim to work with one fermion only, and the irrelevant constant. The one excitation part of the Hamiltonian has the three-diagonal form and reads

$$H_{oe} = -\frac{1}{2} \begin{pmatrix} -J_{1(N-1)} & J_{1(N-1)} & 0 & \dots & 0 & 0 \\ J_{1(N-1)} & -J_{1(N-1)} - J_{2(N-2)} & J_{2(N-2)} & \dots & 0 & 0 \\ 0 & J_{2(N-2)} & -J_{2(N-2)} - J_{3(N-3)} & \dots & 0 & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & 0 & \dots & -J_{1(N-1)} - J_{2(N-2)} & J_{1(N-1)} \\ 0 & 0 & 0 & \dots & J_{1(N-1)} & -J_{1(N-1)} \end{pmatrix} \quad (6)$$

The first step is to separate the odd and even sectors with a unitary operation

$$U = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 0 & \dots & 0 & 1 \\ 0 & 1 & \dots & 1 & 0 \\ \dots & \dots & \dots & \dots & \dots \\ 0 & 1 & \dots & -1 & 0 \\ 1 & 0 & \dots & 0 & -1 \end{pmatrix} \quad (7)$$

for N even and

$$U = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 0 & \dots & \dots & \dots & 0 & 1 \\ 0 & 1 & \dots & \dots & \dots & 1 & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & 0 & \sqrt{2} & 0 & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & 1 & \dots & \dots & \dots & -1 & 0 \\ 1 & 0 & \dots & \dots & \dots & 0 & -1 \end{pmatrix} \quad (8)$$

for N odd. The one excitation part of the Hamiltonian transforms into two three-diagonal matrices, which we will call H_e and H_o :

$$U \cdot H_{oe} \cdot U^\dagger = \begin{pmatrix} H_e & 0 \\ 0 & H_o \end{pmatrix}. \quad (9)$$

If the Hamiltonian expresses a periodic evolution, it can always be rescaled so that all eigenvalues are integers. Having done that, we are looking such the values of $J_{i(N-i)}$ s, which satisfy SPMC that all the eigenvalues of H_e are even and all the eigenvalues of H_o are odd. The first hint is obtained from the comparison of the traces.

$$\text{Tr}(H_e) - \text{Tr}(H_o) = J_{i_0(N-i_0)}, \quad (10)$$

with $i_0 = N/2$ or $(N-1)/2$, depending on the parity of N . $J_{i_0(N-i_0)}$ is hence an integer, the parity of which is dependent on $N \pmod{4}$.

In the next step we compare the determinants of H_e and H_o . The even part is known to have 0 as an eigenvalue associated with the N -partite W state, $\frac{1}{\sqrt{N}}(|10\dots\rangle + |01\dots\rangle + \dots)$. The Hamiltonian commutes with the total angular momentum operator, which in this subspace distinguishes the W state from the rest. Instead of $\text{Det}(H_e)$, we should rather consider the product of non-zero eigenvalues, $\text{Det}'(H_e) = \lim_{\epsilon \rightarrow 0} \text{Det}(H_e + \epsilon)/\epsilon$.

First we consider N odd. We have

$$\text{Det}'(H_e) = \frac{N}{2^{N/2-1}} \prod_{i=1}^{(N-1)/2} J_{i(N-i)}, \quad (11)$$

$$\text{Det}(H_o) = \frac{1}{2^{N/2-1}} \prod_{i=1}^{(N-1)/2} J_{i(N-i)}. \quad (12)$$

The second quantity is supposedly odd. However, if we multiply it by N , we are expected to get something even, namely $\text{Det}'(H_e)$. Hence for nearest neighbor Heisenberg chains of odd N qubits, there cannot be natural state transfer.

Next, let us take $N = 4n + 2$. The determinants are

$$\text{Det}'(H_e) = \frac{2n+1}{2^{2n}} \prod_{i=1}^{2n} J_{i(4n+2-i)}, \quad (13)$$

$$\text{Det}(H_o) = \frac{1}{2^{2n}} \prod_{i=1}^{2n+1} J_{i(4n+2-i)}. \quad (14)$$

To make these two expressions equations, we need to multiply the determinant of H_o by $2n+1$ and the determinant of the reversible part of H_e by an integer $J_{(2n+1)^2}$. But again, with multiplying two odd numbers, we expect to get an even one. The natural state transfer is possible neither for $N = 4n + 2$.

The only exception is, of course, $N = 2$, where there is a single gap between the singlet and the triplet.

The argument for the remaining subset of $N = 4n$ is a bit less obvious:

$$\text{Det}'(H_e) = \frac{2n}{2^{2n-1}} \prod_{i=1}^{2n-1} J_{i(4n-i)}, \quad (15)$$

$$\text{Det}(H_o) = \frac{1}{2^{2n-1}} \prod_{i=1}^{2n} J_{i(4n-i)}. \quad (16)$$

We cannot simply multiply $\text{Det}(H_o)$ by $2n$ as we would have lost the oddity. Instead, we notice that $\text{Det}'(H_e)$ is supposed to be a product of $N/2 - 1$ even numbers, hence it should involve factor 2^{2n-1} . Then $2n$ could be rewritten as $2^p(2q+1)$, where p and q are integers. One has $p \leq N/2 - 1$ (the equality holds for $N = 4$). Thus we are sure that if the chain realizes natural transfer, $2^{-p} \text{Det}'(H_e) = \frac{2q+1}{2^{2n-1}} \prod_{i=1}^{2n} J_{i(4n-i)}$ is an integer. However, one has

$$(2q+1) \text{Det}(H_o) = 2^{-p} J_{4n^2} \text{Det}'(H_e), \quad (17)$$

and since J_{4n^2} is even (such is the number of eigenvalues of H_o), the left-hand side cannot be odd.

In summary, we have shown a fundamental limitation on Heisenberg chains of qubits with nearest neighbor modulated coupling, which prevents them from being mirroring systems. This makes the natural perfect state transfer impossible in those systems, unless one is allowed to introduce local magnetic fields. This is even though it was demonstrated that arrays of Heisenberg interacting qubits are suitable for quantum computing [30]. Our result increases the importance of the transfer protocols described above [8, 20, 21], and others alike (e.g. [22, 23]). Which transfer protocol is most feasible, will, naturally, depend on the specific system realizing a quantum computer and its relation to the environment.

It remains an interesting question if it is possible to naturally transfer a state in a Heisenberg wire, when one applies the magnetic field only to the extreme sites [31]. This is well motivated from the experimental point of view, as the ends of the chain are close to heads-devices, which write in and read out the transferred state. Also, one can ask if including non-nearest interactions can improve the situation. We recall that Kay [32] had studied the non-nearest neighbors interactions in the xx model and failed to observe the perfect transfer, unless local magnetic fields are allowed.

This work is supported by the National Research Foundation and Ministry of Education, Singapore.

-
- [1] D. Deutsch and R. Jozsa, *Proc. R. Soc. London, Ser. A* **439**, 553 (1992).
- [2] P.W. Shor, *Proc. 35th Ann. Symp. Found. Comp. Sc., Los Alamos* (1994).
- [3] L. Grover, *Procc. 28th Ann. ACM Symp. Theor. Comp. (SOTC)* 212 (1996), quant-ph/9605043 @ xxx.lanl.gov.
- [4] R. Raussendorf, D.E. Browne, and H. J. Briegel, *Phys. Rev. A* **68**, 022312 (2003).
- [5] C.H. Bennett, G. Brassard, C. Crépeau, R. Jozsa, A. Peres, and W.K. Wootters, *Phys. Rev. Lett.* **70**, 1895(1993).
- [6] R. Alicki, arXiv:0807.2609 @ www.arxiv.org
- [7] S. Bose, *Phys. Rev. Lett.* **91**, 207901 (2003).
- [8] D. Burgarth and S. Bose, *Phys. Rev. A* **71**, 052315 (2005).
- [9] J.-M. Cai, Z.-W. Zhou, G.-C. Guo, *Phys. Rev. A* **74**, 022328 (2006).
- [10] A. Kay, *Phys. Rev. Lett.* **98**, 010501 (2007).
- [11] D. Tsomokos, M. Hartmann, S. Huelga, and M.B. Plenio, *New. J. Phys.* **9**, 79 (2007).
- [12] M. Wieśniak, arXiv:0711.2357 @ www.arxiv.org
- [13] C.H. Bennett, G. Brassard, S. Popescu, B. Schumacher, J.A. Smolin, and W.K. Wootters, *Phys. Rev. Lett.* **76**, 722 (1996).
- [14] M. Christandl, N. Datta, A. Ekert, and A. J. Landahl, *Phys. Rev. Lett.* **92**, 187902 (2004).
- [15] G.M. Nikolopoulos, D. Petrosyan, and P. Lambropoulos, *Eurphys. Lett.* **65**, 297 (2004).
- [16] G.M. Nikolopoulos, D. Petrosyan, and P. Lambropoulos, *Jour. Phys.:Cond. Matt.* **16**, 4991 (2004).
- [17] Y. Li *et al.*, *Phys. Rev. A* **71** 022301 (2005).
- [18] B. Shi, Y. Li, Z. Song, and C. P. Sun, *Phys. Rev. A* **71**, 032309 (2005).
- [19] M. Christandl, N. Datta, T.C. Dorlas, A. Ekert, A. Kay, and A. J. Landahl, *Phys. Rev. A* **71**, 032312 (2005).
- [20] D. Burgarth and S. Bose, *New Jour. Phys.* **7**, 135 (2005).
- [21] D. Burgarth, V. Giovannetti, and S. Bose, *Phys. Rev. A* **75**, 062327 (2007).
- [22] T. Oshima, A. Ekert, D.K.L. Oi, D. Kaszlikowski, and L.C. Kwek, quant-ph/0702019 @ www.arxiv.org.
- [23] C. Di Franco, M. Paternostro, and M.S. Kim, arXiv:0805.4365 @ www.arxiv.org.
- [24] C. Albanese, M. Christandl, N. Datta, and A. Ekert, *Phys. Rev. Lett.* **93**, 230502 (2004).
- [25] D. G. Angelakis, M. F. Santos, and S. Bose, *Phys. Rev. A* **76**, 031805(R) (2007).
- [26] D. Loss and D. P. DiVincenzo, *Phys. Rev. A* **57**, 120 (1998).
- [27] B. E. Krane, *Nature* **393**, 133 (1998),
- [28] R. Vrijen *et al.*, *Phys. Rev. A*, **62**, 012306 (2000).
- [29] S. Katsura, *Phys. Rev* **127**, 1508 (1962).
- [30] S.C. Benjamin and S. Bose, *Phys. Rev. Lett.* **90**, 247901 (2003).
- [31] D. Burgart, private communication.
- [32] A. Kay, *Phys. Rev. A*, **73**, 032306 (2006).