

Semi-inclusive processes at low and high transverse momentum

Alessandro Bacchetta¹, Daniel Boer², Markus Diehl³, and Piet J. Mulders²

¹*Jefferson Lab, 12000 Jefferson Ave, Newport News, VA 23606, USA*

²*Vrije Universiteit Amsterdam, 1081 HV Amsterdam, The Netherlands*

³*Deutsches Elektronen-Synchrotron DESY, 22603 Hamburg, Germany*

This talk reports on recent work where we studied the connection between the description of semi-inclusive DIS at high transverse momentum (based on collinear factorization) and low transverse momentum (based on transverse-momentum-dependent factorization). We used power counting to determine the leading behavior of the structure functions at intermediate transverse momentum in the two descriptions. When the power behaviors are different, two distinct mechanisms are present and there can be no matching between them. When the power behavior is the same, the two descriptions must match. An explicit calculation however shows that for some observables this is not the case, suggesting that the transverse-momentum-dependent-factorization description beyond leading twist is incomplete.

1. Introduction

The cross section for polarized semi-inclusive DIS can be written in terms of 18 structure functions.¹ Each of them depends, among other variables, on the square of the transverse momentum of the outgoing hadron, $P_{h\perp}^2$, with respect to the virtual photon direction. For theoretical considerations, it is often preferable to consider the transverse momentum $q_T^2 \approx P_{h\perp}^2/z^2$. The problem involves three scales, namely the scale of nonperturbative QCD dynamics, which we represent by the nucleon mass M , the transverse momentum q_T , and the photon virtuality Q , which we require to be large compared with M .

At high q_T ($q_T \gg M$) the structure functions can be described using collinear factorization, i.e., in terms of collinear distribution and fragmentation functions together with perturbative radiation. At low- q_T ($q_T \ll Q$) the structure functions can be described using Collins–Soper TMD factorization,^{2,3} i.e., in terms transverse-momentum-dependent (TMD) parton distribution and fragmentation functions. The low- and high- q_T domains

overlap for $M \ll q_T \ll Q$ (intermediate q_T), where both descriptions can hence be applied.

Studying the relation between the high- q_T and low- q_T regimes is important both from the theoretical and phenomenological point of view. We observe that in some cases the calculations in the two regimes have to give the same result at intermediate q_T , i.e., they have to match. If this does not occur, we can make the important conclusion that there is some flaw in the formalism. We observe in other cases that the two calculations describe different mechanisms and therefore do not have to match. Both of them have to be taken into consideration independently in the overlap region.

2. Matches and mismatches: general discussion

To assess whether the high- q_T and low- q_T calculations have to match or not in the intermediate- q_T region it is sufficient to study the power behavior of the structure functions in the two regimes.

It is important to realize that the power expansions are done in two different ways in the two descriptions. At low q_T , first we expand in $(q_T/Q)^{n-2}$ and neglect terms with n bigger than a certain value (so far, analyses have been carried out only up to $n = 3$, i.e., twist-3). To study the behavior at intermediate q_T we further expand in $(M/q_T)^k$. Conversely, at high q_T we first expand in $(M/q_T)^n$ (also in this case, analyses are available up to $n = 3$, i.e., twist-3). To study the intermediate- q_T region, we further expand in $(q_T/Q)^{k-2}$.

We can encounter two different situations. For simplicity, we will refer to them as type-I and type-II observables.

2.1. Type-I observables

Consider, e.g., a structure functions described by two contributions

$$F = A \frac{M^2}{M^2 + q_T^2} + B \frac{q_T^2}{Q^2} \frac{M^2}{M^2 + q_T^2}. \quad (1)$$

At low transverse momentum, term B is neglected from the very beginning because it is of order $(q_T/Q)^2$ (twist-4). The remaining term is

$$F_{\text{low}}^{\text{twist-2}} = A \frac{M^2}{M^2 + q_T^2} = A \frac{M^2}{q_T^2} + \mathcal{O}\left(\frac{M^4}{q_T^4}\right), \quad (2)$$

where the second step identifies the leading term at $q_T \gg M$.

At high transverse momentum, both terms A and B are twist-2 and are taken into consideration. However, the second term is neglected if a

further expansion in q_T/Q is performed, to study the regime of intermediate transverse momentum, i.e.,

$$F_{\text{high}}^{\text{twist-2}} = A \frac{M^2}{q_T^2} + B \frac{M^2}{Q^2} = A \frac{M^2}{q_T^2} + \mathcal{O}\left(\frac{q_T^2}{Q^2}\right). \quad (3)$$

Therefore, the *leading* terms in the two expansions are the same. In this case, the calculations at high and low transverse momentum must yield exactly the same result at intermediate transverse momentum.^{4,5} If a mismatch occurs, it means that one of the calculations is incorrect or incomplete.

2.2. Type-II observables: expected mismatches

Consider the example of a structure functions composed by two terms

$$F = A \frac{M^4}{M^4 + q_T^4} + B \frac{q_T^2}{Q^2} \frac{M^2}{M^2 + q_T^2}. \quad (4)$$

At low transverse momentum, term B is neglected from the very beginning because it is of order $(q_T/Q)^2$ (twist-4). What is left is

$$F_{\text{low}}^{\text{twist-2}} = A \frac{M^4}{M^4 + q_T^4} = A \frac{M^4}{q_T^4} + \mathcal{O}\left(\frac{M^8}{q_T^8}\right), \quad (5)$$

where in the second step we expanded in M/q_T .

At high transverse momentum, the term A is now twist-4 and it is usually neglected. Only the second term is kept and gives

$$F_{\text{high}}^{\text{twist-2}} = B \frac{q_T^2}{Q^2} \frac{M^2}{q_T^2} \quad (6)$$

In this case, if the calculations at high and low transverse momentum are performed at their respective leading twist, they correspond to two different contributions to the cross section and will not lead to the same result at intermediate transverse momentum. In order to “match”, the calculations would have to be carried out in both regimes up to the sub-subleading order. We could call this situation an “expected mismatch”, since it is simply due to the difference between the two expansions.

3. Matches and mismatches: semi-inclusive DIS case

In Tab. 1 we list the power behavior of the structure functions at intermediate transverse momentum, as obtained from the limits of the low- q_T and high- q_T calculation. For details of the calculation, we refer to Ref. 6. In the names of the structure functions, the first and second subscript respectively

Table 1. Behavior of SIDIS structure functions in the region $M \ll q_T \ll Q$. Empty fields indicate that no calculation is available (in this case, twist 4 indicates observables that are zero when calculated up to twist-three accuracy). Yes/no in parentheses: expected answers based on analogy, rather than actual calculation.

observable	low- q_T calculation		high- q_T calculation		power match	exact match
	twist	power	twist	power		
$F_{UU,T}$	2	$1/q_T^2$	2	$1/q_T^2$	yes	yes
$F_{UU,L}$	4		2	$1/Q^2$		
$F_{UU}^{\cos \phi_h}$	3	$1/(Q q_T)$	2	$1/(Q q_T)$	yes	no
$F_{UU}^{\cos 2\phi_h}$	2	$1/q_T^4$	2	$1/Q^2$	no	
$F_{LU}^{\sin \phi_h}$	3	$1/(Q q_T)$	2	$1/(Q q_T)$	yes	(no)
$F_{UL}^{\sin \phi_h}$	3	$1/(Q q_T)$			(yes)	(no)
$F_{UL}^{\sin 2\phi_h}$	2	$1/q_T^4$			(no)	
F_{LL}	2	$1/q_T^2$	2	$1/q_T^2$	yes	yes
$F_{LL}^{\cos \phi_h}$	3	$1/(Q q_T)$	2	$1/(Q q_T)$	yes	no
$F_{UT,T}^{\sin(\phi_h - \phi_S)}$	2	$1/q_T^3$	3	$1/q_T^3$	yes	yes
$F_{UT,L}^{\sin(\phi_h - \phi_S)}$	4		3	$1/(Q^2 q_T)$		
$F_{UT}^{\sin(\phi_h + \phi_S)}$	2	$1/q_T^3$	3	$1/q_T^3$	yes	(yes)
$F_{UT}^{\sin(3\phi_h - \phi_S)}$	2	$1/q_T^3$	3	$1/(Q^2 q_T)$	no	
$F_{UT}^{\sin \phi_S}$	3	$1/(Q q_T^2)$	3	$1/(Q q_T^2)$	yes	(no)
$F_{UT}^{\sin(2\phi_h - \phi_S)}$	3	$1/(Q q_T^2)$	3	$1/(Q q_T^2)$	yes	(no)
$F_{LT}^{\cos(\phi_h - \phi_S)}$	2	$1/q_T^3$			(yes)	(yes)
$F_{LT}^{\cos \phi_S}$	3	$1/(Q q_T^2)$			(yes)	(no)
$F_{LT}^{\cos(2\phi_h - \phi_S)}$	3	$1/(Q q_T^2)$			(yes)	(no)

specifies the polarization of the beam and the target. When present, the third subscript refers to the polarization of the photon.

In summary, the calculation at high q_T is done using standard collinear factorization, as done in, e.g., Ref. 7,8 and in Ref. 9 for the subleading-twist sector. To obtain the power behavior at intermediate q_T , we need to perform an expansion in q_T/Q . The calculation has no fundamental difficulties and allows us to fill in the third column of Tab. 1. The blank entries correspond to the structure functions that have not yet been computed in the high-transverse-momentum regime.

The calculation at low q_T is done using TMD factorization.^{2,3} The behavior of the TMD functions at intermediate transverse momentum can be calculated perturbatively by considering diagrams as the ones depicted in Fig. 1. We calculated the power behavior of all twist-2 and twist-3 TMD functions, which allowed us to fill in the second column of Tab. 1. Two struc-




Fig. 1. Example diagrams for the calculation of the high- p_T behavior TMD parton distribution functions. Φ_A^α represent the quark-gluon-quark correlator. The dashed lines represent the final-state cut.

ture functions cannot be calculated as they require twist-4 contributions, which are beyond the current limits of the TMD factorization framework.

The fourth column of Tab. 1 is obtained by comparing the second and third column. The structure functions with a “yes” are type-I observables, those with a “no” are type-II. The values in parentheses are expectations based on analogy with similar structure functions, since the high- q_T calculations are not available.

Beside studying the power behavior, we also calculated the explicit form of some of the TMD functions at intermediate transverse momentum, namely the ones requiring only the evaluation of diagrams analogous to that of Fig. 1(a). A calculation of the Sivers function, requiring the evaluation of diagrams like that of Fig. 1(b), was already performed in Refs. 5,10.

The explicit calculations allows us to check if for type-I observables the explicit expressions obtained from high and low transverse momentum exactly match or not. The results are listed in the fifth column of Tab. 1. The entries in parentheses are conjectures based on analogy rather than actual calculation.

3.1. Type-I structure functions

For type-I structure functions (“yes” in the column “power match”), we know from power counting that the two calculations describe the same physics and should therefore exactly match. In these cases, the high- q_T calculation corresponds to the perturbative tail of the low- q_T effect. The two mechanisms need not be distinguished. Using resummation it should be possible to construct expressions for these observables that are valid at any q_T , as was done for the Drell-Yan analog of $F_{UU,T}$ in Ref. 4.

Only five of these structure functions have been calculated explicitly: $F_{UU,T}$, F_{LL} and $F_{UT,T}^{\sin(\phi_h - \phi_S)}$ (Sivers structure function) present an exact

matching,^{5,10} while in our work we showed that $F_{UU}^{\cos \phi_h}$ and $F_{LL}^{\cos \phi_h}$ do not match. In analogy to these results, we expect that also $F_{UT}^{\sin(\phi_h + \phi_S)}$ (Collins structure function) and $F_{LT}^{\cos(\phi_h - \phi_S)}$ will match exactly, while problems will occur with all the others, since they are twist-3 in the low- q_T regime, and the TMD factorization formalism is probably complete only at twist 2.

The structure function $F_{UT,T}^{\sin(\phi_h - \phi_S)}$, related to the Sivers function, is an example of a match between high- and low- q_T . Some of the consequences of the calculation are:

- the leading (twist-3, in this case) contribution of the high- q_T calculation corresponds to the tail of the Sivers function at intermediate q_T , it is not a competing effect and should not be summed to the Sivers function;
- it is conceivable to construct an expression that extends the high- q_T calculation to $q_T \approx M$, through a smooth merging into the Sivers function;
- since the structure function falls as $1/q_T^3$, it is safe to use q_T -weighted asymmetries to extract the Sivers function.

As an example of a mismatch we consider the structure function $F_{UU}^{\cos \phi_h}$, related to the Cahn effect. We show in this case the main steps of the calculation to explain the nature of the problem. In the low- q_T formalism, the expression for this observable is¹

$$F_{UU}^{\cos \phi_h} = \frac{2M}{Q} \mathcal{C} \left[-\frac{\hat{\mathbf{h}} \cdot \mathbf{k}_T}{M_h} \left(xh H_1^\perp + \frac{M_h}{M} f_1 \frac{\tilde{D}^\perp}{z} \right) - \frac{\hat{\mathbf{h}} \cdot \mathbf{p}_T}{M} \left(x f^\perp D_1 + \frac{M_h}{M} h_1^\perp \frac{\tilde{H}}{z} \right) \right], \quad (7)$$

where the convolution means

$$\mathcal{C}[wfD] = \sum_a x e_a^2 \int d^2 \mathbf{p}_T d^2 \mathbf{k}_T d^2 \mathbf{l}_T \delta^{(2)}(\mathbf{p}_T - \mathbf{k}_T + \mathbf{l}_T + \mathbf{q}_T) \times w(\mathbf{p}_T, \mathbf{k}_T) f^a(x, p_T^2) D^a(z, k_T^2) U(l_T^2). \quad (8)$$

The term U denotes the so-called soft factor. It is obtained in the factorization proof for twist-two observables. Here we assume we can use it also for twist-three observables. The terms with h_1^\perp and H_1^\perp fall off as $1/p_T^3$ or $1/k_T^3$ and are power suppressed compared to the terms with f^\perp and \tilde{D}^\perp when $q_T \gg M$. For intermediate q_T we therefore have

$$F_{UU}^{\cos \phi_h} = -\frac{2q_T}{Q} \sum_a x e_a^2 \left[x f^{\perp a}(x, q_T^2) \frac{D_1^a(z)}{z^2} - f_1^a(x) \frac{\tilde{D}^{\perp a}(z, q_T^2)}{z} \right] \quad (9)$$

at leading power. In this case there is no leading contribution from the soft factor taken at large transverse momentum. The tail of the functions at $q_T \gg M$ can be calculated perturbatively and yields

$$\begin{aligned}
x f^{\perp q}(x, p_T^2) &= \frac{\alpha_s}{2\pi^2} \frac{1}{2\mathbf{p}_T^2} \left[\frac{L(\eta^{-1})}{2} f_1^q(x) + (P'_{qq} \otimes f_1^q + P'_{gq} \otimes f_1^g)(x) \right], \\
\frac{\tilde{D}^{\perp q}(z, k_T^2)}{z} &= -\frac{\alpha_s}{2\pi^2} \frac{1}{2z^2 \mathbf{k}_T^2} \left[\frac{L(\eta_h^{-1})}{2} D_1^q(z) - 2C_F D_1^q(z) \right. \\
&\quad \left. + (D_1^q \otimes P'_{qq} + D_1^g \otimes P'_{gq})(z) \right], \quad (10)
\end{aligned}$$

where $L(y) = 2C_F \ln y - 3C_F$, and P'_{qq} , P'_{gq} , P'_{gg} are kernels specific to the functions under consideration. The parameters η and η_h are related to the choice of a nonlightlike gauge (Wilson line) in the calculation of the functions and fulfill the relation $\sqrt{\eta\eta_h} = q_T^2/Q^2$. Putting these ingredients together we arrive at

$$\begin{aligned}
F_{UU}^{\cos \phi_h} &= -\frac{1}{Q q_T} \frac{\alpha_s}{2\pi^2 z^2} \sum_a x e_a^2 \left[f_1^a(x) D_1^a(z) L\left(\frac{Q^2}{q_T^2}\right) \right. \\
&\quad \left. + f_1^a(x) (D_1^a \otimes P'_{qq} + D_1^a \otimes P'_{gq})(z) \right. \\
&\quad \left. + (P'_{qq} \otimes f_1^a + P'_{gq} \otimes f_1^g)(x) D_1^a(z) - 2C_F f_1^a(x) D_1^a(z) \right]. \quad (11)
\end{aligned}$$

This expression differs from the one obtained at high q_T by the last term $2C_F f_1^a(x) D_1^a(z)$. At this point we are forced to conclude that the description of twist-3 structure functions is incomplete in the TMD-factorization formalism. However, it is interesting to note that adding a term $f_1^a(x) D_1^a(z) z^{-2} U(q_T^2)/2$ within brackets in Eq. (9) would be sufficient to cure this problem. It is however not clear how such an expression would be obtained from a factorized formula.

3.2. Type-II structure functions

For type-II structure functions (“no” in the column “power match”) the low- q_T and high- q_T calculations at leading order pick up two different components of the full structure function. They therefore describe two different mechanisms and do not match.

An example of a type-II observable is the structure function $F_{UU}^{\cos 2\phi_h}$, related at low q_T to the Boer–Mulders function.¹¹ Some studies of this structure functions have recently appeared.^{12,13} However, some considerations have to be kept in mind:

- the leading contribution from the high- q_T calculation (often referred to as a pQCD or radiative correction) is a competing effect that has to be taken into account;
- it is at present not possible to construct an expression that extends the high- q_T calculation to $q_T \approx M$, since this requires a smooth merging into unknown twist-4 contributions in TMD factorization;
- Using q_T -weighted asymmetries to extract the Boer–Mulders function is *not* a good idea, since the high- q_T mechanism dominates the observable;
- a solution to the above problems could be to consider observables that are least sensitive to the effect of radiative corrections, for instance by considering specific combinations of structure functions.

We stress that the above considerations apply not only to semi-inclusive DIS, but also to Drell–Yan and e^+e^- annihilation.¹⁴ Drell–Yan data have been already used to extract¹⁵ the Boer–Mulders function, without taking into account radiative corrections, while the extraction¹⁶ of the Collins function from e^+e^- relies on the cancellation of radiative effects through the construction of suitable experimental observables.^{14,17}

References

1. A. Bacchetta, M. Diehl, K. Goeke, A. Metz, P. J. Mulders and M. Schlegel, *JHEP* **02**, 093 (2007), hep-ph/0611265.
2. J. C. Collins and D. E. Soper, *Nucl. Phys.* **B193**, 381 (1981).
3. X. Ji, J.-P. Ma and F. Yuan, *Phys. Rev.* **D71**, 034005 (2005), hep-ph/0404183.
4. J. C. Collins, D. E. Soper and G. Sterman, *Nucl. Phys.* **B250**, 199 (1985).
5. X. Ji, J.-W. Qiu, W. Vogelsang and F. Yuan, *Phys. Rev. Lett.* **97**, 082002 (2006), hep-ph/0602239.
6. A. Bacchetta, D. Boer, M. Diehl and P. J. Mulders, *JHEP* **08**, 023 (2008), arXiv:0803.0227 [hep-ph].
7. A. Mendez, *Nucl. Phys.* **B145**, 199 (1978).
8. Y. Koike, J. Nagashima and W. Vogelsang, *Nucl. Phys.* **B744**, 59 (2006), arXiv:hep-ph/0602188.
9. H. Eguchi, Y. Koike and K. Tanaka, *Nucl. Phys.* **B763**, 198 (2007), arXiv:hep-ph/0610314.
10. Y. Koike, W. Vogelsang and F. Yuan, *Phys. Lett.* **B659**, 878 (2008), arXiv:0711.0636 [hep-ph].
11. D. Boer and P. J. Mulders, *Phys. Rev.* **D57**, 5780 (1998), hep-ph/9711485.
12. V. Barone, A. Prokudin and B.-Q. Ma (2008), arXiv:0804.3024 [hep-ph].
13. B. Zhang, Z. Lu, B.-Q. Ma and I. Schmidt (2008), arXiv:0807.0503 [hep-ph].
14. D. Boer (2008), arXiv:0804.2408 [hep-ph].
15. B. Zhang, Z. Lu, B.-Q. Ma and I. Schmidt, *Phys. Rev.* **D77**, 054011 (2008), arXiv:0803.1692 [hep-ph].
16. M. Anselmino *et al.*, *Phys. Rev.* **D75**, 054032 (2007), hep-ph/0701006.
17. K. Abe *et al.*, *Phys. Rev. Lett.* **96**, 232002 (2006), hep-ex/0507063.