

# COLOURED QUIVER MUTATION FOR HIGHER CLUSTER CATEGORIES

ASLAK BAKKE BUAN AND HUGH THOMAS

ABSTRACT. We define mutation on coloured quivers associated to tilting objects in higher cluster categories. We show that this operation is compatible with the mutation operation on the tilting objects. This gives a combinatorial approach to tilting in higher cluster categories and especially an algorithm to determine the Gabriel quivers of tilting objects in such categories.

## INTRODUCTION

A cluster category is a certain 2-Calabi-Yau orbit category of the derived category of a hereditary abelian category. Cluster categories were introduced in [BMRRT] in order to give a categorical model for the combinatorics of Fomin-Zelevinsky cluster algebras [FZ]. They are triangulated [K] and admit (cluster-)tilting objects, which model the clusters of a corresponding (acyclic) cluster algebra [CK]. Each cluster in a fixed cluster algebra comes together with a finite quiver, and in the categorical model this quiver is in fact the Gabriel quiver of the corresponding tilting object [BMRT].

A principal ingredient in the construction of a cluster algebra is quiver mutation. It controls the exchange procedure which gives a rule for producing a new cluster variable and hence a new cluster from a given cluster. Exchange is modeled by cluster categories in the acyclic case [BMR] in terms of a mutation rule for tilting objects, i.e. a rule for replacing an indecomposable direct summand in a tilting object with another indecomposable rigid object, to get a new tilting object. Quiver mutation describes the relation between the Gabriel quivers of the corresponding tilting objects.

Analogously to the definition of the cluster category, for a positive integer  $m$ , it is natural to define a certain  $m + 1$ -Calabi-Yau orbit category of the derived category of a hereditary abelian category. This is called the  $m$ -cluster category. Implicitly,  $m$ -cluster categories was first studied in [K], and their (cluster-)tilting objects have been studied in [ABST, F, HJ1, HJ2, IY, KR1, KR2, T, W, Z, ZZ]. Combinatorial descriptions of  $m$ -cluster categories in Dynkin type  $A_n$  and  $D_n$  are given in [BM1, BM2].

In cluster categories the mutation rule for tilting objects is described in terms of certain triangles called *exchange triangles*. By [IY] the existence of exchange triangles generalizes to  $m$ -cluster categories. It was shown in [ZZ, W] that there are

---

Both authors were supported by a STORFORSK-grant 167130 from the Norwegian Research Council; the second author was also supported by an NSERC Discovery Grant.

exactly  $m + 1$  non-isomorphic complements to an almost complete tilting object, and that they are determined by the  $m + 1$  exchange triangles defined in [IY].

The aim of this paper is to give a combinatorial description of mutation in  $m$ -cluster categories. *A priori*, one might expect to be able to do this by keeping track of the Gabriel quivers of the tilting objects. However, it is easy to see that the Gabriel quivers do not contain enough information.

We proceed to associate to a tilting object a quiver each of whose arrows has an associated colour  $c \in \{0, \dots, m\}$ . The arrows with colour 0 form the Gabriel quiver of the tilting object. We then define a mutation operation on coloured quivers and show that it is compatible with mutation of tilting objects. A consequence is that the effect of an arbitrary sequence of mutations on a tilting object in an  $m$ -cluster category can be calculated by a purely combinatorial procedure.

Our definition of a coloured quiver associated to a tilting object makes sense in any  $m + 1$ -Calabi-Yau category, such as for example those studied in [IY]. We hope that our constructions may shed some light on mutation of tilting objects in this more general setting.

In section 1, we review some elementary facts about higher cluster categories. In section 2, we explain how to define the coloured quiver of a tilting object, we define coloured quiver mutation, and we state our main theorem. In sections 3 and 4, we state some further lemmas about higher cluster categories, and we prove certain properties of the coloured quivers of tilting objects. We prove our main result in sections 5 and 6. In sections 7 and 8 we point out some applications. In section 9 we interpret our construction in terms of  $m$ -cluster complexes. In section 10, we give an alternative algorithm for computing coloured quiver mutation. Section 11 discusses the example of  $m$ -cluster categories of Dynkin type  $A_n$ , using the model developed by Baur and Marsh [BM1].

We would like to thank Idun Reiten, in conversation with whom the initial idea of this paper took shape.

## 1. HIGHER CLUSTER CATEGORIES

Let  $K$  be an algebraically closed field, and let  $\Gamma$  be a finite acyclic quiver with  $n = n_\Gamma$  vertices. Then the path algebra  $H = K\Gamma$  is a hereditary finite dimensional basic  $K$ -algebra

Let  $\text{mod } H$  be the category of finite dimensional left  $H$ -modules. Let  $\mathcal{D} = D^b(H)$  be the bounded derived category of  $H$ , and let  $[i]$  be the  $i$ 'th shift functor on  $\mathcal{D}$ . We let  $\tau$  denote the Auslander-Reiten translate, which is an autoequivalence on  $\mathcal{D}$  such that we have a bifunctorial isomorphism in  $\mathcal{D}$

$$\text{Hom}(A, B[1]) \simeq D \text{Hom}(B, \tau A). \quad (1)$$

In other words  $\nu = [1]\tau$  is a Serre functor.

Let  $G = \tau^{-1}[m]$ . The  $m$ -cluster category is the orbit category  $\mathcal{C} = \mathcal{C}_m = \mathcal{D} / \tau^{-1}[m]$ . The objects in  $\mathcal{C}$  are the objects in  $\mathcal{D}$ , and two objects  $X, Y$  are isomorphic in  $\mathcal{C}$  if and only if  $X \simeq G^i Y$  in  $\mathcal{D}$ . The maps are given by  $\text{Hom}_{\mathcal{C}_m}(X, Y) = \coprod_{i \in \mathbb{Z}} \text{Hom}_{\mathcal{D}}(X, G^i Y)$ . By [K], the category  $\mathcal{C}$  is triangulated and the canonical functor  $\mathcal{D} \rightarrow \mathcal{C}$  is a triangle functor. We denote therefore by  $[1]$  the suspension in  $\mathcal{C}$ .

The  $m$ -cluster category is also Krull-Schmidt and has an AR-translate  $\tau$  inherited from  $\mathcal{D}$ , such that the formula (1) still holds in  $\mathcal{C}$ . It follows that  $\nu = [1]\tau$  is a Serre functor for  $\mathcal{C}$  and that  $\mathcal{C}$  is  $m+1$ -Calabi-Yau, since  $\nu \simeq [m+1]$ .

The indecomposable objects in  $\mathcal{D}$  are of the form  $M[i]$ , where  $M$  is an indecomposable  $H$ -module and  $i \in \mathbb{Z}$ . We can choose a fundamental domain for the action of  $G = \tau^{-1}[m]$  on  $\mathcal{D}$ , consisting of the indecomposable objects  $M[i]$  with  $0 \leq i \leq m-1$ , together with the objects  $M[m]$  with  $M$  an indecomposable projective  $H$ -module. Then each indecomposable object in  $\mathcal{C}$  is isomorphic to exactly one of the indecomposables in this fundamental domain. We say that  $M[d]$  has degree  $d$ , denoted  $\delta(M[d]) = d$ . Furthermore, for an arbitrary object  $X = \coprod X_i$  in  $\mathcal{C}_m$ , we let  $\Delta_d(X) = \coprod_j X_j[-d]$  be the  $H$ -module which is the (shifted) direct sum of all summands  $X_j$  of  $X$  with  $\delta(X_j) = d$ .

In the following theorem the equivalence between (i) and (ii) is shown in [ZZ, W] and the equivalence between (i) and (iii) is shown in [Z].

**Theorem 1.1.** *Let  $T$  be an object in  $\mathcal{C}$  satisfying  $\text{Hom}_{\mathcal{C}}(T, T[i]) = 0$  for  $i = 1, \dots, m$ . Then the following are equivalent*

- (i) *If  $\text{Hom}_{\mathcal{C}}(T, U[i]) = 0$  for  $i = 1, \dots, m$  then  $U$  is in  $\text{add } T$ .*
- (ii) *If  $\text{Hom}_{\mathcal{C}}(U \amalg T, U[i]) = 0$  for  $i = 1, \dots, m$  then  $U$  is in  $\text{add } T$ .*
- (iii)  *$T$  has  $n$  indecomposable direct summands, up to isomorphism.*

Here  $\text{add } T$  denotes the additive closure of  $T$ . A (cluster-)tilting object  $T$  in an  $m$ -cluster is an object satisfying the conditions of the above Theorem. For a tilting object  $T = \coprod_{i=1}^v T_i$ , with each  $T_i$  indecomposable, and  $T_k$  an indecomposable direct summand, we call  $\bar{T} = T/T_k$  an almost complete tilting object. We let  $\text{Irr}_{\mathcal{A}}(X, Y)$  denote the  $K$ -space of irreducible maps  $X \rightarrow Y$  in a Krull-Schmidt  $K$ -category  $\mathcal{A}$ . The following crucial result is proved in [ZZ] and [W].

**Proposition 1.2.** *There are, up to isomorphism,  $m+1$  complements of an almost complete tilting object.*

Let  $T_k$  be an indecomposable direct summand in an  $m$ -cluster tilting object  $T = \bar{T} \amalg T_k$ . The complements of  $\bar{T}$  are denoted  $T_k^{(c)}$  for  $c = 0, 1, \dots, m$ , where  $T_k = T_k^{(0)}$ . By [IY], there are  $m+1$  exchange triangles

$$T_k^{(c)} \xrightarrow{f_k^{(c)}} B_k^{(c)} \xrightarrow{g_k^{(c+1)}} T_k^{(c+1)} \xrightarrow{h_k^{(c+1)}} .$$

Here the  $B_k^{(c)}$  are in  $\text{add}(T/T_k)$  and the maps  $f_k^{(c)}$  (resp.  $g_k^{(c)}$ ) are minimal left (resp. right)  $\text{add}(T/T_k)$ -approximations, and hence not split mono or split epi. Note that by minimality, the maps  $f_k^{(c)}$  and  $g_k^{(c)}$  have no proper zero summands.

## 2. COLOURED QUIVER MUTATION

We first recall the definition of quiver mutation, formulated in [FZ] in terms of skew-symmetric matrices. Let  $Q = (q_{ik})$  be a quiver with vertices  $1, \dots, n$  and with no loops or oriented two-cycles, where  $q_{ik}$  denotes the number of arrows from  $i$  to  $k$ . Let  $j$  be a vertex in  $Q$ . Then, a new quiver  $\mu_j(Q) = \tilde{Q} = (\tilde{q}_{ik})$  is defined by the following data

$$\tilde{q}_{ik} = \begin{cases} q_{ki} & \text{if } j = k \text{ or } j = i \\ \max\{0, q_{ik} - q_{ki} + q_{ij}q_{jk} - q_{kj}q_{ji}\} & \text{if } i \neq j \neq k \end{cases}$$

It is easily verified that this definition is equivalent to the one of Fomin-Zelevinsky.

Now we consider coloured quivers. Let  $m$  be a positive integer. An  $m$ -coloured (multi-)quiver  $Q$  consists of vertices  $1, \dots, n$  and coloured arrows  $i \xrightarrow{(c)} j$ , where  $c \in \{0, 1, \dots, m\}$ . Let  $q_{ij}^{(c)}$  denote the number of arrows from  $i$  to  $j$  of colour  $(c)$ .

We will consider coloured quivers with the following additional conditions.

- (I) No loops:  $q_{ii}^{(c)} = 0$  for all  $c$ .
- (II) Monochromaticity: If  $q_{ij}^{(c)} \neq 0$ , then  $q_{ij}^{(c')} = 0$  for  $c \neq c'$ .
- (III) Skew-symmetry:  $q_{ij}^{(c)} = q_{ji}^{(m-c)}$ .

We will define an operation on a coloured quiver  $Q$  satisfying the above conditions. Let  $j$  be a vertex in  $Q$  and let  $\mu_j(Q) = \tilde{Q}$  be the coloured quiver defined by

$$\tilde{q}_{ik}^{(c)} = \begin{cases} q_{ik}^{(c+1)} & \text{if } j = k \\ q_{ik}^{(c-1)} & \text{if } j = i \\ \max\{0, q_{ik}^{(c)} - \sum_{t \neq c} q_{ik}^{(t)} + (q_{ij}^{(c)} - q_{ij}^{(c-1)})q_{jk}^{(0)} + q_{ij}^{(m)}(q_{jk}^{(c)} - q_{jk}^{(c+1)})\} & \text{if } i \neq j \neq k \end{cases}$$

In an  $m$ -cluster category  $\mathcal{C}$ , for every tilting object  $T = \coprod_{i=1}^n T_i$ , with the  $T_i$  indecomposable, we will define a corresponding  $m$ -coloured quiver  $Q_T$ , as follows.

Let  $T_i, T_j$  be two non-isomorphic indecomposable direct summands of the  $m$ -cluster tilting object  $T$  and let  $r_{ij}^{(c)}$  denote the multiplicity of  $T_j$  in  $B_i^{(c)}$ . We define the  $m$ -coloured quiver  $Q_T$  of  $T$  to have vertices  $i$  corresponding to indecomposable direct summands  $T_i$ , and  $q_{ij}^{(c)} = r_{ij}^{(c)}$ . Note, in particular, that the  $(0)$ -coloured arrows are the arrows from the Gabriel quiver for the endomorphism ring of  $T$ .

By definition,  $Q_T$  satisfies condition (I). We show in Section 3 that (II) is satisfied (this also follows from [ZZ]), and in Section 4 that (III) is also satisfied.

The aim of this paper is to prove the following theorem, which is a generalization of the main result of [BMR].

**Theorem 2.1.** *Let  $T = \coprod_{i=1}^n T_i$  and  $T' = T/T_j \amalg T_j^{(1)}$  be  $m$ -tilting objects, where there is an exchange triangle  $T_j \rightarrow B_j^{(0)} \rightarrow T_j^{(1)} \rightarrow$ . Then  $Q_{T'} = \mu_j(Q_T)$ .*

In the case  $m = 1$  the coloured quiver of a tilting object  $T$  is given by  $q_{ij}^{(0)} = \bar{q}_{ij}$  and  $q_{ij}^{(1)} = \bar{q}_{ji}$  where  $\bar{q}_{ij}$  denotes the number of arrows in the Gabriel quiver of  $T$ . Then coloured mutation of the coloured quiver corresponds to FZ-mutation of the Gabriel quiver.

**Example:**  $A_3$ ,  $m = 2$

Let  $\Gamma$  be  $A_3$  with linear orientation, i.e. the quiver  $1 \leftarrow 2 \rightarrow 3$ .



(b) *We have that*

$$\delta(T_i^{(c+1)}) - \delta(T_i^{(c)}) \begin{cases} = 1 & \text{if } \delta(T_i^{(c)}) = m \\ \leq 1 & \text{if } \delta(T_i^{(c)}) \notin \{m-1, m\} \\ \leq 2 & \text{if } \delta(T_i^{(c)}) = m-1 \end{cases}$$

- (c) *The distribution of degrees of complements is one of the following*
- *there is exactly one complement of each degree, or*
  - *there is no complement of degree  $m$ , two complements in one degree  $d \neq m$ , and exactly one complement in all degrees  $\neq d, m$ .*
- (d) *If  $\text{Hom}(T_i^{(c)}, T_i^{(c')}) \neq 0$ , then  $c' \in \{c, c+1, c+2\}$ .*
- (e) *For  $t \in \{1, \dots, m\}$  we have*

$$\text{Hom}(T_i^{(c)}, T_i^{(c')}[t]) = \begin{cases} K & \text{if } c' - c + t = 0 \pmod{m+1} \\ 0 & \text{else} \end{cases}$$

*Proof.* (a) follows from the fact that  $\text{Hom}_H(X, X) = K$  for exceptional objects and the definition of maps in a  $m$ -cluster category.

(b) follows from the fact that  $\text{Hom}(T_i^{(c+1)}, T_i^{(c)}[1]) \neq 0$ , since in the exchange triangles, the  $f_i^{(c)}$  are not split mono and (c) follows from (b).

Considering the two different possible distributions of complements, we obtain from (c) that if  $m \geq 3$  and  $c' \geq c+3$  and  $c' \neq c-1$ , then  $\text{Hom}(T_i^{(c)}, T_i^{(c')}) = 0$ . Consider the case  $c' = c-1$ . We can assume  $m > 2$ , since else the statement is void. Hence we can clearly assume that  $\delta(T_i^{(c)}) = \delta(T_i^{(c-1)})$ . There is an exchange triangle induced from an exact sequence in  $\text{mod } H$ ,

$$T_i^{(c-1)} \rightarrow B_i^{(c-1)} \rightarrow T_i^{(c)} \rightarrow T_i^{(c-1)}[1].$$

It is clear that  $\text{Hom}(T_i^{(c-1)}[1], T_i^{(c-1)}) = 0$ , since  $m > 2$ . We claim that also  $\text{Hom}(B_i^{(c-1)}, T_i^{(c-1)}) = 0$ . This holds since  $B_i^{(c-1)} \amalg T_i^{(c-1)}$  is a partial tilting object in  $H$ , and so there are no cycles in the endomorphism ring, by Lemma 3.1. Hence also  $\text{Hom}(T_i^{(c)}, T_i^{(c-1)}) = 0$  follows, and this finishes the proof for (d).

For (e) we first apply  $\text{Hom}(T_i^{(c+1)}, \cdot)$  to the exchange triangle

$$T_i^{(c)} \rightarrow B_i^{(c)} \rightarrow T_i^{(c+1)} \rightarrow$$

and consider the corresponding long-exact sequence, to obtain that

$$\text{Hom}(T_i^{(c+1)}, T_i^{(c)}[t]) = \begin{cases} K & \text{if } t = 1 \\ 0 & \text{if } t = 0 \text{ or } t \in \{2, \dots, m\} \end{cases}.$$

Now consider  $\text{Hom}(T_i^{(c+u)}, T_i^{(c)}[v])$ . When  $0 < v \leq u \leq m$ , we have that

$$\begin{aligned} \text{Hom}(T_i^{(c+u)}, T_i^{(c)}[v]) &\simeq \text{Hom}(T_i^{(c+u+1)}, T_i^{(c)}[v+1]) \simeq \\ &\text{Hom}(T_i^{(c-1)}, T_i^{(c)}[v+m-u]) \simeq \text{Hom}(T_i^{(c)}, T_i^{(c-1)}[1+u-v]). \end{aligned}$$

When  $m \geq v > u \geq 0$ , we have that

$$\mathrm{Hom}(T_i^{(c+u)}, T_i^{(c)}[v]) \simeq \mathrm{Hom}(T_i^{(c+u-1)}, T_i^{(c)}[v-1]) \simeq \mathrm{Hom}(T_i^{(c)}, T_i^{(c)}[v-u]).$$

Combining these facts, (e) follows.  $\square$

**Lemma 3.3.** *The following statements are equivalent*

- (a)  $\mathrm{Hom}(T_i^{(1)}, T_j^{(1)}[1]) = 0$
- (b)  $T_j$  is not a direct summand in  $B_i^{(m)}$
- (c)  $T_i$  is not a direct summand in  $B_j^{(0)}$

Furthermore,  $\mathrm{Hom}(T_i^{(c)}, T_j^{(1)}[1]) = 0$  for  $c \neq 1$ .

*Proof.* Note that  $r_{ji}^{(0)} = r_{ij}^{(m)} = \dim \mathrm{Irr}_{\mathrm{add}T}(T_j, T_i)$ , so (b) and (c) are equivalent. Consider the exact sequence

$$\begin{aligned} \mathrm{Hom}(T_i^{(c)}, T_j^{(0)}[1]) \rightarrow \mathrm{Hom}(T_i^{(c)}, B_j^{(0)}[1]) \rightarrow \\ \mathrm{Hom}(T_i^{(c)}, T_j^{(1)}[1]) \rightarrow \mathrm{Hom}(T_i^{(c)}, T_j^{(0)}[2]) \rightarrow \end{aligned}$$

coming from applying  $\mathrm{Hom}(T_i^{(c)}, \cdot)$  to the exchange triangle

$$T_j^{(0)} \rightarrow B_j^{(0)} \rightarrow T_j^{(1)}.$$

The first and fourth terms are always zero. Using 3.2(e) we get that the second term (and hence the third) is non-zero if and only if  $c = 1$  and  $T_i$  is a direct summand in  $B_j^{(0)}$ .  $\square$

**Lemma 3.4.** ([IY, ZZ]) *For  $0 \leq l \leq m$ , the composition*

$$\gamma_k^{(v,l)} = h_k^{(v)} \circ h_k^{(v-1)}[1] \circ h_k^{(v-2)}[2] \circ \dots \circ h_k^{(v-l+1)}[l-1]: T_k^{(v)} \rightarrow T_k^{(v-l)}[l]$$

*is non-zero and a basis for  $\mathrm{Hom}(T_k^{(v)}, T_k^{(v-l)}[l])$ .*

*Proof.* For  $m = 1$ , see [BMRRT]. Assume  $m \geq 2$ . For the first claim see [IY], while the second claim then follows from Lemma 3.2(e).  $\square$

We include an independent proof of the following crucial property.

**Proposition 3.5.** [ZZ]  $B_k^{(u)}$  and  $B_k^{(v)}$  has no common non-zero direct summands whenever  $u \neq v$ .

*Proof.* When  $m = 1$ , this is proved in [BMR]. Assume  $m > 1$ . We consider two cases,  $|u - v| = 1$  or  $|u - v| > 1$ .

Consider first the case  $|u - v| = 1$ . Without loss of generality we can assume  $u = 0$  and  $v = 1$ , and that  $\delta(T_k^{(0)}) = 0$ . Assume that there exists a (non-zero) indecomposable  $T_x$ , which is a direct summand in  $B_k^{(0)}$  and in  $B_k^{(1)}$ . We have that  $\delta(T_k^{(1)}) \in \{0, 1\}$  by Lemma 3.2(b). Assume first  $\delta(T_k^{(1)}) = 0$ . Then the exchange triangle

$$T_k^{(0)} \rightarrow B_k^{(0)} \rightarrow T_k^{(1)} \rightarrow$$

is induced from the degree 0 part of the derived category, and hence from an exact sequence in  $\text{mod } H$ . Then the endomorphism ring of the partial tilting module  $T_x \amalg T_k^{(1)}$  has a cycle, which is a contradiction to Lemma 3.1. Assume now that  $\delta(T_k^{(1)}) = 1$ . Then  $\delta(T_k^{(2)}) \in \{0, 1, 2\}$ , where 0 can only occur if  $m = 2$ . If  $\delta(T_k^{(2)}) \in \{1, 2\}$ , then clearly  $\delta(T_x) = 1$ , and hence the partial tilting module  $T_k^{(1)} \amalg T_x$  contains a cycle, which is a contradiction. Assume that  $\delta(T_k^{(2)}) = 0$  (and hence  $m = 2$ ). Then  $\delta(T_x) \in \{0, 1\}$ . If  $\delta(T_x) = 1$ , we get a contradiction as in the previous case. If  $\delta(T_x) = 0$ , consider the exchange triangle

$$T_k^{(2)} \rightarrow B_k^{(2)} \rightarrow T_k^{(0)} \rightarrow$$

which is induced from an exact sequence in  $\text{mod } H$ . Hence there is a *non-zero* map  $T_x \rightarrow B_k^{(2)}$  obtained by composing  $T_x \rightarrow T_k^{(2)}$  with the monomorphism  $T_k^{(2)} \rightarrow B_k^{(2)}$ , and thus there are cycles in the endomorphism ring of the partial tilting module  $T_x \amalg B_k^{(2)} \amalg T_k^{(0)}$ , a contradiction. This finishes the case with  $|u - v| = 1$ .

Assume now that  $|u - v| > 1$ . Then we have  $m > 2$ . Since  $\text{Hom}(T_k^{(v)}, T_x) \neq 0$  and  $\text{Hom}(T_k^{(u)}, T_x) \neq 0$ , we have by Lemma 3.2(c) that  $|v - u| \leq 2$ . So without loss of generality we can assume  $v = u - 2$ . Assume that  $\delta(T_k^{(u)}) = 0$ . Then  $\delta(T_k^{(v)}) = m - 1$  using Lemma 3.2(c) and the fact that  $\text{Hom}(T_k^{(v)}, T_x) \neq 0$ . Then also  $\delta(T_k^{(v)}) \leq m$ . But  $\text{Hom}(T_x, T_k^{(v+1)}) \neq 0$ , so  $\delta(T_x) \leq m$ , contradicting the fact that  $\text{Hom}(T_k^{(u)}, T_x) \neq 0$ .  $\square$

**Corollary 3.6.**  $Q_T$  satisfies condition (II).

#### 4. SYMMETRY

Let  $T = \tilde{T} \amalg T_i \amalg T_j$  be a tilting object. In this section we show that the coloured quiver  $Q_T$  satisfies condition (III).

**Proposition 4.1.** *With the notation of the previous section, we have  $r_{ji}^{(c)} = r_{ij}^{(m-c)}$ .*

*Proof.* By Lemma 3.3 we only need to consider the case  $c \notin \{0, m\}$ . It is enough to show that  $r_{ji}^{(c)} \leq r_{ij}^{(m-c)}$ .

We first prove

**Lemma 4.2.** *Let  $\alpha: T_j^{(c)} \rightarrow T_i$  be irreducible in  $\text{add}((T/T_j) \amalg T_j^{(c)})$ . Then the composition  $\alpha[-c] \circ \gamma_i^{(0,c)}[-c]: T_j^{(c)}[-c] \rightarrow T_i^{(m-c+1)}$  is non-zero.*

*Proof.* We have already assumed  $c \neq 0$ . Assume

$$\alpha[-c] \circ h_i^{(0)}[-c]: T_j^{(c)}[-c] \rightarrow T_i^{(0)}[-c] \rightarrow T_i^{(m)}[-c+1]$$

is zero. This means that  $T_j^{(c)} \rightarrow T_i$  must factor through  $B_i^{(m)} \xrightarrow{g_i^{(0)}} T_i$ . Since  $T_i$  is by assumption a summand in  $B_j^{(c)}$ , we have that  $T_i$  is not a summand in  $B_j^{(0)}$  by Proposition 3.5. Since  $r_{ij}^{(m)} = r_{ji}^{(0)} = 0$ , we have that  $T_j$  is not a direct summand in

$B_i^{(m)}$ . This means that  $\alpha$  is not irreducible in  $\text{add}((T/T_j) \amalg T_j^{(c)})$ , a contradiction. So  $\alpha[-c] \circ h_i^{(0)}[-c]: T_j^{(c)}[-c] \rightarrow T_i^{(m)}[-c+1]$  is non-zero.

Assume  $c > 1$ . If the composition  $\alpha[-c] \circ h_i^{(0)}[-c] \circ h_i^{(m)}[-c+1]$  is zero, then  $\alpha[-c] \circ h_i^{(0)}[-c]$  factors through

$$B_i^{(m-1)}[-c+1] \rightarrow T_i^{(m)}[-c+1].$$

We claim that  $\text{Hom}(T_j^{(c)}[-c], B_i^{(m-1)}[-c+1]) \simeq \text{Hom}(T_j^{(c)}, B_i^{(m-1)}[1]) = 0$ . This clearly holds if  $T_j$  is not a summand of  $B_i^{(m-1)}$ . In addition we have that  $\text{Hom}(T_j^{(c)}, T_j[1]) = 0$  since  $c > 1$ , using Lemma 3.2(e). This is a contradiction, and this argument can clearly be iterated to see that  $\alpha[-c] \circ \gamma_i^{(0,c)}[-c]: T_j^{(c)}[-c] \rightarrow T_i^{(m-c+1)}$  is non-zero, using Lemma 3.2(e).  $\square$

We now show that any irreducible map  $\alpha: T_j^{(c)} \rightarrow T_i$  gives rise to an irreducible map  $\delta: T_i^{(m-c)} \rightarrow T_j$ .

Consider the composition

$$B_j^{(c-1)}[-c] \xrightarrow{g_j^{(c)}[-c]} T_j^{(c)}[-c] \longrightarrow T_i^{(m-c+1)}.$$

Since  $T_i$  is a summand in  $B_j^{(c)}$  by assumption, it is not a summand in  $B_j^{(c-1)}$ . Thus,  $B_j^{(c-1)}$  is in  $\text{add } \tilde{T}$ . Since  $\text{Hom}(X, T_i^{(m-c+1)}[c]) = 0$  for any  $X$  in  $\text{add } \tilde{T}$ , the composition vanishes.

Using the exchange triangle

$$B_j^{(c-1)}[-c] \xrightarrow{g_j^{(c)}[-c]} T_j^{(c)}[-c] \xrightarrow{h_j^{(c)}[-c]} T_j^{(c-1)}[-c+1],$$

we see that  $\alpha[-c] \circ \gamma_i^{(0,c)}[-c]: T_j^{(c)}[-c] \rightarrow T_i^{(m-c+1)}$  factors through the map  $T_j^{(c)}[-c] \xrightarrow{h_j^{(c)}[-c]} T_j^{(c-1)}[-c+1]$ , i.e. there is a commutative diagram

$$\begin{array}{ccccc} B_j^{(c-1)}[-c] & \xrightarrow{g_j^{(c)}[-c]} & T_j^{(c)}[-c] & \xrightarrow{h_j^{(c)}[-c]} & T_j^{(c-1)}[-c+1] & \longrightarrow \\ & & \downarrow & \swarrow \phi_1 & & \\ & & T_i^{(m-c+1)} & & & \end{array}$$

Similarly, using the exchange triangle

$$B_j^{(c-2)}[-c+1] \xrightarrow{g_j^{(c-1)}[-c+1]} T_j^{(c-1)}[-c+1] \xrightarrow{h_j^{(c-1)}[-c+1]} T_j^{(c-2)}[-c+2]$$

we obtain a map  $\phi_2: T_j^{(c-2)}[-c+2] \rightarrow T_i^{(m-c+1)}$

Repeating this argument  $c$  times we obtain a map  $\phi_c: T_j \rightarrow T_i^{(m-c+1)}$ , such that  $\gamma_j^{(c,c)}[-c] \circ \phi_c = \alpha[-c] \circ \gamma_i^{(0,c)}$ .

$$\begin{array}{ccc}
T_j^{(c)}[-c] & \xrightarrow{\quad} & T_i^{(m-c+1)} \\
\downarrow h_j^{(c)}[-c] & \nearrow \phi_1 & \\
T_j^{(c-1)}[-c+1] & \nearrow \phi_2 & \\
\downarrow h_j^{(c-1)}[-c+1] & \nearrow \phi_c & \\
T_j^{(c-2)}[-c+2] & & \\
\downarrow h_j^{(c-2)}[-c+2] & & \\
\vdots & & \\
T_j & & 
\end{array}$$

We claim that

**Lemma 4.3.** *There is a map  $\beta: T_j \rightarrow T_i^{(m-c+1)}$ , such that  $\gamma_j^{(c,c)}[-c] \circ \beta = \alpha[-c] \circ \gamma_i^{(0,c)}$ , and such that  $\beta$  is irreducible in  $\text{add}((T/T_i) \amalg T_i^{(m-c+1)})$ .*

*Proof.* Let

$$T_j \begin{pmatrix} \psi' & \psi'' \\ \longrightarrow & \end{pmatrix} (T_i^{(m-c+1)})' \amalg \tilde{T}'$$

be a minimal left  $\text{add}(T_i^{(m-c+1)} \amalg \tilde{T})$ -approximation, with  $\tilde{T}'$  in  $\text{add} \tilde{T}$  and  $(T_i^{(m-c+1)})'$  in  $\text{add} T_i^{(m-c+1)}$ . Let  $\phi_c$  be as above, and factor it as

$$T_j \begin{pmatrix} \psi' & \psi'' \\ \longrightarrow & \end{pmatrix} (T_i^{(m-c+1)})' \amalg \tilde{T}' \xrightarrow{\begin{pmatrix} \epsilon' \\ \epsilon'' \end{pmatrix}} T_i^{(m-c+1)}.$$

Since  $\gamma_j^{(c,c)}$  factors through  $T_j^{(1)}[-1]$ , we have that  $\gamma_j^{(c,c)}[-c]\psi'' = 0$ , so we have

$$\gamma_j^{(c,c)}[-c](\psi'\epsilon' + \psi''\epsilon'') = \gamma_j^{(c,c)}[-c]\psi'\epsilon'.$$

Hence, let us let  $\beta = \psi'\epsilon'$  and since the summands in  $\epsilon'$  are isomorphisms, it is clear that  $\beta$  is irreducible.  $\square$

Next, assume  $\{\alpha_t\}$  is a basis for the space of irreducible maps from  $T_j^{(c)}$  to  $T_i$ . Then, by Lemma 4.2 the set  $\{\alpha_t \circ \gamma_i^{(0,c)}\}$  is also linearly independent. For each  $\alpha_t$ , consider the corresponding map  $\beta_t$ , such that  $\gamma_j^{(c,c)}[-c] \circ \beta_t = \alpha_t[-c] \circ \gamma_i^{(0,c)}$ , and which we by Lemma 4.3 can assume is irreducible. Assume a non-trivial linear

combination  $\sum k_t \beta_t$  is zero. Then also  $\sum k_t (\gamma_j^{(c,c)}[-c] \circ \beta_t) = \sum k_t \alpha_t \circ \gamma_i^{(0,c)} = 0$ . But this contradicts Lemma 4.2 since  $\sum k_t \alpha_t$  is irreducible. Hence it follows that  $\{\beta_t\}$  is also linearly independent. Hence, in the exchange triangle  $T_i^{(m-c)} \rightarrow B_i^{(m-c)} \rightarrow T_i^{(m-c+1)}$ , we have that  $T_j$  appears with multiplicity at least  $r_{ji}^{(c)}$  in  $B_i^{(m-c)}$ . So, we have that  $r_{ji}^{(c)} \leq r_{ij}^{(m-c)}$ , and the proof of the proposition is complete.  $\square$

## 5. COMPLEMENTS AFTER MUTATION

In this section we show how mutation in the vertex  $j$  affects the complements of the almost complete tilting object  $T/T_i$ . As before, let  $T = \tilde{T} \amalg T_i \amalg T_j$  be an  $m$ -tilting object, and let  $T' = T/T_j \amalg T_j^{(1)}$ .

We need to consider

$$T_i \begin{array}{c} \xrightarrow{(e)} T_j \xrightarrow{(d)} T_k \\ \xrightarrow{(c)} \end{array}$$

for all possible values of  $c, d, e$ . However, we have the following restriction on the colour of arrows.

**Proposition 5.1.** *Assume  $q_{ij}^{(e)} > 0, q_{jk}^{(0)} > 0$  and  $q_{ik}^{(c)} > 0$ . Then  $c \in \{e, e+1\}$ .*

*Proof.* Consider the exchange triangle  $T_i^{(e)} \rightarrow T_j \amalg X' \rightarrow T_i^{(e+1)} \rightarrow$ . Note that  $T_j$  is a direct summand in the middle term  $B_i^{(e)}$  by the assumption that  $q_{ij}^{(e)} > 0$ . Consider also the exchange triangle  $T_i^{(c)} \rightarrow T_k \amalg Z \rightarrow T_i^{(c+1)} \rightarrow$ . Pick an arbitrary non-zero map  $h: T_j \rightarrow T_k$ , and consider the map  $\begin{pmatrix} h & 0 \\ 0 & 0 \end{pmatrix}: T_j \amalg X' \rightarrow T_k \amalg Z$ . It suffices to show that whenever  $c \notin \{e, e+1\}$ , then  $h$  is not irreducible in  $\text{add } T$ . So assume that  $c \notin \{e, e+1\}$ . We claim that there is a commutative diagram

$$\begin{array}{ccccccc} T_i^{(e)} & \longrightarrow & T_j \amalg X' & \longrightarrow & T_i^{(e+1)} & \longrightarrow & \\ \downarrow & & \downarrow \begin{pmatrix} h & 0 \\ 0 & 0 \end{pmatrix} & & \downarrow & & \\ T_i^{(c)} & \longrightarrow & T_k \amalg Z & \longrightarrow & T_i^{(c+1)} & \longrightarrow & \end{array}$$

where the rows are the exchange triangles. The composition  $T_i^{(e)} \rightarrow T_j \xrightarrow{h} T_k \rightarrow T_i^{(c+1)}$  is zero since

- if  $c \neq e-1$   $\text{Hom}(T_i^{(e)}, T_i^{(c+1)}) = 0$  by using  $c \notin \{e, e+1\}$  and Lemma 3.2(e)
- if  $c = e-1$ , there is no non-zero composition  $T_i^{(e)} \rightarrow T_j \rightarrow T_k \rightarrow T_i^{(c+1)} = T_i^{(e)}$

Hence the leftmost vertical map exists, and then the rightmost map exists, using that  $\mathcal{C}$  is a triangulated category. Then, since  $\text{Hom}(T_i^{(e+1)}[-1], T_i^{(c)}) = 0$  by Lemma 3.2(e), there is a map  $T_j \amalg X' \rightarrow T_i^{(c)}$ , such that  $T_i^{(e)} \rightarrow T_i^{(c)} = T_i^{(e)} \rightarrow T_j \amalg X' \rightarrow T_i^{(c)}$ . Hence there is map  $T_i^{(e+1)} \rightarrow T_k \amalg Z$  such that  $T_j \amalg X' \rightarrow T_k \amalg Z = (T_j \amalg X' \rightarrow$

$T_i^{(c)} \rightarrow T_k \amalg Z) + (T_j \amalg X' \rightarrow T_i^{(e+1)} \rightarrow T_k \amalg Z)$ . By restriction we get

$$h: T_j \rightarrow T_k = (T_j \rightarrow T_i^{(c)} \rightarrow T_k) + (T_j \rightarrow T_i^{(e+1)} \rightarrow T_k). \quad (2)$$

Under the assumption  $c \notin \{e, e+1\}$  we have that  $T_i^{(e+1)} \rightarrow T_k$  cannot be irreducible in  $\text{add}((T/T_i) \amalg T_i^{(e+1)})$ . Hence  $T_i^{(e+1)} \rightarrow T_k = T_i^{(e+1)} \rightarrow B_i^{(e+1)} \rightarrow T_k$ , where  $T_k$  is not summand in  $B_i^{(e+1)}$ . Also, by Proposition 3.5 we have that  $T_j$  is not a summand in  $B_i^{(e+1)}$ . If  $T_j \rightarrow T_i^{(c)}$  was irreducible in  $\text{add}((T/T_i) \amalg T_i^{(c)})$ , then there would be an irreducible map  $T_i^{(c-1)} \rightarrow T_j$  in  $\text{add}((T/T_i) \amalg T_i^{(c-1)})$ , and since  $c \neq e+1$ , this does not hold, by Proposition 3.5. Hence,  $T_j \rightarrow T_i^{(c)} = T_j \rightarrow B_i^{(c-1)} \rightarrow T_i^{(c)}$ , where  $T_j$  is not a direct summand of  $B_i^{(c-1)}$ . Also by Proposition 3.5 we have that  $T_k$  is not a summand of  $B_i^{(c-1)}$ . By (2), this shows that  $h: T_j \rightarrow T_k$  is not irreducible in  $\text{add} T$ .  $\square$

Let  $T' = (T/T_j) \amalg T_j^{(1)}$ . For  $i \neq j$ , let  $(T_i^{(u)})'$  denote the complements of  $T'/T_i$ , where there are exchange triangles

$$(T_i^{(u)})' \rightarrow (B_i^{(u)})' \rightarrow (T_i^{(u+1)})' \rightarrow$$

We first want to compare  $(T_i^{(u)})'$  with  $(T_i^{(u)})$ .

**Lemma 5.2.** *Assume that  $q_{ij}^{(u)} = 0$  for  $0 \leq u < c$  and that  $q_{ij}^{(m)} = 0$ .*

- (a) *For  $u = 0, 1, \dots, c-1$ , the minimal left  $\text{add}(T/T_i)$ -approximation  $T_i^{(u)} \rightarrow B_i^{(u)}$  is also an  $\text{add}(T'/T_i)$ -approximation.*
- (b) *For  $u = 0, 1, \dots, c$ , we have  $(T_i^{(u)})' = T_i^{(u)}$ .*

*Proof.* By assumption  $T_j$  is not a direct summand in any of the  $B_i^{(u)}$ . Assume there is a map  $T_i^{(u)} \rightarrow T_j^{(1)}$  and consider the diagram

$$\begin{array}{ccccccc} T_i^{(u+1)}[-1] & \longrightarrow & T_i^{(u)} & \longrightarrow & B_i^{(u)} & \longrightarrow & \\ & & \downarrow & & & & \\ & & T_j^{(1)} & & & & \end{array}$$

Since  $\text{Hom}(T_i^{(u+1)}, T_j^{(1)}[1]) = 0$  by Lemma 3.3, we see that the map  $T_i^{(u)} \rightarrow T_j^{(1)}$  factors through  $T_i^{(u)} \rightarrow B_i^{(u)}$ . Hence the minimal left  $\text{add}(T/T_i)$ -approximation  $T_i^{(u)} \rightarrow B_i^{(u)}$  is also an  $\text{add}(T'/T_i)$ -approximation, so we have proved (a). Then (b) follows directly.  $\square$

**Lemma 5.3.** *Assume that  $e \neq m$  and there are exchange triangles*

$$T_i^{(e)} \rightarrow (T_j)^p \amalg X \rightarrow T_i^{(e+1)} \rightarrow \quad (3)$$

and

$$T_j \rightarrow (T_k)^q \amalg Y \rightarrow T_j^{(1)} \rightarrow, \quad (4)$$

where  $p = q_{ij}^{(e)} > 0$  and  $q = q_{jk}^{(0)} \geq 0$ , i.e.  $B_i^{(e)} = (T_j)^p \amalg X$  and  $B_j^{(0)} = (T_k)^q \amalg Y$ , where  $T_k$  is not isomorphic to any direct summand in  $Y$ .

- (a) The composition  $T_i^{(e)} \rightarrow (T_j)^p \amalg X \rightarrow (T_k)^{pq} \amalg Y^p \amalg X$  is a left  $\text{add}(T'/T_i)$ -approximation.  
 (b) There is a triangle

$$T_i^{(e)} \rightarrow (T_k)^{pq} \amalg Y^p \amalg X \rightarrow (T_i^{(e+1)})' \amalg C' \rightarrow$$

with  $C'$  in  $\text{add}(T/(T_i \amalg T_j))$  and  $T_i^{(e)} = (T_i^{(e)})'$ .

- (c) There is a triangle  $T_i^{(e+1)} \rightarrow (T_i^{(e+1)})' \amalg C' \rightarrow (T_j^{(1)})^p \rightarrow$ .

*Proof.* Consider an arbitrary map  $f: T_i^{(e)} \rightarrow U$  with  $U$  in  $\text{add}(T'/T_i)$ . We have that  $\text{Hom}(T_i^{(e+1)}, T_j^{(1)}[1]) = 0$ , by Lemma 3.3. Hence, by applying  $\text{Hom}(\cdot, U)$  to the triangle (3) we get that  $f$  factors through  $T_i^{(e)} \rightarrow (T_j)^p \amalg X$ . By applying  $\text{Hom}(\cdot, U)$  to the triangle (4), and using that  $\text{Hom}(T_j^{(1)}, T_j^{(1)}[1]) = 0$ , we get that  $f$  factors through  $T_i^{(e)} \rightarrow (T_j)^p \amalg X \rightarrow Y^p \amalg (T_k)^{pq} \amalg X$ . This proves (a). For (b) and (c) we use the exchange triangles (3) and (4) and the octahedral axiom to obtain the commutative diagram of triangles

$$\begin{array}{ccccc} T_i^{(e)} & \longrightarrow & (T_j)^p \amalg X & \longrightarrow & T_i^{(e+1)} & \longrightarrow \\ \parallel & & \downarrow & & \downarrow & \\ T_i^{(e)} & \longrightarrow & (T_k)^{pq} \amalg Y^p \amalg X & \longrightarrow & C & \longrightarrow \\ & & \downarrow & & \downarrow & \\ & & (T_j^{(1)})^p & \xlongequal{\quad} & (T_j^{(1)})^p & \longrightarrow \end{array}$$

By (a) the map  $T_i^{(e)} \rightarrow (T_k)^{pq} \amalg Y^p \amalg X$  is a left  $\text{add}(T'/T_i)$ -approximation, and by Lemma 5.2 we have that  $(T_i^{(e)})' = T_i^{(e)}$ . Hence  $C = (T_i^{(e+1)})' \amalg C'$ , where  $C'$  is in  $\text{add}((T_k)^{pq} \amalg Y^p \amalg X) \subset \text{add}(T/(T_i \amalg T_j))$ , and with no copies isomorphic to  $T_k$  in  $Y$ .  $\square$

Note that the induced  $\text{add}(T'/T_i)$ -approximation is in general not minimal.

**Lemma 5.4.** Assume  $e \neq m$  and  $q_{ij}^{(e)} > 0$ .

- (a) Then there is a triangle

$$(T_i^{(e+1)})' \amalg C' \xrightarrow{\alpha} B_i^{(e+1)} \amalg (T_j^{(1)})^p \rightarrow T_i^{(e+2)} \rightarrow$$

where  $\alpha$  is a minimal left  $\text{add}(T'/T_i)$ -approximation, and  $C'$  is as in Lemma 5.3.

- (b) There is an induced exchange triangle

$$(T_i^{(e+1)})' \rightarrow \frac{B_i^{(e+1)} \amalg (T_j^{(1)})^p}{\alpha(C')} \rightarrow T_i^{(e+2)} \rightarrow$$

where  $\alpha(C') \simeq C'$ .  
(c)  $(T_i^{(e+2)})' \simeq T_i^{(e+2)}$ .

*Proof.* Consider the exchange triangle

$$T_i^{(e+2)}[-1] \rightarrow T_i^{(e+1)} \rightarrow B_i^{(e+1)} \rightarrow$$

and the triangle from Lemma 5.3 (b)

$$T_i^{(e+1)} \rightarrow (T_i^{(e+1)})' \amalg C' \rightarrow (T_j^{(1)})^p \rightarrow . \quad (5)$$

Apply the octahedral axiom, to obtain the commutative diagram of triangles

$$\begin{array}{ccccccc} T_i^{(e+2)}[-1] & \longrightarrow & T_i^{(e+1)} & \longrightarrow & B_i^{(e+1)} & \longrightarrow & \\ \parallel & & \downarrow & & \downarrow & & \\ T_i^{(e+2)}[-1] & \longrightarrow & (T_i^{(e+1)})' \amalg C' & \longrightarrow & G & \longrightarrow & \\ & & \downarrow & & \downarrow & & \\ & & (T_j^{(1)})^p & \xlongequal{\quad} & (T_j^{(1)})^p & \longrightarrow & \end{array}$$

Since  $T_j$  does not occur as a summand in  $B_i^{(e+1)}$  by Proposition 3.5, we have that  $\text{Hom}(T_j^{(1)}, B_i^{(e+1)}[1]) = 0$ . Hence the rightmost triangle splits, so we have a triangle

$$T_i^{(e+2)}[-1] \rightarrow (T_i^{(e+1)})' \amalg C' \rightarrow B_i^{(e+1)} \amalg (T_j^{(1)})^p \rightarrow \quad (6)$$

By Lemma 3.3 we have that  $\text{Hom}(T_i^{(e+2)}, T_j^{(1)}[1]) = 0$ . By Lemma 3.2(e) we get that  $\text{Hom}(T_i^{(e+2)}, T_i[1]) = 0$ , and clearly  $\text{Hom}(T_i^{(e+2)}, T_l[1]) = 0$ , for  $l \neq i$ . We hence get that all maps  $(T_i^{(e+1)})' \amalg C' \rightarrow U$ , with  $U$  in  $\text{add } T'$ , factor through  $(T_i^{(e+1)})' \amalg C' \rightarrow B_i^{(e+1)} \amalg (T_j^{(1)})^p$ . Minimality is clear from the triangle (6). This proves (a), and (b) follows from the fact that  $C'$  contains no copies of  $T_j$ , and hence splits off. (c) is a direct consequence of (b).  $\square$

**Proposition 5.5.** (a) If  $q_{ij}^{(u)} = 0$  for  $u = 0, \dots, m$ , then  $(T_i^{(v)})' \simeq T_i^{(v)}$  for all  $v$ .

(b) If  $e \neq m$  and  $q_{ij}^{(e)} > 0$ , then  $(T_i^{(v)})' \simeq T_i^{(v)}$  for  $v \neq e + 1$ .

*Proof.* (a) is a direct consequence of 5.2. For (b) note that by Lemmas 5.2 and 5.4 we have  $(T_i^{(v)})' \simeq T_i^{(v)}$  for  $v = 0, \dots, e$  and  $v = e + 2$ . For  $v \geq e + 2$  consider the exchange triangles

$$T_i^{(v)} \rightarrow B_i^{(v)} \rightarrow T_i^{(v+1)} \rightarrow .$$

Since  $\text{Hom}(T_i^{(v+1)}, T_j^{(1)}[1]) = 0$  by Lemma 3.3 and  $q_{ij}^{(v)} = 0$ , it is clear that the map  $T_i^{(v)} \rightarrow B_i^{(v)}$  is a left  $\text{add } T'/T_i$ -approximation. Hence (b) follows.  $\square$

## 6. PROOF OF THE MAIN RESULT

This section contains the proof of the main result, Theorem 2.1. As before, let  $T = \tilde{T} \amalg T_i \amalg T_j$  be an  $m$ -tilting object, and let  $T' = T/T_j \amalg T_j^{(1)}$ .

We will compare the numbers of  $(c)$ -coloured arrows from  $i$  to  $k$ , in the coloured quivers of  $T$  and  $T'$ , i.e. we will compare  $q_{ik}^{(c)}$  and  $\tilde{q}_{ik}^{(c)}$ .

We need to consider an arbitrary  $T$  whose coloured quiver locally looks like

$$T_i \xrightarrow{(e)} T_j \xrightarrow{(d)} T_k$$

(c)

for any possible value of  $c, d, e$ . Our aim is to show that the formula

$$\tilde{q}_{ik}^{(u)} = \begin{cases} q_{ik}^{(u+1)} & \text{if } j = k \\ q_{ik}^{(u-1)} & \text{if } j = i \\ \max\{0, q_{ik}^{(u)} - \sum_{t \neq u} q_{ik}^{(t)} + (q_{ij}^{(u)} - q_{ij}^{u-1})q_{jk}^{(0)} + q_{ij}^{(m)}(q_{jk}^{(u)} - q_{jk}^{(u+1)})\} & \text{if } i \neq j \neq k \end{cases}$$

holds. The case where  $j = k$  is directly from the definition. The case where  $j = i$  follows by condition (II) for  $Q_{T'}$ . For the rest of the proof we assume  $j \notin \{i, k\}$ . We will divide the proof into four cases, where  $p \geq 0$  denotes the number of arrows from  $i$  to  $j$ , and  $q = q_{jk}^{(0)}$ .

- I.  $p = 0$
- II.  $p \neq 0, e \neq m$  and  $q = 0$
- III.  $p \neq 0, e \neq m$  and  $q \neq 0$ .
- IV.  $p \neq 0$  and  $e = m$

Note that in the three first cases, the formula reduces to

$$\tilde{q}_{ik}^{(u)} = \max\{0, q_{ik}^{(u)} - \sum_{t \neq u} q_{ik}^{(t)} + (q_{ij}^{(u)} - q_{ij}^{(u-1)})q_{jk}^{(0)}\},$$

and in the first two cases it further reduces to

$$\tilde{q}_{ik}^{(u)} = q_{ik}^{(u)}.$$

CASE I. We first consider the situation where there is no coloured arrow  $i \rightarrow j$ , i.e.  $q_{ij}^{(u)} = 0$  for all  $u$ . That is, we assume  $Q_T$  locally looks like this

$$T_i \xrightarrow{\quad} T_j \xrightarrow{(d)} T_k$$

(c)

with  $c, d$  arbitrary. It is a direct consequence of Proposition 5.5 that  $q_{ik}^{(u)} = \tilde{q}_{ik}^{(u)}$  for all  $u$  which shows that the formula holds.

CASE II. We consider the setting where we assume  $Q_T$  locally looks like this

$$T_i \xrightarrow{(e)} T_j \xrightarrow{(d)} T_k$$

(c)

with  $e \neq m$  and  $q = 0$ .

We then claim that we have the following, which shows that the formula holds.

**Lemma 6.1.** *In the above setting  $q_{ik}^{(u)} = \tilde{q}_{ik}^{(u)}$  for all  $u$ .*

*Proof.* It follows directly from Proposition 5.5 that  $q_{ik}^{(u)} = \tilde{q}_{ik}^{(u)}$  for  $u = 0, \dots, e-1$ . We claim that  $q_{ik}^{(e)} = \tilde{q}_{ik}^{(e)}$ .

By Lemma 5.3 we have the (not necessarily minimal) left  $\text{add}(T'/T_i)$ -approximation

$$T_i^{(e)} \rightarrow (T_k)^{pq} \amalg Y^p \amalg X = Y^p \amalg X.$$

First, assume that  $T_k$  does not appear as a summand in  $B_i^{(e)} = (T_j)^p \amalg X$ , then the same holds for  $Y^p \amalg X$ , and hence for  $(B_i^{(e)})'$  which is a direct summand in  $Y^p \amalg X$ .

Next, assume  $T_k$  appears as a summand in  $B_i^{(e)}$ , and hence in  $X$ . Then  $T_k$  is by Proposition 3.5 not a summand in  $B_i^{(e+1)}$ , and by Lemma 5.4 we have that  $T_k$  is also not a summand in  $C'$ . Therefore  $T_k$  appears with the same multiplicity in  $B_i^{(e)}$  as in  $(B_i^{(e)})'$ , also in this case.

We now show that  $q_{ik}^{(u)} = \tilde{q}_{ik}^{(u)}$  for  $u > e$ .

If  $q_{ik}^{(e)} \neq 0$ , then  $q_{ik}^{(u)} = \tilde{q}_{ik}^{(u)} = 0$  for  $u > e$  and we are finished. So assume  $q_{ik}^{(e)} = 0$ , i.e.  $T_k$  does not appear as a direct summand of  $X$ .

Consider the map

$$(T_i^{(e+1)})' \amalg C' \rightarrow B_i^{(e+1)} \amalg (T_j^{(1)})^p.$$

We have that  $(B_i^{(e+1)})' \simeq \frac{B_i^{(e+1)} \amalg (T_j^{(1)})^p}{C'}$ . By assumption,  $T_k$  is not a direct summand in  $(T_k)^{pq} \amalg Y^p \amalg X = Y^p \amalg X$ , and thus not in  $C'$ . Hence it follows that  $q_{ik}^{(e+1)} = \tilde{q}_{ik}^{(e+1)}$ .

Since, by Proposition 5.5 we have for  $u = e+2, \dots, m$ , that  $(T_i^{(u)})' = T_i^{(u)}$  and the left  $\text{add}(T/T_i)$ -approximation coincide with the left  $\text{add}(T'/T_i)$ -approximations of  $T_i^{(u)}$ , it now follows that  $q_{ik}^{(u)} = \tilde{q}_{ik}^{(u)}$  for all  $u$ .  $\square$

CASE III. We now consider the setting with  $p$  non-zero,  $q \neq 0$  and  $e \neq m$ . That is, we assume  $Q_T$  locally looks like this

$$\begin{array}{ccccc} T_i & \xrightarrow{(e)} & T_j & \xrightarrow{(0)} & T_k \\ & \searrow & \nearrow & \searrow & \\ & & & (c) & \end{array}$$

where  $c \in \{e, e+1\}$  by Proposition 5.1, and where there are  $z = q_{ik}^{(c)} \geq 0$  arrows from  $T_i$  to  $T_k$ .

**Lemma 6.2.** *In the above setting, we have that  $Q_{T'}$  is given by*

$$\tilde{q}_{ik}^{(e)} = \begin{cases} q_{ij}^{(e)} q_{jk}^{(0)} + q_{ik}^{(e)} & \text{if } c = e \\ q_{ij}^{(e)} q_{jk}^{(0)} - q_{ik}^{(e+1)} & \text{if } c = e + 1 \text{ and } q_{ik}^{(e+1)} \leq q_{ij}^{(e)} q_{jk}^{(0)} \\ 0 & \text{otherwise} \end{cases} \quad (7)$$

$$\tilde{q}_{ik}^{(e+1)} = \begin{cases} -q_{ij}^{(e)} q_{jk}^{(0)} + q_{ik}^{(e+1)} & \text{if } c = e + 1 \text{ and } q_{ik}^{(e+1)} > q_{ij}^{(e)} q_{jk}^{(0)} \\ 0 & \text{otherwise} \end{cases} \quad (8)$$

and

$$\tilde{q}_{ik}^{(u)} = 0 \text{ for } u \notin \{e, e + 1\} \quad (9)$$

*Proof.* We first deal with the case where  $c = e$  and  $z > 0$ . By assumption  $X$  in the triangle (3) has  $z$  copies of  $T_k$ , so  $(T_k)^{pq} \amalg Y^p \amalg X$  has  $pq + z$  copies of  $T_k$ . Hence to show (7) it is sufficient to show that  $C'$  in the triangle

$$T_i^{(e+1)} \rightarrow (T_i^{(e+1)})' \amalg C' \rightarrow (T_j^{(1)})^p \rightarrow$$

has no copies of  $T_k$ . This follows directly from the Lemma 5.4 and the fact that  $T_k$  (by the assumption that  $z > 0$  and Proposition 3.5) is not a summand in  $B_i^{(e+1)}$ . In this case (8) and (9) follow directly from Proposition 3.5.

Consider the case with  $c = e + 1$  and  $0 \leq z \leq pq$ . We have that  $X$  in the triangle (3) does not have  $T_k$  as a direct summand. Assume  $T_k$  appears as a direct summand of  $C'$  with multiplicity  $z'$ . We claim that  $z' = z$ . Assume first  $z' < z$ , then on one hand  $T_k$  appears with multiplicity  $z - z' > 0$  in  $(B_i^{(e+1)})'$ . On the other hand  $T_k$  appears with multiplicity  $pq - z' > 0$  in  $(B_i^{(e)})'$ . This contradicts Proposition 3.5. Hence  $z' = z$ .

Therefore  $(B_i^{(e)})'$  has  $pq - z$  copies of  $T_k$  and (7) and (8) hold. If  $pq \neq z$ , then (9) follows directly from the above and Proposition 3.5. In the case  $pq = z$ , we also need to show that  $T_k$  does not appear as a summand in  $(B_i^{(u)})'$  for  $u \neq e + 1$ . Since  $pq \neq 0$ , we have  $z \neq 0$ , and the result follows from Proposition 5.5.

Now assume  $c = e + 1$  and  $z > pq$ . Assume  $C' = (T_k)^l \amalg C''$ , where  $T_k$  is not a summand in  $C''$ . Now since

$$(T_i^{(e+1)})' \amalg T_k^l \amalg C'' \rightarrow (T_j^{(1)})^p \amalg T_k^z \amalg Y,$$

with  $T_k$  not a summand in  $Y$ , is a minimal left  $\text{add } T'$ -approximation, we have that  $l \leq pq < z$  and  $T_k$  appears with multiplicity  $z - l > 0$  in the minimal left  $\text{add } T' / (T_i)$ -approximation of  $(T_i^{(e+1)})'$ , hence  $T_k$  cannot appear as a summand in the minimal left  $\text{add } T' / T_i$ -approximation of  $(T_i^{(e)})'$ . Hence  $l = pq$ , and we have completed the proof of (7) and (8) in this case. The case (9), i.e.  $u \neq e, e + 1$  follows from Proposition 3.5.  $\square$

CASE IV. We now consider the case with  $q_{ij}^{(m)} \neq 0$ . Assume first there are no arrows

from  $j$  to  $k$ . Then we can use the symmetry proved in Proposition 4.1 and reduce to case I. The formula is easily verified in this case.

Assume  $d \neq 0$ , again we can use the symmetry, this time to reduce to case III. It is straightforward to verify that the formula holds also in this case.

Assume now that  $d = 0$ , i.e. we need to consider the following case

$$\begin{array}{ccccc}
 & & \xrightarrow{(m)} & & \xrightarrow{(0)} \\
 T_i & \xleftrightarrow{\quad} & T_j & \xleftrightarrow{\quad} & T_k \\
 & \xleftarrow{(0)} & & \xleftarrow{(m)} & \\
 & \searrow & \text{---} & \swarrow & \\
 & & (c) & & \\
 & \searrow & \text{---} & \swarrow & \\
 & & (m-c) & & 
 \end{array}$$

Now by Proposition 5.1 we have that  $c$  is in  $\{m, 0\}$ . Assume there are  $z \geq 0$   $(c)$ -coloured arrows

The coloured quiver of  $T'$  is of the form

$$\begin{array}{ccccc}
 & & \xrightarrow{(0)} & & \xrightarrow{(m)} \\
 T_i & \xleftrightarrow{\quad} & T_j^{(1)} & \xleftrightarrow{\quad} & T_k \\
 & \xleftarrow{(m)} & & \xleftarrow{(0)} & \\
 & \searrow & \text{---} & \swarrow & \\
 & & (c') & & \\
 & \searrow & \text{---} & \swarrow & \\
 & & (m-c') & & 
 \end{array}$$

and applying the symmetry of Proposition 4.1 we have that if  $z > 0$ , then  $c' \in \{0, m\}$  by Proposition 5.1. Hence for all  $u \notin \{0, m\}$  we have that  $\tilde{q}_{ik}^{(u)} = q_{ik}^{(u)} = 0$ . Therefore it suffices to show that  $\tilde{q}_{ik}^{(u)} = q_{ik}^{(u)}$ , for  $u \in \{0, m\}$ . This is a direct consequence of the following.

**Lemma 6.3.** *Assume we are in the above setting. A map  $T_i \rightarrow T_k$  or  $T_k \rightarrow T_i$  is irreducible in  $\text{add } T$  if and only if it is irreducible in  $\text{add } T'$ .*

*Proof.* Assume  $T_i \rightarrow T_k$  is not irreducible in  $\text{add } T'$ , and that  $T_i \rightarrow T_k = T_i \rightarrow U \rightarrow T_k$  for some  $U = \coprod_t U_t \in \text{add } T'$ , with  $U_t$  the indecomposable direct summands of  $U$ . Note that by Lemma 3.2(a), we can assume that all  $T_i \rightarrow U_t$  and all  $U_t \rightarrow T_k$  are non-isomorphisms. If there is some index  $t$  such that  $U_t \simeq T_j^{(1)}$ , the map  $U_t \rightarrow T_k$  factors through some  $U'$  in  $\text{add}(T/(T_i \amalg T_k))$ , since there are no  $(1)$ -coloured arrows  $j \rightarrow i$  or  $j \rightarrow k$  in the coloured quiver of  $T$ . This shows that  $T_i \rightarrow T_k$  is not irreducible in  $\text{add } T$ .

Assume  $T_i \rightarrow T_k$  is not irreducible in  $\text{add } T$ , and that  $T_i \rightarrow T_k = T_i \rightarrow V \rightarrow T_k$  for some  $V = \coprod_t V_t \in \text{add } T$ , with  $V_t$  the indecomposable direct summands of  $V$ . If there is some index  $t$  such that  $V_t \simeq T_j$ , the map  $T_i \rightarrow V_t$  factors through  $B_j^{(m)}$ , which is in  $\text{add}(T/(T_i \amalg T_j \amalg T_k)) \subset \text{add } T'$ , since there are no  $(0)$ -coloured arrows  $i \rightarrow j$  or  $k \rightarrow j$  in the coloured quiver of  $T$ . This shows that  $T_i \rightarrow T_k$  is not irreducible in  $\text{add } T'$ .

By symmetry, the same property holds for maps  $T_k \rightarrow T_i$ .  $\square$

Thus we have proven that the formula holds in all four cases, and this finishes the proof of Theorem 2.1.

## 7. $m$ -CLUSTER-TILTED ALGEBRAS

An  $m$ -cluster-tilted algebra is an algebra given as  $\text{End}_{\mathcal{C}}(T)$  for some tilting object  $T$  in an  $m$ -cluster category  $\mathcal{C} = \mathcal{C}_m$ . Obviously, the subquiver of the coloured quiver of  $T$  given by the (0)-coloured maps is the Gabriel quiver of  $\text{End}_{\mathcal{C}}(T)$ .

An application of our main theorem is that the quivers of the  $m$ -cluster-tilted algebras can be combinatorially determined via repeated (coloured) mutation. For this one needs transitivity in the tilting graph of  $m$ -tilting objects. More precisely, we need the following, which is also pointed out in [ZZ].

**Proposition 7.1.** *Any  $m$ -tilting object can be reached from any other  $m$ -tilting object via iterated mutation.*

*Proof.* We sketch a proof for the convenience of the reader. Let  $T'$  be a tilting object in an  $m$ -cluster category  $\mathcal{C}$  of the hereditary algebra  $H = KQ$ , and let  $\mathcal{C}_1$  be the 1-cluster category of  $H$ . By [Z], there is a tilting object  $T$  of degree 0, i.e. all direct summands in  $T$  have degree 0, such that  $T$  can be reached from  $T'$  via mutation. It is sufficient to show that the canonical tilting object  $H$  can be reached from  $T$  via mutation. Since  $T$  is of degree 0, it is induced from a  $H$ -tilting module. Especially  $T$  is a tilting object in  $\mathcal{C}_1$ . Since  $T$  and  $H$  are tilting objects in  $\mathcal{C}_1$ , by [BMRRT] there are  $\mathcal{C}_1$ -tilting objects  $T = T_0, T_1, \dots, T_r = H$ , such that  $T_i$  mutates to  $T_{i+1}$  (in  $\mathcal{C}_1$ ) for  $i = 0, \dots, r-1$ . Now each  $T_i$  is induced by a tilting module for some  $Q_i$  where all  $KQ_i$  are derived equivalent to  $KQ$ . Hence, each  $T_i$  is easily seen to be an  $m$ -cluster tilting object. Since  $T_{i+1}$  differs from  $T_i$  in only one summand the mutations in  $\mathcal{C}_1$  are also mutations in  $\mathcal{C}$ . This concludes the proof.  $\square$

A direct consequence of the transitivity is the following.

**Corollary 7.2.** *For an  $m$ -cluster category  $\mathcal{C} = \mathcal{C}_m$  of the acyclic quiver  $Q$ , all quivers of  $m$ -cluster-tilted algebras are given by repeated coloured mutation of  $Q$ .*

## 8. COMBINATORIAL COMPUTATION

In this section, we discuss concrete computation with tilting objects in an  $m$ -cluster tilting category.

An exceptional indecomposable object in  $\text{mod } H$  is uniquely determined by its image  $[T]$  in the Grothendieck group  $K_0(\text{mod } H)$ . There is a map from  $\mathcal{D}^b(\text{mod } H)$  to  $K_0(\text{mod } H)$  which, for  $T \in \text{mod } H$ , takes  $T[i]$  to  $(-1)^i[T]$ . An exceptional indecomposable in  $\mathcal{D}^b(\text{mod } H)$  can be uniquely specified by its class in  $K_0(\text{mod } H)$  together with its degree.

The map from  $\mathcal{D}^b(\text{mod } H)$  to  $K_0(\text{mod } H)$  does not descend to  $\mathcal{C}$ . However, if we fix our usual choice of fundamental domain in  $\mathcal{D}^b(\text{mod } H)$ , then we can identify the indecomposable objects in it as above.

Let us define the combinatorial data corresponding to a tilting object  $T$  to be  $Q_T$  together with  $([T_i], \deg T_i)$  for  $1 \leq i \leq n$ .

**Theorem 8.1.** *Given the combinatorial data for a tilting object  $T$  in  $\mathcal{C}$ , it is possible to determine, by a purely combinatorial procedure, the combinatorial data for the tilting object which results from an arbitrary sequence of mutations applied to  $T$ .*

*Proof.* Clearly, it suffices to show that, for any  $i$ , we can determine the class and degree for  $T_i^{(j)}$ . If we can do that then, by the coloured mutation procedure, we can determine the coloured quiver for  $(T/T_i) \amalg T_i^{(j)}$ , and by applying this procedure repeatedly, we can calculate the result of an arbitrary sequence of mutations.

Since we are given  $Q_T$ , we know  $B_i^{(0)}$ , and we can calculate  $[B_i^0]$ . Now we have the following lemma:

**Lemma 8.2.**  *$[T_i^{(1)}] = [B_i^{(0)}] - [T_i^{(0)}]$ , and  $\deg(T_i^{(1)}) = \deg T_i$  or  $\deg T_i + 1$ , whichever is consistent with the sign of the class of  $[T_i^{(1)}]$ , unless this yields a non-projective indecomposable object in degree  $m$ , or an indecomposable of degree  $m + 1$ .*

*Proof.* The proof is immediate from the exchange triangle  $T_i \rightarrow B_i^{(0)} \rightarrow T_i^{(1)} \rightarrow$ .  $\square$

Applying this lemma, and supposing that we are not in the case where its procedure fails, we can determine the class and degree  $T_i^{(1)}$ . By the coloured mutation procedure, we can also determine the coloured quiver for  $\mu_i(T)$ . We therefore have all the necessary data to apply Lemma 8.2 again. Repeatedly applying the lemma, there is some  $k$  such that we can calculate the class and degree of  $T_i^{(j)}$  for  $1 \leq j \leq k$ , and the procedure described in the lemma fails to calculate  $T_i^{(k+1)}$ .

We also have the following lemma:

**Lemma 8.3.**  *$[T_i^{(m)}] = [B_i^{(m)}] - [T_i^{(0)}]$ , and  $\deg T_i^{(m)} = \deg T_i$  or  $\deg T_i - 1$ , whichever is consistent with the sign of  $[T_i^{(m)}]$ , unless this yields an indecomposable in degree  $-1$ .*

Applying this lemma, starting again with  $T$ , we can obtain the degree and class for  $T_i^{(m)}$ . We can then determine the coloured quiver for  $\mu_i^{-1}(T)$ , and we are now in a position to apply Lemma 8.3 again. The last complement which Lemma 8.3 will successfully determine is  $T_i^{(k+1)}$ . It follows that we can determine the degree and class of any complement to  $T/T_i$ .  $\square$

## 9. THE $m$ -CLUSTER COMPLEX

In this section, we discuss the application of our results to the study of the  $m$ -cluster complex, a simplicial complex defined in [FR] for a finite root system  $\Phi$ . We shall begin by stating our results for the  $m$ -cluster complex in purely combinatorial language, and then briefly describe how they follow from the representation-theoretic perspective in the rest of the paper. For simplicity, we restrict to the case where  $\Phi$  is simply laced.

Number the vertices of the Dynkin diagram for  $\Phi$  from 1 to  $n$ . The  $m$ -coloured almost positive roots,  $\Phi_{\geq -1}^m$ , consist of  $m$  copies of the positive roots, numbered 1 to  $m$ , together with a single copy of the negative simple roots. We refer to an element of the  $i$ -th copy of  $\Phi^+$  as having colour  $i$ , and we write such an element as  $\beta^{(i)}$ .

Since the Dynkin diagram for  $\Phi$  is a tree, it is bipartite; we fix a bipartition  $\{1, \dots, n\} = I_+ \cup I_-$ .

The  $m$ -cluster complex,  $\Delta_m$ , is a simplicial complex on the ground set  $\Phi_{\geq -1}^m$ . Its maximal faces are called  $m$ -clusters. The definition of  $\Delta_m$  is combinatorial; we refer the reader to [FR]. The  $m$ -clusters each consist of  $n$  elements of  $\Phi_{\geq -1}^m$  [FR, Theorem 2.9]. Every codimension 1 face of  $\Delta_m$  is contained in exactly  $m+1$  maximal faces [FR, Proposition 2.10]. There is a certain combinatorially-defined bijection  $R_m: \Phi_{\geq -1}^m \rightarrow \Phi_{\geq -1}^m$ , which takes faces of  $\Delta_m$  to faces of  $\Delta_m$  [FR, Theorem 2.4].

It will be convenient to consider *ordered  $m$ -clusters*. An ordered  $m$ -cluster is just a  $n$ -tuple from  $\Phi_{\geq -1}^m$ , the set of whose elements forms an  $m$ -cluster. Write  $\Sigma_m$  for the set of ordered  $m$ -clusters.

For each ordered  $m$ -cluster  $C = (C_1, \dots, C_n)$ , we will define a coloured quiver  $Q_C$ . We will also define an operation  $\mu_j: \Sigma_m \rightarrow \Sigma_m$ , which takes ordered  $m$ -clusters to ordered  $m$ -clusters, changing only the  $j$ -th element.

We will define both operations inductively. The set  $-\Pi$  of negative simple roots forms an  $m$ -cluster. Its associated quiver is defined by drawing, for each edge  $ij$  in the Dynkin diagram, a pair of arrows. Suppose  $i \in I_+$  and  $j \in I_-$ . Then we draw an arrow from  $i$  to  $j$  with colour 0, and an arrow from  $j$  to  $i$  with colour  $m$ .

Suppose now that we have some ordered  $m$ -cluster  $C$ , together with its quiver  $Q_C$ . We will now proceed to define  $\mu_j(C)$ . Write  $q_{jk}^{(0)}$  for the number of arrows in  $Q_C$  of colour 0 from  $j$  to  $k$ . Define:

$$\beta = -C_j + \sum_{k \neq j} q_{jk}^{(0)} C_k.$$

Let  $c$  be the colour of  $C_j$ . We define  $\mu_j(C)$  by replacing  $C_j$  by some other element of  $\Phi_{\geq -1}^m$ , according to the following rules:

- If  $C_j$  is positive and  $\beta$  is positive, replace  $C_j$  by  $\beta^{(c)}$ .
- If  $C_j$  is positive and  $\beta$  is negative, replace  $C_j$  by  $R_m(-\beta^{(c)})$ .
- If  $C_j$  is negative simple  $-\alpha_i$ , define  $\gamma$  by  $\gamma^{(0)} = R_m(-\alpha_i)$ , and then replace  $C_j$  by  $\beta + C_j - \gamma$ , with colour zero.

Define the quiver for the  $m$ -cluster  $\mu_j(C)$  by the coloured quiver mutation rule from Section 2. Since any  $m$ -cluster can be obtained from  $-\Pi$  by a sequence of mutations, the above suffices to define  $\mu_j(C)$  and  $Q_C$  for any ordered  $m$ -cluster  $C$ .

**Proposition 9.1.** *The operation  $\mu_j$  defined above takes  $m$ -clusters to  $m$ -clusters, and the  $m$ -clusters  $\mu_j^i(C)$  for  $0 \leq i \leq m$  are exactly those containing all the  $C_i$  for  $i \neq j$ .*

The connection between the combinatorics discussed here and the representation theory in the rest of the paper is as follows.  $\Phi_{\geq -1}^m$  corresponds to the indecomposable objects of (a fundamental domain for)  $\mathcal{C}_m$ . The cluster tilting objects in  $\mathcal{C}_m$  correspond to the  $m$ -clusters. The operation  $R_m$  corresponds to [1]. For further details on the translation, the reader is referred to [T, Z]. The above proposition then follows from the approach taken in Section 8.

## 10. AN ALTERNATIVE ALGORITHM FOR COLOURED MUTATION

Here we give an alternative description of coloured quiver mutation at vertex  $j$ .

- (1) For each pair of arrows

$$i \xrightarrow{(c)} j \xrightarrow{(0)} k$$

with  $i \neq k$ , the arrow from  $i$  to  $j$  of arbitrary colour  $c$ , and the arrow from  $j$  to  $k$  of colour 0, add a pair of arrows: an arrow from  $i$  to  $k$  of colour  $c$ , and one from  $k$  to  $i$  of colour  $m - c$ .

- (2) If the graph violates property II, because for some pair of vertices  $i$  and  $k$  there are arrows from  $i$  to  $k$  which have two different colours, cancel the same number of arrows of each colour, until property II is satisfied.
- (3) Add one to the colour of any arrow going into  $j$  and subtract one from the colour of any arrow going out of  $j$ .

**Proposition 10.1.** *The above algorithm is well-defined and correctly calculates coloured quiver mutation as previously defined.*

*Proof.* Fix a quiver  $Q$  and a vertex  $j$  at which the mutation is being carried out.

To prove that the algorithm is well-defined, we must show that at step 2, there are only two colours of arrows running from  $i$  to  $k$  for any pair of vertices  $i, k$ . (Otherwise there would be more than one way to carry out the cancellation procedure of step 2.)

Since in the original quiver  $Q$ , there was only one colour of arrows from  $i$  to  $k$ , in order for this problem to arise, we must have added two different colours of arrows from  $i$  to  $k$  at step 1. Two colours of arrows will only be added from  $i$  to  $k$  if, in  $Q$ , there are both (0)-coloured arrows from  $j$  to  $k$  and from  $j$  to  $i$ . In this case, by property III, there are  $(m)$ -coloured arrows from  $i$  to  $j$  and from  $k$  to  $j$ . It follows that in step 1, we will add both (0)-coloured and  $(m)$ -coloured arrows. Applying Proposition 5.1, we see that any arrows from  $i$  to  $k$  in  $Q$  are of colour 0 or  $m$ . Thus, as desired, after step 1, there are only two colours of arrows in the quiver, so step 2 is well-defined.

We now prove correctness. Let  $\tilde{Q} = \mu_j(Q)$ . Write  $q_{ij}^{(c)}$  for the number of  $c$ -coloured arrows from  $i$  to  $j$  in  $Q$ , and similarly  $\tilde{q}_{ij}^{(c)}$  for  $\tilde{Q}$ . Write  $\hat{Q}$  and  $\hat{q}_{ij}^{(c)}$  for the result of applying the above algorithm.

It is clear that only the final step of the algorithm is relevant for  $\hat{q}_{ik}$  where one of  $i$  or  $k$  coincides with  $j$ , and therefore that in this case  $\hat{q}_{ij}^{(c)} = \tilde{q}_{ij}^{(c)}$  as desired.

Suppose now that neither  $i$  nor  $k$  coincides with  $j$ . Suppose further that in  $Q$  there are no (0)-coloured arrows from either  $i$  or  $k$  to  $j$ , and therefore also no  $m$ -coloured arrows from  $k$  to  $i$  or  $j$ . In this case,  $\tilde{q}_{ik}^{(c)} = q_{ik}^{(c)}$ . In the algorithm, no arrows will be added between  $i$  and  $k$  in step 1, and therefore no further changes will be made in step 2. Thus  $\hat{q}_{ik}^{(c)} = q_{ik}^{(c)} = \tilde{q}_{ik}^{(c)}$ , as desired.

Suppose now that there are (0)-coloured arrows from  $j$  to both  $i$  and  $k$ . In this case,  $\tilde{q}_{ik}^{(c)} = q_{ik}^{(c)}$ . In this case, as discussed in the proof of well-definedness, an equal

number of (0)-coloured and ( $m$ )-coloured arrows will be introduced at step 1. They will therefore be cancelled at step 2. Thus  $\hat{q}_{ik}^{(c)} = q_{ik}^{(c)} = \tilde{q}_{ik}^{(c)}$  as desired.

Suppose now that there is a (0)-coloured arrow from  $j$  to  $k$ , but not from  $j$  to  $i$ . Let the arrows from  $i$  to  $j$ , if any, be of colour  $c$ . At step 1 of the algorithm, we will add  $q_{ij}^{(c)} q_{jk}^{(0)}$  arrows of colour  $c$  to  $Q$ . By Proposition 5.1, the arrows in  $Q$  from  $i$  to  $k$  are of colour  $c$  or  $c + 1$ . One verifies that the algorithm yields the same result as coloured quiver mutation, in the three cases that the arrows from  $i$  to  $k$  in  $Q$  are of colour  $c$ , that they are of colour  $c + 1$  but there are fewer than  $q_{ij}^{(c)} q_{jk}^{(0)}$ , and that they are of colour  $c + 1$  and there are at least as many as  $q_{ij}^{(c)} q_{jk}^{(0)}$ .

The final case, that there is a (0)-coloured arrow from  $j$  to  $i$  but not from  $j$  to  $k$ , is similar to the previous one.  $\square$

## 11. EXAMPLE: TYPE $A_n$

In [BM1], a certain category  $\mathcal{C}_{BM}$  is constructed, which is shown to be equivalent to the  $m$ -cluster category of Dynkin type  $A_n$ . The description of  $\mathcal{C}_{BM}$  is as follows. Take an  $nm + 2$ -gon  $\Pi$ , with vertices labelled clockwise from 1 to  $nm + 2$ . Consider the set  $X$  of diagonals  $\gamma$  of  $\Pi$  with the property that  $\gamma$  divides  $\Pi$  into two polygons each having a number of sides congruent to 2 modulo  $m$ . For each  $\gamma \in X$ , there is an object  $A_\gamma$  in  $\mathcal{C}_{BM}$ . These objects  $A_\gamma$  form the indecomposables of the additive category  $\mathcal{C}_{BM}$ . We shall not recall the exact definition of the morphisms, other than to note that they are generated by the morphisms  $p_{ijk} : A_{ij} \rightarrow A_{ik}$  which exist provided that  $ij$  and  $ik$  are both diagonals in  $X$ , and that, starting at  $j$  and moving clockwise around  $\Pi$ , one reaches  $k$  before  $i$ .

A collection of diagonals in  $X$  is called non-crossing if its elements intersect pairwise only on the boundary of the polygon. An inclusion-maximal such collection of diagonals divides  $\Pi$  into  $m + 2$ -gons; we therefore refer to such a collection of diagonals as an  $m + 2$ -angulation. If we remove one diagonal  $\gamma$  from an  $m + 2$ -angulation  $\Delta$ , then the two  $m + 2$ -gons on either side of  $\gamma$  become a single  $2m + 2$ -gon. We say that  $\gamma$  is a *diameter* of this  $2m + 2$ -gon, since it connects vertices which are diametrically opposite (with respect to the  $2m + 2$ -gon). If  $\delta$  is another diameter of this  $2m + 2$ -gon, then  $(\Delta \setminus \gamma) \cup \delta$  is another maximal noncrossing collection of diagonals from  $X$ . (In particular,  $\delta \in X$ .)

For  $\Delta$  an  $m + 2$ -angulation, let  $A_\Delta = \prod_{\gamma \in \Delta} A_\gamma$ . Then we have that  $A_\Delta$  is a basic ( $m$ -cluster-)tilting object for  $\mathcal{C}_{BM}$ , and all basic tilting objects of  $\mathcal{C}_{BM}$  arise in this way. It follows from the previous discussion that if  $T = A_\Delta$  is a basic tilting object, and  $\gamma \in \Delta$ , then the complements to  $A_{\Delta \setminus \gamma}$  will consist of the objects  $A_\delta$  where  $\delta$  is a diameter of the  $2m + 2$ -gon obtained by removing  $\gamma$  from the  $m + 2$ -angulation determined by  $\Delta$ . In fact, we can be more precise. Define  $\delta_{(i)}$  to be the diameter of the  $2m + 2$ -gon obtained by rotating the vertices of  $\gamma$  by  $i$  steps counterclockwise (within the  $2m + 2$ -gon). Then  $A_\gamma^{(i)} = A_{\delta_{(i)}}$ .


Define  $Q_\Delta$  to be the coloured quiver for the tilting object  $A_\Delta$ , with vertex set  $\Delta$ . Using the setup of [BM1], it is straightforward to verify:

**Proposition 11.1.** *The coloured quiver  $Q_\Delta$  of  $T = A_\Delta$  has an arrow from  $\gamma$  to  $\delta$  if and only if  $\gamma$  and  $\delta$  both lie on some  $m + 2$ -gon in the  $m + 2$ -angulation defined by  $\Delta$ . In this case, the colour of the arrow is the number of edges forming the segment of the boundary of the  $m + 2$ -gon which lies between  $\gamma$  and  $\delta$ , counterclockwise from  $\gamma$  and clockwise from  $\delta$ .*

Given the proposition above, it is straightforward to verify directly that  $Q_\Delta$  satisfies conditions (I), (II), and (III), and that mutation is indeed given by the mutation of coloured quivers.

**Example:**  $A_3$ ,  $m = 2$

We return to the example from Section 2. The quadrangulation of a decagon corresponding to the tilting object  $T$  is on the left. The quadrangulation corresponding to  $T'$  is on the right. Passing from the figure on the left to the figure on the right, the diagonal 27 (which corresponds to the summand  $I_2$ ) has been rotated one step counterclockwise within the hexagon with vertices 1,2,3,4,7,10.



#### REFERENCES

- [ABST] I. Assem, T. Brüstle, R. Schiffler, G. Todorov, *m-cluster categories and m-replicated algebras*, J. Pure Appl. Algebra 212 (2008), no. 4, 884–901
- [BM1] K. Baur, R. J. Marsh, *A geometric description of m-cluster categories*, Trans. Amer. Math. Soc. 360 (2008), 5789–5803
- [BM2] K. Baur, R. J. Marsh, *A geometric description of the m-cluster categories of type  $D_n$* , Int. Math. Res. Notices (2007) Vol. 2007, article ID rnm011, 19 pages
- [BMR] A. B. Buan, R. J. Marsh, I. Reiten, *Cluster mutation via quiver representations*, Commentarii Mathematici Helvetici, Volume 83 no.1, 143–177
- [BMRRT] A. B. Buan, R. J. Marsh, M. Reineke, I. Reiten, G. Todorov, *Tilting theory and cluster combinatorics*, Advances in Mathematics 204 (2) (2006), 572–618
- [BMRT] A. B. Buan, R. J. Marsh, I. Reiten, G. Todorov, *Clusters and seeds in acyclic cluster algebras*, Proc. Amer. Math. Soc. 135, No. 10 (2007), 3049–3060
- [CK] P. Caldero, B. Keller, *From triangulated categories to cluster algebras II*, Annales Scientifiques de l’École Normale Supérieure, 4ème série, 39, (2006), 983–1009
- [F] E. Fernandez, *On m-clusters and iterated tilted algebras*, talk at ICRA XIII, Torun, August 2007

- [FR] S. Fomin, N. Reading, *Generalized cluster complexes and Coxeter combinatorics*, Int. Math. Res. Notices (2005), 2709–2757
- [FZ] S. Fomin, A. Zelevinsky, *Cluster algebras. I. Foundations*, J. Amer. Math. Soc. 15 (2002), no. 2, 497–529
- [HR] D. Happel, C. M. Ringel, *Tilted algebras*, Trans. Amer. Math. Soc. 274 (1982), no. 2, 399–443
- [HJ1] T. Holm, P. Jørgensen, *Cluster categories, selfinjective algebras, and stable Calabi-Yau dimensions: type A* arXiv:math.RT/0610728
- [HJ2] T. Holm, P. Jørgensen, *Cluster categories, selfinjective algebras, and stable Calabi-Yau dimensions: types D and E* arXiv:math.RT/0612451
- [IR] O. Iyama, I. Reiten, *Fomin-Zelevinsky mutation and tilting modules over Calabi-Yau algebras*, Amer. J. Math. 139 (2008), no.4, 1087–1148
- [IY] O. Iyama, Y. Yoshino, *Mutation in triangulated categories and rigid Cohen-Macaulay modules*, Inv. Math. Volume 172, Number 1, (2008), 117–168
- [K] B. Keller, *On triangulated orbit categories*, Doc. Math. 10 (2005), 551–581
- [KR1] B. Keller, I. Reiten, *Cluster tilted algebras are Gorenstein and stably Calabi-Yau*, Adv. Math. 211 (2007), 123–151
- [KR2] B. Keller, I. Reiten, *Acyclic Calabi-Yau categories*, arXiv: math/0605594, to appear in Compositio Math.
- [T] H. Thomas, *Defining an  $m$ -cluster category*, Journal of Algebra, Volume 318, Issue 1 (2007), 37–46
- [W] A. Wraalsen, *Rigid objects in higher cluster categories*, preprint, arXiv: 0712.2970
- [Z] B. Zhu, *Generalized cluster complexes via quiver representations*, Journal of Algebraic Combinatorics, Volume 27, Number 1 (2008), 35–54
- [ZZ] Y. Zhou, B. Zhu, *Cluster combinatorics of  $d$ -cluster categories*, preprint, arXiv: 0712.1381

INSTITUTT FOR MATEMATISKE FAG, NORGES TEKNISK-NATURVITENSKAPELIGE UNIVERSITET, N-7491 TRONDHEIM, NORWAY

*E-mail address:* `aslakb@math.ntnu.no`

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF NEW BRUNSWICK, FREDERICTON NB, E3B 1J4 CANADA

*E-mail address:* `hthomas@unb.ca`