# On principal fibrations associated with one algebra

# Maria Trnková<sup>1,2</sup>

email: Maria-Trnkova@seznam.cz

<sup>1</sup>Department of Algebra and Geometry, Faculty of Science, Palacky University in Olomouc, Tomkova 10, Olomouc, Czech Republic <sup>2</sup>Department of Geometry, NIMM, Kazan State University, prof. Nuzhina 1/37, Kazan, Russia

To the memory of B.N. Shapukov

**Abstract** In this paper we study two types of fibrations associated with a 3-dimensional unital associative irreducible algebra and their basic properties. We investigate trivial principal fibrations of degenerate semi-Euclidean sphere and their semi-conformal and projective models. We use Norden normalization method for constructing second model.

#### 1. Principal fibrations of G by its subgroups

Let us denote by  $\mathfrak{A}$  a unital associative n-dimensional algebra with multiplication xy, by  $G \subset \mathfrak{A}$  the set of invertible elements. It is well known that G is a Lie group with the same multiplication rule. Let  $\mathfrak{B}$  be a unital subalgebra of algebra  $\mathfrak{A}$  and  $H \subset G$  the set of invertible elements. So H is a Lie subgroup of group G. We consider the factor-space G/H of right cosets. A fibration  $(G, \pi, M = G/H)$  is a principal fibration with the structure group H, where  $\pi$  is a canonical projection [1]. Here we have a fibration:

$$H \to G \to G/H$$
.

We consider all 3-dimensional unital associative irreducible algebras up to isomorphism. It is known that there exist only tree types of such algebras [2], [3]. In special choice of basic units they have the following multiplication rules:

I)		1	$e_1$	$e_2$
	1	1	$e_1$	$e_2$
	$e_1$	$e_1$	$e_2$	0
	$e_2$	$e_2$	0	0

II) 
$$\begin{array}{c|ccccc}
 & 1 & e_1 & e_2 \\
\hline
 & 1 & 1 & e_1 & e_2 \\
\hline
 & e_1 & e_1 & 1 & e_2 \\
\hline
 & e_2 & e_2 & -e_2 & 0
\end{array}$$

III) 
$$\begin{array}{c|cccc}
 & 1 & e_1 & e_2 \\
\hline
 & 1 & 1 & e_1 & e_2 \\
\hline
 & e_1 & e_1 & 0 & 0 \\
\hline
 & e_2 & e_2 & 0 & 0
\end{array}$$

For these algebras N.Belova found all subalgebras with unit and principal fibrations [4]. Only in algebras of type II and III exist the associated conjugation  $x = x_0 + x^i e_i \to \overline{x} = x_0 - x^i e_i$  which has the property  $\overline{xy} = \overline{y} \, \overline{x}$ .

We consider the bilinear form

$$(x,y) = \frac{1}{2}(x\overline{y} + y\overline{x}). \tag{1}$$

For algebras of type II and III this form takes the real values and it determines a degenerate scalar product on them:

II)
$$(x, y) = x_0 y_0 - x_1 y_1$$
, III) $(x, y) = x_0 y_0$ .

So these algebras have a structure of semi-Euclidean vector spaces with rank 2 and 1, respectively. In our discussion we concentrate on the algebra of type II, because it is less degenerate than type III.

This is non-Abelian algebra. Multiplication rule for algebra elements is:

$$xy = (x_0 + x_1e_1 + x_2e_2)(y_0 + y_1e_1 + y_2e_2) =$$
 (2)

$$x_0y_0 + x_1y_1 + (x_0y_1 + x_1y_0)e_1 + (x_2(y_0 - y_1) + (x_0 + x_1)y_2)e_2.$$

An inverse element of x is:

$$x^{-1} = \frac{x_0 - x_1 e_1 - x_2 e_2}{(x_0)^2 - (x_1)^2}. (3)$$

The set of invertible elements  $G = \{x \in \mathfrak{A} \mid (x_0)^2 - (x_1)^2 \neq 0\}$  is a non-Abelian Lie group. Its underlying manifold is  $\mathbb{R}^3$  without two transversal 2-planes, hence it consists from 4 connected components.

**Proposition**(N.Belova) Any 2-dimensional subalgebra of algebra of type II with an algebra unit is isomorphic to double or dual numbers algebras.

We consider a subalgebra  $R(e_1)$  with basis  $\{1, e_1\}$ , it is an algebra of double numbers, and a subalgebra  $R(e_2)$  with basis  $\{1, e_2\}$ , it is an algebra of dual numbers. The set of their invertible elements  $H_1 = \{x_0 + x_1 e_1 \in R(e_1) \mid x_0^2 - x_1^2 \neq 0\}$  and  $H_2 = \{x_0 + x_2 e_2 \in R(e_2) \mid x_0 \neq 0\}$  are Lie subgroups of the Lie group G. First we take the space of right cosets by  $R(e_1)$  subalgebra and fibration by it. Then we have the following proposition:

**Proposition** (N.Belova) Fibration  $(G, \pi, M = G/H_1)$  determined by formula

$$\pi(x) = \frac{x_2}{x_0 - x_1} \tag{4}$$

is a trivial principal fibration over the real line  $\mathbb{R}$ . The typical fiber, it is a plane without two transversal lines. The structure group is  $H_1$ .

Therefore, the manifold of the group G is diffeomorphic to direct sum  $\mathbb{R} \times H_1$ . The equation of fibers is:

$$u(x_0 - x_1) - x_2 = 0, \ u \in R. \tag{5}$$

This is 1-parametric family of planes with common axis:  $x_0 - x_1 = 0$ ,  $x_2 = 0$ . Of course it is necessary to remove the intersection points of these planes with planes:  $x_0 \pm x_1 = 0$ .

We consider the left multiplications x' = ax on the group G. They form the 3-parametric Lie group of linear transformations:

$$L(a) = \begin{pmatrix} a_0 & a_1 & 0 \\ a_1 & a_0 & 0 \\ a_2 & -a_2 & a_0 + a_1 \end{pmatrix}, \tag{6}$$

where determinant  $\det L(a) = (a_0^2 - a_1^2)(a_0 + a_1) \neq 0$ . The multiplications preserve fibers if and only if when  $a \in H_1$ . This group consists of four connected components.

The right multiplications x' = xb on the group G forms the 3-parametric Lie group of linear transformations:

$$R(b) = \begin{pmatrix} b_0 & b_1 & 0 \\ b_1 & b_0 & 0 \\ b_2 & b_2 & b_0 - b_1 \end{pmatrix}. \tag{7}$$

They preserve the fibration and induce the 2-parametric group of affine transformations on the base:

$$u' = \alpha u + \beta$$
,

here  $\alpha = \frac{b_0 - b_1}{b_0 + b_1} \neq 0, \ \beta = \pi(b).$ 

The right multiplications include involutions of hyperbolic type with matrices:

$$R(b_2) = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ b_2 & b_2 & -1 \end{pmatrix}, \quad R^2 = \text{Id.}$$

They induce the 1-parametric family involutions on the base:

$$u' = -u + b_2.$$

Let us consider rotations and anti-rotations of this semi-Euclidean space. Multiplications x' = ax and x' = xa, where  $|a|^2 = \pm 1$ , are rotations and anti-rotations because of the relations:

$$|x'|^2 = |a|^2 |x|^2 = \pm |x|^2.$$

Each element  $a \in \mathfrak{A}$ ,  $|a|^2 = \pm 1$ , can be represented as:

$$a = \cosh \varphi + a_0 \sinh \varphi$$
 or  $a = \sinh \varphi + a_0 \cosh \varphi$ ,

where  $a_0 = -\overline{a}_0$ ,  $|a_0|^2 = -1$ . The bilinear form (1) in the algebra  $\mathfrak{A}$  takes real values, therefore it is possible to present it as:  $(x,y) = \frac{1}{2}(x\overline{y} + y\overline{x}) = \frac{1}{2}(\overline{x}y + \overline{y}x)$ . Consequently, the hyperbolic cosines or sines of angle between x and x' is equal:

$$\frac{(x,ax)}{|x||ax|} = \frac{1/2(x\overline{ax} + ax\overline{x})}{|x|^2} = \frac{1/2(x\overline{x}\,\overline{a} + ax\overline{x})}{|x|^2} = \frac{1}{2}(\overline{a} + a),$$

$$\frac{(x,xa)}{|x||xa|} = \frac{1/2(\overline{x}xa + \overline{x}\overline{a}x)}{|x|^2} = \frac{1/2(\overline{x}xa + \overline{a}\,\overline{x}x)}{|x|^2} = \frac{1}{2}(a + \overline{a}).$$

In the last two equations we get  $\cosh \varphi$  if  $|a|^2 = 1$  and we get  $\sinh \varphi$  if  $|a|^2 = -1$ . It means that this angle does not depend on x.

Transformations

$$x' = axb, (8)$$

where  $|a|^2 = \pm 1$ ,  $|b|^2 = \pm 1$ , are compositions of rotations or anti-rotations x' = ax and x' = xb. So, there are proper rotations or anti-rotations.

Transformations

$$x' = a\overline{x}b\tag{9}$$

are compositions of (8) and reflection  $x' = \overline{x}$ . So, there are improper rotations or anti-rotations.

**Lemma** Any proper or improper rotation and anti-rotation of this semi-Euclidean space can be represented by (8) or (9).

**Proof** These rotations and anti-rotations are compositions of odd and even numbers of reflections of planes passing through the origin. To each plane corresponds its orthonormal vector n. If vectors  $x_1$  and n are collinear, then  $\overline{x}_1 n = \overline{n}x_1$  and  $x'_1 = -n\overline{x}_1 n = -n\overline{n}x_1 = -x_1$ . If vectors  $x_2$  and n are orthogonal, then  $\overline{x}_2 n + \overline{n}x_2 = 0$  and  $x'_2 = -n\overline{x}_2 n = n\overline{n}x_2 = x_2$ . On the other hand any vector x can be represented as a sum of vectors  $x_1$  and  $x_2$ . It means, that a reflection of plane is:  $x' = -n\overline{x}n$ . So, composition of odd and even numbers of planes reflections are transformation (8) or (9).  $\square$ 

Let us introduce adapted coordinates  $(u, \lambda, \varphi)$  to fibration in semi-Euclidean space, here u is a basic coordinate,  $\lambda, \varphi$  are fiber coordinates. If  $|x|^2 > 0$ , we denote  $\lambda = \pm \sqrt{x_0^2 - x_1^2} \neq 0$ , the sign of  $\lambda$  is equal to the sign of  $x_0$ . The adapted coordinates of fibration in this case are:

$$x_0 = \lambda \cosh \varphi, \quad x_1 = \lambda \sinh \varphi, \quad x_2 = u\lambda \exp \varphi,$$
 (10)

where  $\lambda \in \mathbb{R}_0$ ,  $u, \varphi \in \mathbb{R}$ .

If  $|x|^2 < 0$ , then we write  $\lambda = \pm \sqrt{x_1^2 - x_0^2}$ , the sign of  $\lambda$  is equal to the sign of  $x_1$ :

$$x_0 = \lambda \sinh \varphi, \quad x_1 = \lambda \cosh \varphi, \quad x_2 = u\lambda \exp \varphi.$$
 (11)

The structure group acts as follows:

$$u' = u, \quad \lambda' = \lambda \rho, \quad \varphi' = \varphi + \psi,$$
 (12)

where the element  $a(0, \rho, \psi)$  of structure group acts on the element  $x(u, \lambda, \varphi) \in G$ . This group consists of 4 connected components.

# 2. The principal subfibration of the fibration $(G, \pi, M = G/H_1)$

As we already said, the scalar product in the algebra  $\mathfrak{A}$  of type II is:  $(x,y) = x_0y_0 - x_1y_1$ . So, the algebra  $\mathfrak{A}$  is a 3-dimensional semi-Euclidean space with

rank 2. We call semi-Euclidean sphere with unit radius the set of all elements of algebra  $\mathfrak{A}$  whose square is equal to one,

$$S^{2}(1) = \{ x \in \mathfrak{A} \mid x_{0}^{2} - x_{1}^{2} = 1 \}.$$

It looks like a hyperbolic cylinder. The set of elements with imaginary unit module  $|x|^2 = -1$  we call semi-Euclidean sphere with imaginary unit radius  $S^2(-1)$ . One of these spheres can be obtained up from another one by rotation.

We consider the subfibration of the fibration  $(G, \pi, M = G/H_1)$  to semi-Euclidean sphere  $S^2(1)$ , i.e. the fibration  $\pi: S^2(1) \to M$ . The fibers of new fibration are intersections of  $S^2(1)$  and planes (5). The restriction of the group of double numbers  $H_1$  to  $S^2(1)$  is a Lie subgroup  $S_1$  of double numbers with unit module

$$S_1 = \{a_0 + a_1 e_1 \in H_1 \mid a_0^2 - a_1^2 = 1\}.$$

This group consists of two connected components.

**Proposition** The fibration  $(S^2(1), \pi, M)$  is principal fibration of the group  $S^2(1)$  by the Lie subgroup  $S_1$  to right cosets.

**Proof.** Let x and y be two sphere points from one fiber. They are also from the same fiber of the principal fibration  $(G, \pi, M)$ . So, there exists a unique element  $a \in H_1$  such that y = ax. Then  $|y|^2 = |a|^2|x|^2$  and hence  $|a|^2 = 1$ . It means that  $a \in S_1$ .  $\square$ 

We define coordinates adapted to the fibration on semi-Euclidean sphere  $S^2(1)$ . If  $x \in S^2(1)$  then by (10) we get  $\lambda = \varepsilon$ ,  $\varepsilon = \pm 1$ . The parametric equation of semi-Euclidean sphere in the adapted coordinates  $(u, \varphi)$  is:

$$\mathbf{r}(u,\varphi) = \varepsilon(\cosh\varphi, \sinh\varphi, u\exp\varphi),\tag{13}$$

where u is a basis coordinate,  $\varphi$  is a fiber coordinate. Different values of  $\varepsilon$  correspond to different connected components of semi-Euclidean sphere  $S^2(1)$ .

Let us define the action of the structure group  $S_1$  on semi-Euclidean sphere. By (12) and the adapted coordinates of elements  $a(0, \varepsilon_1, \psi), x(u, \varepsilon, \varphi) \in S^2(1)$  we get:

$$u' = u, \quad \varepsilon' = \varepsilon \varepsilon_1, \qquad \varphi' = \varphi + \psi.$$

This group also consists of two connected components.

The metric tensor for semi-Euclidean sphere has the matrix representation:

$$(g_{ij}) = \begin{pmatrix} 0 & 0 \\ 0 & -1 \end{pmatrix}, \quad \operatorname{rank}(g_{ij}) = 1$$

So the linear element of metric is:

$$ds_1^2 = -d\varphi^2. (14)$$

## 3. Semi-conformal model of the sphere fibration $(S^2(1), \pi, \mathbb{R})$

We consider the semi-conformal model of the fibration  $(S^2(1), \pi, \mathbb{R})$ . We project stereographically the sphere  $S^2(1)$  from the point  $N(1,0,0) \in S^2(1)$  (pole) to the equatorial plane  $\mathbb{R}^2$  with equation  $x_0 = 0$ . We write the stereographic map in the coordinate form. For this we find the equation of line which pass through the pole and an arbitrary point of the sphere  $X(x_0, x_1, x_2)$ :  $\tilde{x}_0 = 1 + (x_0 - 1)t$ ,  $\tilde{x}_1 = x_1t$ ,  $\tilde{x}_2 = x_2t$ . When this line crosses the equatorial plane, we get the formulas of stereographic map  $f: S^2(1) \to \mathbb{R}^2$  where  $x_0 \neq 1$ :

$$x = \frac{x_1}{1 - x_0}, \quad y = \frac{x_2}{1 - x_0}, \tag{15}$$

here  $(x,y) \in \mathbb{R}^2$ ,  $(x_0,x_1,x_2)$  are coordinates on  $S^2(1)$ . An inverse map  $f^{-1}: \mathbb{R}^2 \to S^2(1)$  when  $x \neq \pm 1$  is:

$$x_0 = -\frac{1+x^2}{1-x^2}, \quad x_1 = \frac{2x}{1-x^2}, \quad x_2 = \frac{2y}{1-x^2}.$$
 (16)

If we put formulas (15) into (13) then we obtain the relations between coordinates x, y and adapted coordinates  $u, \varphi$  which are on semi-Euclidean sphere:

$$f: \quad x = \frac{\sinh \varphi}{\varepsilon - \cosh \varphi}, \quad y = \frac{u \exp \varphi}{\varepsilon - \cosh \varphi}.$$

Then the inverse map is:

$$\varphi = \ln\left(\varepsilon \frac{x-1}{x+1}\right), \quad u = -\frac{2y}{(1-x)^2}.$$
(17)

If we expand  $\mathbb{R}^2$  to the semi-conformal plane  $C^2$  by infinitely distant point and ideal line crossing this point, then the stereographic map f becomes diffeomorphism  $\widetilde{f}$ . This infinitely distant point is the image of point N. The ideal line is the image of straight line belonging to  $S^2(1)$  and crossing the pole:  $x_0 = 1$ ,  $x_1 = 0$ . Do not forget, that we need to exclude lines  $x \neq \pm 1$  from the the plane  $\mathbb{R}^2$ .

Let us now consider the commutative diagram:

The map  $p = \pi \circ \widetilde{f}^{-1}: C^2 \to R$  is defined by this diagram. We find the coordinate form of this map:

$$u = -\frac{2y}{(1-x)^2} \,.$$

So, the map  $p: \mathbb{C}^2 \to \mathbb{R}$  defines the trivial principal fibration with the base  $\mathbb{R}$  and the structure group  $S_1$ .

**Proposition.** The map  $\widetilde{f}: S^2(1) \to C^2$  is conformal.

**Proof.** The metric on G induces the metric on  $C^2$ . In the coordinates x, y it has the form:

$$d\tilde{s}^2 = -dx^2. (18)$$

Let us find the metric of semi-Euclidean sphere from the metric on  $C^2$ . By (17) we get  $d\varphi = \frac{2}{x^2-1}$ . So, by (14) and (18), we find:

$$ds_1^2 = \frac{4}{(x^2 - 1)^2} d\tilde{s}^2.$$

Hence, the linear element of semi-Euclidean sphere differs from the linear element of semi-plane by conformal factor and so, the map  $\tilde{f}$  is conformal.  $\square$ 

We find the fibers equations on  $C^2$ . The 1-parametric fibers family of the fibration  $(S^2(1), \pi, \mathbb{R})$  in the adaptive coordinates (13) is:  $u = c, c \in \mathbb{R}$ . By (17) we get the image of this family under map  $\widetilde{f}$ :

$$y = -c/2 \cdot (x-1)^2. \tag{19}$$

The semi-conformal plane is also fibred by this 1-parametric family of curves. It is parabolas with axis x = 1 and top (1, 0).

# 4. The projective semi-conformal model of the fibration of sphere $S^2(1)$

Now we construct the projective semi-conformal model of the sphere  $S^2(1)$  and the principal fibration on it. We use normalization method of A.P.Norden [5]. A. P. Shirokov in his work [6] constructed conformal models of Non-Euclidean spaces with this method.

In a projective space  $P_n$  a hypersurface  $X_{n-1}$  as an absolute is called *normalized* if with every point  $Q \in X_{n-1}$  is associated:


- 1) line  $P_I$  which has the point Q as the only intersection with the tangent space  $T_{n-1}$ ,
- 2) linear space  $P_{n-2}$  that belongs to  $T_{n-1}$ , but it does not contain the point Q. We call them normals of first and second types,  $P_I$  and  $P_{II}$ .

In order that normalization be polar,  $P_I$  and  $P_{II}$  must be polar with respect to absolute  $X_{n-1}$ .

We expand the semi-Euclidean space  $_2E_1^3$  to a projective space  $P^3$ . Here  $_kE_l^n$  denotes an n-dimensional semi-Euclidean space with metric tensor of rank k, l is the number of negative inertia index in a quadric form. We consider homogeneous coordinates  $(y_0:y_1:y_2:y_3)$  in  $P^3$ , where  $x_i=\frac{y_i}{y_3}$ , i=0,1,2. Thus  $S^2(1):x_0^2-x_1^2=1$  describes the hyperquadric in  $P^3$ :

$$y_0^2 - y_1^2 - y_3^2 = 0. (20)$$

Here the projective basis  $(E_0, E_1, E_2, E_3)$  is chosen in the following way. The vertex  $E_0$  of basis is inside the hyperquadric. The other vertices  $E_1, E_2, E_3$  are on its polar plane,  $y_0 = 0$ . The line  $E_0E_3$  cross the hyperquadric at poles N(1):



0:0:1), N'(1:0:0:-1). Vertices  $E_1, E_2$  lie on the polar of the line  $E_0E_3$ . The vertex of the hyperquadric coincides with the vertex  $E_2$ .

The stereographic map of projective plane  $P^2: y_0 = 0$  to the hyperquadric (20) from the pole N(1:0:0:1) is shown on the picture. Let  $U(0:y_1:y_2:y_3) \in P^2$ . If  $y_3 = 0$ , then the line UN belongs to the tangent plane  $T_N: y_0 - y_3 = 0$  of the hyperquadric (20) at the point N and in this case the intersection point of the line UN with the hyperquadric is not uniquely determined. If  $y_3 \neq 0$ , then the intersection point of line UN with the hyperquadric is unique. So, we choose the line  $E_1E_2: y_3 = 0$  as the line at infinity. In the area  $y_3 \neq 0$  we consider the Cartesian coordinates  $x_1 = \frac{y_1}{y_3}, x_2 = \frac{y_2}{y_3}$ . Then the plane  $\alpha: y_0 = 0, y_3 \neq 0$  becomes a plane with an affine structure  $A^2$ . It is possible to introduce the structure of semi-Euclidean plane  ${}_1E^2$  with the linear element

$$ds_0^2 = dx_1^2. (21)$$

The hyperquadric and the plane  $\alpha$  do not intersect or intersect in two imaginary parallel lines

$$x_1^2 = -1. (22)$$

The restriction of the stereographic projection to the plane  $\alpha$  maps the point  $U(0:x_1:x_2:1)$  into the point  $X_1$ 

$$X_1(-1-x_1^2:2x_1:2x_2:1-x_1^2)$$
. (23)

So, the Cartesian coordinates  $x_i$  can be used as the local coordinates at the hyperquadric except the point of its intersection with the tangent plane  $T_N$ .

We construct an autopolar normalization of the hyperquadric. As a normal of first type we take lines with fixed center  $E_0$  and as a normal of second type we

take their polar lines which belong to the plane  $\alpha$  and cross the vertex  $E_2$  of the hyperquadric. The line  $E_0X_1$  intersects the plane  $\alpha$  at the point

$$X(0:2x_1:2x_2:1-x_1^2)$$
.

In this normalization the polar of the point X intersects the plane  $\alpha$  on normal  $P_{II}$ . Thus for any point X in the plane  $\alpha$  there corresponds a line which does not cross this point. It means that the plane  $\alpha$  is also normalized. Normalization of  $\alpha$  is defined by an absolute quadric (22).

We consider the derivative equations for this normalization. If we take normals of the first type with fixed center  $E_0$ , then the derivative equations ([5], p.204) have the form:

$$\partial_i X = Y_i + l_i X, \nabla_j Y_i = l_j Y_i + p_{ji} X.$$
(24)

The points X,  $Y_i$ ,  $E_0$  define a family of projective frames. Here  $Y_i$  are generating points of normal  $P_{II}$ .

We can calculate the values (X, X),  $(X, Y_i)$  on the plane  $\alpha$  using the quadric form, which is in the left part of equation (20). So,  $(X, X) = -(1 + x_1^2)^2$ .

Let us find coordinates of metric tensor on plane  $\alpha$ . So, we take the Weierstrass standartization

$$(\widetilde{X}, \widetilde{X}) = -1, \quad \widetilde{X} = \frac{X}{1 + x_1^2}.$$

Then the coordinates of metric tensor are scalar product of partial derivatives  $g_{ij} = -(\partial_i \widetilde{X}, \partial_j \widetilde{X})$ :

$$(g_{ij}) = \begin{pmatrix} \frac{4}{(1+x_1^2)^2} & 0\\ 0 & 0 \end{pmatrix}.$$

So, we get the conformal model of polar normalized plane  $\alpha: y_0 = 0, y_3 \neq 0$  with a linear element

$$ds^2 = \frac{dx_1^2}{(1+x_1^2)^2} \,. {25}$$

It means that this non-Euclidean plane is conformal to semi-Euclidean plane  $_1E^2$ . The points X and  $Y_i$  are polarly conjugated,  $(X, Y_i) = 0$ . From this equation and the derivative equations (24) we can get the non-zero connection coefficients:

$$\Gamma_{11}^1 = \Gamma_{12}^2 = \Gamma_{21}^2 = -\frac{2x_1}{1+x_1^2}, \quad \Gamma_{11}^2 = \frac{2x_2}{1+x_1^2}.$$

The sums  $\Gamma_{ks}^s = \partial_k \ln \frac{c}{(1+x_1^2)^2}$  (c = const) are gradient, so the connection is equiaffine. Curvature tensor has the following form:

$$R_{121}^2 = -R_{211}^2 = -\frac{4}{(1+x_1^2)^2}.$$

Ricci curvature tensor  $R_{sk} = R_{isk}^{i}$  is symmetric:  $R_{11} = \frac{4}{(1+x_1^2)^2}$ . Metric  $g_{ij}$  and curvature  $R_{rsk}^{i}$  tensors are covariantly constant in this connection:  $\nabla_k g_{ij} = 0$ ,  $\nabla_l R_{rsk}^{i} = 0$ .

Geodesic curves in this connection satisfy the following equations:

$$x_2 = A(x_1^2 - 1) + Bx_1$$
 and  $x_1 = 0$ ,

where A and B are arbitrary constants. There are parabolas and lines orthogonal to axis  $Ox_1$ .

Let us consider the fibration of this plane by double numbers subalgebra. We write the equations of fibers of semi-Euclidean sphere  $S^2(1)$  in homogeneous coordinates:

$$\begin{cases} (y_0 - y_1)v - y_2 = 0, \\ y_0^2 - y_1^2 - y_3^2 = 0. \end{cases}$$
 (26)

This 1-parametric family of curves fibers the hyperquadric and it defines a fibration on it. The stereographic projection of these fibers from the pole N to the plane  $\alpha$  is:

$$x_2 = -v/2 \cdot (x_1 + 1)^2.$$

It is 1-parametric family of parabolas.

#### Remark

We would obtain the similar results for the space of right cosets by Lie subgroup  $H_2$  (it is the subgroup of invertible dual numbers) and the fibration of the group G by  $H_2$ . However,  $H_2$  is a normal divisor of the group G. Therefore, the spaces of right and left cosets coincide. We have the proposition:

**Proposition** The fibration  $(G, \pi', M' = G/H_2)$  determined by the formula

$$\pi'(x) = \frac{x_1}{x_0} \tag{27}$$

is a trivial principal fibration over the real line  $\mathbb{R}\setminus\{0\}$ . The typical fiber is a plane without a line. The structure group is  $H_2$ .

The equations of fibers are:

$$ux_0 - x_1 = 0, \ u \in \mathbb{R}. \tag{28}$$

This is a 1-parametric family of planes with common axis  $Ox_2$ . It is necessary to remove intersection points of these planes with planes:  $x_0 \pm x_1 = 0$ ,  $x_0 = 0$ .

The left multiplications x' = ax on the group G preserve fibres if and only if when  $a \in H_2$ .

The right multiplications: x' = xb on group G forms the 3-parametric Lie group of linear transformations that preserve the fibration and induce a 1-parametric group of hyperbolic transformations on the base with two fixed points  $u = \pm 1$ :

$$u' = \frac{u + \alpha}{\alpha \cdot u + 1},$$

here  $\alpha = \pi(b)$ .

The right multiplications include involutions of hyperbolic type:

$$R(b) = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ b_2 & b_2 & -1 \end{pmatrix}, R^2 = \text{Id}.$$

They induce the involution on the base:

$$u' = \frac{1}{u}.$$

The similar properties has the fibration of semi-Euclidean sphere by subgroup  $S_2 = \{x \in H_2 | x_0^2 - x_1^2 = 1\}$  and its models.

### ACKNOWLEDGEMENT

I would like to thank Professor Jiri Vanzura for fruitful discussions and support with writing this paper. This work was supported in part by grant No. 201/05/2707 of The Czech Science Foundation and by the Council of the Czech Government MSM 6198959214.

#### References

- [1] B. N. Shapukov, Exercises on Lie groups and its applications, Moskow: RXD (2002).
- [2] Study E., Cartan E. Nombres complexes, Encyclopedie des sciences mathematigues pures et appliquees, t.1. Vol.1. (1908) 329-468.
- [3] V. V. Vishnevskii, A.P. Shirokov, V.V. Shurygin, Spaces over algebras, Izdat. Kazan. univ. (1985).
- [4] N. E. Belova, various articles and works published in Kazan, for example: Mat. vseros. molod. nauc. shkoli-konf. po mat. mod., geom. i algebre. Kazan.(1998) 169-174.
- [5] A. P. Norden, Spaces with affine connection, Moscow: Nauka (1976).
- [6] A. P. Shirokov, Non-Euclidean spaces, Izdat. Kazan. univ. (1997).
- [7] B. A. Rosenfeld, Geometry of Lie Groups, Kluwer (1991).