Low-resolution spectroscopy and spectral energy distributions of selected sources towards σ Orionis

J. A. Caballero¹, L. Valdivielso², E. L. Martín^{2,3}, D. Montes¹, S. Pascual¹, and P. G. Pérez-González¹

¹ Dpto. de Astrofísica y Ciencias de la Atmósfera, Facultad de Física, Universidad Complutense de Madrid, E-28040 Madrid, Spain

² Instituto de Astrofísica de Canarias, Avenida Vía Láctea, 38200 La Laguna, Tenerife, Islas Canarias, Spain

³ University of Central Florida, Dept. of Physics, PO Box 162385, Orlando, FL 32816-2385, USA

Received 15 July 2008 / Accepted 2 September 2008

ABSTRACT

Aims. We investigated in detail nine sources in the direction of the young σ Orionis cluster, which is considered a unique site for studying stellar and substellar formation. The nine sources were selected because of some peculiar properties, such as extremely red infrared colours or too strong H α emission for their blue optical colours.

Methods. We took high-quality, low-resolution spectroscopy (R ~ 500) of the nine targets with ALFOSC at the Nordic Optical Telescope. We also re-analyzed [24]-band photometry from MIPS/Spitzer and compiled the best photometry available at the $ViJHK_s$ passbands and the four IRAC/Spitzer channels for constructing accurate spectral energy distributions covering from 0.55 to $24 \mu m$.

Results. The nine targets were classified into: one Herbig Ae/Be star with a scatterer edge-on disc, two G-type stars, one X-ray flaring, early-M, young star with chromospheric H α emission, one very low-mass, accreting, young spectroscopic binary, two young objects at the brown dwarf boundary with the characteristics of classical T Tauri stars, and two emission-line galaxies, one undergoing star formation, and another one whose spectral energy distribution is dominated by an active galactic nucleus. Besides, we discover three infrared sources associated to overdensities in a cold cloud in the cluster centre.

Conclusions. Low-resolution spectroscopy and spectral energy distributions are a vital tool for measuring the physical properties and the evolution of young stars and candidates in the σ Orionis cluster.

Key words. stars: emission-line, Be – stars: low mass, brown dwarfs – stars: pre-main sequence – Galaxy: open clusters and associations: individual: σ Orionis – galaxies: quasars: emission lines

1. Introduction

The σ Orionis cluster is one of the very few star-forming regions within 1 kpc with massive stars (B2 and earlier – Zinnecker & Yorke 2007). In particular, the Trapezium-like system σ Ori, that gives the name to the cluster, contains five known stars with masses above ~7 M_{\odot} (A, B, D, E, and the recently confirmed F component - Caballero 2007a and references therein; D. M. Peterson et al., in prep.). The nearby Horsehead Nebula, which mane is illuminated by the Trapezium-like system, acts as a retaining wall to the east, while approximately 80% of the cluster members are contained in a circular pool of only 20 arcmin radius, where the extinction is very low (Lee 1968; Béjar et al. 2004; Caballero 2008a). Besides, although there are still uncertainties on its actual heliocentric distance ($d \sim 350-440 \,\mathrm{pc}$ – Caballero 2008b; Mayne & Naylor 2008; Sherry et al. 2008), σ Orionis has been classified as the closest massive-star forming region (Zinnecker & Yorke 2007). Furthermore, with ~3 Ma (Zapatero Osorio et al. 2002a; Oliveira et al. 2006; Caballero 2008b), the cluster has an age intermediate between those of extremely young, highly embedded (~1 Ma: Orion Nebula Cluster, NGC 2024) and relatively cleared, still very young clusters (~5-10 Ma: λ Orionis, NGC 2023) at similar heliocentric distances $(d \sim 400 \,\mathrm{pc}).$

The low absorption, relative closeness, and youth make σ Orionis to be one of the most fruitful hunting grounds for

substellar objects (Béjar et al. 1999, 2001, 2004; Scholz & Eislöffel 2004; Kenyon et al. 2005). Notably, the cluster accounts for more than a half of all known *confirmed* isolated planetary-mass objects in the literature (Zapatero Osorio et al. 2000, 2002b, 2002c, 2007; Barrado y Navascués et al. 2001; Martín et al. 2001a; González-García et al. 2006; Caballero et al. 2007; Scholz & Jayawardhana 2008; Luhman et al. 2008). The reader can find in Caballero (2008c) additional cluster data and useful references for going deeply into all subjects that have been investigated in σ Orionis (from X-ray emission, through Herbig-Haro objects, to brown dwarfs with discs).

For a deeper understanding of the processes that take place in the cluster, Caballero (2008c) tabulated, in his Mayrit catalogue, 241 σ Orionis stars and brown dwarfs with known feaures of youth (e.g. OB spectral types, Li $\scriptstyle 1\lambda$ 6707.8Å in absorption, abnormal strength of other alkali lines because of low surface gravity, H α λ 6562.8 Å in strong emission due to accretion, infrared flux excess due to surrounding disc), 97 candidate cluster members, and 115 back- and foreground sources. Of the 338 cluster members and member candidates, 54 are fainter than the stellar/substellar boundary at $J \approx 14.5$ mag for null extinction (Caballero et al. 2007). The Mayrit catalogue is to date the most comprehensive database of σ Orionis members, and is being used as an input to investigate the cluster spatial distribution, the frequency of discs at different mass intervals, the X-ray emission, the occurrence of wide binaries, or the initial mass function from about $18 M_{\odot}$ to a few Jupiter masses (Caballero 2008b; Luhman et al. 2008; López-Santiago & Caballero 2008). Any

Send offprint requests to: José A. Caballero, investigador Juan de la Cierva at the UCM (e-mail: caballero@astrax.fis.ucm.es).

Fig. 1. False-colour composite images roughly centred on σ Ori at different wavelengths. From left to right: optical photographic $B_J + R_F + I_N$, 2MASS $J + H + K_s$, and *Spitzer* [3.6] + [8.0] + [24] (blue + green + red). Approximate size is 30 × 30 arcmin²; north is up, east is left. The nine analyzed targets are encircled. Note the relative dimming of the Trapezium-like star system and the appearence of a large dust cloud eastwards of it at the reddest wavelengths (see Sect. A). Colour version of all our figures are available in the electronic publication. **Note: the images are available only in the A&A version.**

systematic error in the catalogue (e.g. incompleteness, contamination) may have a pernicious influence on the previous results. It is therefore necessary to confirm or discard membership in the σ Orionis cluster of many Mayrit sources that lack spectroscopy. Some of these disputable sources have been spectroscopically followed-up in the very recent works by Maxted et al. (2008), Gatti et al. (2008), and Sacco et al. (2008). However, there still remain dozens cluster members and member candidates without spectroscopy.

2. Observations and analysis

2.1. The target sample

Here we show low-resolution spectroscopy and spectral energy distributions (from 0.55 to $24 \,\mu$ m) of nine selected sources towards the σ Orionis cluster. We selected the nine targets based on a variety of properties derived from existing catalogues (peculiar H α emission, possible extended point spread function, abnormally red infrared colours, X-ray flares). The earliest and latest *expected* spectral types were B to late M, covering a magnitude interval of $\Delta V \approx 9$ mag (which represents a factor ~4000 in flux, or a factor ~30 in mass). We provide in Table 1 the names, coordinates, and references of the nine targets. False-colour composite images centred on each target, with different wavelength combinations, can be provided upon request to the authors.

2.2. Low-resolution optical spectroscopy with ALFOSC

We used the Andalucía Faint Object Spectrograph and Camera (ALFOSC) at the 2.56 m Nordic Optical Telescope (NOT) in the Spanish Observatorio del Roque de Los Muchachos during five nights in Oct 2006. The ALFOSC detector was the CCD#8, a nimo back illuminated E2V 2k×2k 42–40 with a plate scale of 0.19 arcsec/pixel (2×2 binning). With the grism #5 and the 1.0 arcsec-size slit, we got a dispersion of 3.1 Å/pixel and a resolution of 14.2 Å at 7000 Å (R $\equiv \lambda/\Delta\lambda = 494$).

The total wavelength coverage was 5350–9910Å. The central wavelength (7630Å) was red-shifted with respect to the effective blaze wavelength (6500Å) to maximise the use of the reddest part of the optical spectrum, where late M stars (our faintest objects) emit most of their energy. With the grism #5, the second order (corresponding to bandpasses *B* and *V*) is noticeable at the 20% level only at wavelengths $\lambda > 9700$ Å (no blocking filter was therefore used). Unfortunately, the peak-topeak fringe levels start at 6700Å and increase in intensity redwards (8% at 7500Å, 18% at 8000Å, >30% at >9000Å). The fringing, together with the low number of counts of flux standard stars at the extremes of the total wavelength coverage due to the lower quantum efficiency of the detector, made the final useful wavelength coverage to shrink to 5500–9300Å.

The weather during our run was in general fair, with high thin cirruses only on the last night. The seeing ranged in the interval $\sim 0.7-1.0$ arcsec, which justified our slit size choice. Wind speed was low and humidity maintained roughly stable within

each night (with maximum variations of 15 %). The dates of observation and the exposure times of the nine (combined) spectra are provided in Table 1. Total exposure times ranged from only 30 s for the brightest target to 1 h 20 min for the faintest cluster member. Bias images for posterior calibration were taken at sunsets, while flats and arcs were obtained just before and after each scientific spectrum. The spectra of the five faintest targets are actually the combination of two to four individual spectra, sometimes taken on different nights (e.g. the final spectrum of Mayrit 264077 is the combination of two 900 s-long spectra taken on 2006 Oct 11 and 12). We also got the spectra of GJ 3517 (M9V) during nighttime. It was used as a flux, spectral type, and radial velocity standard/comparison star.

Reduction of the raw images (bias subtraction and flatfielding), spectrum extraction, flux calibration, and combination of the spectra (in this order), was carried out using standard procedures within the IRAF environment (daophot.onedspec). We applied a tiny Doppler shift of 2 Å to the spectra taken during the first two nights due to a systematic error in the calibration. Since the resolution power was 14.2 Å (measured from the arcs and sky lines), this shift had no implications on further analysis steps. Before combination, we checked that there were no significant differences between individual spectra (e.g. no appreciable variation of the H α emission).

The nine final (combined) spectra are shown in Figs. 2 and 3. Note the strong telluric absorption features, especially the O₂ A and B bands at 6850–6960 and 7550–7710 Å and another patent water vapour absorption feature at $\lambda = 7150-7340$ Å (there is other strong feature at >9360 Å, not visible with our display).

2.3. Multi-wavelenght photometry

2.3.1. ViJHK_s[3.6][4.5][5.8][8.0]-band photometry

We collected optical, red optical, near-infrared, and mid-infrared magnitudes from the literature for the nine sources. The *i* and *JHK*_s magnitudes were homogeneously retrieved from the Deep Near Infrared Survey of the Southern Sky (DENIS; Epchtein et al. 1997) and the Two Micron All Sky Survey (2MASS; Skrutskie et al. 2006). The magnitudes in the four channels of the Infrared Array Camera (IRAC; 3.8–8.0 μ m) on the *Spitzer Space Telescope* were taken either from Caballero et al. (2008) or from Hernández et al. (2007), depending on availability. Finally, we compiled the most accurate available visual magnitudes for each target from a heterogeneous collection of sources. We give in Table 2 all the necessary information, including the magnitudes and their origin.

2.3.2. [24]-band photometry with MIPS/*Spitzer* and spectral energy distributions

Hernández et al. (2007) tabulated $24 \mu m$ Multiband Imaging Photometer for *Spitzer* (MIPS) magnitudes ([24]) for only a few of our targets. To expand the wavelength coverage of our spectral energy distributions, we downloaded with the Leopard tool the most recent post-Basic Calibrated Data [24]-band MIPS mosaic



Fig. 4. Spectral energy distributions of the nine targets in Tables 1 and 2. From the shortest to the longest wavelengths, the data points correspond to the passbands Johnson *V*, DENIS *i*, 2MASS *J*, *H*, and K_s , IRAC/*Spitzer* [3.6], [4.5], [5.8], and [8.0], and MIPS/*Spitzer* [24]. The lower triangle indicates an upper limit for the flux at the [24] passband.

covering the whole σ Orionis cluster (P.I.: G. Rieke, programme: Evolution and Lifetimes of Protoplanetary Disks).

First, we looked for sources suitable for studying their observed [24]-band flux. As can be seen in Caballero et al. (2008) and Hernández et al. (2007), intracluster clouds (some of them associated to the Horsehead Nebula), hardly visible in the optical and in the near-infrared, are apparent at wavelengths redwards of $\sim 8 \,\mu m$ (see Sect. A). To avoid contamination by spurious detections, we imposed a relatively high threshold for the source finding at 10σ over the noise background. Next, we performed standard aperture and point-spread-function (PSF) photometry on the mosaic with the IRAF package digiphot, using an aperture radius of 13 arcsec and a background annulus from 39 to 48 arcsec (which leaded to an aperture correction of 1.175 according to the *Spitzer* Science Centre¹). We used $F_{v0} =$ 7.17±0.11 Jy and $\lambda_0 = 23.675 \,\mu\text{m}$ for transforming PSF fluxes into apparent magnitudes. In the previous-to-the-last column in Table 2, we give the measured [24] magnitudes and their errors, computed from the quadratic sum of the error in the PSF photometry and the errors in the aperture correction and the absolute calibration (~7 %; Engelbracht et al. 2007). The 10 σ threshold made the completeness limit of our [24]-band survey to be \sim 9.2 mag.

Only the magnitudes for five of our nine main targets were obtained. Besides, the tabulated magnitude for one of them (Mayrit 102101AB) actually corresponds to a bright knot in a cloud in the north-east vicinity of σ Ori, spatially-coincident with our star (see again Sect. A). The other four stars without MIPS measurement have, therefore, magnitudes $[24] \ge 9.2$ mag.

Using the magnitudes in Table 2 and the zero-point values for conversion between magnitudes and flux densities compiled in the NASA/IPAC/MSC Star and Exoplanet Database², we constructed the spectral energy distributions shown in Fig. 4.

3. Results and discussion

We have classified our nine targets into three categories, depending on their astrophysical nature. The classes are early-type and solar-like stars (3), late-type young stars and brown dwarfs (4), and emission-line galaxies at intermediate redshifts (2). In the last columns of Table 1, we list the spectral types, equivalent widths of $H\alpha$, and cluster membership measured by us of

¹ http://ssc.spitzer.caltech.edu/mips/apercorr/.

² http://nsted.ipac.caltech.edu/NStED/docs/parhelp/Photometry.h



Fig. 2. Flux-calibrated, low-resolution ALFOSC/NOT spectra of the stars and brown dwarfs Mayrit 459340, Mayrit 968292, Mayrit 1064167, Mayrit 797272, Mayrit 102101AB, Mayrit 264077, and Mayrit 358154, from top to bottom. Vertical dotted lines indicate strong telluric absorption features. Beyond ~8000 Å, ripples show regions where fringing contributes to the spectra.



Fig. 3. Same as Fig. 2, but for the emission-line galaxies UCM 0537–0227 (top window) and UCM 0536–0239 (bottom window).

the seven young object candidates. Actually, we measured the pseudo-equivalent width (pEW) of the lines with respect to the pseudo-continuum of the four M-type stars and brown dwarfs. This definition extends to other lines different from the Balmer series (e.g. Li 1). Hereafter, we use the term equivalent width (EW) for simplicity. The error in the EWs comes from the uncertainty in the determination of the (pseudo-)continuum.

The seven young object candidates and the two galaxies are discussed in detail in the following sections.

3.1. Early-type and solar-like stars

3.1.1. Mayrit 459340

Mayrit 459340 (StHA 50) was a new H α emission star found above 10 deg galactic latitude in Stephenson (1986), who investigated a large number of red-sensitive objective prism plates. Soon afterwards, Downes & Keyes (1988) classified Mayrit 459340 as a Be star from a spectrum taken with an image tube scanner. It went unnoticed until Caballero (2007a) listed it as one of the (30) brightest stars of the σ Orionis cluster (Mayrit 459340, with $M \gtrsim 1.4 M_{\odot}$, is also one the most massive cluster stars). According to him, Mayrit 459340 deviates at the level of 3σ to the blue of the spectro-photometric cluster sequence in a V vs. B - V colour-magnitude diagram from Tycho-2 magnitudes. Although its early (Be) spectral type matches, within uncertainties, its blue colours in the optical $(B_T - V_T)$ $= -0.06 \pm 0.11$ mag), Mayrit 459340 is $\Delta B_T = 1.96 \pm 0.06$ mag fainter than the faintest B-type star in σ Orionis (HD 294275 [Mayrit 1227243, B9V]; Caballero 2007a). Its V_T magnitude better matches that of an early- or mid-F spectral-type star in the cluster, or a late-B spectral-type star at an unlikely heliocentric distance thrice larger than to σ Orionis. The first hypothesis would easily explain its H α emission, while the second hypothesis would locate Mayrit 459340 at about 300 pc below the Galactic plane (the Galactic vertical scale height of OBA-type stars is only about 100 pc; e.g. Bahcall & Soneira 1980).

Likewise, Caballero (2007a) also reported that Mayrit 459340 has a disc based on 3.6-8.0µm IRAC/Spitzer photometry from Caballero et al. (2008). Our MIPS/Spitzer [24] photometry (tabulated in Table 2) corroborated his statement: we derived colours $[3.6] - [24] = 1.5 \pm 0.6 \text{ mag}$ and $[3.6] - [8.0] = 0.52 \pm 0.05$ mag, that deviates about 3σ and 10σ with respect the expected values for a disc-less dwarf $([3.6] - [24] \sim [3.6] - [8.0] \sim 0 \text{ mag})$. From the ALFOSC spectrum, shown in the top of Fig. 2, we measured a very intense H α emission of -10 ± 1 Å and derived a spectral type in the interval A2-6V (that corresponds to a most probable mass in the interval 3–4 M_{\odot}). This spectral type approximately matches the blue $B_T - V_T$ colour of Mayrit 459340 and the previous determination by Downes & Keyes (1988). For the classification, we investigated the depth of the blended doublets He I D₃ $\lambda\lambda$ 5875.6.5876.0Å and Na I $\lambda\lambda$ 5890.0.5895.6Å and of the line Ca I λ 5857.5 Å and compared the slope of the continuum of its spectrum with those of early-type standard stars from Valdes et al. (2004). The strong H α emission seems to persist in time, since 1978–1984 (Stephenson 1986), through 1986–1987 (Downes & Keyes 1988), to the present (this work). Besides, the absorption feature of the diffuse interstellar band at about 6280 Å, typical in early-type stars in young clusters, is quite clear. To our knowledge, there is only one scenario that simultaneously explain the A spectral type, the H α emission, the abnormal blue colour for its magnitude (or, alternatively, its abnormal faintness for its colour), and the flux excess at $8-24\,\mu\text{m}$: the *blueing* effect in a Herbig Ae/Be (HAeBe) star with a scatterer edge-on disc (Thé et al. 1994 and references therein).

There are grounds to consider Mayrit 459340 a (long-scale) photometric variable, which is another characteristic typical of HAeBe stars. From data of the All Sky Automated Survey (ASAS-3; Pojmański 2002), we measured $\overline{V} = 11.312 \text{ mag}$, $\sigma_V = 0.045 \text{ mag}$, and $\overline{\delta V} = 0.032 \text{ mag}$ (N = 248), consistent with the Tycho-2 V_T measurement and with Mayrit 459340 lacking large-amplitude photometric variations ($\Delta V \gg 0.045 \text{ mag}$)

Table 1. Names, coordinates, and basic spectroscopic data of observed targets.

Name ^{<i>a</i>}	Alternative name	α (J2000)	δ (J2000)	Reference(s) ^{b}	Date	t_{\exp} (s)	Sp. type	EW(Hα) [Å]	Cluster member
Mayrit 459340	StHA 50	05 38 34.45	-02 28 47.6	St86, DK88, Ca07a	2006 Oct 15	30	A2–6 V e	-10±1	Yes
Mayrit 968292	GSC 04771-00962	05 37 44.92	-02 29 57.3	Ca06, Ca07a	2006 Oct 14	60	G8–K0 V	$+2.5\pm0.5$	Yes?
Mayrit 1064167	2MASS J05390088-0253164	05 39 00.88	-02 53 16.4		2006 Oct 12	150	G2–5 V	$+3.5\pm0.5$	No?
Mayrit 797272	[W96] rJ053751-0235	05 37 51.61	-02 35 25.7	Sh04, Fr06, Ca06	2006 Oct 13	300	M1–3 V e	-4.5 ± 0.5	Yes
Mayrit 102101AB	[W96] rJ053851-0236	05 38 51.45	-02 36 20.6	Wo96, Fr06, He07, Sa08	2006 Oct 11, 13	300+500	M2–4 V e	-32 ± 2	Yes
Mayrit 264077	S Ori J053902.1-023501	05 39 01.94	-02 35 02.9	He07, Ca07	2006 Oct 11, 12	900+900	M2–4 V e	-72 ± 4	Yes
Mayrit 358154	S Ori J053855.4–024121	05 38 55.42	-02 41 20.8	He07, Ca07	2006 Oct 13, 15	1200+3×1200	M4–6 V e	-190 ± 20	Yes
UCM 0537-0227	Mayrit 926051	05 39 32.70	-02 26 15.4	Ca08	2006 Oct 11, 12	1200 + 1200	^c	c	No
UCM 0536-0239	[HHM2007] 668	05 38 41.24	-02 37 37.7	He07, Ca07b	2006 Oct 11, 12	1200 + 1200	^c	c	No

^{*a*} Mayrit and UCM designations follow the nomenclatures introduced in Caballero (2007b) and Zamorano et al. (1994), respectively (truncated coordinates in the UCM HHMM+DDMMW designation are for the 1950.0 equinox).

^b References – St86: Stephenson (1986); DK88: Downes & Keyes (1988); Sh04: Sherry, Walter & Wolk (2004); Fr06: Franciosini, Pallavicini & Sanz-Forcada (2006); He07: Hernández et al. (2007); Ca07a: Caballero (2007a); Ca07b: Caballero (2007b). Ca07: Caballero et al. (2007); Ca08: Caballero (2008c); Sa08: Sacco et al. (2008).

^c Not applicable for galaxies UCM 0537–0227 and UCM 0536–0239.

 Table 2. Photometry of observed targets.

Name	$V \pm \delta V$ [mag]	$i \pm \delta i$ [mag]	$J \pm \delta J$ [mag]	$H \pm \delta H$ [mag]	$\begin{array}{c} K_{\rm s} \pm \delta K_{\rm s} \\ [\rm mag] \end{array}$	[3.6] [mag]	[4.5] [mag]	[5.8] [mag]	[8.0] [mag]	[24.0] [mag]	Sources of photometry ^{<i>a</i>}
Mayrit 459340	11.28 ± 0.05	11.004 ± 0.040	10.666 ± 0.024	10.614 ± 0.024	10.545 ± 0.024	10.40 ± 0.03	10.29 ± 0.03	10.07 ± 0.03	9.88 ± 0.04	8.90 ± 0.55	A122244440
Mayrit 968292	11.06 ± 0.07	10.747 ± 1.000	10.202 ± 0.026	9.960 ± 0.022	9.905 ± 0.022	9.91±0.03	9.90 ± 0.03	9.91±0.03	9.89 ± 0.03	$\gtrsim 9.2^{b}$	A122233330
Mayrit 1064167	12.65 ± 0.12	11.912 ± 0.030	11.451±0.024	11.184 ± 0.023	11.101 ± 0.024	11.04 ± 0.04	10.97 ± 0.04	11.01 ± 0.05	10.95 ± 0.07	≳9.2	B122244440
Mayrit 797272	15.64 ± 0.01	13.446 ± 1.000	11.894 ± 0.026	11.172 ± 0.023	10.977 ± 0.022	10.81 ± 0.03	10.80 ± 0.03	10.79 ± 0.03	10.72 ± 0.03	≳9.2	3122233330
Mayrit 102101AB	17.68 ± 0.09	14.248 ± 0.030	12.438±0.027	11.813±0.023	11.552 ± 0.025	11.23 ± 0.03	11.16±0.03	11.12 ± 0.03	10.96±0.03	3.44 ± 0.27^{c}	3122233330
Mayrit 264077	19.08 ± 0.13	16.380 ± 0.080	14.445 ± 0.035	13.378 ± 0.030	12.607 ± 0.030	11.45 ± 0.04	10.99 ± 0.04	10.52 ± 0.04	9.75 ± 0.04	6.37±0.14	C122255550
Mayrit 358154	≥20.0	18.063 ± 0.250	15.622 ± 0.096	14.842 ± 0.050	13.968 ± 0.061	12.88 ± 0.08	12.52 ± 0.08	12.08 ± 0.09	11.15 ± 0.08	$\gtrsim 9.2^d$	D122255550
UCM 0537-0227	~18.5	17.622 ± 0.250	16.141±0.098	15.099 ± 0.071	14.345 ± 0.092	13.40 ± 0.10	13.21±0.11	12.47 ± 0.10	9.72 ± 0.04	6.23±0.12	D122244440
UCM 0536-0239	18.90 ± 0.10	17.693 ± 0.170	16.375 ± 0.094	15.253 ± 0.080	14.040 ± 0.058	11.88 ± 0.03	11.11±0.03	10.44 ± 0.03	9.39 ± 0.03	5.89 ± 0.12	E122233330

^{*a*} Sources of photometry – 0: this work; 1: DENIS; 2: 2MASS; 3: Hernández et al. (2007); 4: Caballero et al. (2008); 5: Caballero et al. (2007); A: transformed from V_T Tycho-2 (Høg et al. 2000) to *V* Johnson using relations in ESA (1997); B: ASAS-3 (Pojmański 2002); C: Wolk (1996); D: estimated from photographic B_J and R_F magnitudes; E: measured using unpublished *V* data obtained with the Wide Field Camera at the Isaac Newton Telescope, calibrated into the Johnson system through Tycho-2 comparison stars in an overlapping short exposure *V* image in Caballero (2007b).

^b Hernández et al. (2007) measured [24] = 9.80 ± 0.05 mag for Mayrit 968292; it is fainter than the 10σ threshold in our data (note their under estimated errors).

^c The [24] emission of Mayrit 102101AB actually comes from a spatially-coincident intracluster cloud described in the text.

^d Hernández et al. (2007) measured [24] = 8.58 ± 0.03 mag for Mayrit 264077; this emission actually comes from a nearby source (which magnitude we re-determine at [24] = 8.32 ± 0.37 mag).

during years 2000 to 2005. There is not evidence of photometric variability in the USNO-B1.0 catalogue, either (Monet et al. 2003). However, from the spectral energy distribution, the DENIS *i* magnitude, measured in 1996 Feb, seems to be ~ 0.6 -0.7 magnitudes brighter than expected. The value $i = 11.00 \pm$ 0.04 mag is still far away from the $i \sim 10.0$ mag threshold below which brighter DENIS stars in σ Orionis are affected by saturation and non-linear effects (Caballero 2008c). From the ratio between our flux-calibrated spectra, we also estimated that during the observations (2006 Oct), Mayrit 459340 was $\sim 0.9 \text{ mag brighter}$ at 6200–6300 Å than the solar-like, young star candidate Mayrit 968292. However, from the compiled magnitudes in Table 2, Mayrit 459340 is only brighter than Mayrit 968292 at $\lambda \gtrsim 8 \,\mu m$ (where the flux excess by the disc gets more important). Furthermore, the tabulated V_T magnitudes are contemporaries (early 1990s), and Mayrit 459340 was 0.17±0.12 mag *fainter* than Mayrit 968292 at the corresponding effective wavelength of 5320 Å. This apparent brightening at recent epochs, if accompanied by a suitable colour change, would make Mayrit 459340 to lie on the right spectro-photometric sequence of σ Orionis. Recalling the scatterer edge-on disc scenario of Thé et al. (1994), "the variations in brightness and in colour [of a HAeBe star] are caused by a dust cloud in a Keplerian orbit revolving in the outer part of a star's circumstellar disc". Most of the time, the disc of Mayrit 459340 is oriented almost edge on with respect to the observer, and the star is well obscured by the dust cloud. It is at this stage when the dimming is maximum, and scattered blue light is more easily detected. It is expected that when the dust cloud passes away from the front of the HAeBe star, it gets brighter and redder. This effect has been clearly detected in the A3e-type, variable star (with rapid variations) UX Ori (Bibo & Thé 1991). Mayrit 459340 is about 2.6 mag fainter than UX Ori, which probably had made it to get unnoticed until now. High-resolution spectroscopy would enable better constraints on membership through measurments of radial velocity.

3.1.2. Mayrit 968292 and Mayrit 1064167

On the one hand, the Tycho-2 star Mayrit 968292 (GSC 04771-00962) was the brightest (in the optical) σ Orionis member candidate in the Mayrit catalogue without spectroscopy (Caballero 2008c). This was a per se reason for first obtaining an optical spectrum with ALFOSC. With $V \approx 11.1$ mag, it was necessary to exposure only during 1 min. Besides, Mayrit 968292 is the brightest component of a visual binary with a relatively small angular separation ($\rho \approx 9.8 \operatorname{arcmin}, \theta \approx 97 \operatorname{deg}, \Delta J = 0.72 \pm$ 0.03 mag). The secondary, Mayrit 958292, displayed an apparent lithium absorption feature in a spectrum taken by Caballero (2006), which confirmed the star to be a very young member of the Ori OB1b association. If both stars formed a physical pair, then it would be one of the very few wide binaries ($r \approx 3800 \text{ AU}$) detected so far in the σ Orionis cluster (several wide binary candidates were listed in Caballero [2007b]). See also Caballero (2007a) for a consideration on the possible X-ray emission of Mayrit 968292.

On the other hand, Mayrit 1064167 (2MASS J05390088– 0253164) was an anonymous star whose position in the *i* vs. $i - K_s$ diagram made it to fall in the side of the candidate foreand background stars, but still very close to the sequence of the cluster (see fig. 4 in Caballero 2008c). It is the brightest one in a trio of stars separated by 20–40 arcsec. One of the other two components of the asterism has also optical colours and magnitudes typical of solar-like cluster members (e.g. Sherry et al. 2004). However, Caballero (2006) found no trace of lithium in a high signal-to-noise, mid-resolution spectrum of the target. The remaining star in the trio is a foreground late-type dwarf or giant based on a relatively large proper motion and quite red optical/near-infrared colours for its magnitude $(J - K_s)$ 1.30 ± 0.04 mag; Caballero 2008c). Far for extrapolating the non-membership also to Mayrit 1064167, we found three Xray events tabulated by the Second ROSAT Position-Sensitive Proportional Counter (PSPC) Catalogue, 2RXP (ROSAT 2000), and the White-Giommi-Angelini version of the ROSAT PSPC Catalogue, WGACAT (White et al. 2000), at 10–30 arcsec to the star. Although the X-ray events could also be associated to a background extragalactic source, we decided to test photometric selection criteria for cluster members in the literature (in particular: Sherry et al. 2004; Caballero 2008c) with the possible X-ray emitter Mayrit 1064167.

At a first glance at the ALFOSC spectra and the spectral energy distributions of Mayrit 968292 and Mayrit 1064167, the two stars seemed to be G-type stars with no indication of accretion or infrared flux excess. We could only impose lower limits to the [24]-band magnitudes. A careful comparison of the blended Na I D_1D_2 doublets and H α lines indicated that Mayrit 968292 is cooler than Mayrit 1064167 (the absorption of the Na I doublets and the H α line increases and decreases, respectively, towards later spectral types). By comparison with several spectraltype standard stars taken from Jacoby et al. (1984), Montes et al. (1997), and Valdes et al. (2004), we classified Mayrit 968292 as G8–K0V and Mayrit 1064167 as G2–G5V. We obtained a similar classification using the lines ratio H α /Fe I λ 6495.0 Å (Danks & Dennefeld 1994). Other lines typical of solar-like stars that we observed in the spectra of Mayrit 968292 and Mayrit 1064167 were the calcium features Ca 1 λ 6163.8Å and, especially, the infrared triplet Ca II λλλ8498.0,8542.1,8662.1 Å (partially embedded in the fringing region). Unfortunately, our spectra have not enough spectral resolution to clearly detect any other spectroscopic feature of youth (especially lithium).

Although Mayrit 1064167 has an earlier spectral type than Mayrit 968292, the first star is about 1.2 mag fainter than the second one at all passbands. Since such a large reversal of brightness is improbable, we consider that at least one of the two stars is actually a contaminant. The interloper is probably Mayrit 1064167, because Mayrit 968292 follows the cluster sequence and Mayrit 1064167 does not (it has roughly the same magnitudes of mid-K-type young stars in σ Orionis). Albeit Mayrit 968292 remains as a good solar-like cluster member candidate, it is necessary to obtain higher-resolution spectroscopy to confirm its membership. If its youth is verified, its membership in the Mayrit 968292–958292 binary will even so remain unknown for a longer time: because of of the location of σ Orionis in the solar antapex, its cluster members have very low proper motions ($\mu < 10$ mas a⁻¹; Caballero 2007a)

3.2. Late-type young stars and brown dwarfs

We mostly used the PC3 index (Martín, Rebolo & Zapatero Osorio 1996; Martín et al. 1999) for the spectral type classification of the four M-type objects, although we also took into account the information provided by the I1, I2, and I3 indices (indicators of the strength of several CaH and TiO absorption bands; Martín & Kun 1996) and some indices associated to metallic hydrides and oxides (CrH, FeH, H₂O, TiO, and VO). We also compared the indices values with those of spectral standards in the

literature and of GJ 3517, observed with the same instrumental configuration.

3.2.1. Mayrit 797272

Mayrit 797272 ([W96] rJ053751-0235) is a moderate X-ray emitter, early-M star in σ Orionis. It was first discovered by Wolk (1996), who classified it as a ROSAT M0 star with lithium in absorption (EW(Li I) = +0.32 Å) and H α in faint chromospheric emission (EW(H α) = -6.4 Å). A decade later, Franciosini et al. (2006) measured its X-ray count rate with *XMM-Newton* and estimated an M2 spectral type from the R - I, V - I, and I - J colours using the relations by Kenyon & Hartmann (1995) and Leggett et al. (2001). Almost simultaneously, Caballero (2006) took a low signal-to-noise, optical spectrum where the H α line appeared in faint, broadened emission and the Li I line was undetected. Very recently, López-Santiago & Caballero (2008) have found Mayrit 797272 to be an X-ray flaring star from new XMM-Newton data. Finally, Mayrit 797272 forms a visual binary ($\rho \approx 10.5 \operatorname{arcsec}, \theta \approx 5 \operatorname{deg}$) with a slightly brighter star that does not seem to belong to the σ Orionis cluster.

We have derived the most probable spectral type of Mayrit 797272 in the range M1–3V from the ALFOSC spectrum shown in Fig. 2. Our determination is consistent with previous analyses. The H α line in our spectrum of Mayrit 797272 is sharp and with an equivalent width similar to that one measured by Wolk (1996). The broad emission detected by Caballero (2006) was possibly due to the low signal-to-noise ratio and background contamination by a nearby fibre (he used a multifibre spectrograph). The chromospheric H α emission and the absence of flux excess at $\lambda \leq 8 \mu$ m suggest that Mayrit 797272 lacks a disc, which would make the star to rotate slower. The X-ray variability is easier to explain for a very young, fast-rotating, disc-free, almost completely-convective object like Mayrit 797272 (Güdel et al. 1995; Feigelson et al. 2002; Stelzer et al. 2004; Robrade & Schmitt 2005).

3.2.2. Mayrit 102101AB

Mayrit 102101AB ([W96] rJ053851–0236) was, like Mayrit 797272, first detected as a *ROSAT* X-ray source by Wolk (1996). He derived a K5 spectral type and measured EW(H α) \approx -86 Å in a faint spectrum where he did not detect the Li I line. The star has later been detected with *XMM-Newton* and *Chandra* by Franciosini et al. (2006), Caballero (2007b), and Skinner et al. (2008). Using IRAC and MIPS/*Spitzer* data, Hernández et al. (2007) claimed that Mayrit 102101AB harbours an "evolved disc" (i.e. the star exhibits smaller IRAC excesses than optically thick disc systems). Remarkably, Mayrit 102101AB is one of the very few low-mass, spectroscopic binaries (SB2) in the σ Orionis cluster whose orbital parameters have been derived yet ($P = 8.72\pm0.02$ d; Sacco et al. 2008 – they also derived a later spectral type, K9.5, and found lithium in both spectra).

Our observations confirm and contradict some of the observables above. First, using the same technique as for Mayrit 797272, we derived an M3±1 spectral type for Mayrit 102101AB, which matches better the spectrophotometric cluster sequence than previous determinations. The H α emission in our ALFOSC spectrum (EW(H α) = -32±2Å) is not so strong as measured by Wolk (1996), although still satisfying the accreting empirical criterion of Barrado y Navascués & Martín (2003 – the White & Basri [2003]'s criterion cannot be applied because of our poor wavelength resolution). The strong (variable?) H α emission might be originated by material interchange between the two binary components or accretion from a disc. As already noticed by Hernández et al. (2007), the IRAC [3.8] – [8.0] colour of Mayrit 102101AB is only of 0.27 ± 0.05 mag. Furthermore, the apparent flux excess at the MIPS [24] passband is due to an overdensity in a large intracluster dust cloud, and not to the star itself (Sect. A). This overdensity may also be responsible of the slight excess at [8.0]. The close separation between Mayrit 102101A and B, of $(a_1+a_2) \sin i$ = $11.6\pm0.3 R_{\odot}$ (i.e. a few objects radii³; Sacco et al. 2008), prevents the formation of inner disc(s) surrounding one (or both) components. From our point of view, the origin of the X-ray and $H\alpha$ emissions is better explained by the mutual interaction of the two binary components rather than by the presence of an "evolved disc". In any case, we do not completely discard the presence of a cool, *circun-binary* disc at several tens solar radii.

3.2.3. Mayrit 264077

Mayrit 264077 (S Ori J053902.1–023501) is an object at the stellar/substellar boundary in σ Orionis, with a most probable mass of 0.061 M_{\odot} (Caballero et al. 2004, 2007). It had, by far, the reddest near-infrared colours among the objects in the Caballero et al. (2007)'s survey ($J - K_s = 1.84 \pm 0.05 \text{ mag}$ [2MASS], [3.6] – [8.0] = 1.70\pm0.06 \text{ mag}) and fell in a differentiated location in the [3.6] – [8.0] vs. $J - K_s$ colour-colour diagram. Simultaneously, Hernández et al. (2007) measured [24] = 6.45\pm0.03 \text{ mag} (consistent with our photometry) and sorted Mayrit 264077 as a class II object. There was no spectroscopic information on it.

The spectral energy distribution in Fig. 4 complements that one shown in Caballero et al. (2007). The presence of a circumstellar disc is obvious only from photometric data; the ALFOSC spectrum powerfully reinforces the disc existence. We determined an M3±1 spectral type for Mayrit 264077, identical to that of Mayrit 102101AB. However, the H α emission was stronger: EW(H α) = -72±4Å. Besides, we also found [O I] λ 6300.3Å and [S II] $\lambda\lambda$ 6716.4,6730.8Å in forbidden emission (the [N II] λ 6583.4Å is blended with the strong H α line).

It is worthy of notice that the *I*-band light curve of Mayrit 264077 displayed short-, mid-, and long-scale variations in the original data of Caballero et al. (2004). Since it was one of the brightest targets in their sample and its photometry was, therefore, close to the non-linear limit of the detector, Mayrit 264077 was conservatively discarded from the detailed analysis. However, coming back to the original light curve, it is quite possible that those variations were intrinsic to the object and not due to seeing variations and non-linear effects. As shown by Caballero et al. (2006), T Tauri-analog brown dwarfs in σ Orionis with strong H α emission, forbidden lines, and near-infrared flux excess also display large amplitude photometric variations at all considered time scales.

Mayrit 264077 might actually be a very low-mass T Tauri star with a large extinction bluewards of $\sim 1.5 \,\mu\text{m}$ accompanied by a flux excess redwards of it (i.e. the central object is a star brighter than $J = 14.44 \pm 0.04 \,\text{mag}$, and not a brown dwarf).

³ The objects are still contracting and have radii much larger than measured for field dwarf of the same spectral type.

3.2.4. Mayrit 358154

Mayrit 358154 (S Ori J053855.4-024121) was another class II, brown dwarf candidate without spectroscopy in the works by Caballero et al. (2007) and Hernández et al. (2007). It went unnoticed during the optical/near-infrared survey in Caballero et al. (2004) because it fell in the glare of the 3.8 mag-V multiple star σ Ori. Since Mayrit 358154 is 1.2 mag fainter than Mayrit 264077, we expect the first object to be a bona fide brown dwarf, even accounting for a possible edge-on disc. To date, there is only one σ Orionis *star* fainter than $J \sim 14.5$ mag, and it is the source of a well, long-time known Herbig-Haro object (HH 446 [Mayrit 633105] - Reipurth et al. 1998; Andrews et al. 2004). However, Mayrit 358154 is even fainter in J than the Herbig-Haro star. Besides, Mayrit 358154 is at 8.8 arcsec to the north of a brighter photometric cluster member candidate from Sherry et al. (2004). The hypothetical primary was not, however, in the comprehensive cluster catalogue by Caballero (2008c).

The ALFOSC spectrum and the spectral energy distribution strenghten the evidence of Mayrit 358154 having a circum(sub)stellar disc. The flux excess at the [8.0] passband is clear, although the spectral energy distribution seems to drop again at redder wavelengths (our upper limit at the [24] passband is still restricting). The distribution has a peculiar feature at the Hband (Mayrit 633105 should be brighter at $1.6\,\mu m$) whose true origin we fail to ascertain; the feature is significative and cannot be explained by photometric error. The H α emission is even larger than for the previous objects (EW(H α) = -190±20 Å) and we could identify the [O I], [S II], and [N II] forbidden emission lines in the studied wavelength interval. The derived spectral type, M5±1, is consistent with Mayrit 358154 being a highmass brown dwarf in σ Orionis (the earlier brown dwarf candidates known to date in the cluster have M5.5 spectral types). Although it lies on the substellar domain, Mayrit 358154 displays most of the phenomena observed in classical T Tauri stars.

3.3. Background galaxies

UCM0537–0227 was the faintest new σ Orionis photometric member candidate in the Mayrit catalogue of Caballero (2008c). He gave the DENIS *i* and 2MASS *JHK*_s magnitudes for the object, from where one could derive a mass in the middle of the brown dwarf domain, *if* it belonged to the cluster. He also noticed that UCM0537–0227 possibly has an extended, nonstellar point spread function. Its extremely red 2MASS (*J* – *K*_s = 2.34±0.11 mag) and IRAC/*Spitzer* colours measured by Caballero et al. (2008) made us to obtain spectroscopy of it. The low-resolution optical spectrum (top panel of Fig. 3) and the spectral energy distribution of UCM0537–0227 correspond to an emission-line, star-forming galaxy at a spectroscopic redshift z_{sp} = 0.1009 ± 0.0002 with strong PAH features and a moderate star formation rate.

UCM0536–0239 ([HHM2007] 668) and its odd spectral energy distribution from the *I* band to [8.0] was first identified by Caballero (2006). Just afterwards, Hernández et al. (2007) measured its [24]-band magnitude and presented it as one of the very few class I object *candidates* in σ Orionis. Caballero (2007b) noticed that UCM0536–0239 (incorrectly named "Mayrit 111208" in his work) had quite blue I-J and extremely red $J-K_s$ colours for its faint magnitude $(J = 16.14\pm0.10 \text{ mag})$ and found it to be an X-ray emitter detected with the ACIS and HRC-I instruments onboard *Chandra*. The (sub)stellar nature of UCM0536–0239 was put into question by Caballero (2008c): he measured a possible extended point spread function in public optical images.

Finally, Skinner et al. (2008) measured the mean photon energy of UCM0536–0239 at $\overline{E} = 3.40 \text{ keV}$ from ACIS/*Chandra* data, which is a quite high value for a normal star, but typical of active galactic nuclei. The ALFOSC spectrum (bottom panel of Fig. 3) and the spectral energy distribution of UCM0536–0239 favour its classification as a Type 1 obscured quasi-stellar object at $z_{sp} = 0.2362 \pm 0.0005$.

4. Summary

We obtained, compiled, and analyzed low-resolution spectroscopy (R ~ 500), taken with ALFOSC at the Nordic Optical Telescope, and spectral energy distributions, covering from the optical V band to the mid-infrared MIPS/Spitzer [24] band, of nine selected sources towards the young σ Orionis cluster. They covered a very wide range in brightness, that was expected to be translated into a wide range in mass and spectral type. Of the nine optical spectra presented, six correspond to objects without previous spectroscopic observations.

Two of the targets in our sample were identified as low redshift ($z \approx 0.10-0.24$) galaxies with prominent emission-lines in the optical and large fluxes at $8-24\,\mu\text{m}$ due to an active galactic nucleus (UCM0536-0239) and a burst of star formation (UCM0537-0227). Another one seemed to be a solar-like star in the field. Other five objects were classified as bona-fide cluster members with spectral types in the interval A2-6 to M4-6. We failed to detect spectroscopic features of youth in the remaining solar-like, cluster member candidate (Mayrit 968292).

Four of the five confirmed σ Orionis members display peculiar features in their optical spectra and energy distributions:

- Mayrit 459340 is an A2–6-type Herbig Ae/Be star with strong H α emission [EW(H α) = -10 ± 1 Å], flux excess at the MIPS/*Spitzer* [24] band, and a probable scatterer edge-on disc, in analogy to UX Ori.
- Mayrit 102101AB is a young, low-mass, spectroscopic binary whose H α emission could be variable and originated in the mass transfer between A and B.
- Mayrit 264077 is an accretor at the substellar boundary with a developed disc and a flux excess that extends longwards of $24 \,\mu\text{m}$.
- Mayrit 358154 remains as the second faintest σ Orionis member with spectroscopic features of youth and DENIS and 2MASS identification. Because of its strong, probably broad, Hα emission (of up to about –200 Å), [O I], [S II], and [N II] forbidden lines in emission, M5±1 spectral type, flux excess at the four IRAC/Spitzer channels, and faintness (*i* = 18.1±0.2 mag), we classified it as a "classical T Tauri brown dwarf".

The fifth cluster member, Mayrit 797272, is an X-ray flaring star without a disc. Besides, we also present three previuosly unknown mid-infrared sources without optical/near-infrared counterpart. The σ Orionis cluster still hides many astonishing surprises; simple low-resolution spectroscopy and analysis of spectral energy distributions are suitable tools for finding them.

Acknowledgements. We thank the anonymous referee, G. Barro Calvo, M. Rego, V. Villar, and J. Zamorano for helpful comments. The data presented here have been taken using ALFOSC, which is owned by the Instituto de Astrofísica de Andalucía (IAA) and operated at the Nordic Optical Telescope under agreement between IAA and the NBIfAFG of the Astronomical Observatory of Copenhagen. Partial financial support was provided by the Universidad Complutense de Madrid, the Spanish Virtual Observatory, the Spanish Ministerio Educación y Ciencia, and the European Social Fund under grants AyA2005–02750, AyA2005–04286, AyA2005–24102–E, and AyA2007–67458 of the

Programa Nacional de Astronomía y Astrofísica and by the Comunidad Autónoma de Madrid under PRICIT project S-0505/ESP-0237 (AstroCAM).

References

- Andrews, S. M., Reipurth, Bo, Bally, J., Heathcote, S. R. 2004, ApJ, 606, 353
- Bahcall, J. N. & Soneira, R. M. 1980, ApJS, 44, 73
- Barrado y Navascués, D., Zapatero Osorio, M. R., Béjar, V. J. S., et al. 2001, A&A, 377, L9
- Barrado y Navascués, D. & Martín, E. L. 2003, AJ, 126, 2997
- Béjar, V. J. S., Zapatero Osorio, M. R., & Rebolo, R. 1999, ApJ, 521, 671
- Béjar, V. J. S., Martín, E. L., Zapatero Osorio, M. R. et al. 2001, ApJ, 556, 830
- Béjar, V. J. S., Zapatero Osorio, M. R., & Rebolo, R. 2004, AN, 325, 705
- Bibo, E. A. & Thé, P. S. 1991, A&AS, 89, 319
- Bouy, H., Huélamo, N., Martín, E. L. et al. 2008, A&A, in press, eprint arXiv:0808.3890
- Caballero, J. A. 2005, AN, 326, 1007
- Caballero, J. A. 2006, PhD thesis, Universidad de La Laguna, Spain
- Caballero, J. A. 2007a, A&A, 466, 917
- Caballero, J. A. 2007b, AN, 328, 917
- Caballero, J. A. 2008a, MNRAS, 383, 375
- Caballero, J. A. 2008b, MNRAS, 383, 750
- Caballero, J. A. 2008c, A&A, 478, 667
- Caballero, J. A., Béjar, V. J. S., Rebolo, R., Zapatero Osorio, M. R. 2004, A&A, 424, 857
- Caballero, J. A., Martín, E. L., Zapatero Osorio, M. R., et al. 2006, A&A, 445, 143
- Caballero, J. A., Béjar, V. J. S., Rebolo, R. et al. 2007, A&A, 470, 903
- Caballero, J. A., Barrado y Navascués, D., Béjar, V. J. S., et al. 2008, 14th Cambridge Workshop on Cool Stars, Stellar Systems, and the Sun ASP Conference Series, Vol. 384, proceedings of the conference held 5-10 November, 2006, at the Spitzer Science Center and Michelson Science Center, Pasadena, California, USA. Ed. G. van Belle. Poster #31 (in DVD)
- Danks, A. C. & Dennefeld, M. 1994, PASP, 106, 382 Downes, R. A., Keyes, C. D. 1988, AJ, 96, 777
- Engelbracht, C. W., Blaylock, M., Su, K. Y. L. et al., 2007, PASP, 119, 994
- Epchtein, N., de Batz, B., Capoani, L. et al. 1997, Msngr, 87, 27
- European Space Agency (ESA), The Hipparcos and Tycho catalogues, ESA SP-1200, 1997
- Feigelson, E. D., Broos, P., Gaffney, J. A., III et al. 2002, ApJ, 574, 258
- Franciosini, E., Pallavicini, R., Sanz-Forcada, J. 2006, A&A, 446, 501
- Gatti, T., Natta, A., Randich, S., Testi, L., Sacco, G. 2008, A&A, 481, 423
- González-García, B. M., Zapatero Osorio, M. R., Béjar, V. J. S., et al. 2006, A&A, 460, 799
- Güdel, M., Schmitt, J. H. M. M., Benz, A. O., Elias, N. M., II 1995, A&A, 301, 201
- Hernández, J., Hartmann, L., Megeath, T. et al. 2007, ApJ, 662, 1067
- Høg, E., Fabricius, C., Makarov, V. V. et al. 2000, A&A, 355, L27
- Jacoby, G. H., Hunter, D. A., & Christian, C. A. 1984, ApJS, 56, 257
- Kenyon, S. J. & Hartmann, L. 1995, ApJS, 101, 117
- Kenyon, M. J., Jeffries, R. D., Naylor, T., Oliveira, J. M., Maxted, P. F. L. 2005, MNRAS, 356, 89
- Lee, T. A. 1968, ApJ, 152, 913
- Leggett, S. K., Allard, F., Geballe, T. R., Hauschildt, P. H., Schweitzer, A. 2001, ApJ, 548, 908
- van Loon, J. Th. & Oliveira, J. M. 2003, A&A, 405, L33
- López-Santiago, J. & Caballero J. A. 2008, A&A, in press, eprint arXiv:0809.0990
- Luhman, K. L., Hernández J., Downes, J. J., Hartmann, L., Briceño, C. 2008, ApJ, in press, eprint arXiv:0808.0471
- Martín, E. L. & Kun, M. 1996, A&AS, 116, 467
- Martín, E. L., Rebolo, R. & Zapatero Osorio, M. R. 1996, ApJ, 469, 706
- Martín, E. L., Zapatero Osorio, M. R., Barrado y Navascués, D., Béjar, V. J. S., Rebolo, R. 2001a, ApJ, 558, L117
- Martín, E. L., Dougados, C., Magnier, E., et al. 2001b, ApJ, 561, L195
- Maxted, P. F. L., Jeffries, R. D., Oliveira, J. M., Naylor, T., Jackson, R. J. 2008, MNRAS, 385, 2210
- Mayne, N. J. & Naylor, T. 2008, MNRAS, 386, 261
- Monet, D. G., Levine, S. E., Canzian, B. et al. 2003, AJ, 125, 984
- Montes, D., Martín, E. L., Fernández-Figueroa, M. J., Cornide, M., & de Castro, E. 1997, A&AS, 123, 473
- Oliveira, J. M., Jeffries, R. D., van Loon, J. Th., Rushton, M. T. 2006, MNRAS, 369.272
- Pojmański, G. 2002, AcA, 52, 397
- Reipurth, Bo, Bally, J., Fesen, R. A., Devine, D. 1998, Nature, 396, 343
- Robrade, J. & Schmitt, J. H. M. M. 2005, A&A, 435, 1073

- ROSAT Consortium 2000, VizieR On-line Data Catalog: IX/30. Originally published in: ROSAT News 72, 25-May-2000 (2000)
- Sacco, G. G., Franciosini, E., Randich, S., Pallavicini, R. 2008, A&A, 488, 167
- Sanz-Forcada, J., Franciosini, E., Pallavicini, R. 2004, A&A, 421, 715
- Scholz, A. & Eislöffel, J. 2004, A&A, 419, 249
- Scholz, A. & Jayawardhana, R. 2008, ApJ, 672, L49
- Sherry, W. H., Walter, F. M., Wolk, S. J. 2004, AJ, 128, 2316
- Sherry, W. H., Walter, F. M., Wolk, S. J., Adams, N. R. 2008, AJ, 135, 1616
- Skinner, S. L., Sokal, K. R., Cohen, D. H., et al. 2008, ApJ, 683, 796
- Skrutskie, M. F., Cutri, R. M., Stiening, R. et al. 2006, AJ, 131, 1163
- Stelzer, B., Micela, G., Neuhäuser, R. 2004, A&A, 423, 1029 Stephenson, C. B. 1986, ApJ, 300, 779
- Thé, P. S., de Winter, D., Pérez, M. R. 1994, A&AS, 104, 315
- Turner, N. H., ten Brummelaar, T. A., Roberts, L. C., et al. 2008, AJ, 136, 554 Valdes, F., Gupta, R., Rose, J. A., Singh, H. P., & Bell, D. J. 2004, ApJS, 152, 251
- White, N. E., Giommi, P., Angelini, L. 2000, VizieR On-line Data Catalog: IX/31. Originally published in: Laboratory for High Energy Astrophysics (LHEA/NASA), Greenbelt (2000)
- White, R. J. & Basri, G. 2003, ApJ, 582, 1109
- Wolk, S. J. 1996, PhD thesis, State University New York at Stony Brook, USA
- Zamorano, J., Rego, M., Gallego, J. G., et al. 1994, ApJS, 95, 387
- Zapatero Osorio, M. R., Béjar, V. J. S., Martín, E. L., et al. 2000, Science, 290, 103
- Zapatero Osorio, M. R., Béjar, V. J. S., Pavlenko, Ya., et al. 2002a, A&A, 384, 937
- Zapatero Osorio, M. R., Béjar, V. J. S., Martín, E. L., Barrado y Navascués, D., Rebolo, R. 2002b, ApJ, 569, L99
- Zapatero Osorio, M. R., Béjar, V. J. S., Martín, E. L., et al. 2002c, ApJ, 578, 536
- Zapatero Osorio, M. R., Caballero, J. A., Béjar, V. J. S., et al. 2007, A&A, 472,

Zinnecker, H. & Yorke, H. W. 2007, ARA&A, 45, 481

Appendix A: The new infrared sources σ Ori IRS2, 3A, and 3B

There exist dust clouds at relatively short separations from the Trapezium-like system σ Ori AF–B. First, van Loon & Oliveira (2003) discovered σ Ori IRS1, a mid-infrared source at just ~3 arcsec to the OB triple stars, that they associated to a dust cloud. It is originated by the photoevaporation of the outer layers of a binary low-mass star that lies on its densest part (Sanz-Forcada et al. 2004; Caballero 2005, 2007a, 2007b; Turner et al. 2008; Skinner et al. 2008; Bouy et al. 2008). Next, Caballero et al. (2008) discovered a "cloud or shell of cool gas surrounding [...] σ Ori". Although the cloud is obvious in the MIPS/Spitzer [24]-band image (right window in Fig. 1; see also Figure 2 in Hernández et al. 2007), it has received no attention in the literature.

The brightest, densest knot in the cloud falls at a couple arcseconds to our spectroscopic binary Mayrit 102101AB, whose [24]-band photometry was not properly extracted because of the knot. We called it σ Ori IRS2, following the nomenclature introduced by van Loon & Oliveira (2003) for unidentified midinfrared sources without (known) optical/near-infrared counterpart. There are other two additional mid-infrared sources associated to a longitudinal arrangement in the cloud, to the northeast of σ Ori AF–B. Since they seem to form part of the same structure, we called them σ Ori IRS3A and 3B.

Approximate central coordinates (J2000) of the new infrared sources are 05 38 51.0 –02 36 24 (σ Ori IRS2), 05 38 47.6 -02 35 14 (σ Ori IRS3A), and 05 38 58.7 -02 35 30 (σ Ori IRS3B). The origin of the dust cloud that harbours σ Ori IRS2, 3A, and 3B is yet unknown.

List of Objects

' σ Orionis cluster' on page 1 ' σ Ori' on page 1

'Horsehead Nebula' on page 1 'Orion Nebula Cluster' on page 1 'NGC 2024' on page 1 ' λ Orionis' on page 1 'NGC 2023' on page 1 'GJ 3517' on page 2 'Mayrit 1227243' on page 4 'Mayrit 459340' on page 5 'Mayrit 968292' on page 5 '2MASS J05390088-0253164' on page 5 'Mayrit 797272' on page 5 'Mayrit 102101' on page 5 'Mayrit 264077' on page 5 'Mayrit 358154' on page 5 "UCM 0537–0227" on page 5 "UCM 0536–0239" on page 5 'Mayrit 958292' on page 7 'HH 446' on page 9 ' σ Ori IRS1' on page 10 ' σ Ori IRS2' on page 10 ' σ Ori IRS3A' on page 10 ' σ Ori IRS3B' on page 10