On canonical transformations between equivalent Hamiltonian formulations of General Relativity

A.M. $Frolov^*$

Department of Chemistry, University of Western Ontario, N6A 5B7, London, Canada

N. Kiriushcheva[†] and S.V. Kuzmin[‡]

Faculty of Arts and Social Science, Huron University College, N6G 1H3 and Department of Applied Mathematics,
University of Western Ontario, N6A 5B7, London, Canada

(Dated: November 15, 2018)

Abstract

Two Hamiltonian formulations of General Relativity, due to Pirani, Schild and Skinner (Phys. Rev. 87, 452, 1952) and Dirac (Proc. Roy. Soc. A 246, 333, 1958), are considered. Both formulations, despite having different expressions for constraints, allow one to derive four-dimensional diffeomorphism invariance. The relation between these two formulations at all stages of the Dirac approach to the constrained Hamiltonian systems is analyzed. It is shown that the complete sets of their phase-space variables are related by a transformation which satisfies the ordinary condition of canonicity known for unconstrained Hamiltonians and, in addition, converts one total Hamiltonian into another, thus preserving form-invariance of generalized Hamiltonian equations for constrained systems.

^{*}Electronic address: afrolov@uwo.ca

[†]Electronic address: nkiriush@uwo.ca

[‡]Electronic address: skuzmin@uwo.ca

I. INTRODUCTION

The Hamiltonian formulation of General Relativity (GR) is an old subject which is still plagued by some long-standing questions. One of the most important problems, related to essence of Einstein's General Relativity, was the disappearance of four-dimensional diffeomorphism

$$\delta g_{\mu\nu} = -\xi_{\mu;\nu} - \xi_{\nu;\mu} \tag{1}$$

in the Hamiltonian formulation of GR "that has worried many people working in geometrodynamics for so long" [1]¹. According to some authors, in the Hamiltonian formulation of GR, it is possible to restore only spatial diffeomorphism [2, 3] or, according to others, the socalled "special diffeomorphism", for which a non-covariant and field-dependent redefinition of gauge parameters is needed, can be derived [4–6].

In fact, the two Hamiltonian formulations that preserve four-dimensional diffeomorphism have been known for a long time. They are the first Hamiltonian formulations of GR due to Pirani, Schild, and Skinner (PSS) [7] and Dirac [8], both of which allow one to derive the full diffeomorphism from their constraint structure [9, 10]. These two formulations both lead to the expected gauge invariance (1). At the same time, they provide an example that allows us to discuss the conditions under which different Hamiltonian formulations of GR are equivalent. The study of the conditions for which a change of phase-space variables preserves the properties of an original Hamiltonian system is of great importance for constrained dynamical systems. It is especially important in the Hamiltonian formulations of General Relativity where it is customary to perform changes of variables or to introduce new variables. The legitimacy of such changes must be verified. Without showing that a change of variables is canonical, it is impossible to assert that the Hamiltonian formulation of a new theory is equivalent to the original one. This what happened with the formulation that prevailed all others for more than fifty years: the ADM gravity [11]. The introduction of "lapse" and "shift" functions: $N = (-g^{00})^{-1/2}$ and $N^i = -\frac{g^{0i}}{g^{00}}$, which is in fact nonlinear and non-covariant transformation, leads to the formulation (ADM) that is not related

¹ The word "diffeomorphism" is often used as equivalent to the transformation (1) (the semicolon ";" means a covariant derivative); in this paper the same meaning is employed.

to Dirac's (and consequently, to PSS) formulation by any canonical transformation. As we demonstrated in [10], one Poisson bracket is enough to prove the non-canonicity of the ADM variables, e.g.

$$\left\{N\left(x\right),\Pi^{ki}\left(x'\right)\right\} = \left\{\left(-g^{00}\right)^{-1/2},p^{ki}\right\}_{g,p} = -\frac{\delta}{\delta g_{ki}}\left(-g^{00}\right)^{-1/2} = \frac{1}{2}\left(-g^{00}\right)^{-3/2}g^{0k}g^{0i} \neq 0$$
(2)

(where Π^{ki} and p^{ki} are momenta in ADM and Dirac's formulations, respectively, conjugate to the space-space components of the metric tensor, g_{ki}).

As the result, the ADM formulation (the only one in which a restoration of gauge invariance from the complete set of first class constraints has even been considered before) does not produce the expected diffeomorphism invariance (1), as it was recently demonstrated in [6] using the method [12]. The derivation of [6] is the most complete one in the literature but it is not new; the gauge transformations of the ADM variables have been discussed in part previously in [4] and [13]. The gauge transformations that follow from the ADM formulation can be presented *in the form* of diffeomorphism only if a *field-dependent* and *non-covariant* redefinitions of gauge parameters are performed

$$\xi_{diff}^{0} = \left(-g^{00}\right)^{1/2} \varepsilon_{ADM}^{\perp} , \quad \xi_{diff}^{k} = \varepsilon_{ADM}^{k} + \frac{g^{0k}}{g^{00}} \left(-g^{00}\right)^{1/2} \varepsilon_{ADM}^{\perp} , \qquad (3)$$

which, according to [6], "demonstrate the unity of the different symmetries involved". The transformations of [6] are consistent with transformations obtained in [14] using the Lagrangian approach of [15]. However, as we have already pointed out in [16], the field-dependent redefinition of gauge parameters contradicts the essence of all known algorithms of restoration of gauge invariance, as all of them start from the assumption that the gauge parameters should be independent of fields (for details see [16]). To prevent the situation where some manipulations are needed to justify non-canonical transformations, it is better not to perform such transformations from the beginning.

To the best of our knowledge, the equivalence of Hamiltonian formulations of GR, which differ from each other by a change of phase-space variables, was never been analyzed. What is only known to us is a brief statement of DeWitt, which he made for PSS formulation: "four so-called primary constraints could, by a phase transformations, be changed into pure momenta" (see [2] where the author refers to his unpublished report). The connection between the linearized versions of the two formulations of [7] and [8] was analyzed in [17] where it was demonstrated that the two formulations of linearized GR are connected by a change of phase-space variables, which is in fact a canonical transformation in the sense of ordinary Classical Mechanics. Moreover, these formulations, despite having different expressions for Hamiltonians and constraints, give an equivalent description, i.e. the corresponding generators built from the first-class constraints allow one to derive the same gauge invariance (the linearized version of diffeomorphism invariance).

The main goal of this paper is to extend this analysis to the two Hamiltonian formulations of full GR [7–10]. We investigate the relation between the corresponding phase-space variables in both formulations and discuss the effects of such a change of variables at all stages of the Dirac procedure. Another aim is to formulate some general conditions that should be imposed on transformations of phase-space variables for singular systems to preserve the equivalence of different Hamiltonian formulations.

II. COMPARISON OF THE TWO HAMILTONIAN FORMULATIONS OF GR

A starting point of the Hamiltonian formulations of GR in both the approaches of [7] and [8] is the "gamma-gamma" part of the Einstein-Hilbert (EH) Lagrangian which is quadratic in first-order derivatives of the metric tensor (for more details see, e.g., [18, 19])

$$L = \sqrt{-g} g^{\alpha\beta} \left(\Gamma^{\mu}_{\alpha\nu} \Gamma^{\nu}_{\beta\mu} - \Gamma^{\nu}_{\alpha\beta} \Gamma^{\mu}_{\nu\mu} \right) = \frac{1}{4} \sqrt{-g} B^{\alpha\beta\gamma\mu\nu\rho} g_{\alpha\beta,\gamma} g_{\mu\nu,\rho} \tag{4}$$

where

$$B^{\alpha\beta\gamma\mu\nu\rho} = g^{\alpha\beta}g^{\gamma\rho}g^{\mu\nu} - g^{\alpha\mu}g^{\beta\nu}g^{\gamma\rho} + 2g^{\alpha\rho}g^{\beta\nu}g^{\gamma\mu} - 2g^{\alpha\beta}g^{\gamma\mu}g^{\nu\rho}.$$
 (5)

To find the momenta $\pi^{\alpha\beta}$, conjugate to the ten components of the metric tensor $g_{\alpha\beta}$, we rewrite Eq. (4) in a form which explicitly contains the time derivatives of the metric tensor, i.e. in terms of "velocities"

$$L = \frac{1}{4}\sqrt{-g}B^{\alpha\beta0\mu\nu0}g_{\alpha\beta,0}g_{\mu\nu,0} + \frac{1}{2}\sqrt{-g}B^{(\alpha\beta0|\mu\nuk)}g_{\alpha\beta,0}g_{\mu\nu,k} + \frac{1}{4}\sqrt{-g}B^{\alpha\betak\mu\nut}g_{\alpha\beta,k}g_{\mu\nu,t} , \quad (6)$$

where the Latin alphabet is used for spatial components and "0" for a temporal one. The brackets $(\alpha\beta)$ indicate symmetrization in two indices, while the notation (... | ...) is used for symmetrization in two groups of indices, i.e.

$$B^{(\alpha\beta\gamma|\mu\nu\rho)} = \frac{1}{2} \left(B^{\alpha\beta\gamma\mu\nu\rho} + B^{\mu\nu\rho\alpha\beta\gamma} \right).$$

Momenta conjugate to the metric tensor are defined in standard way, and (6) gives

$$\pi^{\gamma\sigma} = \frac{\delta L}{\delta g_{\gamma\sigma,0}} = \frac{1}{2} \sqrt{-g} B^{((\gamma\sigma)0|\mu\nu0)} g_{\mu\nu,0} + \frac{1}{2} \sqrt{-g} B^{((\gamma\sigma)0|\mu\nuk)} g_{\mu\nu,k} .$$
(7)

By using (5), one finds the explicit form of the first term of (7)

$$B^{((\gamma\sigma)0|\mu\nu0)}g_{\mu\nu,0} = g^{00}E^{\mu\nu\gamma\sigma}g_{\mu\nu,0} \tag{8}$$

where

$$E^{\mu\nu\gamma\sigma} \equiv e^{\mu\nu}e^{\gamma\sigma} - e^{\mu\gamma}e^{\nu\sigma}, \ e^{\mu\nu} \equiv g^{\mu\nu} - \frac{g^{0\mu}g^{0\nu}}{g^{00}}.$$
(9)

Note that both $e^{\mu\nu}$ and $E^{\mu\nu\gamma\sigma}$ are zero unless all of the μ , ν , γ , and σ indices differ from 0. The notation e^{km} designates the inverse of the spatial components of the metric tensor, i.e. $g_{nk}e^{km} = \delta_n^m$, and $\frac{\delta}{\delta g_{0\alpha}}e^{\mu\nu} = \frac{\delta}{\delta g_{0\alpha}}E^{\mu\nu\gamma\sigma} = 0$. From (8)-(9) it follows that we cannot express some of the velocities in (7) in terms of momenta, therefore, d primary constraints arise (here d is the dimension of space-time); they are

$$\phi^{0\sigma} = \pi^{0\sigma} - \frac{1}{2}\sqrt{-g}B^{((0\sigma)0|\mu\nu k)}g_{\mu\nu,k} \approx 0.$$
 (10)

If γ and δ indices in (7) are space-like, then (7) is invertible and we find

$$g_{mn,0} = I_{mnpq} \frac{1}{g^{00}} \left(\frac{2}{\sqrt{-g}} \pi^{pq} - B^{((pq)0|\mu\nu k)} g_{\mu\nu,k} \right)$$
(11)

where

$$I_{mnpq} \equiv \frac{1}{d-2} g_{mn} g_{pq} - g_{mp} g_{nq}, \ I_{mnpq} E^{pqkl} = \delta_m^k \delta_n^l .$$
 (12)

The appearance of a singularity in (12) for d = 2 corresponds to the fact that in two dimensions none of the components of (7) can be solved for "velocities". The number of primary constraints (three) in this case equals the number of independent components of the metric tensor in two dimensions [20, 21].

The Hamiltonian is defined by $H = \pi^{\alpha\beta}g_{\alpha\beta,0} - L$. After using (11) to eliminate all the "velocities" $g_{mn,0}$, one finds the following total Hamiltonian:

$$H_T = H_c + g_{00,0}\phi^{00} + 2g_{0k,0}\phi^{0k}, \tag{13}$$

where the 'canonical part'² is

$$H_{c} = \frac{1}{\sqrt{-g}g^{00}} I_{mnpq} \pi^{mn} \pi^{pq} - \frac{1}{g^{00}} I_{mnpq} \pi^{mn} B^{(pq0|\mu\nu k)} g_{\mu\nu,k} + \frac{1}{4} \sqrt{-g} \left[\frac{1}{g^{00}} I_{mnpq} B^{((mn)0|\mu\nu k)} B^{(pq0|\alpha\beta t)} - B^{\mu\nu k\alpha\beta t} \right] g_{\mu\nu,k} g_{\alpha\beta,t} .$$
(14)

For the detailed analysis of (13), including the constraint structure and derivation of the corresponding generators and gauge transformations, see [9].

In this Letter we want to compare the Hamiltonian formulation of GR given by (13) with that of Dirac [8, 10]. Dirac's main idea was based on the fact that the Lagrangian, (4) (it is called below L_{PSS}) can be modified in order to simplify the primary constraints by adding a non-covariant combination of spatial and temporal derivatives that does not affect the equations of motion. This modification leads to the following Lagrangian

$$L_D = L_{PSS} - L^* \tag{15}$$

where L^* [8] is taken by Dirac to be

$$L^* = \left[\left(\sqrt{-g} g^{00} \right)_{,k} \frac{g^{0k}}{g^{00}} \right]_{,0} - \left[\left(\sqrt{-g} g^{00} \right)_{,0} \frac{g^{0k}}{g^{00}} \right]_{,k}.$$
 (16)

The explicit form of (16) can be found using the identity $F_{,\gamma} = \frac{\delta F}{\delta g_{\mu\nu}} g_{\mu\nu,\gamma}$ for the metricdependent functional and rewriting the contravariant components of the metric tensor in terms of $e^{\alpha\beta}$ (see (9)). Finally, we find

$$L^* = \frac{1}{2} \sqrt{-g} A^{\alpha\beta 0\mu\nu k} g_{\alpha\beta,0} g_{\mu\nu,k} , \qquad (17)$$

where we have introduced the following notation

$$A^{\alpha\beta0\mu\nu k} = e^{\alpha\beta} e^{k\mu} g^{0\nu} - e^{\mu\nu} e^{k\alpha} g^{0\beta} + e^{k\alpha} \frac{g^{0\mu} g^{0\nu} g^{0\beta}}{g^{00}} - e^{k\mu} \frac{g^{0\alpha} g^{0\nu} g^{0\beta}}{g^{00}}.$$
 (18)

This relation is obtained by taking into account symmetries $\alpha\beta \Leftrightarrow \beta\alpha$ and $\mu\nu \Leftrightarrow \nu\mu$ in (17) due to the presence of $g_{\alpha\beta,0}g_{\mu\nu,k}$. The important property of $A^{\alpha\beta0\mu\nu k}$ is its antisymmetry with respect to interchange of the two pairs of indices

$$A^{\alpha\beta0\mu\nu k} = -A^{\mu\nu0\alpha\beta k}.$$
(19)

² We shall use this standard terminology, or alternatively 'canonical Hamiltonian', both of which however, can be misleading because for the canonical treatment of singular systems the total Hamiltonian, H_T , not its individual parts, is needed to provide the complete description.

Using the explicit form of (5) we can rewrite the coefficients $B^{(\alpha\beta0|\mu\nu k)}$ in terms of the $A^{\alpha\beta0\mu\nu k}$ and $E^{\alpha\beta\mu\nu}$

$$B^{(\alpha\beta0|\mu\nu k)} = A^{\alpha\beta0\mu\nu k} + g^{0k} E^{\alpha\beta\mu\nu} - 2g^{0\mu} E^{\alpha\beta k\nu}.$$
(20)

Now, the relation between Dirac's Lagrangian L_D and the Lagrangian of PSS, L_{PSS} , takes the form

$$L_D = L_{PSS} - \frac{1}{2}\sqrt{-g}A^{\alpha\beta0\mu\nu k}g_{\alpha\beta,0}g_{\mu\nu,k} . \qquad (21)$$

Note that at the Lagrangian level the difference between the PSS and Dirac formulations does not affect the equations of motion and, in this sense, the two formulations are equivalent. Now, we analyze this difference from the point of view of the Hamiltonian formulation. If we have the two Lagrangians, then we can introduce the two corresponding Hamiltonians which, as we know, give the same gauge invariance [9, 10]. Let us find the relation between their phase-space variables and constraints. This will provide a clue about the changes which can be performed at the Hamiltonian level in a constrained system that will preserve its properties.

The two Lagrangians in (21) differ from each other by the terms linear in time derivatives of the metric tensor; this will affect the expression for conjugate momenta in these two Hamiltonian formulations. For PSS we have

$$\pi^{\gamma\sigma} = \frac{\delta L_{PSS}}{\delta g_{\gamma\sigma,0}},\tag{22}$$

while for the Dirac formulation the momentum is

$$p^{\gamma\sigma} = \frac{\delta L_D}{\delta g_{\gamma\sigma,0}} = \frac{\delta L_{PSS}}{\delta g_{\gamma\sigma,0}} + \frac{\delta L^*}{\delta g_{\gamma\sigma,0}}.$$
(23)

To obtain the relation between these two momenta, we subtract the last two equations, which gives

$$\pi^{\gamma\sigma} - p^{\gamma\sigma} = \frac{\delta}{\delta g_{\gamma\sigma,0}} \left(\frac{1}{2} \sqrt{-g} A^{\alpha\beta 0\mu\nu k} g_{\alpha\beta,0} g_{\mu\nu,k} \right)$$
(24)

or

$$p^{\gamma\sigma} = \pi^{\gamma\sigma} - \frac{1}{2}\sqrt{-g}A^{(\gamma\sigma)0\mu\nu k}g_{\mu\nu,k} .$$
(25)

Equation (25) represents the transformation of phase-space variables for two Hamiltonian formulations of GR, [7] and [8].

Thus, we have two Hamiltonians with two sets of phase-space variables, $(g_{\alpha\beta}, \pi^{\alpha\beta})$ and $(g_{\alpha\beta}, p^{\alpha\beta})$; the momenta of these two sets are connected by the transformation of (25) and

the components of the metric tensor are identical in both formulations. The two sets of fundamental Poisson brackets (PB) are:

$$\{g_{\alpha\beta},\pi^{\mu\nu}\} = \frac{1}{2} \left(\delta^{\mu}_{\alpha}\delta^{\nu}_{\beta} + \delta^{\nu}_{\alpha}\delta^{\mu}_{\beta}\right) \equiv \Delta^{\mu\nu}_{\alpha\beta}, \ \{g_{\alpha\beta},g_{\mu\nu}\} = \left\{\pi^{\alpha\beta},\pi^{\mu\nu}\right\} = 0 \tag{26}$$

and

$$\{g_{\alpha\beta}, p^{\mu\nu}\} = \Delta^{\mu\nu}_{\alpha\beta}, \ \{g_{\alpha\beta}, g_{\mu\nu}\} = \{p^{\alpha\beta}, p^{\mu\nu}\} = 0.$$

$$(27)$$

Note that the conjugate momenta have to be introduced for all generalized coordinates irrespective of whether or not the corresponding time derivatives ("velocities") for particular fields are present in the Lagrangian. In fact, in the first order, metric-affine, formulations of GR due to Einstein [22] (for English translation see [23]) some fields enter the Lagrangian with no derivatives, but nevertheless momenta, conjugate to all of fields, are needed [16, 24].

For regular, i.e. non-singular, systems, the two sets of phase-space variables are related to each other by a canonical transformation if and only if the following relations are fulfilled

$$\{g_{\alpha\beta}, p^{\mu\nu}\}_{g,p} = \{g_{\alpha\beta}, p^{\mu\nu}(\pi, g)\}_{g,\pi} = \Delta^{\mu\nu}_{\alpha\beta} ,$$

$$\{g_{\alpha\beta}, g_{\mu\nu}\}_{g,p} = \{g_{\alpha\beta}, g_{\mu\nu}\}_{g,\pi} = 0, \ \{p^{\alpha\beta}, p^{\mu\nu}\}_{g,p} = \{p^{\alpha\beta}, p^{\mu\nu}\}_{g,\pi} = 0.$$
(28)

In our case, which is based on the use of phase-space variables of [7] and [8], we have to explicitly check in detail only one PB (the rest of PBs is obviously fulfilled) to find

$$\left\{p^{\gamma\sigma}, p^{\delta\rho}\right\}_{g,p} = \left\{\pi^{\gamma\sigma} - \frac{1}{2}\sqrt{-g}A^{(\gamma\sigma)0\mu\nu k}g_{\mu\nu,k} , \ \pi^{\delta\rho} - \frac{1}{2}\sqrt{-g}A^{(\delta\rho)0\alpha\beta m}g_{\alpha\beta,m}\right\}_{g,\pi} = 0.$$
(29)

Note that for pairs with at least one temporal index this PB was calculated in [7, 9], where it is just the PB between primary constraints

$$\left\{\phi^{\sigma 0}, \phi^{\gamma 0}\right\} = 0. \tag{30}$$

However, this result is also valid for all indices. The PB of (29) shows that Dirac's modification of L_{PSS} at the Lagrangian level leads to a Hamiltonian formulation in which the phase-space variables are canonically related to those of PSS.

Our next goal is to consider the effect of such a canonical change of phase-space variables on all steps of the Dirac procedure. We can utilize the PB of (29), and, by rearranging terms, present the canonical part of the PSS Hamiltonian in a different, but equivalent form by explicitly creating (extracting) combinations that correspond to a canonical change of variables

$$\phi^{pq} = \pi^{pq} - \frac{1}{2}\sqrt{-g}A^{(pq)0\mu\nu k}g_{\mu\nu,k} \ . \tag{31}$$

This simple rearrangement is very convenient to study canonical transformations and allows us to present two Hamiltonians of [7] and [8] as one expression that will make transparent the effect of such changes on all steps of the Dirac procedure. By substituting Eq. (31) into the canonical part of the PSS Hamiltonian, (14), and using (18) we obtain the total Hamiltonian, (13), where H_c written in terms of ϕ^{pq} takes the form

$$H_{c} = \frac{1}{\sqrt{-g}g^{00}} I_{mnpq} \phi^{mn} \phi^{pq} - \frac{1}{g^{00}} \phi^{mn} \left(g^{0t} g_{mn,t} - 2g^{0\alpha} g_{\alpha n,m} \right) + \frac{1}{4} \sqrt{-g} \left[\frac{1}{g^{00}} \left(g^{0k} E^{(mn)\mu\nu} - 2g^{0\mu} E^{(mn)k\nu} \right) \left(g^{0t} \delta^{\alpha}_{m} \delta^{\beta}_{n} - 2g^{0\alpha} \delta^{t}_{m} \delta^{\beta}_{n} \right) - B^{\mu\nu k\alpha\beta t} \right] g_{\mu\nu,k} g_{\alpha\beta,t} .$$
(32)

Note that (13) and (32) simultaneously represent the total Hamiltonians for both formulations, [7] and [8]. In the Dirac case $\phi^{\alpha\beta} = p^{\alpha\beta}$; while for PSS, $\phi^{\alpha\beta}$ is given by (31). Both equations (13) and (32) manifestly demonstrate the effect of canonical transformations for the total Hamiltonians:

$$H_T^{PSS}\left(g,\pi\right)|_{g_{\mu\nu}=g_{\mu\nu};\ p^{\gamma\sigma}=\pi^{\gamma\sigma}-\frac{1}{2}\sqrt{-g}A^{(\gamma\sigma)0\mu\nu k}g_{\mu\nu,k}}=H_T^D\left(g,p\right);$$
(33)

and for the generalized Hamiltonian equations:

$$g_{\alpha\beta,0} = \left\{ g_{\alpha\beta}, H_T^{PSS} \right\}, \ \pi_{,0}^{\alpha\beta} = \left\{ \pi^{\alpha\beta}, H_T^{PSS} \right\} \Longrightarrow g_{\alpha\beta,0} = \left\{ g_{\alpha\beta}, H_T^D \right\}, \ p_{,0}^{\alpha\beta} = \left\{ p^{\alpha\beta}, H_T^D \right\}.$$

$$(34)$$

In [9] the PSS formulation was analyzed by considering the combinations of different orders in $\pi^{\alpha\beta}$. Here we will work in orders of $\phi^{\alpha\beta}$ which, due to the simple relation $\{\phi^{\alpha\beta}(g,\pi),\phi^{\mu\nu}(g,\pi)\}_{(g,\pi)} = 0$, makes the amount of calculations absolutely the same for the PSS and Dirac formulations. It also makes transparent the effect of the considered canonical transformation.

Now we calculate the time development of the primary constraints,

$$\phi_{,0}^{0\sigma} = \left\{ \phi^{0\sigma}, H_c \right\}.$$

By working with combinations $\phi^{\alpha\beta}$, we can use associative properties of PB, and therefore $\{\phi^{0\sigma}, H_c\} = -\frac{\delta H_c}{\delta g_{0\sigma}}$, where the variation is not performed inside the expression for $\phi^{\alpha\beta}$, since

 $\left\{\phi^{\alpha\beta},\phi^{\mu\nu}\right\} = 0$. The variation $-\frac{\delta H_c}{\delta g_{0\sigma}}$ leads to the following secondary constraint

$$\chi^{0\sigma} = -\frac{1}{2} \frac{1}{\sqrt{-g}} \frac{g^{0\sigma}}{g^{00}} I_{mnpq} \phi^{mn} \phi^{pq} + \delta^{\sigma}_{m} \left[\phi^{mk}_{,k} + \left(\phi^{pk} e^{qm} - \frac{1}{2} \phi^{pq} e^{km} \right) g_{pq,k} \right] + \frac{1}{2} \sqrt{-g} g^{0\sigma} \left[-g_{mn,kt} E^{mnkt} + \frac{1}{4} g_{mn,k} g_{pq,t} \left(-E^{mnpq} e^{kt} + 2E^{ktpn} e^{mq} + E^{pqnt} e^{mk} \right) \right].$$
(35)

This expression coincides with the secondary constraint in Dirac's formulation [10] where $\phi^{mn} = p^{mn}$. In order to show the equivalence of (35) to the secondary constraint of PSS [9], one has to rewrite this result in terms of π^{km} using (31).

Let us continue the Dirac procedure and consider the time development of the secondary constraint

$$\chi_{,0}^{0\sigma} = \left\{ \chi^{0\sigma}, H_c \right\} + \left\{ \chi^{0\sigma}, g_{00,0} \phi^{00} + 2g_{0k,0} \phi^{0k} \right\}.$$
(36)

We have found it more convenient to perform the calculations in different orders of momenta $\phi^{\mu\nu}$, which are indicated by the numbers in brackets. We start from the PB of $\chi^{0\sigma}$ with the primary constraint for which the highest order contribution gives

$$\left\{\chi^{0\sigma}(2),\phi^{0\gamma}\right\} = -\frac{1}{2}\frac{\delta}{\delta g_{0\gamma}}\left(\frac{g^{0\sigma}}{\sqrt{-gg^{00}}}\right)I_{mnpq}\phi^{mn}\phi^{pq} = -\frac{1}{2}g^{\sigma\gamma}\chi^{00}(2).$$
 (37)

Using this higher order result, (37), as a guide, we have to verify by calculation that to all orders of ϕ^{km} this structure is preserved. The explicit calculation confirms that the following PB is valid to all orders of ϕ^{km}

$$\left\{\chi^{0\sigma},\phi^{0\gamma}\right\} = -\frac{1}{2}g^{\gamma\sigma}\chi^{00}.$$
(38)

We obtained this relation for both formulations, [9] and [10], and it demonstrates the forminvariance of the PB among the primary and secondary constraints for canonically related formulations. Now we proceed and find the PB of the secondary constraints with the canonical part of the Hamiltonian $\{\chi^{0\sigma}, H_c\}$. As before, we start from the highest order contribution, which for this part is of third order in ϕ^{ab} ,

$$\left\{\chi^{0\sigma}, H_{c}\right\}(3) = \left\{\chi^{0\sigma}(2), H_{c}(2)\right\} = -\frac{2}{\sqrt{-g}g^{00}}g^{\sigma d}\phi^{ab}I_{abcd}\frac{g^{0c}}{g^{00}}\left(-\frac{1}{2}\frac{1}{\sqrt{-g}}I_{mnpq}\phi^{mn}\phi^{pq}\right);$$
(39)

it can be presented as a term proportional to $\chi^{00}(2)$ or $\chi^{0c}(2)$, or as a linear combination of both. So, we have many possible and non-unique ways to present this result, which requires us to investigate all combinations, to all orders. Such an approach involves a considerable amount of calculation. The wrong choice can lead to the erroneous conclusion that the time development of the secondary constraint $\chi^{0\sigma}$ is not proportional to the secondary constraints and gives rise to tertiary constraints, etc. The approach that allows one to perform unambiguous calculations (sort out the contributions uniquely in terms of secondary constraints) is presented in the Appendix and here we give only the final result:

$$\left\{\chi^{0\sigma}, H_{c}\right\} = -\left[\frac{2}{\sqrt{-g}}I_{pqmk}\frac{g^{\sigma m}}{g^{00}}\phi^{pq} + g^{0\sigma}g_{00,k} + 2g^{\sigma p}g_{0p,k} + g^{\sigma p}\frac{g^{0q}}{g^{00}}\left(g_{pq,k} + g_{qk,p} - g_{pk,q}\right)\right]\chi^{0k} - \delta_{0}^{\sigma}\chi^{0k}_{,k} + \frac{1}{2}g^{\sigma k}g_{00,k}\chi^{00},$$
(40)

which is easy to compare with the results obtained for the Dirac and PSS formulations in [9] and [10]. Note that again the constraints and structure functions are different for the two formulations; but the whole structure of (40) is form-invariant. In fact, (40) can be presented in the following compact form $\{\chi^{0\sigma}, H_e\} = V^{\sigma}_{\gamma}(g, \phi, \partial) \chi^{0\gamma}$ where upon a canonical transformation not only the constraints, but also the structure functionals V^{σ}_{γ} of one formulation transforms into another independently. Equation (40) proves at the same time the closure of the Dirac procedure for both formulations. This equation, along with (38) and (30), is sufficient to find the gauge generators and derive the gauge transformations for both formulations. We do not want to repeat such calculations here, since they are given in detail in [9] and [10] using two different methods described in [13] and [12]. The result of such calculations is the four-dimensional diffeomorphism invariance (1) (for both formulations) that follows directly using each formalism, without any non-covariant and field-dependent redefinition of the gauge parameters; and the gauge transformation can be written in the covariant form (1) for all components of the metric tensor. For completeness we provide the expression for the canonical part of the Hamiltonian

$$H_{c} = -2g_{0\sigma}\chi^{0\sigma} + \left(2g_{0m}\phi^{mk}\right)_{,k} - \left[\sqrt{-g}E^{mnki}g_{mn,i} - \sqrt{-g}g_{\mu\nu,i}\frac{g^{0\mu}}{g^{00}}\left(g^{\nu k}g^{0i} - g^{\nu i}g^{0k}\right)\right]_{,k}.$$
 (41)

In both formulations, H_c is the sum of the term proportional to the secondary constraints, $-2g_{0\sigma}\chi^{0\sigma}$, and the total spatial derivatives, despite having different expressions for $\chi^{0\sigma}$ and ϕ^{mk} (for details see [9] and [10]).

III. CONCLUSION

We have analyzed the relation between the two Hamiltonian formulations of GR, [7] and [8], which allow one to derive four-dimensional diffeomorphism invariance [9, 10]. It has been shown that the full sets of phase-space variables for these two formulations are related to each other by a transformation of (25), which satisfies the condition of canonicity (28) known for the Hamiltonian formulations of non-singular systems. It also preserves the form-invariance of the expressions for the total Hamiltonians (33). These properties are well known for Hamiltonian formulations of systems with regular (i.e., non-singular) Lagrangians. Despite these similarities with regular systems, the analysis of singular systems has a peculiarity; in the former, condition (28) is necessary and sufficient for equivalence of the two formulations, whereas in the latter case (for singular systems) this condition is necessary, but not sufficient. For singular systems it is also important to preserve the form-invariance of the algebra of constraints, as is the case for the PSS and Dirac formulations (see (30), (38), and (40)). As we demonstrated in [10], if one is tempted to convert the ADM variables into canonical ones then, to satisfy (28) all momenta have to be involved, which leads to the relationships between "old" and "new" momenta given by Eqs. (163-165) of [10]

$$p^{00} = -\Pi \frac{1}{2} \left(-g^{00} \right)^{1/2}, \tag{42}$$

$$p^{0m} = \Pi \frac{1}{2} \left(-g^{00} \right)^{-1/2} g^{0m} + \Pi_k \frac{1}{2} e^{km}, \tag{43}$$

$$p^{pq} = -\Pi \frac{1}{2} \left(-g^{00} \right)^{-3/2} g^{0p} g^{0q} + \Pi_k \frac{1}{2} \left(\frac{g^{0p}}{g^{00}} e^{kq} + \frac{g^{0q}}{g^{00}} e^{kp} \right) + \Pi^{pq}, \tag{44}$$

or Eqs. (166-168) for the inverse transformations (see [10], p. 608-609)

$$\Pi = -2 \left(-g^{00}\right)^{-1/2} p^{00},\tag{45}$$

$$\Pi_n = 2g_{mn}p^{0m} + 2g_{0n}p^{00},\tag{46}$$

$$\Pi^{pq} = p^{pq} + \frac{g^{0q}}{g^{00}} \frac{g^{0p}}{g^{00}} p^{00} - \frac{g^{0p}}{g^{00}} p^{0q} - \frac{g^{0q}}{g^{00}} p^{0p}, \tag{47}$$

where $p^{\alpha\beta}$ and $\{\Pi, \Pi_k, \Pi^{pq}\}$ are momenta conjugate to $g_{\alpha\beta}$ and $\{N, N^k, g_{pq}\}$, respectively.

This transformation (Eqs. (163-165) or Eqs. (166-168) of [10]) can be called canonical, if only the PBs (28) among phase-space variables are concerned. However, as we have shown for the example of ADM gravity, even if momenta conjugate to the ADM variables are introduced according to (28), which is a canonical transformation in the ordinary sense, then these transformations nevertheless destroy the form-invariance of the total Hamiltonian; the space-space "velocities", which were eliminated at the first step of the Dirac procedure, reappear again, and it is not clear what to do with them at this stage of the Hamiltonian analysis (see Sec. 4.4 of [10] for the detail discussion).

For non-singular systems, the canonicity condition (28) is actually independent of the particular form of the unconstrained Hamiltonian; and the canonical transformations automatically convert the Hamiltonian, written in terms of one set of phase-space variables, into another Hamiltonian. Whereas for singular systems the Hamiltonian (that is, the total Hamiltonian) consists of two distinct parts namely, the 'canonical Hamiltonian' and a linear combination of primary constraints, both of which play different roles. In particular, the number of primary constraints corresponds to the number of "velocities" that cannot be solved in terms of momenta, and, for systems with first-class constraints, it defines the number of gauge parameters, an important intrinsic characteristic of a theory. Thus, in the case of singular systems, the explicit form of the total Hamiltonian becomes crucial and only transformations that preserve this form (as in (33)) keep different formulations equivalent. This additional condition makes the canonical transformations for singular systems much more restrictive in comparison to non-singular systems.

We have considered two equivalent Hamiltonian formulations of GR, connected by a relatively simple transformation that involves only a change of momenta. But even from this simple case, it is possible to make a conjecture that the equivalence of different Hamiltonian formulations of singular systems (at least restricted to systems with only first-class constraints) is preserved if the complete set of their phase-space variables is related by a canonical, in ordinary sense, transformation, which in addition, must preserve the forminvariance of the *total* Hamiltonian and the algebra of constraints. The transformation that does not satisfy these conditions will not lead to a Hamiltonian formulation equivalent to the original theory (Einstein's GR, as well as any other singular system).

Let us demonstrate our conjecture by the simple example, or the toy model recently

analyzed in [25], which illustrates the importance of the compliance of both additional conditions: the preservation of the form-invariance of the *total* Hamiltonian and the algebra of constraints. The following model is considered in [25]:

$$L_1 = -\frac{1}{2}\frac{a\dot{a}^2}{N} + \frac{1}{2}Na , \qquad (48)$$

which after the change of variables ("parametrization", according to [25])

$$N = \sqrt{\mu} , \qquad a = a \tag{49}$$

becomes

$$L_2 = -\frac{1}{2}\frac{a\dot{a}^2}{\sqrt{\mu}} + \frac{1}{2}\sqrt{\mu}a \ . \tag{50}$$

The Hamiltonian formulations for these two Lagrangians, (48) and (50), are presented in [25] (Sec. 3) including the restoration of gauge symmetries using the algorithm of [13]. The two total Hamiltonians are:

$$H_T^{(1)} = \dot{N}P - NT_1 , \qquad H_T^{(2)} = \dot{\mu}\pi - 2\mu T_2 , \qquad (51)$$

where P and π are momenta conjugate to N and μ , respectively; and T_1 and T_2 are the secondary first class constraints of two formulations (all details and explicit form of constraints can be found in [25]). The PBs among primary and secondary constraints for these two formulations are:

$$\{P, T_1\} = 0$$
, $\{\pi, T_2\} = \frac{1}{2\mu}T_2$. (52)

This difference in the algebra of constraints is responsible for distinct transformations of these two formulations, because the generators are built from all the first class constraints (this is Dirac's conjecture [27] realized as the algorithm in [13]). Consequently, they depend on the algebra of PBs among all first class constraints. From the canonicity condition, which reads for this model

$$\dot{N}P = \frac{1}{2\sqrt{\mu}}\dot{\mu}P = \dot{\mu}\left(\frac{1}{2\sqrt{\mu}}P\right) = \dot{\mu}\pi , \qquad (53)$$

and using (49) (in addition to (53)), we obtain the relation between the momenta:

$$P = 2\sqrt{\mu}\pi.\tag{54}$$

This phase-space transformation and its inverse leads to

$$\{N, P\}_{\mu,\pi} = \{\mu, \pi\}_{N,P} = 1.$$
(55)

(Note that in this simple model, the transformations of N and P are decoupled from the transformations of a and p.)

The change of phase-space variables (49) and (54) can be called canonical because of (55), but this is not enough for there to be equivalence of the two Hamiltonians (51) because the algebra of constraints (52) is different, contrary to the preservation of the constraint algebra for PSS and Dirac formulations (see the discussion after (40)) that guarantees the equivalence of gauge transformations. In [25, 26] great attention is paid to the condition of canonicity (28) and verification of it for different parametrizations is emphasized. However, the example in [25] provides simple and explicit illustration of our point that the notion of canonicity is more complicated for constrained Hamiltonian systems and the ordinary condition on PBs is necessary, but not sufficient. For this simple example the variables (49) and (54) have canonical PBs (55) and there is no need to use the effective Lagrangian (Sec. 4 of [25]) if only the PBs are concerned. The canonical transformation between Dirac's variables (metric and the corresponding momentum) and ADM is calculated in the extended phase-space approach, and it was shown that it preserves the canonical structure of PBs. However, we would like to mention that these transformations, Eqs. (12) and (18) of [26]and Eqs. (9) and (73) of [25], do not touch the variables from the gauge and ghost sectors of the extended phase space and are in fact the same as we constructed before for ADM gravity (see Eqs. (166-168) of |10|) without reference to any extension of a phase space. In [10] this was done to illustrate that it is possible to make changes from the Dirac to ADM phase-space variables in such a way that the fundamental PBs preserve a canonical form, but this change destroys the constraint structure and as such are not sufficient to ensure the equivalence of the two formulations (see discussion on p. 609 of [10]). In the extended phase-space approach, presented in [25, 26], this question cannot be analyzed because the Hamiltonian for the effective action is presented neither in [26], nor in [25]. This was done only for the simple model described above, where the PBs (55) are canonical. The extended phase-space Hamiltonian is written and the gauge transformations are obtained, but they are different from the transformations that the author calls "correct" [25]; and both are different from the transformations that should be called "canonical" [28].

The motivation for the approach of [25, 26] was based on the results of the ADM Hamiltonian formulation that is not related to the Dirac [8, 10] and PSS [7, 9] formulations by a canonical transformation [10, 29, 30], and consequently, leads to the gauge transformations which differ from diffeomorphism. An answer in the affirmative to the question posed by Shestakova in [25] "Would not it be better to restrict ourself by transformations in phase space of original canonical variables in the sense of Dirac?" (i.e. metric) cannot be reconciled with the statement of [25] "the ADM parametrization is more preferable" (but it does not lead to diffeomorphism when the Dirac method is applied). The solution of this dilemma is proposed in [25]: "we cannot consider the Dirac approach as fundamental and undoubted", and one ought to conclude (based on this choice) that the ADM parametrization is, at least, more fundamental and less doubted. So, as an alternative to the Dirac parametrizationdependent method, another approach is proposed: the extended phase space; by switching from the original Lagrangian to another, effective, Lagrangian the equivalence of which to the original was not proven; further many other questions, related to this proposal, remained unanswered in [25, 26], e.g. "how to construct a generator of gauge transformations in phase space" [26]. It is difficult to discuss in detail the alternative approach before, for example, the equivalence of the EH and effective action is demonstrated. We argue that because the Dirac approach allows unique selection of the canonical variables, it is more preferable than approaches (if they exist) that lead to many canonical formulations. Our answer to the above question (in italic) is "YES" and, because of this, we doubt that the approach proposed in [25, 26] or any other approach "will restore a legitimate status of the ADM parametrization" [25]. The detailed discussion of the many attempts to modify the Dirac method to eliminate, what is, in our opinion, its the most attractive feature, the uniqueness of a canonical formulation, is left for another paper.

IV. ACKNOWLEDGMENT

The authors are grateful to P.G. Komorowski, D.G.C. McKeon and A.V. Zvelindovsky for helpful discussions. The partial support of The Huron University College faculty of Arts and Social Science Research Grant Fund is gratefully acknowledged.

V. APPENDIX

In this Appendix we describe the calculation of $\{\chi^{0\sigma}, H_c\}$, (40), that proves the closure of the Dirac procedure. This result is also needed to find generators and gauge transformations which are discussed in detail for both formulations in [9] and [10].

As we have already pointed out, it is more convenient to perform the calculations in different orders of momenta $\phi^{\mu\nu}$, which are indicated by the numbers in brackets. Comparison of contributions of the highest order in ϕ^{km} into the constraints $\chi^{0\sigma}$, (35), leads to a simple relation: $\chi^{0m}(2) - \frac{g^{0m}}{g^{00}}\chi^{00}(2) = 0$. A generalization of this result to all orders gives

$$\chi^{0m} - \frac{g^{0m}}{g^{00}}\chi^{00} = \psi^{0m} \tag{56}$$

where

$$\psi^{0m} = \phi_{,k}^{mk} + \left(\phi^{pk}e^{qm} - \frac{1}{2}\phi^{pq}e^{km}\right)g_{pq,k} .$$
(57)

If the Dirac procedure is closed in terms of the constraints (χ^{00}, χ^{0k}) , then it is also closed in terms of (χ^{00}, ψ^{0k}) , and vice versa, which follows directly from (56). However, working with the (χ^{00}, ψ^{0k}) – pair allows us to sort terms unambiguously because we have only the following non-zero contributions in both constraints: $\chi^{00}(2)$, $\chi^{00}(0)$, and $\psi^{0k}(1)$, so separating terms of different order in ϕ^{km} simplifies calculations (note that in [9] this procedure could not be simplified to such an extent).

Let us start with $\{\chi^{00}, H_c\}$. In the highest order, this PB, (39), unambiguously gives (there are no derivatives of ϕ^{km} in this expression, so it cannot be proportional to ψ^{0k}):

$$\left\{\chi^{00}, H_c\right\}(3) = \left\{\chi^{00}(2), H_c(2)\right\} = -\frac{2}{\sqrt{-g}} \frac{g^{0d}g^{0c}}{g^{00}g^{00}} \phi^{ab} I_{abcd} \chi^{00}(2).$$
(58)

In the next order, we have contributions with and without derivatives of the momenta ϕ^{km} :

$$\left\{\chi^{00}, H_c\right\}(2) = \left\{\chi^{00}(2), H_c(1)\right\}(\phi\partial\phi) + \left\{\chi^{00}(2), H_c(1)\right\}(\phi\phi),$$
(59)

which we consider separately starting from the terms proportional to $\phi \partial \phi$. Such terms might be presented as proportional to the corresponding orders of the (χ^{00}, ψ^{0k}) – pair through either derivatives of χ^{00} (2) or $\phi \phi_{,k}^{mk}$, both of which have a particular structure in the indices. Again this allows us to sort such terms uniquely:

$$\left\{\chi^{00}(2), H_{c}(1)\right\}(\phi\partial\phi) = -\frac{g^{0k}}{g^{00}}\chi^{00}_{,k}(2)(\phi\partial\phi) - \frac{2}{\sqrt{-g}}I_{mnpq}\phi^{mn}\frac{g^{0p}}{g^{00}}\psi^{0q}(\partial\phi)$$
(60)

where

$$\chi_{,k}^{00}(2) = \chi_{,k}^{00}(2) \left(\phi \partial \phi\right) + \chi_{,k}^{00}(2) \left(\phi \phi\right) = -\frac{1}{\sqrt{-g}} I_{mnpq} \phi^{mn} \phi_{,k}^{pq} - \frac{1}{2} \phi^{mn} \phi^{pq} \left(\frac{1}{\sqrt{-g}} I_{mnpq}\right)_{,k},$$
(61)
(62)

$$\psi^{0m} = \psi^{0m} \left(\partial\phi\right) + \psi^{0m} \left(\phi\right) = \phi_{,k}^{mk} + \left(\phi^{pk} e^{qm} - \frac{1}{2} \phi^{pq} e^{km}\right) g_{pq,k} .$$
 (62)

By performing the completion of (60) to full expressions $\chi^{00}_{,k}(2)$ and $\psi^{0q}(1)$, and combining them with the second term of (59), we obtain

$$\left\{\chi^{00}(2), H_{c}(1)\right\}(\phi\phi) + \frac{g^{0k}}{g^{00}}\chi^{00}_{,k}(2)(\phi\phi) + \frac{2}{\sqrt{-g}}I_{mnpq}\phi^{mn}\frac{g^{0p}}{g^{00}}\psi^{0q}(\phi)$$

that, in case of closure, can be proportional only to $\chi^{00}(2)$. For the second order, we finally have

$$\left\{\chi^{00}, H_{c}\right\}(2) = -\frac{g^{0k}}{g^{00}}\chi^{00}_{,k}(2) - \frac{2}{\sqrt{-g}}I_{mnpq}\phi^{mn}\frac{g^{0p}}{g^{00}}\psi^{0q} + \left[-\frac{g^{0k}g^{0\alpha}g^{0\beta}}{\left(g^{00}\right)^{2}}g_{\alpha\beta,k} + \frac{1}{2}g^{0k}g_{00,k} - \left(\frac{g^{0k}}{g^{00}}\right)_{,k}\right]\chi^{00}(2).$$
(63)

Next, the first order

$$\left\{\chi^{00}, H_c\right\}(1) = \left\{\chi^{00}(2), H_c(0)\right\} + \left\{\chi^{00}(0), H_c(2)\right\} - (Eq.(58))\left(\chi^{00}(2) \to \chi^{00}(0)\right)$$
(64)

can be proportional to only ψ^{0k} , where the last term comes from the completion of (58) to the full (all orders) constraint χ^{00} . Here and in what follows the equations (Eq.(#)) inside the formulae are used to indicate that the right-hand side of (Eq.(#)) must be substituted with the change indicated by " \rightarrow ".

In the last order, by using a similar compensation from the second order, we have

$$\left\{\chi^{00}, H_c\right\}(0) = \left\{\chi^{00}(0), H_c(1)\right\} - \left(Eq.(63)\right)\left(\chi^{00}(2) \to \chi^{00}(0), \psi^{0k} \to 0\right)$$
(65)

which, in the case of closure, should give zero. This is confirmed by explicit calculation.

By calculating (64) and combining it with the results of (58) and (63), we obtain:

$$\left\{\chi^{00}, H_c\right\} = -\frac{2}{\sqrt{-g}} I_{mnpq} \phi^{pq} \frac{g^{0m} g^{0n}}{g^{00} g^{00}} \chi^{00} - \frac{g^{0k}}{g^{00}} \chi^{00}_{,k} - \frac{2}{\sqrt{-g}} I_{mnpq} \phi^{mn} \frac{g^{0p}}{g^{00}} \psi^{0q}$$

$$+\left[-\frac{g^{0k}g^{0\alpha}g^{0\beta}}{\left(g^{00}\right)^2}g_{\alpha\beta,k} + \frac{1}{2}g^{0k}g_{00,k} - \left(\frac{g^{0k}}{g^{00}}\right)_{,k}\right]\chi^{00} - \psi^{0k}_{,k} - \frac{g^{0\alpha}g^{0\beta}}{g^{00}}g_{\alpha\beta,t}\psi^{0t},\tag{66}$$

and in terms of χ^{00} and χ^{0k} equation (66) gives:

$$\left\{\chi^{00}, H_c\right\} = -\left(\frac{2}{\sqrt{-g}}I_{mnpk}\phi^{mn}\frac{g^{0p}}{g^{00}} + \frac{g^{0\alpha}g^{0\beta}}{g^{00}}g_{\alpha\beta,k}\right)\chi^{0k} - \chi^{0k}_{,k} + \frac{1}{2}g^{0k}g_{00,k}\chi^{00}.$$
 (67)

In a similar way, order by order, we consider the PB of ψ^{0k} with H_c , which leads to

$$\left\{\psi^{0m}, H_{c}\right\} = -\frac{2}{\sqrt{-g}} I_{pqkn} \phi^{pq} e^{mn} \psi^{0k} - \frac{g^{0m}}{g^{00}} \frac{g^{0k}}{g^{00}} \chi^{00}_{,k} + \frac{g^{0m}}{g^{00}} \psi^{0k}_{,k} + 2\left(\frac{g^{0k}}{g^{00}}\right)_{,k} \psi^{0k} - \frac{g^{0p}}{g^{00}} e^{qm} \left(g_{pk,q} - g_{pq,k} - g_{qk,p}\right) \psi^{0k}$$

$$(68)$$

$$+ \left[\frac{g^{0m}}{g^{00}}\frac{g^{0q}}{g^{00}}e^{qk}g_{pq,k} + \frac{1}{g^{00}}\left(g^{0q}e^{km} - g^{0k}e^{qm} - g^{0m}e^{qk}\right)g_{0q,k} + \frac{1}{2}e^{km}g_{00,k}\right]\chi^{00}.$$

Now, using Eq. (56) and Eqs. (66) - (68) we find $\{\chi^{0m}, H_c\}$ in terms of χ^{00} and χ^{0k}

$$\left\{\chi^{0m}, H_c\right\} = \left\{\frac{g^{0m}}{g^{00}}\chi^{00} + \psi^{0m}, H_c\right\} = \frac{g^{0m}}{g^{00}}\left\{\chi^{00}, H_c\right\} + \chi^{00}\left\{\frac{g^{0m}}{g^{00}}, H_c\right\} + \left\{\psi^{0m}, H_c\right\}$$
$$= -\left[\frac{2}{\sqrt{-g}}I_{pqkn}\phi^{pq}g^{mn} + g^{0m}g_{00,k} + 2g^{pm}g_{0p,k} + \frac{g^{0p}}{g^{00}}g^{qm}\left(g_{pk,q} + g_{pq,k} - g_{qk,p}\right)\right]\chi^{0k} + \frac{1}{2}g^{km}g_{00,k}\chi^{00}$$
(69)

Finally, Eqs. (67) and (69) which are written in terms of (χ^{00}, χ^{0k}) can be combined into one covariant expression expression, Eq. (40), given in the main text.

- [1] C.J. Isham and K.V. Kuchar, Ann. Phys. 164, 316 (1985).
- [2] B.S. DeWitt, Phys. Rev. **160**, 1113 (1967).
- [3] J. Pullin, in "Cosmology and Gravitation: Xth Brazilian School of Cosmology and Gravitation; 25th Anniversary (1977-2002)" (eds. M. Novello, S.E.P. Bergliaffa, AIP conference proceedings, 2003), vol. 668, p. 141; arXiv:gr-qc/0209008.
- [4] P.G. Bergmann and A. Komar, Int. J. Theor. Phys. 5, 15 (1972).
- [5] J. M. Pons, Class. Quantum Grav. **20**, 3279 (2003).
- [6] P. Mukherjee and A. Saha, Int. J. Mod. Phys. A 24, No 23, 4305 (2009).
- [7] F.A.E. Pirani, A. Schild and S. Skinner, Phys. Rev. 87, 452 (1952).

- [8] P.A.M. Dirac, Proc. Roy. Soc. A 246, 333 (1958).
- [9] N. Kiriushcheva, S.V. Kuzmin, C. Racknor, S.R. Valluri, Phys. Lett. A 372, 5101 (2008).
- [10] N. Kiriushcheva and S.V. Kuzmin, Central Eur. J. Phys. 9, No 3, 576 (2011).
- [11] R. Arnowitt, S. Deser and C.W. Misner, in "Gravitation: An Introduction to Current Research" (ed. L. Witten, Wiley, New York, 1962), p. 227.
- [12] R. Banerjee, H.J. Rothe and K.D. Rothe, Phys. Lett. B 479, 429 (2000).
- [13] L. Castellani, Ann. Phys. **143**, 357 (1982).
- [14] R. Banerjee, S. Gangopadhyay, P. Mukherjee and D. Roy, JHEP 1002, 075 (2010).
- [15] D.M. Gitman and I.V. Tyutin, Quantization of Fields with Constraints (Springer, Berlin, 1990).
- [16] N. Kiriushcheva, S.V. Kuzmin, Eur. Phys. J. C 70, 389 (2010).
- [17] K.R. Green, N. Kiriushcheva and S.V. Kuzmin, Eur. Phys. J. C 71 (2011); online DOI 10.1140/epjc/s10052-011-1678-2.
- [18] L.D. Landau and E.M. Lifshitz, *The Classical Theory of Fields* (fourth ed., Pergamon Press, Oxford, 1975).
- [19] M. Carmeli, Classical Fields: General Relativity and Gauge Theory (World Scientific, New Jersey, 2001).
- [20] N. Kiriushcheva and S.V. Kuzmin, Mod. Phys. Lett. A 21, 899 (2006).
- [21] R.N. Ghalati, N. Kiriushcheva and S.V. Kuzmin, Mod. Phys. Lett. A 22, 17 (2007).
- [22] A. Einstein, Sitzungsber. preuss. Akad. Wiss., Phys.-Math. 1, 414 (1925); The Complete Collection of Scientific Papers (Nauka, Moscow, 1966) vol. 2, p. 171.
- [23] A. Unzicker, T. Case, arXiv:physics/0503046; http://www.alexander-unzicker.de/ae1930.html>.
- [24] N. Kiriushcheva and S.V. Kuzmin, Ann. Phys. **321**, 958 (2006).
- [25] T.P. Shestakova, Class. Quantum Grav. 28, 055009 (2011).
- [26] T.P. Shestakova, Grav. and Cosm. 17, 67 (2011).
- [27] P.A.M. Dirac, Lectures on Quantum Mechanics (Belfer Graduate School of Sciences, Yeshiva University, New York, 1964).
- [28] N. Kiriushcheva, P.G. Komorowski, S.V. Kuzmin: arXiv:1107.2449 [gr-qc].
- [29] B.A. Fayzullaev, Int. J. Mod. Phys. A 25, No 21, 4101 (2010).
- [30] F. Cianfrani, M. Lulli, G. Montani: arXiv:1104.0140 [gr-qc].