ON THE RADIAL GEODESICS OF KERR'S MANIFOLD

ANGELO LOINGER AND TIZIANA MARSICO

ABSTRACT. The so-called stationary-limit surface of Kerr's manifold represents an insuperable barrier for the arriving material elements and light-rays.

Summary. – Introduction. – 1. Different forms of Kerr's interval ds^2 . – 2. A test-particle or a light-ray, moving along a radial geodesic in the negative direction of the radial coordinate, arrive at the "stationary-limit" surface with a zero velocity and a positive, or zero, acceleration. – 3. Action of Hilbert's repulsive effect on the radial geodesics for which $\vartheta = \pi/2$. – 4. Action of Hilbert's repulsive effect on the radial geodesics for which ϑ has a generic value. – Appendix.

PACS 04.20 – General relativity.

Introduction. – At the end of a paper of ours [1] you find the intent to give in a future work a reasonable explanation of those observational data, reported by Remillard et al. [2] and by McClintock [2], which suppositionally concern the event horizons of Kerr's mass points [3], [4]. Well, we do not give here the promised explanation, we give instead a demonstration that Kerr's horizons are incapable of swallowing anything. The reason of this fact is simple: as we shall see, the stationary-limit surface of Kerr's manifold, which is external to the event horizons, exerts a gravitational repulsion on the arriving material elements and light-rays, that is quite similar to the Schwarzschildian repulsive action [5].

1. – As the most adequate choice from the *physical* standpoint, we shall choose the time $t=x^4$ as evolution parameter [6], and we shall compute velocity $\mathrm{d}r/\mathrm{d}t \equiv \dot{r}(t)$ and acceleration $\mathrm{d}^2r/\mathrm{d}t^2 \equiv \ddot{r}(t)$ of the test-particles and of the light-rays travelling along the radial geodesics of Kerr's manifold.

It is however suitable to recall, first of all, some important properties of the coordinate frames describing the above manifold. With Boyer and Lindquist [3], we start from the following form of Kerr's interval:

$$ds^{2} = dx^{2} + dy^{2} + dz^{2} - dt^{2} +$$

$$(1) + \frac{2mr^{3}}{r^{4} + a^{2}z^{2}} \times \left[\frac{r(xdx + ydy) + a(xdy - ydx)}{r^{2} + a^{2}} + \frac{zdz}{r} + dt \right]^{2} ,$$

where r is defined by

(1')
$$[(x^2 + y^2) / (r^2 + a^2)] + z^2 / r^2 = 1 ;$$

here: c = G = 1; m and a are two constants. The case $m \ge a$, with m and a nonnegative parameters, is particularly interesting. If m = 0, the above ds^2 reduces to the customary expression of Minkowskian interval. If $m \ne 0$ and a = 0, eqs. (1) - (1') reduce to the Eddington-Finkelstein form of the solution to Schwarzschild problem of a gravitating point mass m [7]:

(2)
$$ds^{2} = dr^{2} + r^{2}(d\vartheta^{2} + \sin^{2}\vartheta d\varphi^{2}) - dt^{2} + (2m/r)(dr + dt)^{2}$$

remark that this interval was obtained from the standard (Hilbert, Droste, Weyl) form of ds^2 with a coordinate transformation whose derivative is singular at the value 2m of standard r "in just the appropriate way for providing a transformed metric that is regular there." [8].

A different form of Kerr's ds^2 is obtained by putting, see *e.g.* Boyer and Lindquist [3]:

(3)
$$\begin{cases} x = (r^2 + a^2)^{1/2} \sin \vartheta \cos \left[\varphi - \arctan(a/r)\right] &, \\ y = (r^2 + a^2)^{1/2} \sin \vartheta \sin \left[\varphi - \arctan(a/r)\right] &, \\ z = r \cos \vartheta &; \end{cases}$$

the function r of eqs. (1') has been promoted to the rank of a coordinate. If $\Sigma \equiv r^2 + a^2 \cos^2 \vartheta$, we get:

$$ds^{2} = dr^{2} + 2a\sin^{2}\theta dr d\varphi + (r^{2} + a^{2})\sin^{2}\theta d\varphi^{2} + \Sigma(r,\theta) d\theta^{2} - dt^{2} + (4) + [2mr/\Sigma(r,\theta)] (dr + a\sin^{2}\theta d\varphi + dt)^{2}.$$

If we wish a form of ds^2 such that for a=0 the space coordinates reduce to the usual polar coordinates and the interval reduces to the standard form of solution to Schwarzschild problem, we must perform – with Boyer and Lindquist [3] – the following substitutions, where $\Delta(r) \equiv r^2 - 2mr + a^2$:

(5)
$$\begin{cases} r \to r \; ; \; \vartheta \to \vartheta \; , \\ \mathrm{d}\varphi \to \mathrm{d}\varphi - a\,\mathrm{d}r/\Delta(r) \; , \\ \mathrm{d}t \to \mathrm{d}t + 2\,m\,r\,\mathrm{d}r/\Delta(r) \; ; \end{cases}$$

accordingly:

$$ds^{2} = \Sigma (dr^{2}/\Delta + d\vartheta^{2}) + (r^{2} + a^{2})\sin^{2}\vartheta d\varphi^{2} - dt^{2} + (2mr/\Sigma) (a\sin^{2}\vartheta d\varphi + dt)^{2}.$$
(6)

This $\mathrm{d}s^2$ has two "soft" singularities at $r=m\pm(m^2-a^2)^{1/2}$ ("event horizons"). Two remarks: i) the presence of mass m in formulae (5) is indispensable for the passage from an "Eddington-Finkelstein" coordinate frame (eqs. (1) - (1') and (4)) to a "standard" one (eq. (6)); ii) for m=0 and $a\neq 0$, we have, of course, a Minkowskian $\mathrm{d}s^2\equiv\mathrm{d}s_M^2$:

$$ds_M^2 = \frac{r^2 + a^2 \cos^2 \vartheta}{r^2 + a^2} dr^2 + (r^2 + a^2 \cos^2 \vartheta) d\vartheta^2 + (r^2 + a^2) \sin^2 \vartheta d\varphi^2 - dt^2 ;$$
(7)

it is not difficult to give direct proofs of the Minkowskian character of (7) (see Appendix). For $a \to 0$, $\mathrm{d} s_M^2 \to \mathrm{d} r^2 + r^2 (\mathrm{d} \vartheta^2 + \sin^2 \vartheta \, \mathrm{d} \vartheta^2) - \mathrm{d} t^2$. (Analogous considerations hold for the form of $\mathrm{d} s_M^2$ which follows from eq. (4) when m=0.

The Minkowskian nature of Kerr's ds^2 when m=0 and $a\neq 0$ is not a trivial fact, because it proves that the parameter a does not create a spacetime curvature of its own, but it is only a characteristic of some reference frames. Accordingly, it could be "absorbed" by a suitable coordinate transformation.

2. – Let us consider the form (6) of Kerr's interval. The generic radial geodesic of a test-particle or of a light-ray is characterized by the following formulae for the velocity $dr/dt \equiv \dot{r}(t)$ and the acceleration $d^2r/dt^2 \equiv \ddot{r}(t)$, where the constant A is negative for the test-particles and zero for the light-rays:

$$\dot{r}^{2} = \frac{\left(r^{2} + a^{2}\cos^{2}\vartheta - 2mr\right)\left(r^{2} + a^{2} - 2mr\right)}{\left(r^{2} + a^{2}\cos^{2}\vartheta\right)^{2}} - \left|A\right| \frac{\left(r^{2} + a^{2}\cos^{2}\vartheta - 2mr\right)^{2}\left(r^{2} + a^{2} - 2mr\right)}{\left(r^{2} + a^{2}\cos^{2}\vartheta\right)^{3}};$$

the two roots of $r^2 + a^2 \cos^2 \vartheta - 2mr = 0$ are $r_{1,2} := m \pm (m^2 - a^2 \cos^2 \vartheta)^{1/2}$. We have:

(9)
$$\left[\dot{r}^2\right]_{r=r_{1,2}} = 0$$
 ; $\left[\dot{r}^2\right]_{r=\infty} = 1 - |A|$.

$$\ddot{r} = \frac{(r-m)\left[\left(r^2 + a^2\cos^2\vartheta - 2\,m\,r\right) + \left(r^2 + a^2 - 2\,m\,r\right)\right]}{(r^2 + a^2\cos^2\vartheta)^2} - \frac{2\,r\,\left(r^2 + a^2\cos^2\vartheta - 2\,m\,r\right)\left(r^2 + a^2 - 2mr\right)}{(r^2 + a^2\cos^2\vartheta)^3} - |A|\,\left\{\frac{2\,(r-m)\,(r^2 + a^2\cos^2\vartheta - 2\,m\,r)(r^2 + a^2 - 2mr)}{(r^2 + a^2\cos^2\vartheta)^3} + \frac{(r-m)\,(r^2 + a^2\cos^2\vartheta - 2\,m\,r)^2}{(r^2 + a^2\cos^2\vartheta)^3} - \frac{3\,r\,(r^2 + a^2\cos^2\vartheta - 2\,m\,r)^2(r^2 + a^2 - 2mr)}{(r^2 + a^2\cos^2\vartheta)^4}\right\}.$$

$$(10) \qquad -\frac{3\,r\,(r^2 + a^2\cos^2\vartheta - 2\,m\,r)^2(r^2 + a^2 - 2mr)}{(r^2 + a^2\cos^2\vartheta)^4}.$$

For the acceleration $\ddot{r}(t)$ we have:

(11)
$$[\ddot{r}]_{r=r_{1,2}} = \pm \frac{a^2 \sin^2 \vartheta \left(m^2 - a^2 \cos^2 \vartheta \right)^{1/2}}{\left\{ 2m \left[m \pm \left(m^2 - a^2 \cos^2 \vartheta \right)^{1/2} \right] \right\}^2} ;$$

$$[\ddot{r}]_{r=\infty} = 0 \quad .$$

A test-particle or a light-ray, which arrives from the exterior region, travelling in the negative direction of r, arrives at the surface $r = r_1 \equiv m + (m^2 - a^2 \cos^2 \vartheta)^{1/2}$; now:

(12)
$$[\ddot{r}]_{r=r_1} = \frac{a^2 \sin^2 \vartheta \left(m^2 - a^2 \cos^2 \vartheta \right)^{1/2}}{\left\{ 2m \left[m + (m^2 - a^2 \cos^2 \vartheta)^{1/2} \right] \right\}^2} ;$$

this acceleration is positive (or zero for $\vartheta = 0$ and $\vartheta = \pi$); *i.e.*, at $r = r_1$ we have a gravitational repulsion.

Thus, with a zero velocity and a positive (or zero) acceleration no test-particle and no light-ray can cross the barrier $r = r_1$.

Kerr's mass-point – exactly as Schwarzschild's mass point – does not possess any marvellous property. By virtue, *in particular*, of the Hilbertian repulsive effect [5].

3. The surface $r = r_1 \equiv m + (m^2 - a^2 \cos^2 \vartheta)^{1/2}$ is tangent to the surface $r = r^* \equiv m + (m^2 - a^2)^{1/2}$ at $\vartheta = 0$ and $\vartheta = \pi$. For $\vartheta = \pi/2$, we have $r_1[\vartheta = \pi/2] = 2m$; this case is particularly interesting.

With a rather tedious computation, we find that $(\dot{r}^2)_b$, the value of \dot{r}^2 which corresponds to a value zero of the acceleration \ddot{r} (attraction and repulsion balance each other), is given by

(13)
$$(\dot{r}^2)_b \left[\vartheta = \pi/2\right] = \frac{m \left(r - 2m\right) \left(r^2 + a^2 - 2mr\right)^2}{r^3 \left[3 m r^2 - (6m^2 + a^2) r + 4ma^2\right]} ;$$

for a = 0, the right side of eq. (13) yields Hilbert's result [5]: (1/3) [(r -

(-2m)/r]². Where $\dot{r}^2 < (\dot{r}^2)_b$ we have a gravitational attraction. Where $\dot{r}^2 > (\dot{r}^2)_b$

For the light-rays we get from eq. (8) – with A = 0 – that

$$(14) \qquad \dot{r}^{2} \left[\vartheta = \pi/2\right] = \frac{\left(r-2m\right)\left(r^{2}+a^{2}-2mr\right)}{r^{3}} \, \geq \, \left(\dot{r}^{2}\right)_{b} \, \left[\vartheta = \pi/2\right] \quad ;$$

i.e., the light-rays are repulsed for all values of $r \geq 2m$. An analogous result was found by Hilbert [5] for Schwarzschild manifold. (The region r < 2m is quite unphysical).

- **4.** For a generic value of ϑ we have results quite similar to those of sect.
- 3. We write here the expression of $(\dot{r}^2)_b$:

$$(\dot{r}^2)_b = \left[m \left(r^2 + a^2 - 2mr \right)^2 \left(r^2 + a^2 \cos^2 \vartheta - 2mr \right) \left(r^2 - a^2 \cos^2 \vartheta \right) \right] \times \\ \times \left\{ \left(r^2 + a^2 \cos^2 \vartheta \right)^2 \left[a^4 r \cos^2 \vartheta \left(\cos^2 \vartheta - 1 \right) - \right. \\ \left. - a^4 m \cos^2 \vartheta \left(\cos^2 \vartheta + 2 \right) + \left(a^2 r^3 - 4a^2 r^2 m \right) \left(\cos^2 \vartheta - 1 \right) + \right. \\ \left. + 6a^2 m^2 r \cos^2 \vartheta + 3r^4 m - 6r^3 m^2 \right] \right\}^{-1} ;$$

for a=0, we obtain Hilbert's result [5], i.e. $(1/3)[(r-2m)/r]^2$. Where $\dot{r}^2 < (\dot{r}^2)_b$ we have a gravitational attraction. Where $\dot{r}^2 > (\dot{r}^2)_b$ we have a gravitational repulsion. The light-rays are repulsed for all values of $r \geq$ $m + (m^2 + a^2 \cos^2 \vartheta)^{1/2}$.

There is an analogous result for Schwarzschild's manifold for $r \geq 2m$ [5]. A last remark. The repulsive effect makes evident that for Schwarzschild's and Kerr's manifolds the question of qeodesic completeness, though interesting from a geometric standpoint, is destitute of a physical importance.

Acknowledgment. We are grateful to Dr. Giovanni Rastelli, who has obtained with a computer program formulae (13) and (15), and the analytical proof of the Appendix.

APPENDIX

The metric tensor of eq. (7) is regular, time independent, and satisfies $R_{jk} = 0$. Consequently, by virtue of Serini's theorem, it satisfies also $R_{jklm} = 0$, i.e. the ds_M^2 of eq. (7) has a Minkowskian character.

 $R_{jklm}=0$, *i.e.* the $\mathrm{d}s_M^2$ of eq. (7) has a Minkowskian character. An analytical proof: we start from $\mathrm{d}s_M^2=\mathrm{d}x^2+\mathrm{d}y^2+\mathrm{d}z^2-\mathrm{d}t^2$; the transformations

(A1)
$$\begin{cases} x = r' \sin \vartheta' \cos \varphi' & , \\ y = r' \sin \vartheta' \sin \varphi' & , \\ z = r' \cos \vartheta' & \end{cases}$$

give, as it is well known:

(A2)
$$ds_M^2 = dr'^2 + r'^2 (d\vartheta'^2 + \sin^2 \vartheta' d\varphi'^2) - dt^2 ;$$

the transformations:

(A3)
$$\begin{cases} x = (r^2 + a^2)^{1/2} \sin \vartheta \cos \varphi &, \\ y = (r^2 + a^2)^{1/2} \sin \vartheta \sin \varphi &, \\ z = r \cos \vartheta &, \end{cases}$$

give eq. (7), *i.e.*:

$$ds_M^2 = \frac{r^2 + a^2 \cos^2 \vartheta}{r^2 + a^2} dr^2 + (r^2 + a^2 \cos^2 \vartheta) d\vartheta^2 + (r^2 + a^2) \sin^2 \vartheta d\varphi^2 - dt^2 .$$

$$Q.e.d.$$

The relation between $(r', \vartheta', \varphi')$ and (r, ϑ, φ) is:

(A5)
$$\begin{cases} r'^2 = r^2 + a^2 \sin^2 \vartheta & ,\\ \cos^2 \vartheta' = \left(r^2 \cos^2 \vartheta\right) / \left(r^2 + a^2 \sin^2 \vartheta\right) & ,\\ \tan \varphi' = \tan \varphi & \end{cases}$$

(The Minkowskian character of Kerr's interval for m=0 and $a\neq 0$ is quite evident from eqs. (1)-(1).)

References

- [1] A. Loinger and T. Marsico: a) arXiv:0706.3891 v3 [physics.gen-ph] 16 Jul 2007.
- [2] R. Remillard et al., arXiv:astro-ph/0208402 v1 (August 21tst, 2002); J. E. McClintock, Astrophys. J., 652518 (2006-06-04).
- [3] See, in particular: R.P. Kerr, Phys. Rev. Letters, 11 (1963) 237; R.H. Boyer and R.W Lindquist, J. Math. Phys., 8 (1967) 265; C. Møller, The Theory of Relativity, Second Edition (Clarendon Press, Oxford) 1972, p. 482 sqq.; S. Chandrasekhar, The Observatory, 92 (1972) 160.

- [4] A. Loinger, arXiv:gr-qc/9911077 (January 5th, 2000); it is proved here that simple charges of the radial coordinate leave no room for the attribution of extraordinary properties to the field of a Kerr's corpuscle.
- [5] D. Hilbert, Mathem. Annalen, 92 (1924) 1; also in Gesammelte Abhandlungen, Dritter Band (J. Springer, Berlin) 1935, p.258.
- [6] Cf., e.g., V. Fock, The Theory of Space, Time and Gravitation, Second Revised Edition (Pergamon Press, Oxford, etc) 1964, pp. 134 and 135.
- [7] A.S. Eddington, Nature, 113 (1924) 192; D. Finkelstein, Phys. Rev., 110 (1958) 965.
- [8] S. Antoci and D.-E. Liebscher, Astr. Nachrichten, 322 (2001) 137.
- A.L. Dipartimento di Fisica, Università di Milano, Via Celoria, 16 20133 Milano (Italy)
- $\rm T.M.-Liceo$ Classico "G. Berchet", Via della Commenda, 26 20122 Milano (Italy)

 $E\text{-}mail\ address: \verb|angelo.loinger@mi.infn.it||$

E-mail address: martiz64@libero.it