

THE RELATIVE BURNSIDE KERNEL - THE ELEMENTARY ABELIAN CASE

ERIC B. KAHN

ABSTRACT. We give a conjectural description for the kernel of the map assigning to each finite \mathbb{Z}_p -free $G \times \mathbb{Z}_p$ -set its rational permutation module where G is a finite p -group. We prove that this conjecture is true when G is an elementary abelian p -group or a cyclic p -group.

1. INTRODUCTION

Given a prime p and a finite p -group G , there is a natural map between the Burnside ring $A(G)$ and the rational representation ring $R(G)$ taking finite G -sets X to their permutation modules $\mathbb{Q}[X]$. It was shown by J. Ritter and G. Segal [5, 6] that when G is a p -group this map is a surjection and we can study the connection between $A(G)$ and $R(G)$ by looking at generators for the kernel denoted $N(G)$ of the map. These generators were shown by Tornehave in [8] to always be induced by “small” subquotients of G .

From a topological viewpoint we have the Segal conjecture proved by G. Carlsson in [3] which states that the ring $A(G)$ is isomorphic up to completion to the stable cohomotopy group of BG . In addition it was shown by M. Atiyah in [2] that the complex representation ring $R_{\mathbb{C}}(G)$ is isomorphic up to completion to the complex K -theory of the classifying space BG . Thus we gain a connection between stable cohomotopy theory and complex K -theory of BG which can be studied by purely algebraically means.

In a generalization of these ideas, we can consider for a finite p -group G , the natural map between the relative Burnside ring $A(G, \mathbb{Z}_p)$ and the relative rational representation ring $R(G, \mathbb{Z}_p)$ which takes finite, \mathbb{Z}_p -free, $G \times \mathbb{Z}_p$ -sets X to the $\mathbb{Q}[\mathbb{Z}_p]$ -free, $G \times \mathbb{Z}_p$ permutation $\mathbb{Q}[G \times \mathbb{Z}_p]$ -modules $\mathbb{Q}[X]$. We give their definition in section 2.2. It was show by M. Anton in [1] that this map is surjective.

The purpose of this paper is to give a conjecture describing the kernel denoted $N(G, \mathbb{Z}_p)$ of the map in the relative case and to prove the conjecture for some specific p -groups G . We begin in section 2 by giving a brief overview of the classical case, including the construction of induction maps. In section 3 we consider the case when G is an elementary abelian p -group. In this case we decompose both the classical and relative Burnside rings into graded modules and compute their ranks. The main theorem is given in section 3.3 and describes the generators of the relative kernel $N(G, \mathbb{Z}_p)$. In section 4 we offer a conjecture for describing the kernel of the relative map for any p -group G and offer proofs in the cases of G a cyclic p -group and an elementary abelian p -group.

Acknowledgments I am grateful to the department at the University of Kentucky for funding over the summer to work on this problem and to Dr. M. Anton for his enlightening discussions.

2. PRELIMINARIES

2.1. Burnside and representation rings. For a finite group G the isomorphism classes of finite G -sets form a semiring S with respect to disjoint union and direct product. The Burnside ring $A(G)$ is defined to be the Grothendieck construction of the semiring S . In fact $A(G)$ is a free \mathbb{Z} -module with a basis given by the set of left coset spaces $[G/L]$ where L runs thru conjugacy class representatives of subgroups $L < G$. For each such subgroup we define an induction map $L \uparrow: A(L) \rightarrow A(G)$ by sending an L -set X to the G -set $G \times_L X$ where $gl \times x = g \times lx$ for all (g, l, x) in $G \times L \times X$. This definition extends to induction maps $L/C \uparrow: A(L/C) \rightarrow A(G)$ via the pullback map $A(L/C) \rightarrow A(L)$ where L/C is a subquotient of G . The induction maps are \mathbb{Z} -linear but do not preserve the product.

Likewise let T be the semiring of isomorphism classes of finitely generated $\mathbb{Q}[G]$ -modules with respect to direct sum and tensor product. The rational representation ring $R(G)$ is defined to be the Grothendieck construction of T . For $L < G$, the induction map $L \uparrow: R(L) \rightarrow R(G)$ sends a $\mathbb{Q}[L]$ -module M to the $\mathbb{Q}[G]$ -module $\mathbb{Q}[G] \otimes_{\mathbb{Q}[L]} M$ and extends to subquotients as in the Burnside ring case.

The Burnside and representation rings are related by a natural ring homomorphism $f: A(G) \rightarrow R(G)$ sending a G -set X to the permutation $\mathbb{Q}[G]$ -module $\mathbb{Q}[X]$. It is immediate that f commutes with the induction maps.

Definition 2.1. *The Burnside kernel $N(G)$ is the kernel of the map f .*

2.2. Relative Burnside and representation modules. If $\tilde{G} = G \times H$ is a direct product of two finite groups then a \tilde{G} -set is thought of with G acting on the left and H on the right. Let S' be the monoid of isomorphism classes of finite H -free \tilde{G} -sets with respect to disjoint union. The relative Burnside module $A(G, H)$ is the Grothendieck construction of the monoid S' . Then $A(G, H) \subset A(\tilde{G})$ is a free \mathbb{Z} -submodule with a basis given by twisted products $[G \times_{\rho} H]$ where ρ runs thru conjugacy class representatives of homomorphisms $\rho: K \rightarrow H$ with $K < G$ and $gk \times h = g \times \rho(k)h$ for all (g, k, h) in $G \times K \times H$.

Similarly, a $\mathbb{Q}[\tilde{G}]$ -module is thought of with $\mathbb{Q}[G]$ acting on the left and $\mathbb{Q}[H]$ on the right. Let T' be the monoid of isomorphism classes of finitely generated $\mathbb{Q}[H]$ -free $\mathbb{Q}[\tilde{G}]$ -modules with respect to direct sum. The relative rational representation module $R(G, H)$ is the Grothendieck construction of the monoid T' . Then the natural ring homomorphism $f: A(\tilde{G}) \rightarrow R(\tilde{G})$ will restrict to a module homomorphism $f': A(G, H) \rightarrow R(G, H)$.

Definition 2.2. *The relative Burnside kernel $N(G, H)$ is the kernel of f' .*

2.3. Relative induction. The relative induction maps $L/C \uparrow: \tilde{A}(L/C) \rightarrow A(G, H)$ are defined by the usual induction $L/C \uparrow$ restricted to the submodule $\tilde{A}(L/C)$ made of those elements of $A(L/C)$ that land in $A(G, H)$ where L/C is a subquotient of \tilde{G} . The same observation applies to the relative induction map $L/C \uparrow: \tilde{R}(L/C) \rightarrow R(G, H)$. It is immediate that the natural map f' from the relative Burnside module to the relative representation module commutes with the relative induction maps. We conclude by proving from scratch the following

Proposition 2.3. *All induction maps are injective.*

Proof. Let M be any of the monoids S, T, S', T' defining the Burnside and representation modules and their relative versions. The induction map is defined by a

homomorphism $L/C \uparrow: M \rightarrow N$ of monoids extended to the Grothendieck constructions where L/C is a subquotient of \tilde{G} . The Grothendieck construction of M consists of formal differences $[X] - [Y]$ of elements in M such that $[X] - [Y] = [X'] - [Y']$ if and only if

$$[X + Y' + Z] = [X' + Y + Z]$$

for some $[Z]$ in M . In particular, if $L/C \uparrow [X] - L/C \uparrow [Y] = 0$ then

$$[L/C \uparrow X + V] = [L/C \uparrow Y + V]$$

for some $[V]$ in N . By restricting the G -structure to an L -structure we have a restriction map $L \downarrow$ such that $L \downarrow (L/C \uparrow [X]) = [\tilde{G} : L][X]$. In particular,

$$[\tilde{G} : L][X] + L \downarrow [V] = [\tilde{G} : L][Y] + L \downarrow [V].$$

Since each element of M has a unique decomposition into a sum of irreducible elements, we conclude that $[X] = [Y]$ proving the injectivity of the induction map. \square

2.4. The p -group case. For G a finite p -group it was shown by Tornehave [8] that $N(G)$ is generated by the induced kernels $L/C \uparrow N(L/C)$ where L/C runs thru all subquotients of G that are isomorphic to the elementary abelian group $\mathbb{Z}_p \times \mathbb{Z}_p$, the dihedral group, or the nonabelian group of order p^3 and exponent p . Combining this with the Ritter-Segal [5, 6] proof for the surjectivity of f we get a well understood short exact sequence:

$$(1) \quad 0 \rightarrow N(G) \rightarrow A(G) \xrightarrow{f} R(G) \rightarrow 0.$$

In the abelian case $G = \mathbb{Z}_p \times \mathbb{Z}_p$ for instance, it is shown in [4] that $N(G)$ is the free cyclic group generated by

$$(2) \quad [G] - \sum [G/C] + p[G/G]$$

where C runs thru all proper cyclic subgroups of G . In the relative case $\tilde{G} = G \times H$ with G a finite p -group and $H = \mathbb{Z}_p$ it is known by [1] only that we have a short exact sequence

$$(3) \quad 0 \rightarrow N(G, H) \rightarrow A(G, H) \xrightarrow{f'} R(G, H) \rightarrow 0$$

and the purpose of this paper is to study $N(G, H)$.

2.5. A useful trick.

Lemma 2.4. *Consider the chain complex of finitely generated free \mathbb{Z} -modules*

$$0 \rightarrow A \xrightarrow{\alpha} B \xrightarrow{\beta} C \rightarrow 0$$

with α injective and β surjective. If the cokernel of α is a free module and the rank of the image of α equals the rank of the kernel of β , then the sequence is exact.

Proof. Since $Im(\alpha) \subset Ker(\beta)$ and $Coker(\alpha)$ is free, we have the free \mathbb{Z} -submodule $Ker(\beta)/Im(\alpha) \subset B/Im(\alpha)$. But the rank of the image of α equals the rank of the kernel of β so that $Ker(\beta)/Im(\alpha)$ is torsion. Therefore $Ker(\beta)/Im(\alpha) = 0$. \square

3. THE RELATIVE BURNSIDE KERNEL FOR ELEMENTARY ABELIAN GROUPS

3.1. Notations. In this section let $G = \mathbb{Z}_p^n$ and $H = \mathbb{Z}_p$ so that $\tilde{G} = G \times H = \mathbb{Z}_p^{n+1}$. Also, we denote

$$\begin{aligned} A &= A(\tilde{G}), \quad R = R(\tilde{G}), \quad N = N(\tilde{G}) \\ A' &= A(G, H), \quad R' = R(G, H), \quad N' = N(G, H) \end{aligned}$$

and define $A_k \subset A$ to be generated by all $[\tilde{G}/L]$ with $L \subset \tilde{G}$ of dimension k . Thus,

$$A = A_0 \oplus A_1 \oplus \dots \oplus A_{n+1}$$

and a similar decomposition holds for A' with $A'_k = A_k \cap A'$.

3.2. Rank calculations. Let $G(k, n)$ denote the number of k -dimensional subspaces of the vector space \mathbb{Z}_p^n . Then by [7, p. 28] we have

$$G(k, n) = \prod_{j=1}^k \frac{p^{n-j+1} - 1}{p^j - 1}$$

Proposition 3.1. *The ranks a_k and a'_k of A_k and A'_k are given by the formulas*

$$a_k = G(k, n+1), \quad a'_k = p^k G(k, n).$$

Proof. The basis elements $[\tilde{G}/L]$ for A_k are in one-to-one correspondence with the k -dimensional subspaces $L < \tilde{G} = \mathbb{Z}_p^{n+1}$. Hence, we get the first formula.

The basis elements $[G \times_\rho H]$ for A'_k are in one-to-one correspondence with pairs (K, ρ) with $K < G$ a k -dimensional subspace and $\rho : K \rightarrow H$ a homomorphism. Given K , ρ is uniquely determined by its kernel and an automorphism of its image. If K is k -dimensional, the kernel of ρ is either K or any $(k-1)$ -dimensional subspace of K . In the later case the image admits $(p-1)$ automorphisms. Hence, for a given k -dimensional K there are $(p-1)G(k-1, k) + 1$ different ρ 's. For a given dimension k the number of pairs (K, ρ) is thus given by the formula:

$$G(k, n)[(p-1)G(k-1, k) + 1] = G(k, n)[(p-1) \sum_{i=0}^{k-1} p^i + 1] = p^k G(k, n).$$

□

Let ζ denote a primitive p -root of unity and $F = \mathbb{Q}[\zeta]$ be the associated cyclotomic field. For each $s \in \mathbb{Z}_p^{n+1}$ let F_s be the $\mathbb{Q}[\tilde{G}]$ -module F obtained by letting the i^{th} canonical generator of \tilde{G} act on F via the automorphism sending ζ to ζ^{s_i} where s_i is the i^{th} coordinate of s .

Proposition 3.2. *The ranks r and r' of R and R' are given by the formulas*

$$r = G(1, n+1) + 1, \quad r' = G(1, n+1).$$

Proof. With the above notations, two isomorphism classes are equal $[F_s] = [F_t]$ if and only if $t = s = 0$ or $t = us$ for some unit u in \mathbb{Z}_p . In the later case we say that s and t represent the same point $[s] = [t]$ in the projective n -space P^n over \mathbb{Z}_p . With this observation $[F_s]$ indexed by $[s] \in P^n$ and the trivial module $[\mathbb{Q}]$ form a basis for R . Thus we get the first formula.

For the second formula we claim that a basis for R' is given by the elements

$$[F_{s' \times 1}] + [\mathbb{Q}], \quad [F_{t \times 0}] - (p-1)[\mathbb{Q}]$$

indexed by $s' \in \mathbb{Z}_p^n$ and $[t] \in P^{n-1}$. Let \mathcal{B} denote the set of these elements and \mathcal{M} the \mathbb{Z} -module generated by \mathcal{B} . Since $F_{0 \times 1} + \mathbb{Q} = \mathbb{Q}[H]$ it follows that by forgetting the G -action, the elements:

$$[F_{s' \times 1}] + [\mathbb{Q}], [F_{t \times 0}] + (p-1)[F_{0 \times 1}], \text{ and } (p-1)[F_{0 \times 1}] + (p-1)[\mathbb{Q}]$$

are all represented by the $\mathbb{Q}[H]$ -free modules $\mathbb{Q}[H]$ or $(p-1)\mathbb{Q}[H]$. Thus $\mathcal{M} \subset R'$ and it is immediate that \mathcal{B} is a linearly independent set. Now by inspection R/\mathcal{M} is the free module generated by $[\mathbb{Q}]$ and $m[\mathbb{Q}] \in R'$ implies $m = 0$. Thus the rank of R' equals the rank of \mathcal{M} . In particular Lemma 2.4 applies to the sequence:

$$0 \rightarrow \mathcal{M} \rightarrow R \rightarrow R/R' \rightarrow 0$$

implying that $R' = \mathcal{M}$. □

From the Propositions 3.1 and 3.2 and the short exact sequences (1) and (3) we deduce the following result.

Corollary 3.3. *The ranks b and b' of N and N' are given by the formulas*

$$b = \sum_{k=0}^{n-1} G(k, n+1), \quad b' = \sum_{k=0}^n p^k G(k, n) - G(1, n+1).$$

3.3. The main theorem. It is convenient to identify each basis element $[\tilde{G}/L]$ of A where $L < \tilde{G}$ with the projective subspace $(L) \subset P^n$ generated by L . Also, let e denote the distinguished vector $(0, \dots, 0, 1) \in \mathbb{Z}_p^{n+1}$. Then we have the following characterization for the basis elements of A'_k :

Lemma 3.4. *The submodule $A'_k \subset A'$ is the free abelian group on the set of projective subspaces $(L) \subset P^n$ with $L < \tilde{G}$ of dimension k not containing e .*

Proof. It is easy to see that the basis elements $[G \times_\rho H]$ of A' associated with a pair (K, ρ) is of the form $[\tilde{G}/L]$ where $K < G$, $\rho : K \rightarrow H$ is a homomorphism, and $L = \{(k, \rho(k)) \mid k \in K\}$ is a linear subspace of \tilde{G} not containing e .

Conversely, let $(L) \subset P^n$ with $L < \tilde{G}$ of dimension k not containing e and define K to be the image of the canonical projection $\tilde{G} \rightarrow G$. If (g, h) is an element in L which maps to 0 under the projection, then $g = 0$. This would imply $he \in L$ so $h = 0$. Thus the projection induces an isomorphism $L \cong K$. Let $\alpha : K \rightarrow L$ be the inverse and define $\rho : K \rightarrow H$ by composing α with the canonical projection $\tilde{G} \rightarrow H$. We can then check that $[\tilde{G}/L] = [G \times_\rho H]$. □

Given L a subspace of codimension at least 2 in \tilde{G} we define $L^* < \tilde{G}$ to be a distinguished subspace such that the following two conditions are both satisfied:

- (1) L^* contains L and L^*/L has rank 2
- (2) If L does not contain e and has codimension at least 3 then L^* does not contain e

Now we observe that L^* always exists subject to the two conditions. In particular, if L has codimension exactly 2 then $L^* = \tilde{G}$ is the only choice without violating condition 2.

Definition 3.5. *For each such L define*

$$t(L) = (L) - \sum (C) + p(L^*)$$

where the sum is over all proper subspaces $L < C < L^$.*

In particular, define M_{n-1} to be the set of all (L) with $L < \tilde{G}$ an $(n-1)$ -dimensional subspace where $e \notin L$. By Lemma 3.4, this set is also a basis for A'_{n-1} .

Definition 3.6. Let A''_{n-1} be the submodule of A'_{n-1} generated by all those differences $(L) - (L')$ of elements in M_{n-1} that are subject to the relation

$$(L + \mathbb{Z}_p e) = (L' + \mathbb{Z}_p e).$$

Theorem 3.7. The rank of A''_{n-1} is $G(1, n)(p^{n-1} - 1)$ and we have the following commutative diagram of short exact sequences:

$$\begin{array}{ccccccccc} 0 & \longrightarrow & A'_0 \oplus A'_1 \oplus \dots \oplus A'_{n-2} \oplus A''_{n-1} & \xrightarrow{t'} & A' & \xrightarrow{f'} & R' & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & A_0 \oplus A_1 \oplus \dots \oplus A_{n-2} \oplus A_{n-1} & \xrightarrow{t} & A & \xrightarrow{f} & R & \longrightarrow & 0 \end{array}$$

where the vertical arrows are all inclusions and t' , f' are the restrictions of t , f .

Proof. Given (L) in M_{n-1} ,

$$t(L) = (L) - (C_0) - \sum (C) + p(\tilde{G})$$

where $C_0 = L + \mathbb{Z}_p e$ and the sum is taken over all $L < C < \tilde{G}$ not containing e . By Definition 3.6, if $(L) - (L')$ is a generator of A''_{n-1} then

$$(L + \mathbb{Z}_p e) = (L' + \mathbb{Z}_p e) = (C_0)$$

and we deduce that $t((L) - (L'))$ is in A' so all maps in the diagram are well defined. Also it was shown respectively in [5, 6] and [1] that f and f' are surjective.

From subsection 2.4 we know that the kernel of f is generated by the induced kernels $L/C \uparrow N(L/C)$ where $L/C \cong \mathbb{Z}_p \times \mathbb{Z}_p$. In particular, by applying $L/C \uparrow$ to equation (2) of Section 2.4 with $G = L/C$ we deduce that $N(\tilde{G})$ is generated by elements of the form

$$[\tilde{G}/C] - \sum [\tilde{G}/D] + p[\tilde{G}/L]$$

where L/C is any subquotient of \tilde{G} isomorphic to $\mathbb{Z}_p \times \mathbb{Z}_p$ and the sum runs over all proper subgroups $C < D < L$. By Definition 3.5 the above elements with L/C replaced by L^*/L generate the image of t so that the composition $f \circ t$ and $f' \circ t'$ are both zero.

We are left to prove the injectivity of the map t and the inclusions of the kernels of f and f' inside the images of t and t' respectively. Under the map t , each basis element (L) of A_i is mapped to an element inside $A_i \oplus A_{i+1} \oplus A_{i+2}$ whose first component is again (L) . Therefore the matrix representation of t is upper triangular with cokernel $A_n \oplus A_{n+1}$ which is free. Hence t , and therefore t' , are injective.

Regarding the exactness at A observe that by Proposition 3.1 and the injectivity of t it follows that the rank of t is the sum $G(k, n+1)$ for $k = 0, 1, \dots, n-1$. The same sum by Corollary 3.3 is the rank of the kernel of f . Since the cokernel of t is a free module we conclude by Lemma 2.4 that the bottom sequence is exact at A .

To determine exactness at A' we must first determine the rank of A'_{n-1} . As calculated in Proposition 3.1, the rank of A'_{n-1} which is also the order of M_{n-1} is equal to $a'_{n-1} = p^{n-1}G(n-1, n)$. We observe that M_{n-1} breaks into $G(n-1, n)$ equivalence classes relative to the equivalence relation $(L) \sim (L')$ if and only if


$$L + \mathbb{Z}_p e = L' + \mathbb{Z}_p e.$$

By Lemma 3.4 and Proposition 3.1 with n replaced by $n - 1$, each n -subspace containing e contains $p^{n-1}G(n-1, n-1)$ subspaces of dimension $n-1$ not containing e . This is the number of elements in any of the equivalence classes. Hence, since each equivalence class produces $p^{n-1} - 1$ basis elements for A''_{n-1} and there are $a'_{n-1}p^{n-1}$ equivalence classes, we conclude that the rank of A''_{n-1} is given by the formula

$$a'_{n-1}p^{n-1}(p^{n-1} - 1) = G(1, n)(p^{n-1} - 1).$$

Combining this with the fact that t' is injective it follows that the image of t' has rank equal to the kernel of f' . Moreover when considering the generators of A''_{n-1} , if we allow any given basis element $(K) \in A(n-1)$ to play the role of an (L) in the difference $(L) - (L')$ at most once, then we see that the matrix of t' will be upper triangular as t' maps a difference $(L) - (L')$ to an element inside $A_{n-1} \oplus A_n \oplus A_{n+1}$ with first component $(L) - (L')$. Therefore the cokernel of t' is a free module and by Lemma 2.4, the top row is exact. □

3.4. An illustration for $n = 2$ and $p = 2$. Order \mathbb{Z}_2 such that $0 < 1$ and order \mathbb{Z}_2^3 lexicographically. Then for $n = p = 2$ we gain a labeling of the basis of $A(\tilde{G})$ $\{e_i\}$ such that $e_1 < e_2 < \dots < e_{16}$. With this labeling of the basis of $A(\tilde{G})$, the subgroup lattice of \tilde{G} can be represented by the graph E below and offers a visual description of the relationship between basis elements e_i and e_j .



Theorem 3.7 implies that we have the commutative diagram of short exact sequences:

$$\begin{array}{ccccccccc} 0 & \longrightarrow & A'_0 \oplus A'_1 & \xrightarrow{t'} & A' & \xrightarrow{f'} & R' & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & A_0 \oplus A_1 & \xrightarrow{t} & A & \xrightarrow{f} & R & \longrightarrow & 0 \end{array}$$


We see using our basis that

$$A_0 = A'_0 = \mathbb{Z}e_1, A_1 = \sum_{i=2}^8 \mathbb{Z}e_i \text{ and } A'_1 = \mathbb{Z}(e_3 - e_4) + \mathbb{Z}(e_5 - e_6) + \mathbb{Z}(e_7 - e_8).$$

Hence t is well defined on A_1 while we define:

$$t(e_1) = e_1 - e_3 - e_5 - e_7 + 2e_{12}.$$

Define the subgraph E_i to be the full subgraph of E where the vertices are the terms occurring in $t(e_i)$. Then the image $t'(e_i - e_j)$ is associated with the subgraph $E_i - E_j$ whose vertices are those in E_i and E_j . For example, if $i = 3$ and $j = 4$ these subgraphs are:



Conversely, given a subgraph E_i the image $t(e_i)$ is uniquely determined by taking a weighted sum of the vertices of E_i . Moreover, given a subgraph $E_i - E_j$, the image $t(e_i - e_j)$ is also uniquely determined by the vertices of $E_i - E_j$. It follows that the kernel of f is generated by all of the subgraphs E_i for $i = 1, 2, \dots, 8$ and the kernel of f' is generated by all the non-singular subgraphs $E_1, E_3 - E_4, E_5 - E_6, E_7 - E_8$.

4. FINAL REMARKS

We would like to develop a description for the kernel $N(G, H)$ with $H \cong \mathbb{Z}_p$ similar to that given by Tornehave in [8] for $N(G)$ with G an arbitrary finite p -groups G . Define $\tilde{N}(L/C)$ for L/C a subquotient of $\tilde{G} = G \times H$ to be the intersection of $N(L/C)$ with the submodule $\tilde{A}(L/C)$ of $A(L/C)$ that lands inside $A(G, H)$ under the induction $L/C \uparrow$ of Section 2.3.

Conjecture 4.1. *Let p be a prime, G any finite p -group, and $H \cong \mathbb{Z}_p$. Then*

$$N(G, H) = \sum L/C \uparrow \tilde{N}(L/C)$$

where the sum is taken over subquotients L/C of \tilde{G} isomorphic to $T \times H$ where T is the elementary abelian group $\mathbb{Z}_p \times \mathbb{Z}_p$, the dihedral group, or the nonabelian group of order p^3 and exponent p .

For G elementary abelian or cyclic this conjecture can readily be checked using Theorem 3.7 and rank arguments.

Proposition 4.2. *Let p be any prime, G be an elementary abelian p -group, and $H \cong \mathbb{Z}_p$. Then*

$$N(G, H) = \sum L/C \uparrow \tilde{N}(L/C)$$

with the sum taken over all subquotients $L/C \cong \mathbb{Z}_p^3$.

Proof. From Theorem 3.7 we know that the image of t generates the kernel $N(G, H)$. If $(L) \in A'_i$ with $0 \leq i \leq n - 2$, then there exists subgroups $L < L^* < B < \tilde{G}$ such that $B/L \cong \mathbb{Z}_p^3$ where L^* is the distinguished element used to define t in Definition 3.5. In addition, regardless of our choice of B ,

$$t((L)) \in B/L \uparrow \tilde{N}(B/L).$$

If $(L) - (L') \in \tilde{A}_{n-1}$, let

$$C = L + \mathbb{Z}_p e = L' + \mathbb{Z}_p e, D = L \cap L'.$$

We see immediately that $\tilde{G}/D \cong \mathbb{Z}_p^3$ and also that $t((L) - (L'))$ is an element of $\tilde{G}/D \uparrow \tilde{N}(\tilde{G}/D)$. Hence we conclude that

$$N(G, H) \subset \sum L/C \uparrow \tilde{N}(L/C).$$

The converse is immediate. \square

Proposition 4.3. *Let p be any prime, G the cyclic p -group with order p^k , and $H \cong \mathbb{Z}_p$. Then f' is an isomorphism between $A(G, H)$ and $R(G, H)$.*

Proof. Let $\tilde{G} = G \times H$ and ξ be the primitive p^k -root of unity. Since G is cyclic, easily the rank of $A(G, H)$ is equal to $kp + 1$ as G has $k + 1$ subgroups and for a nontrivial subgroup $K < G$, there are p homomorphisms $\rho : K \rightarrow H$. Let $F_{\nu, \phi} = \mathbb{Q}(\xi^{p^{k-\nu}}, \xi^{\phi p^{k-1}})$ be the $\mathbb{Q}[\tilde{G}]$ -module with the generators of G and H acting by multiplication by $\xi^{p^{k-\nu}}$ and $\xi^{\phi p^{k-1}}$ respectively where $\nu = 0, 1, \dots, k$ and $\phi = 0, 1, \dots, p - 1$. Then the irreducible $\mathbb{Q}[\tilde{G}]$ -modules as seen from the decomposition of the group ring $\mathbb{Q}[\tilde{G}]$ are:

$$\begin{aligned} F_{0,0} &= \mathbb{Q} \\ F_{0,1} &= \mathbb{Q}(\xi^{\phi p^{k-1}}) \\ F_{\nu, \phi} &\text{ with } \nu = 1, \dots, k \text{ and } \phi = 0, 1, \dots, p - 1. \end{aligned}$$

This implies the rank of $R(\tilde{G})$ is $kp + 2$.

For $[M], [M'] \in R(\tilde{G})$, define $[M] \equiv [M']$ if we have $[M] - [M'] \in R(G, H)$. Using this relation we immediately gain the following equivalences from [1]:

$$\begin{aligned} [F_{\nu, \phi}] &\equiv -p^{\nu-1}[\mathbb{Q}] && \text{for } \nu = 1, \dots, k \text{ and } \phi = 1, \dots, p - 1, \\ [F_{0,1}] &\equiv -[\mathbb{Q}], \\ [F_{\nu,0}] &\equiv p^{\nu-1}(p-1)[\mathbb{Q}] && \text{for } \nu = 1, \dots, k. \end{aligned}$$

The equivalences imply that the rank of $R(\tilde{G})/R(G, H)$ is less than or equal to 1. In addition, since f' is surjective, the rank of $A(G, H)$ is $kp + 1$, and the rank of $R(\tilde{G})$ is $kp + 2$, we see that the rank of $R(\tilde{G})/R(G, H)$ is at least 1. Thus the rank of $R(\tilde{G})/R(G, H)$ is exactly 1 which implies the rank of $R(G, H) = kp + 1$. As $A(G, H)$ is a free module, the rank of $A(G, H)$ is equal to the rank of $R(G, H)$, and f' is a surjection, we see that f' is an isomorphism. \square

As a corollary, Conjecture 4.1 is true for G a cyclic p -group.

REFERENCES

- [1] Marian F. Anton. The double burnside ring and and rational representations. *Revue Roumaine de Mathématiques Pures et Appliquées*, 2006.
- [2] Michael F. Atiyah. Characters and cohomology of finite groups. *Publications mathématiques de l'I.H.E.S.*, 9, 1961.
- [3] Gunnar Carlsson. Equivariant stable homotopy and segal's burnside ring conjecture. *The Annals of Mathematics*, 120, 1984.
- [4] Erkki Laitinen. On the burnside ring and stable comotopy of a finite group. *Mathematica Scandinavica*, 44, 1979.
- [5] Jürgen Ritter. Ein induktion für rationale charaktere von nilpotenten gruppen. *J. Reine Angew. Math.*, 254, 1972.
- [6] Graeme Segal. Permutation representations of finite p-groups. *Quarterly Journal of Mathematics*, 1972.

- [7] Richard P. Stanley. *Enumerative Combinatorics*. Wadsworth and Brooks/Cole, 1986.
- [8] J. Tornehave. Relations among permutation representations of p-groups. preprint, 1984.