

# Matter Collineations of Plane Symmetric Spacetimes

M. Sharif \*and Nousheen Ilyas

Department of Mathematics, University of the Punjab,  
Quaid-e-Azam Campus, Lahore-54590, Pakistan.

## Abstract

This paper is devoted to the study of matter collineations of plane symmetric spacetimes (for a particular class of spacetimes) when the energy-momentum tensor is non-degenerate. There exists many interesting cases where we obtain proper matter collineations. The matter collineations in these cases are *four*, *five*, *six*, *seven* and *ten* with some constraints on the energy-momentum tensor. We have solved some of these constraints to obtain solutions of the Einstein field equations.

**Keywords** : Matter symmetries, Plane Symmetric spacetimes

**PACS**: 04.20Gz, 02.40Ky

## 1 Introduction

It has been an interesting subject to use the symmetry group of a spacetime in constructing the solution of the Einstein field equation (EFEs) given by

$$G_{ab} \equiv R_{ab} - \frac{1}{2}Rg_{ab} = \kappa T_{ab}, \quad (1)$$

where  $G_{ab}$  are components of the Einstein tensor,  $R_{ab}$  are components of the Ricci tensor and  $T_{ab}$  are components of the matter (energy-momentum) tensor,  $R$  is the Ricci scalar and  $\kappa$  is the gravitational constant. Further, these

---

\*msharif@math.pu.edu.pk

solutions are classified according to the Lie algebra structure generated by these symmetries. Curvature and the Ricci tensors are the important quantities which play a significant role in understanding the geometric structure of spacetime. A pioneer study of curvature collineations (CCs) and Ricci collineations (RCs) has been carried out by Katzin, et al [1] and a further classification of CCs and RCs has been obtained by different authors [2,3].

Symmetries of the Ricci tensor and, in particular, energy-momentum tensor have recently been studied. The Einstein tensor quite naturally arises in the theory of General Relativity and is related, via EFEs, to the material content of the spacetime. Thus it has an important role in this theory. In this paper, we shall analyze the properties of a vector field along which the Lie derivative of the energy-momentum tensor vanishes, i.e.,  $\mathcal{L}_\xi T_{ab} = 0$ .

A smooth vector field  $\xi$  is said to preserve a matter symmetry [4] on  $M$  if, for each smooth local diffeomorphism  $\phi_t$  associated with  $\xi$ , the tensors  $T$  and  $\phi_t^*T$  are equal on the domain  $U$  of  $\phi_t$ , i.e.,  $T = \phi_t^*T$ . Equivalently, a vector field  $\xi^a$  is said to generate a matter collineation if it satisfies the following equation

$$\mathcal{L}_\xi T_{ab} = 0, \quad \text{or} \quad T_{ab,c}\xi^c + T_{ac}\xi_{,b}^c + T_{cb}\xi_{,a}^c = 0, \quad (a, b, c = 0, 1, 2, 3), \quad (2)$$

where  $\mathcal{L}$  is the Lie derivative operator,  $\xi^a$  is the symmetry or collineation vector. Every Killing vector (KV) is a matter collineation (MC) but the converse is not true, in general. A proper MC is an MC which is not a KV, or a homothetic vector.

There is a growing interest in the study of MCs [5-7 and references therein]. Hall, et al [5], in the discussion of RC and MC, have argued that the symmetries of the energy-momentum tensor may also provide some extra understanding of the subject which has not been provided by KVs, Ricci and CCs. Carot, et al [6,7] have discussed MCs from the point of view of the Lie algebra of vector fields generating them and, in particular, he discussed spacetimes with a degenerate  $T_{ab}$ . Some recent investigations [8-20] show keen interest in the study of matter collineations. In the papers [13-15], the study of MCs has been taken for spherically symmetric, static plane symmetric and cylindrically symmetric spacetimes and some interesting results have been obtained. Recently, the proper MCs of non-static plane symmetric spacetimes have been found [19] when the energy-momentum tensor is degenerate for spacetimes satisfying  $T_{01} = 0$ . It turns out that this admits an MC Lie algebra of 4, 6, 10 dimensions apart from the infinite dimensional algebras. This paper extends the above study to the non-degenerate case.

The distribution of the paper is as follows. In the next section, we write down the MC equations for plane symmetric spacetimes. In section **3**, we calculate MCs by solving MC equations for the non-degenerate case. Section **4** carries some examples satisfying the constraints. Finally, a discussion of the results is given in the last section.

## 2 Matter Collineation Equations

The most general form of the plane symmetric metric is given in the form [21]

$$ds^2 = e^{\nu(t,x)} dt^2 - e^{\lambda(t,x)} dx^2 - e^{\mu(t,x)} (dy^2 + dz^2), \quad (3)$$

where  $\nu$ ,  $\lambda$  and  $\mu$  are arbitrary functions of  $t$  and  $x$ . The surviving components of the energy-momentum tensor are given as

$$\begin{aligned} T_{00} &= \frac{1}{4}(\dot{\mu}^2 + 2\dot{\mu}\dot{\lambda}) - \frac{1}{4}e^{v-\lambda}(4\mu'' + 3\mu'^2 - 2\mu'\lambda'), \\ T_{01} &= -\frac{1}{2}(2\dot{\mu}' + \dot{\mu}\mu' - \dot{\mu}\nu' - \mu'\dot{\lambda}), \\ T_{11} &= \frac{1}{4}(\mu'^2 + 2\mu'\nu') - \frac{1}{4}e^{\lambda-\nu}(4\ddot{\mu} + 3\dot{\mu}^2 - 2\dot{\mu}\dot{\nu}), \\ T_{22} &= \frac{1}{4}e^{\mu-\lambda}(2\mu'' + \mu'^2 - \mu'\lambda' + \mu'\nu' - \lambda'\nu' + \nu'^2 + 2\nu'') \\ &\quad - \frac{1}{4}e^{\mu-\nu}(2\ddot{\mu} + \dot{\mu}^2 - \dot{\mu}\dot{\nu} + \dot{\mu}\dot{\lambda} - \dot{\nu}\dot{\lambda} + \dot{\lambda}^2 + 2\ddot{\lambda}), \\ T_{33} &= T_{22}. \end{aligned} \quad (4)$$

Here dot and prime indicate differentiation w.r.t. time and  $x$  coordinate respectively. The MC equations can be written as follows

$$2T_{01}\xi_{,0}^1 + T_{00,0}\xi^0 + T_{00,1}\xi^1 + 2T_{00}\xi_{,0}^0 = 0, \quad (5)$$

$$T_{01,0}\xi^0 + T_{01}\xi_0^0 + T_{00}\xi_{,1}^0 + T_{01,1}\xi^1 + T_{11}\xi_{,0}^1 + T_{01}\xi_{,1}^1 = 0, \quad (6)$$

$$T_{00}\xi_{,2}^0 + T_{01}\xi_{,2}^1 + T_{22}\xi_{,0}^2 = 0, \quad (7)$$

$$T_{00}\xi_{,3}^0 + T_{01}\xi_{,3}^1 + T_{22}\xi_{,0}^3 = 0, \quad (8)$$

$$T_{11,0}\xi^0 + 2T_{01}\xi_0^0 + T_{11,1}\xi^1 + 2T_{11}\xi_{,1}^1 = 0, \quad (9)$$

$$T_{01}\xi_{,2}^0 + T_{11}\xi_{,2}^1 + T_{22}\xi_{,1}^2 = 0, \quad (10)$$

$$T_{01}\xi_{,3}^0 + T_{11}\xi_{,3}^1 + T_{22}\xi_{,1}^3 = 0, \quad (11)$$

$$T_{22,0}\xi^0 + T_{22,1}\xi^1 + 2T_{22}\xi_{,2}^2 = 0, \quad (12)$$

$$T_{22}(\xi_{,3}^2 + \xi_{,2}^3) = 0, \quad (13)$$

$$T_{22,0}\xi^0 + T_{22,1}\xi^1 + 2T_{22}\xi_{,3}^3 = 0. \quad (14)$$

These are the first order non-linear partial differential equations in four variables  $\xi^a(x^b)$ . We restrict ourselves with the assumption  $T_{01} = 0$  for the sake of simplicity. For this purpose, either we need to make transformations such that  $T_{01} = 0$  or we take such spacetimes which satisfy this condition. Here we take the following class of metrics satisfying this assumption.

$E(2) \otimes \mathbf{R}$ , where  $\mathbf{R} = \partial_x$  if and only if  $\nu = \nu(t)$ ,  $\lambda = \lambda(t)$ ,  $\mu = 2 \ln t$ .

For the sake of simplicity, we use the notation  $T_{aa} = T_a$  when  $T_{01} = 0$ . The solution of these equations for the degenerate case, i.e.,  $\det(T_{ab}) = 0$  for the above class of metrics has already been completed [19]. In this paper, we are giving the complete classification for the non-degenerate case with  $T_{01} = 0$ .

### 3 Solution for the Non-Degenerate Case

This section explicitly provides a complete classification of plane symmetric spacetimes for the non-degenerate case for the above mentioned case.

For this case, MC Eqs.(5)-(14) reduce to

$$T_{0,0}\xi^0 + 2T_0\xi_{,0}^0 = 0, \quad (15)$$

$$T_0\xi_{,1}^0 + T_1\xi_{,0}^1 = 0, \quad (16)$$

$$T_0\xi_{,2}^0 + T_2\xi_{,0}^2 = 0, \quad (17)$$

$$T_0\xi_{,3}^0 + T_2\xi_{,0}^3 = 0, \quad (18)$$

$$T_{1,0}\xi^0 + 2T_1\xi_{,1}^1 = 0, \quad (19)$$

$$T_1\xi_{,2}^1 + T_2\xi_{,1}^2 = 0, \quad (20)$$

$$T_1\xi_{,3}^1 + T_2\xi_{,1}^3 = 0, \quad (21)$$

$$T_{2,0}\xi^0 + 2T_2\xi_{,2}^2 = 0, \quad (22)$$

$$T_2(\xi_{,3}^2 + \xi_{,2}^3) = 0, \quad (23)$$

$$T_{2,0}\xi^0 + 2T_2\xi_{,3}^3 = 0. \quad (24)$$

When we solve Eqs.(15)-(24) simultaneously, after some algebraic computations, we obtain

$$\begin{aligned}
\xi^0 &= -\frac{T_2}{T_0} \left\{ \frac{1}{2} \dot{A}_1 (y^2 + z^2) + \dot{A}_2 z + \dot{A}_3 y \right\} + A_4, \\
\xi^1 &= -\frac{T_2}{T_1} \left\{ \frac{1}{2} A'_1 (y^2 + z^2) + A'_2 z + A'_3 y \right\} + A_5, \\
\xi^2 &= c_2 y z + \frac{c_3}{2} (z^2 - y^2) + c_4 z + A_1 y + A_3, \\
\xi^3 &= \frac{c_2}{2} (z^2 - y^2) - (c_3 z + c_4) y + A_1 z + A_2
\end{aligned} \tag{25}$$

subject to the constraints

$$\dot{T}_2 \dot{A}_1 = 0, \tag{26}$$

$$\frac{\dot{T}_1}{T_0} \dot{A}_i + 2A''_i = 0, \tag{27}$$

$$\frac{\dot{T}_2}{T_2} A_4 + 2A_1 = 0, \tag{28}$$

$$\frac{\dot{T}_2}{T_2 \sqrt{T_0}} g_2(x) - 2c_2 = 0, \tag{29}$$

$$\frac{\dot{T}_2}{T_2 \sqrt{T_0}} g_3(x) + 2c_3 = 0, \tag{30}$$

$$\frac{\dot{T}_2}{T_2 \sqrt{T_0}} g_4(x) + 2A_1 = 0, \tag{31}$$

$$\frac{\dot{T}_1}{T_2 \sqrt{T_0}} g_i(x) + 2A''_i = 0, \tag{32}$$

$$A'_i = \sqrt{\frac{T_1}{T_2}} f_i(x), \quad \dot{A}_i = \frac{\sqrt{T_0}}{T_2} g_i(x), \quad (i = 1, 2, 3) \tag{33}$$

$$A'_5 = -\frac{\dot{T}_1}{2T_1 \sqrt{T_0}} g_4(x), \quad \dot{A}_5 = -\frac{\sqrt{T_0}}{T_1} g'_4(x), \tag{34}$$

where  $c_2, c_3, c_4$  are arbitrary constants and  $A_\mu = A_\mu(t, x)$  ( $\mu = 1, 2, 3, 4, 5$ ),  $f_i(x), g_i(x), g_4(x)$  are integration functions. Thus the problem is reduced to solving the set of Eqs.(15)-(24) subject to the constraint Eqs.(26)-(34). From Eqs.(29) and (30), there arise the following two cases:

$$(1) \quad \left( \frac{\dot{T}_2}{T_2 \sqrt{T_0}} \right) \neq 0, \quad (2) \quad \left( \frac{\dot{T}_2}{T_2 \sqrt{T_0}} \right) = 0.$$

**Case (1):** In this case, we have  $\dot{T}_2 \neq 0$  and Eq.(26) yields  $A_1 = A_1(x)$ . Then it follows from Eq.(31) that

$$\frac{\dot{T}_2}{T_2\sqrt{T_0}}g_4(x) + 2A_1(x) = 0 \quad (35)$$

which is possible only if  $g_4 = 0 = A_1$ . Also, from Eqs.(28) and (34), we have  $A_4 = 0$ ,  $A_5 = c_0$ . Now Eqs.(29) and (30) yield that  $g_2 = 0 = g_3$ ,  $c_2 = 0 = c_3$  and from Eqs.(32) and (33), we have  $f_j(x) = a_j$ , ( $j = 2, 3$ ), where  $a_j$  are arbitrary constants. Also, it follows from Eq.(33) that

$$\dot{A}_j = 0, \quad A'_j = \sqrt{\frac{T_1}{T_2}}a_j$$

which gives

$$A_j = \sqrt{\frac{T_1}{T_2}}a_jx + h_j(t), \quad (36)$$

$$\left(\sqrt{\frac{T_1}{T_2}}\right)'a_jx + \dot{h}_j(t) = 0.$$

This implies that either

$$(i) \quad \left(\sqrt{\frac{T_1}{T_2}}\right)' = 0, \quad \text{or} \quad (ii) \quad \left(\sqrt{\frac{T_1}{T_2}}\right)' \neq 0.$$

For both the cases  $h_j(t) = b_j$ , where  $b_j$ , ( $j = 2, 3$ ) are arbitrary constants and hence Eq.(36) takes the form

$$A_j = \sqrt{\frac{T_1}{T_2}}a_jx + b_j \quad (37)$$

**Case 1(i):** Here we have  $\left(\frac{T_1}{T_2}\right)' = 0$  which implies that  $T_1 = dT_2$ , where  $d$  is an arbitrary constant. Thus Eq.(37) reduces to

$$A_j = d_jx + b_j, \quad (j = 2, 3) \quad (38)$$

where  $d_j$  ( $j = 2, 3$ ) are arbitrary constants. Consequently, we obtain six independent MCs out of which three are the minimal KVs of plane symmetric and three are proper MCs. These are given by

$$\begin{aligned} \xi_{(1)} &= \partial_y, & \xi_{(2)} &= \partial_z, & \xi_{(3)} &= z\partial_y - y\partial_z, \\ \xi_{(4)} &= \partial_x, & \xi_{(5)} &= x\partial_z - \frac{T_2}{T_1}z\partial_x, & \xi_{(6)} &= x\partial_y - \frac{T_2}{T_1}y\partial_x. \end{aligned} \quad (39)$$

The corresponding Lie algebra is given by

$$\begin{aligned}
[\xi_{(1)}, \xi_{(3)}] &= -\xi_{(2)}, & [\xi_{(1)}, \xi_{(6)}] &= -\frac{T_2}{T_1}\xi_{(4)}, & [\xi_{(2)}, \xi_{(3)}] &= \xi_{(1)}, \\
[\xi_{(2)}, \xi_{(5)}] &= -\frac{T_2}{T_1}\xi_{(4)}, & [\xi_{(3)}, \xi_{(5)}] &= \xi_{(6)}, & [\xi_{(3)}, \xi_{(6)}] &= -\xi_{(5)}, \\
[\xi_{(4)}, \xi_{(5)}] &= \xi_{(2)}, & [\xi_{(4)}, \xi_{(6)}] &= \xi_{(1)}, & [\xi_{(5)}, \xi_{(6)}] &= -\frac{T_2}{T_1}\xi_{(3)}, \\
[\xi_{(i)}, \xi_{(j)}] &= 0, & & & & \text{(otherwise)}.
\end{aligned}$$

**Case 1(ii):** Here  $(\sqrt{\frac{T_1}{T_2}}) \cdot \neq 0$  which implies that  $a_j = 0$  and then Eq.(37) reduces to  $A_j = b_j$ . Solving the MCs equations, we obtain three usual plane symmetry KVs and one proper MC.

**Case 2.** This case yields  $(\frac{\dot{T}_2}{T_2\sqrt{T_0}}) = \alpha$ , where  $\alpha$  is an arbitrary constant which can be zero or non-zero.

**Case 2(i):** First we take  $\alpha \neq 0$  which implies  $\dot{T}_2 \neq 0$ . It follows from Eqs.(26), (29) and (30) that

$$g_2 = \frac{2c_2}{\alpha}, \quad g_3 = -\frac{2c_3}{\alpha}, \quad A_1 = A_1(x) \quad (40)$$

and hence Eq.(33) yields

$$\left(\frac{T_2}{T_1}\right) \cdot A'_i = 0. \quad (41)$$

This implies further two possibilities:

$$(a) \quad \left(\frac{T_2}{T_1}\right) \cdot \neq 0, \quad (b) \quad \left(\frac{T_2}{T_1}\right) \cdot = 0.$$

For the **case 2i(a)**, Eq.(40) implies that  $A_1 = c_5$  and consequently Eqs.(27), (28) and (34) respectively yield

$$\dot{T}_1 \dot{A}_i = 0, \quad A_4 = -\frac{2c_5}{\alpha\sqrt{T_0}}, \quad (42)$$

$$\dot{A}_5 = 0, \quad A'_5 = \frac{\dot{T}_1}{\alpha T_1 \sqrt{T_0}} c_5, \quad (43)$$

$$A_5 = \frac{\dot{T}_1}{\alpha T_1 \sqrt{T_0}} c_5 x + c_0. \quad (44)$$

Eq.(43) further gives two possibilities

$$(*) \quad \left(\frac{\dot{T}_1}{T_1\sqrt{T_0}}\right)' \neq 0, \quad (**) \quad \left(\frac{\dot{T}_1}{T_1\sqrt{T_0}}\right)' = 0.$$

In the **case 2ia**(\*),  $\dot{T}_1 \neq 0$ , we obtain three usual plane symmetry KVs and one proper MC.

For the **case 2ia**(\*\*), we have  $\frac{\dot{T}_1}{T_1\sqrt{T_0}} = \beta$ , where  $\beta$  is an arbitrary constant which can be zero or non-zero. When  $\beta \neq 0$ , the proper MCs are given by

$$\xi_{(4)} = \partial_x, \quad \xi_{(5)} = -\frac{2}{\alpha\sqrt{T_0}}\partial_t + \frac{\beta}{\alpha}x\partial_x + y\partial_y + z\partial_z. \quad (45)$$

The Lie algebra will become

$$[\xi_{(1)}, \xi_{(3)}] = -\xi_{(2)}, \quad [\xi_{(1)}, \xi_{(5)}] = \xi_{(1)}, \quad [\xi_{(2)}, \xi_{(3)}] = \xi_{(1)},$$

$$[\xi_{(2)}, \xi_{(5)}] = \xi_{(2)}, \quad [\xi_{(4)}, \xi_{(5)}] = \frac{\beta}{\alpha}\xi_{(4)}, \quad [\xi_{(i)}, \xi_{(j)}] = 0, \quad (\text{otherwise}).$$

For  $\beta = 0$ , the proper MCs are

$$\begin{aligned} \xi_{(4)} &= \partial_x, \\ \xi_{(5)} &= -\frac{2}{\alpha\sqrt{T_0}}z\partial_t + yz\partial_y + \left\{\frac{1}{2}(z^2 - y^2) - \frac{2}{\alpha^2 T_2}\right\}\partial_z, \\ \xi_{(6)} &= \frac{2}{\alpha\sqrt{T_0}}y\partial_t + \left\{\frac{1}{2}(z^2 - y^2) + \frac{2}{\alpha^2 T_2}\right\}\partial_y - yz\partial_z, \\ \xi_{(7)} &= -\frac{2}{\alpha\sqrt{T_0}}\partial_t + y\partial_y + z\partial_z. \end{aligned} \quad (46)$$

The Lie algebra is

$$[\xi_{(1)}, \xi_{(3)}] = -\xi_{(2)}, \quad [\xi_{(1)}, \xi_{(5)}] = \xi_{(3)}, \quad [\xi_{(1)}, \xi_{(6)}] = -\xi_{(7)},$$

$$[\xi_{(1)}, \xi_{(7)}] = \xi_{(1)}, \quad [\xi_{(2)}, \xi_{(3)}] = \xi_{(1)}, \quad [\xi_{(2)}, \xi_{(5)}] = \xi_{(7)},$$

$$[\xi_{(2)}, \xi_{(6)}] = \xi_{(3)}, \quad [\xi_{(2)}, \xi_{(7)}] = \xi_{(2)}, \quad [\xi_{(3)}, \xi_{(5)}] = \xi_{(6)},$$

$$[\xi_{(3)}, \xi_{(6)}] = -\xi_{(5)}, \quad [\xi_{(5)}, \xi_{(7)}] = -\xi_{(5)}, \quad [\xi_{(6)}, \xi_{(7)}] = -\xi_{(6)},$$



$$[\xi_{(i)}, \xi_{(j)}] = 0, \quad (\text{otherwise}).$$

For the **case 2i(b)**,  $T_2 = \gamma T_1$ , where  $\gamma$  is non-zero arbitrary constant, the proper MCs are

$$\begin{aligned}
\xi_{(4)} &= \partial_x, \\
\xi_{(5)} &= \frac{x}{\sqrt{\gamma}}\partial_z - \sqrt{\gamma}z\partial_x, \\
\xi_{(6)} &= -\frac{2}{\alpha\sqrt{T_0}}z\partial_t + zx\partial_x + yz\partial_y + \frac{1}{2}\left(z^2 - y^2 - \frac{x^2}{\gamma} - \frac{4}{\alpha^2 T_2}\right)\partial_z, \\
\xi_{(7)} &= \frac{2}{\alpha\sqrt{T_0}}y\partial_t - yx\partial_x + \frac{1}{2}\left(z^2 - y^2 + \frac{x^2}{\gamma} + \frac{4}{\alpha^2 T_2}\right)\partial_y - yz\partial_z, \\
\xi_{(8)} &= -\frac{2}{\alpha\sqrt{T_0}}x\partial_t + \frac{\gamma}{2}\left(\frac{x^2}{\gamma} - y^2 - z^2 - \frac{4}{\alpha^2 T_2}\right)\partial_x + xy\partial_y + xz\partial_z, \\
\xi_{(9)} &= -\frac{2}{\alpha\sqrt{T_0}}\partial_t + x\partial_x + y\partial_y + z\partial_z, \\
\xi_{(10)} &= \frac{x}{\sqrt{\gamma}}\partial_y - \sqrt{\gamma}y\partial_x.
\end{aligned} \tag{47}$$

The Lie algebra is given by

$$\begin{aligned}
[\xi_{(1)}, \xi_{(3)}] &= -\xi_{(2)}, & [\xi_{(1)}, \xi_{(6)}] &= \xi_{(3)}, & [\xi_{(1)}, \xi_{(7)}] &= -\xi_{(9)}, \\
[\xi_{(1)}, \xi_{(8)}] &= \sqrt{\gamma}\xi_{(10)}, & [\xi_{(1)}, \xi_{(9)}] &= \xi_{(1)}, & [\xi_{(1)}, \xi_{(10)}] &= -\sqrt{\gamma}\xi_{(4)}, \\
[\xi_{(2)}, \xi_{(3)}] &= \xi_{(1)}, & [\xi_{(2)}, \xi_{(5)}] &= -\sqrt{\gamma}\xi_{(4)}, & [\xi_{(2)}, \xi_{(6)}] &= \xi_{(9)}, \\
[\xi_{(2)}, \xi_{(7)}] &= \xi_{(3)}, & [\xi_{(2)}, \xi_{(8)}] &= \sqrt{\gamma}\xi_{(5)}, & [\xi_{(2)}, \xi_{(9)}] &= \xi_{(2)}, \\
[\xi_{(3)}, \xi_{(5)}] &= -\xi_{(10)}, & [\xi_{(3)}, \xi_{(6)}] &= \xi_{(7)}, & [\xi_{(3)}, \xi_{(7)}] &= -\xi_{(6)}, \\
[\xi_{(3)}, \xi_{(10)}] &= \xi_{(5)}, & [\xi_{(4)}, \xi_{(5)}] &= \frac{1}{\sqrt{\gamma}}\xi_{(2)}, & [\xi_{(4)}, \xi_{(6)}] &= -\frac{1}{\sqrt{\gamma}}\xi_{(5)}, \\
[\xi_{(4)}, \xi_{(7)}] &= \frac{1}{\sqrt{\gamma}}\xi_{(10)}, & [\xi_{(4)}, \xi_{(8)}] &= \xi_{(9)}, & [\xi_{(4)}, \xi_{(9)}] &= \xi_{(4)},
\end{aligned}$$

$$[\xi_{(4)}, \xi_{(10)}] = \frac{1}{\sqrt{\gamma}}\xi_{(1)}, \quad [\xi_{(5)}, \xi_{(6)}] = \frac{1}{\sqrt{\gamma}}\xi_{(8)}, \quad [\xi_{(5)}, \xi_{(8)}] = -\sqrt{\gamma}\xi_{(6)},$$

$$[\xi_{(5)}, \xi_{(10)}] = -\xi_{(3)}, \quad [\xi_{(6)}, \xi_{(9)}] = -\xi_{(6)}, \quad [\xi_{(7)}, \xi_{(9)}] = -\xi_{(7)},$$

$$[\xi_{(7)}, \xi_{(10)}] = \frac{1}{\sqrt{\gamma}}\xi_{(8)}, \quad [\xi_{(8)}, \xi_{(9)}] = -\xi_{(8)}, \quad [\xi_{(8)}, \xi_{(10)}] = -\sqrt{\gamma}\xi_{(7)},$$

$$[\xi_{(i)}, \xi_{(j)}] = 0, \quad (\text{otherwise}).$$

**Case 2(ii):** Here  $\alpha = 0$  and we have  $T_2 = \text{constant}$ . Eqs.(29), (30) and (31) yield  $c_2 = 0 = c_3$ ,  $A_1 = 0$  and Eq.(33) gives

$$\left(\frac{(\sqrt{T_1})^\cdot}{\sqrt{T_0}}\right)^\cdot f_j(x) = 0. \quad (48)$$

This leads to two possibilities

$$(a) \quad \left(\frac{(\sqrt{T_1})^\cdot}{\sqrt{T_0}}\right)^\cdot = 0, \quad (b) \quad \left(\frac{(\sqrt{T_1})^\cdot}{\sqrt{T_0}}\right)^\cdot \neq 0.$$

For the **case 2ii(a)**, we have  $\frac{(\sqrt{T_1})^\cdot}{\sqrt{T_0}} = \delta$ , where  $\delta$  is an arbitrary constant which can be zero or non-zero.

In the **case 2iia(\*)**, when  $\delta \neq 0$ , the proper MCs are

$$\begin{aligned} \xi_{(4)} &= \partial_x, \\ \xi_{(5)} &= \frac{1}{\sqrt{T_0}} \cos \delta x \partial_t - \frac{1}{\sqrt{T_1}} \sin \delta x \partial_x, \\ \xi_{(6)} &= \frac{1}{\sqrt{T_0}} \sin \delta x \partial_t + \frac{1}{\sqrt{T_1}} \cos \delta x \partial_x, \\ \xi_{(7)} &= -\frac{z}{\sqrt{T_0}} \cos \delta x \partial_t + \frac{z}{\sqrt{T_1}} \sin \delta x \partial_x + \frac{\sqrt{T_1}}{\delta T_2} \cos \delta x \partial_z, \\ \xi_{(8)} &= -\frac{z}{\sqrt{T_0}} \sin \delta x \partial_t - \frac{z}{\sqrt{T_1}} \cos \delta x \partial_x + \frac{\sqrt{T_1}}{\delta T_2} \sin \delta x \partial_z, \\ \xi_{(9)} &= -\frac{y}{\sqrt{T_0}} \cos \delta x \partial_t + \frac{y}{\sqrt{T_1}} \sin \delta x \partial_x + \frac{\sqrt{T_1}}{\delta T_2} \cos \delta x \partial_y, \\ \xi_{(10)} &= -\frac{y}{\sqrt{T_0}} \sin \delta x \partial_t - \frac{y}{\sqrt{T_1}} \cos \delta x \partial_x + \frac{\sqrt{T_1}}{\delta T_2} \sin \delta x \partial_y. \end{aligned} \quad (49)$$

The Lie algebra is given by

$$\begin{aligned}
[\xi_{(1)}, \xi_{(3)}] &= -\xi_{(2)}, & [\xi_{(1)}, \xi_{(9)}] &= -\xi_{(5)}, & [\xi_{(1)}, \xi_{(10)}] &= -\xi_{(6)}, \\
[\xi_{(2)}, \xi_{(3)}] &= \xi_{(1)}, & [\xi_{(2)}, \xi_{(7)}] &= -\xi_{(5)}, & [\xi_{(2)}, \xi_{(8)}] &= -\xi_{(6)}, \\
[\xi_{(3)}, \xi_{(7)}] &= -\xi_{(9)}, & [\xi_{(3)}, \xi_{(8)}] &= -\xi_{(10)}, & [\xi_{(3)}, \xi_{(9)}] &= \xi_{(7)}, \\
[\xi_{(3)}, \xi_{(10)}] &= \xi_{(8)}, & [\xi_{(4)}, \xi_{(5)}] &= -\delta\xi_{(6)}, & [\xi_{(4)}, \xi_{(6)}] &= \delta\xi_{(5)}, \\
[\xi_{(4)}, \xi_{(7)}] &= -\delta\xi_{(8)}, & [\xi_{(4)}, \xi_{(8)}] &= \delta\xi_{(7)}, & [\xi_{(4)}, \xi_{(9)}] &= -\delta\xi_{(10)}, \\
[\xi_{(4)}, \xi_{(10)}] &= \delta\xi_{(9)}, & [\xi_{(5)}, \xi_{(7)}] &= \frac{1}{T_2}\xi_{(2)}, & [\xi_{(5)}, \xi_{(9)}] &= \frac{1}{T_2}\xi_{(1)}, \\
[\xi_{(6)}, \xi_{(8)}] &= \frac{1}{T_2}\xi_{(2)}, & [\xi_{(6)}, \xi_{(10)}] &= \frac{1}{T_2}\xi_{(1)}, & [\xi_{(7)}, \xi_{(9)}] &= -\frac{1}{T_2}\xi_{(3)}, \\
[\xi_{(8)}, \xi_{(10)}] &= \frac{1}{T_2}y\xi_{(2)}, & [\xi_{(9)}, \xi_{(10)}] &= -\frac{1}{T_2}y\xi_{(1)}, \\
[\xi_{(i)}, \xi_{(j)}] &= 0, & & & & \text{(otherwise)}.
\end{aligned}$$

For the **case 2iia(\*\*)**, i.e.,  $\delta = 0$ ,  $T_1 = \text{constant}$  and the proper MCs are given by

$$\begin{aligned}
\xi_{(4)} &= \partial_x, \\
\xi_{(5)} &= -\frac{T_2}{T_1}z\partial_x + x\partial_z, \\
\xi_{(6)} &= -\frac{T_2}{T_1}y\partial_x + x\partial_y, \\
\xi_{(7)} &= -\frac{T_2}{\sqrt{T_0}}z\partial_t + \int \sqrt{T_0}dt\partial_z, \\
\xi_{(8)} &= -\frac{T_2}{\sqrt{T_0}}y\partial_t + \int \sqrt{T_0}dt\partial_y, \\
\xi_{(9)} &= \frac{1}{\sqrt{T_0}}\partial_t, \\
\xi_{(10)} &= \frac{x}{\sqrt{T_0}}\partial_t - \frac{1}{T_1} \int \sqrt{T_0}dt\partial_x.
\end{aligned} \tag{50}$$

The Lie algebra is

$$[\xi_{(1)}, \xi_{(3)}] = -\xi_{(2)}, \quad [\xi_{(1)}, \xi_{(6)}] = -\frac{T_2}{T_1}\xi_{(4)}, \quad [\xi_{(1)}, \xi_{(8)}] = -T_2\xi_{(9)},$$

$$[\xi_{(2)}, \xi_{(3)}] = \xi_{(1)}, \quad [\xi_{(2)}, \xi_{(5)}] = -\frac{T_2}{T_1}\xi_{(4)}, \quad [\xi_{(2)}, \xi_{(7)}] = -T_2\xi_{(9)},$$

$$[\xi_{(3)}, \xi_{(5)}] = -\xi_{(6)}, \quad [\xi_{(3)}, \xi_{(6)}] = \xi_{(5)}, \quad [\xi_{(3)}, \xi_{(7)}] = -\xi_{(8)},$$

$$[\xi_{(3)}, \xi_{(8)}] = \xi_{(7)}, \quad [\xi_{(4)}, \xi_{(5)}] = \xi_{(2)}, \quad [\xi_{(4)}, \xi_{(6)}] = \xi_{(1)},$$

$$[\xi_{(4)}, \xi_{(10)}] = \xi_{(9)}, \quad [\xi_{(5)}, \xi_{(6)}] = -\frac{T_2}{T_1}\xi_{(3)}, \quad [\xi_{(5)}, \xi_{(7)}] = -T_2\xi_{(10)},$$

$$[\xi_{(5)}, \xi_{(10)}] = \frac{1}{T_1}\xi_{(7)}, \quad [\xi_{(6)}, \xi_{(8)}] = -T_2\xi_{(10)}, \quad [\xi_{(6)}, \xi_{(10)}] = \frac{1}{T_1}\xi_{(8)},$$

$$[\xi_{(7)}, \xi_{(8)}] = -T_2\xi_{(3)}, \quad [\xi_{(7)}, \xi_{(10)}] = -\xi_{(5)}, \quad [\xi_{(8)}, \xi_{(10)}] = -\xi_{(6)},$$

$$[\xi_{(9)}, \xi_{(10)}] = -\frac{1}{T_1}\xi_{(4)}, \quad [\xi_{(i)}, \xi_{(j)}] = 0, \quad (\text{otherwise}).$$

For the **case 2ii(b)**, Eqs.(27) and (33) yield  $A_j = a_j$  and hence Eq.(34) gives

$$g_4''(x) = \frac{T_1}{2\sqrt{T_0}} \left( \frac{\dot{T}_1}{T_1\sqrt{T_0}} \right) g_4(x). \quad (51)$$

This leads to the following two possibilities

$$(*) \quad \left( \frac{T_1}{2\sqrt{T_0}} \left( \frac{\dot{T}_1}{T_1\sqrt{T_0}} \right) \right) = 0, \quad (**) \quad \left( \frac{T_1}{2\sqrt{T_0}} \left( \frac{\dot{T}_1}{T_1\sqrt{T_0}} \right) \right) \neq 0.$$

In the **case 2iib(\*)**, we have  $\frac{T_1}{2\sqrt{T_0}} \left( \frac{\dot{T}_1}{T_1\sqrt{T_0}} \right) = \eta$ , where  $\eta$  is an integration constant which gives further three cases according as  $\eta > 0$ , or  $\eta = 0$  or  $\eta < 0$ .

When  $\eta = 0$ , i.e., **case 2iib\*(+)**, we have  $\frac{\dot{T}_1}{T_1\sqrt{T_0}} = \epsilon$ , where  $\epsilon$  is non-zero arbitrary constant, the proper MCs are

$$\begin{aligned}\xi_{(4)} &= \partial_x, \\ \xi_{(5)} &= \frac{x}{\sqrt{T_0}}\partial_t + \left(\frac{1}{\epsilon T_1} - \frac{\epsilon}{4}x^2\right)\partial_x, \\ \xi_{(6)} &= \frac{1}{\sqrt{T_0}}\partial_t - \frac{\epsilon}{2}x\partial_x.\end{aligned}\tag{52}$$

The Lie algebra is given by

$$[\xi_{(1)}, \xi_{(3)}] = -\xi_{(2)}, \quad [\xi_{(2)}, \xi_{(3)}] = \xi_{(1)}, \quad [\xi_{(4)}, \xi_{(5)}] = \xi_{(6)},$$

$$[\xi_{(4)}, \xi_{(6)}] = -\frac{\epsilon}{2}\xi_{(4)}, \quad [\xi_{(5)}, \xi_{(6)}] = \frac{\epsilon}{2}\xi_{(5)}, \quad [\xi_{(i)}, \xi_{(j)}] = 0, \quad (\text{otherwise}).$$

For the **case 2iib\*(++)**, we have the following proper MCs

$$\begin{aligned}\xi_{(4)} &= \partial_x, \\ \xi_{(5)} &= \frac{\cosh\sqrt{\eta}x}{\sqrt{T_0}}\partial_t - \left(\frac{\dot{T}_1}{2T_1\sqrt{T_0}}\right)\frac{\sinh\sqrt{\eta}x}{\sqrt{\eta}}\partial_x, \\ \xi_{(6)} &= \frac{\sinh\sqrt{\eta}x}{\sqrt{T_0}}\partial_t - \left(\frac{\dot{T}_1}{2T_1\sqrt{T_0}}\right)\frac{\cosh\sqrt{\eta}x}{\sqrt{\eta}}\partial_x\end{aligned}\tag{53}$$

and the corresponding Lie algebra is

$$[\xi_{(1)}, \xi_{(3)}] = -\xi_{(2)}, \quad [\xi_{(2)}, \xi_{(3)}] = \xi_{(1)}, \quad [\xi_{(4)}, \xi_{(5)}] = \sqrt{\eta}\xi_{(6)},$$

$$[\xi_{(4)}, \xi_{(6)}] = \sqrt{\eta}\xi_{(5)}, \quad [\xi_{(5)}, \xi_{(6)}] = -\frac{1}{\sqrt{\eta}}\left\{\frac{\eta}{T_1} + \left(\frac{\dot{T}_1}{2T_1\sqrt{T_0}}\right)^2\right\}\xi_{(4)},$$

$$[\xi_{(i)}, \xi_{(j)}] = 0, \quad (\text{otherwise}).$$

In the **case 2iib\*(+++)**, we take  $\psi = -\eta$ , where  $\psi$  is also an arbitrary constant and the proper MCs turn out to be

$$\begin{aligned}\xi_{(4)} &= \partial_x, \\ \xi_{(5)} &= \frac{\cos\sqrt{\psi}x}{T_0}\partial_t + \left(\frac{\dot{T}_1}{2T_1\sqrt{T_0}}\right)\frac{\sin\sqrt{\psi}x}{\sqrt{\psi}}\partial_x, \\ \xi_{(6)} &= \frac{\sin\sqrt{\psi}x}{T_0}\partial_t - \left(\frac{\dot{T}_1}{2T_1\sqrt{T_0}}\right)\frac{\cos\sqrt{\psi}x}{\sqrt{\psi}}\partial_x.\end{aligned}\tag{54}$$

The Lie algebra is

$$\begin{aligned} [\xi_{(1)}, \xi_{(3)}] &= -\xi_{(2)}, & [\xi_{(2)}, \xi_{(3)}] &= \xi_{(1)}, & [\xi_{(4)}, \xi_{(5)}] &= -\sqrt{\psi}\xi_{(6)}, \\ [\xi_{(4)}, \xi_{(6)}] &= \sqrt{\psi}\xi_{(5)}, & [\xi_{(5)}, \xi_{(6)}] &= -\frac{1}{\sqrt{\psi}}\left\{\frac{\psi}{T_1} - \left(\frac{\dot{T}_1}{2T_1\sqrt{T_0}}\right)^2\right\}\xi_{(4)}, \\ [\xi_{(i)}, \xi_{(j)}] &= 0, & & \text{(otherwise)}. \end{aligned}$$

The **case 2iib(\*\*)** leads to the four MCs out of which three are usual plane symmetry KVs and one is proper MC.

## 4 Examples

In this section, we shall present some examples which satisfy the constraints.

1. The following metric satisfy the constraints of the case 1(i)

$$ds^2 = e^{t^a}(dt^2 - dx^2 - dy^2 - dz^2) \quad (55)$$

which has six independent MCs.

2. The solution of the constraint of the case 2ia\* \* (+) turns out to be

$$ds^2 = dt^2 - e^{\frac{2t}{a}}dx^2 - e^{\frac{2t}{b}}(dy^2 + dz^2), \quad (56)$$

where  $a$  and  $b$  are non-zero constants such that  $a \neq b$ . This admits five independent MCs.

3. When we solve the constraints of the case 2ia\* \* (++) , we obtain the following solution

$$ds^2 = dt^2 - dx^2 - e^{\frac{2t}{a}}(dy^2 + dz^2), \quad (a \neq 0) \quad (57)$$

which admits seven independent MCs.

4. The metrics given below satisfy the constraints of the case 2ii(a)

$$ds^2 = t^2(dt^2 - dx^2 - dy^2 - dz^2), \quad (58)$$

$$ds^2 = dt^2 - e^{\frac{2t}{a}}(dx^2 + dy^2 + dz^2), \quad (59)$$

$$ds^2 = dt^2 - \left(\frac{t}{t_0}\right)^{2a}(dx^2 + dy^2 + dz^2) \quad (a \neq 0) \quad (60)$$

which have ten independent MCs.

## 5 Discussion and Conclusion

In a recent paper [14], a classification of static plane symmetric spacetimes has been obtained according to their energy-momentum tensor. We have extended this classification to non-static plane symmetric spacetimes satisfying  $T_{01} = 0$  for the degenerate case only [19]. It was found four, six and ten independent MCs out of which four are the usual KVs and the rest are the proper MCs. This paper has been focussed to extend the classification to the non-degenerate case along with  $T_{01} = 0$ . The results are summarized in the form of table 1.

**Table 1.** MCs for the Non-Degenerate Case

Cases	MCs	Constraints
1(i)	6	$(\frac{\dot{T}_2}{T_2\sqrt{T_0}}) \cdot \neq 0, (\sqrt{\frac{T_1}{T_2}}) \cdot = 0$
1(ii)	4	$(\frac{\dot{T}_2}{T_2\sqrt{T_0}}) \cdot \neq 0, (\sqrt{\frac{T_1}{T_2}}) \cdot \neq 0$
2ia(*)	4	$(\frac{\dot{T}_2}{T_2\sqrt{T_0}}) \cdot = 0, \frac{\dot{T}_2}{T_2\sqrt{T_0}} \neq 0, (\frac{T_2}{T_1}) \cdot \neq 0, (\frac{T_1}{T_1\sqrt{T_0}}) \cdot \neq 0$
2ia** (+)	5	$(\frac{\dot{T}_2}{T_2\sqrt{T_0}}) \cdot = 0, \frac{\dot{T}_2}{T_2\sqrt{T_0}} \neq 0, (\frac{T_2}{T_1}) \cdot \neq 0,$ $(\frac{\dot{T}_1}{T_1\sqrt{T_0}}) \cdot = 0, \frac{\dot{T}_1}{T_1\sqrt{T_0}} \neq 0$
2ia** (++)	7	$(\frac{\dot{T}_2}{T_2\sqrt{T_0}}) \cdot = 0, \frac{\dot{T}_2}{T_2\sqrt{T_0}} \neq 0, (\frac{T_2}{T_1}) \cdot \neq 0,$ $(\frac{\dot{T}_1}{T_1\sqrt{T_0}}) \cdot = 0, \frac{\dot{T}_1}{T_1\sqrt{T_0}} = 0$
2i(b)	10	$(\frac{\dot{T}_2}{T_2\sqrt{T_0}}) \cdot = 0, \frac{\dot{T}_2}{T_2\sqrt{T_0}} \neq 0, (\frac{T_2}{T_1}) \cdot = 0,$
2iia(*)	10	$(\frac{\dot{T}_2}{T_2\sqrt{T_0}}) \cdot = 0, \frac{\dot{T}_2}{T_2\sqrt{T_0}} = 0, (\frac{\sqrt{T_1}}{\sqrt{T_0}}) \cdot = 0,$ $\frac{\sqrt{T_1}}{\sqrt{T_0}} \cdot \neq 0$
2iia(**)	10	$(\frac{\dot{T}_2}{T_2\sqrt{T_0}}) \cdot = 0, \frac{\dot{T}_2}{T_2\sqrt{T_0}} = 0, (\frac{\sqrt{T_1}}{\sqrt{T_0}}) \cdot = 0,$ $\frac{\sqrt{T_1}}{\sqrt{T_0}} \cdot = 0$
2iib(*)	6	$(\frac{\dot{T}_2}{T_2\sqrt{T_0}}) \cdot = 0, \frac{\dot{T}_2}{T_2\sqrt{T_0}} = 0, (\frac{\sqrt{T_1}}{\sqrt{T_0}}) \cdot \neq 0,$ $(\frac{T_1}{2T_0} (\frac{\dot{T}_1}{T_1\sqrt{T_0}}) \cdot) = 0$
2iib(**)	4	$(\frac{\dot{T}_2}{T_2\sqrt{T_0}}) \cdot = 0, \frac{\dot{T}_2}{T_2\sqrt{T_0}} = 0, (\frac{\sqrt{T_1}}{\sqrt{T_0}}) \cdot \neq 0,$ $(\frac{T_1}{2T_0} (\frac{\dot{T}_1}{T_1\sqrt{T_0}}) \cdot) \neq 0$

We have obtained four, five, six, seven and ten independent MCs out of which three are the usual plane symmetry KVs and the rest are the proper. Further, it is mentioned here that each case has different constraints on the

energy-momentum tensor. Solving these constraints may lead to some interesting solutions to the EFEs. We have found some examples satisfying the constraints in some cases which help to find the exact number of proper MCs. It would be interesting to classify plane symmetric spacetimes according to their MCs for the degenerate and non-degenerate cases by removing the assumption  $T_{01} = 0$ . This would lead to provide a complete classification of non-static plane symmetric spaces according to their MCs.

## References

- [1] Katzin, G.H., Levine, J. and Davis, H.R.: J. Math. Phys. **10**(1969)617.
- [2] Bokhari, A.H., Kashif, A.R. and Qadir, A.: J. Math. Phys. **41**(2000)2167.
- [3] Hall, G.S. and da Costa, J.: J. Math. Phys. **32**(1991)2848; *ibid* 2854.
- [4] Hall, G.S.: Gen. Rel. and Grav. **30**(1998)1099; *Symmetries and Curvature Structure in General Relativity* (World Scientific, 2004).
- [5] Hall, G.S., Roy, I. and Vaz, L.R.: Gen. Rel. and Grav. **28**(1996)299.
- [6] Carot, J., da Costa, J. and Vaz, E.G.L.R.: J. Math. Phys. **35**(1994)4832.
- [7] Carot, J. and da Costa, J.: *Procs. of the 6th Canadian Conf. on General Relativity and Relativistic Astrophysics*, Fields Inst. Commun. 15, Amer. Math. Soc. WC Providence, RI(1997)179.
- [8] Tsamparlis, M., and Apostolopoulos, P.S.: J. Math. Phys. **41**(2000)7543.
- [9] Sharif, M.: Nuovo Cimento **B116**(2001)673; Astrophys. Space Sci. **278**(2001)447.
- [10] Camci, U. and Barnes, A.: Class. Quantum Grav. **19**(2002)393.
- [11] Camci, U. and Sharif, M.: Gen. Rel. and Grav. **35**(2003)97.
- [12] Camci, U. and Sharif, M.: Class. Quantum Grav. **20**(2003)2169.
- [13] Sharif, M. and Aziz, S.: Gen. Rel. and Grav. **35**(2003)1091; Sharif, M.: J. Math. Phys. **44**(2003)5141.
- [14] Sharif, M.: J. Math. Phys. **45**(2004)1518.



- [15] Sharif, M.: J. Math. Phys. **45**(2004)1532.
- [16] Sharif, M.: Int. J. Mod. Phys. **D14**(2005)1675.
- [17] Sharif, M.: Int. J. Mod. Phys. **A21**(2006)2645.
- [18] Sharif, M. and Ismaeel, T.: Commun. Theore. Phys. **47**(2007)829.
- [19] Sharif, M. and Ismaeel, T.: Mod. Phys. Lett. **A22**(2007)1813.
- [20] Sharif, M.: Acta Physica Polonica **B38**(2007)2003.
- [21] Stephani, H., Kramer, D., MacCallum, M.A.H., Hoenselaers, C. and Hearlt, E.: *Exact Solutions of Einstein's Field Equations* (Cambridge University Press, 2003).