

Finite size effects for the gap in the excitation spectrum of the one-dimensional Hubbard model

M. Colomé-Tatché¹, S. I. Matveenko^{1,2} and G. V. Shlyapnikov^{1,3}

¹ *Laboratoire de Physique Théorique et Modèles Statistiques, Université Paris Sud, CNRS, 91405 Orsay, France.*

² *Landau Institute for Theoretical Physics, Kosygina Str. 2, 119334, Moscow, Russia.*

³ *Van der Waals-Zeeman Institute, University of Amsterdam, Valckenierstraat 65/67, 1018 XE Amsterdam, The Netherlands.*

(Dated: June 21, 2024)

Abstract

We study finite size effects for the gap of the quasiparticle excitation spectrum in the one-dimensional Hubbard model with on-site attraction. Two type of corrections to the result of the thermodynamic limit are obtained. First of all, there is a power law correction due to gapless excitations which behaves as $1/N_a$, where N_a is the number of lattice sites. Second, we obtain an exponential correction related to the existence of gapped excitations. In the weakly interacting regime this correction behaves as $\sim \exp(-N_a\Delta_\infty/4t)$ in the extreme limit of $N_a\Delta_\infty/t \gg 1$, where t is the hopping amplitude and Δ_∞ is the gap in the thermodynamic limit. For not very large $N_a\Delta_\infty/t$ it significantly increases and becomes comparable with the power law corrections. For sufficiently small number of particles, which can be of the order of thousands in the weakly interacting regime, the gap is dominated by finite size effects.

PACS numbers: 03.75.Sc, 71.10.Pm, 03.75.Mn

I. INTRODUCTION

The one-dimensional (1D) Hubbard model with only on-site interaction is exactly solvable by the Bethe Ansatz, and the properties of this model have been widely studied [1]. Low lying excitations of the Hubbard model with attraction are gapless charge excitations and gapped spin excitations, whereas for the repulsive model the gap exists only at half filling in the charge sector.

Finite size corrections to the ground state energy E_0 , due to the gapless part of the spectrum, follow from conformal field theory [2, 3, 4, 5] and have the form: $E_0 - \epsilon_0 L = -\pi v \hbar / 6L$, where $L = N_a a$ is the size of the system, N_a is the number of lattice sites, a is the lattice constant, ϵ_0 is the energy per unit length in the thermodynamic limit, and v is the velocity of gapless excitations. Finite size corrections to the energies of low-lying gapless excitations are also proportional to v/N_a , and the proportionality coefficient depends on the scaling dimensions of the primary fields. However, finite size corrections originating from the gapped sector remained unknown for the Hubbard model. Finite size effects are expected to be important for sufficiently small systems, such as cold atoms in a 1D optical lattice, where the number of particles and lattice sites ranges from several tens to several hundreds [6, 7].

In this paper we solve the Bethe Ansatz equations for a finite number of particles and calculate finite size corrections to the gap for the attractive Hubbard model. As expected, there are power law $1/N_a$ corrections due to gapless excitations, and we also find contributions related to the existence of gapped excitations. In the weakly interacting regime this contribution behaves as $\sim \exp(-N_a \Delta_\infty / 4t)$ in the extreme limit of $N_a \Delta_\infty / t \gg 1$, where t is the hopping amplitude and Δ_∞ is the gap in the thermodynamic limit. For not very large $N_a \Delta_\infty / t$ it significantly increases and can become comparable with power law corrections. We find that the value of the gap increases with decreasing the system size and show how the gap becomes dominated by finite size effects in sufficiently small systems.

The paper is organized as follows. In section II we introduce the Hubbard model together with related Bethe Ansatz equations, and discuss the thermodynamic limit. In section III we present a general approach for finding finite size corrections to the ground state energy and to the gap. Section IV contains our results for (exponential) corrections due to the gapped sector at half filling, and section V the results for power law corrections. In Section VI we show a perturbative approach for solving the Bethe Ansatz equations in the limiting case of $L \ll at/U$. In Section VII we present our numerical results and in Section VIII we conclude.

II. GENERAL EQUATIONS, THERMODYNAMIC LIMIT

The Hubbard model for a system of interacting spin-1/2 fermions on a lattice is described by the Hamiltonian

$$H = -t \sum_{\sigma=\uparrow,\downarrow;j+1}^{N_a} (c_{j,\sigma}^\dagger c_{j+1,\sigma} + c_{j+1,\sigma}^\dagger c_{j,\sigma}) + U \sum_{j=1}^{N_a} n_{j,\uparrow} n_{j,\downarrow}, \quad (1)$$

where the subscript j labels the lattice sites, and U is the on-site interaction. The index σ labels the spin projection, $c_{j,\sigma}^\dagger$ and $c_{j,\sigma}$ are the creation and annihilation fermion operators, and $n_{j,\sigma} = c_{j,\sigma}^\dagger c_{j,\sigma}$ are the particle number operators. Below we express all quantities having the dimension of energy in units of t , and quantities having the dimension of length in units of the lattice constant a .

Lieb and Wu have solved the Fermi-Hubbard model by means of the Bethe Ansatz [8]. The corresponding eigenvalue equations read

$$\sum_{j=1}^N 2 \arctan \left(\frac{\lambda_\alpha - \sin k_j}{u} \right) = 2\pi J_\alpha + \sum_{\beta=1}^M 2 \arctan \left(\frac{\lambda_\alpha - \lambda_\beta}{2u} \right), \quad \alpha = 1, \dots, M, \quad (2)$$

$$N_a k_j = 2\pi I_j - \sum_{\beta=1}^M 2 \arctan \left(\frac{\sin k_j - \lambda_\beta}{u} \right), \quad j = 1, \dots, N, \quad (3)$$

where $u = |U|/4t$, M is the number of spin-down fermions, and N is the total number of particles. The energy of a given state is expressed through the charge momenta k_j :

$$E_N = -2 \sum_{j=1}^N \cos k_j, \quad (4)$$

and it depends on the spin rapidities λ_α only implicitly through the coupled equations (2) and (3). For the ground state the quantum numbers I_j and J_α are integers or half-odd integers depending on the parities of N and M :

$$J_\alpha = \frac{N + M + 1}{2} (\text{mod } 1), \quad I_j = \frac{M}{2} (\text{mod } 1). \quad (5)$$

For the Hubbard model with attraction ($U < 0$), there is a gap in the spectrum of spin excitations. Considering a finite number of particles we define the gap Δ as:

$$2\Delta = E_{N+2}(N_\uparrow + 2, N_\downarrow, U) + E_{N-2}(N_\uparrow - 2, N_\downarrow, U) - 2E_N(N_\uparrow, N_\downarrow, U), \quad (6)$$

where $E_N(N_\uparrow, N_\downarrow; U)$ is the ground state energy for a system with N_\uparrow spin-up and $N_\downarrow = N - N_\uparrow$ spin-down fermions in a lattice with N_a lattice sites, at the interaction strength U . This definition is convenient as it does not change the parity of the quantum numbers I_j and J_α . Without loss of generality, we may put $N_\downarrow \leq N_\uparrow$.

In the thermodynamic limit, where $N \rightarrow \infty$ and $N_a \rightarrow \infty$ while keeping constant densities $n = N/L$ and $n_\downarrow = N_\downarrow/L$, Eq. (6) leads to the same result as the definition $\Delta = E_{N_a}(N_\uparrow + 1, N_\downarrow; U) - 2E_{N_a}(N_\uparrow, N_\downarrow; U) + E_{N_a}(N_\uparrow - 1, N_\downarrow; U)$ introduced in Refs. [8, 13].

In the thermodynamic limit the density of momenta k and the density of rapidities λ are defined as $\rho(k) = L^{-1}\partial I/\partial k$ and $\sigma(\lambda) = L^{-1}\partial J/\partial \lambda$, respectively. Then, the Bethe Ansatz equations for the ground state of the repulsive model become:

$$\rho_\infty(k) = \frac{1}{2\pi} + \frac{\cos k}{\pi} \int_{-B}^B \frac{u}{u^2 + (\lambda - \sin k)^2} \sigma_\infty(\lambda) d\lambda, \quad (7)$$

$$\sigma_\infty(\lambda) + \frac{1}{\pi} \int_{-B}^B \frac{2u}{4u^2 + (\lambda - \lambda')^2} \sigma_\infty(\lambda') d\lambda' = \frac{1}{\pi} \int_{-Q}^Q \frac{u}{u^2 + (\lambda - \sin k)^2} \rho_\infty(k) dk, \quad (8)$$

where the constants B and Q are given by

$$\frac{N}{N_a} = n = \int_{-Q}^Q \rho_\infty(k) dk, \quad \frac{N_\downarrow}{N_a} = n_\downarrow = \int_{-B}^B \sigma_\infty(\lambda) d\lambda, \quad (9)$$

and the ground state energy follows from the relation:

$$E_\infty = -2N_a \int_{-Q}^Q \rho_\infty(k) \cos k dk. \quad (10)$$

The value of the spin gap for an arbitrary filling factor has been calculated in the thermodynamic limit in Refs. [13, 14]. In the case of weak attraction the result is

$$\Delta_\infty = \frac{16 \sin^{3/2}(\pi n/2) \sqrt{u}}{\pi} \exp\left(-\frac{\pi \sin(\pi n/2)}{2u}\right), \quad (11)$$

and the validity of Eq. (11) requires large values of the exponent.

The particle-hole symmetry and the symmetry with respect to interchanging spin-up and spin-down fermions allow one to establish relations between ground state energies of the repulsive and attractive Hubbard models [8]:

$$\begin{aligned} E(N_\uparrow, N_\downarrow; U) &= -(N_a - N_\uparrow - N_\downarrow)U + E(N_a - N_\uparrow, N_a - N_\downarrow; U) \\ &= N_\uparrow U + E(N_\uparrow, N_a - N_\downarrow; -U) = N_\downarrow U + E(N_a - N_\uparrow, N_\downarrow; -U). \end{aligned} \quad (12)$$

Using Eqs. (6) and (12) we can express the spin gap for the attractive Hubbard model through the energies of the repulsive model:

$$2\Delta = 2|U| + E_{N_a-2}(M-2, N_a - M, |U|) + E_{N_a-2}(N_a - M - 2, M, |U|) - 2E_{N_a}(M, N_a - M, |U|). \quad (13)$$

For the half-filled case ($N = 2M = N_a$), with $N_\uparrow = N_\downarrow = M$, Eq. (13) takes the form:

$$\Delta = |U| + E_{N_a-2}(N_a/2, N_a/2 - 2, |U|) - E_{N_a}(N_a/2, N_a/2, |U|). \quad (14)$$

Below we calculate the gap for the attractive Hubbard model. For this purpose we first perform calculations of ground state energies for the repulsive Hubbard model, where the momenta k_j are real numbers, and then obtain Δ for the attractive model by using Eq. (13) (Eq. (14) for the half filled case). This allows us to find exponential finite size corrections, which is not possible in direct calculations for the case of attraction where k_j are complex and can be found only with an exponential accuracy.

III. FINITE SIZE CORRECTIONS. GENERAL APPROACH

Thus, in order to calculate finite size corrections to the gap we have to obtain the three energies of a finite size system, entering the right hand side of Eq. (13). We will follow the scheme proposed by de Vega and Woynarovich [9], which introduces a formalism allowing us to use the Bethe Ansatz in order to calculate finite size corrections to the energy of the ground state. The scheme consists of writing the Bethe Ansatz equations (2) and (3) in the form:

$$Z^s(\lambda) = \frac{1}{N_a} \sum_j^N \frac{1}{\pi} \arctan \frac{\lambda - \sin k_j}{u} - \frac{1}{N_a} \sum_\beta^M \frac{1}{\pi} \arctan \frac{\lambda - \lambda_\beta}{2u}, \quad Z^s(\lambda_\alpha) = \frac{J_\alpha}{N_a}, \quad (15)$$

$$Z^c(k) = \frac{k}{2\pi} + \frac{1}{N_a} \sum_\alpha^M \frac{1}{\pi} \arctan \frac{\sin k - \lambda_\alpha}{u}, \quad Z^c(k_j) = \frac{I_j}{N_a}. \quad (16)$$

We then define the densities of momenta k and rapidities λ for a finite size system as

$$\sigma_N(\lambda) \equiv \frac{dZ^s}{d\lambda} = \frac{1}{2\pi N_a} \sum_{j=1}^N K_1(\lambda - \sin k_j) - \frac{1}{2\pi N_a} \sum_{\beta=1}^M K_2(\lambda - \lambda_\beta), \quad (17)$$

$$\rho_N(k) \equiv \frac{dZ^c}{dk} = \frac{1}{2\pi} + \frac{1}{N_a} \frac{\cos k}{2\pi} \sum_{\alpha=1}^M K_1(\sin k - \lambda_\alpha), \quad (18)$$

where $K_1(x) = 2u/(u^2 + x^2)$, and $K_2 = 4u/(4u^2 + x^2)$. The densities satisfy the relations:

$$\int_{\Lambda_-}^{\Lambda_+} \sigma_N(\lambda) d\lambda = \frac{M}{N_a}, \quad \int_{Q_-}^{Q_+} \rho_N(k) dk = \frac{N}{N_a}, \quad (19)$$

where Q_\pm, Λ_\pm are determined from the equations

$$Z^c(Q_+) = \frac{I_+}{N_a} = \frac{I_{max} + 1/2}{N_a}; \quad Z^s(\Lambda_+) = \frac{J_+}{N_a} = \frac{J_{max} + 1/2}{N_a}. \quad (20)$$

We first perform calculations for the half-filled case. According to Eq. (14) we have to calculate $E_{N_a}(N_a/2, N_a/2; |U|)$ and $E_{N_a}(N_a/2, N_a/2 - 2; |U|)$. In the former case we have $N = N_a$ and $N_\uparrow = N_\downarrow = N_a/2 = M$, and the quantum numbers for the ground state are

$$J_\alpha = \left\{ -\frac{M-1}{2}, \dots, -1, 0, 1, \dots, \frac{M-1}{2} \right\},$$

$$I_j = \left\{ -\frac{N-1}{2}, \dots, -\frac{1}{2}, \frac{1}{2}, \dots, \frac{N-1}{2} \right\}. \quad (21)$$

With $N = N_a$ and $M = N_a/2$, from Eq. (21) we have

$$J_{max} = \frac{N_a/2 - 1}{2}; \quad I_{max} = \frac{N_a - 1}{2}. \quad (22)$$

Then, from Eq. (20) we obtain

$$\begin{aligned} J_+ &= \frac{N_a}{4}, & Z_s(\Lambda_+) &= \frac{1}{4}, \\ I_+ &= \frac{N_a}{2}, & Z_c(Q_+) &= \frac{1}{2}, \end{aligned} \quad (23)$$

and Eqs. (15) and (16) lead to

$$\Lambda_+ = \infty; \quad Q_+ = \pi. \quad (24)$$

When calculating the energy $E_{N_a}(N_a/2, N_a/2 - 2; |U|)$ we have $N = N_a - 2$, $N_\uparrow = N_a/2$, and $N_\downarrow = N_a/2 - 2$. It is convenient to introduce two additional particles with momenta $k_h = \{k_1, k_{N_a}\}$ and spin rapidities $\lambda_h = \{\lambda_1, \lambda_{N_a/2}\}$ in order to satisfy the conditions $Q_\pm = \pm\pi$ and $\Lambda_\pm = \pm\infty$. The discrete Bethe Ansatz equations in this case read

$$\begin{aligned} \sum_{j=1}^{N_a} 2 \arctan \left(\frac{\lambda_\alpha - \sin k_j}{u} \right) &= 2\pi J_\alpha + \sum_{\beta=1}^{N_a/2} 2 \arctan \left(\frac{\lambda_\alpha - \lambda_\beta}{2u} \right) \\ &+ \sum_{k_h} 2 \arctan \left(\frac{\lambda_\alpha - \sin k_h}{u} \right) - \sum_{\lambda_h} 2 \arctan \left(\frac{\lambda_\alpha - \lambda_h}{2u} \right), \quad \alpha = 1, \dots, N_a/2, \end{aligned} \quad (25)$$

$$N_a k_j = 2\pi I_j - \sum_{\beta=1}^{N_a/2} 2 \arctan \left(\frac{\sin k_j - \lambda_\beta}{u} \right) + \sum_{\lambda_h} 2 \arctan \left(\frac{\sin k_j - \lambda_h}{u} \right), \quad j = 1, \dots, N_a. \quad (26)$$

Note that we added only 4 additional equations for defining the numbers k_h, λ_h . Other equations are exactly the Bethe Ansatz equations for $N_a - 2$ particles. The sets J_α, I_j are the same as for the ground state of N_a particles (21).

For the case $N = N_a$ we rewrite equations (17) and (18) in the form of integral equations for the densities $\sigma_{N_a}(\lambda), \rho_{N_a}(k)$:

$$\begin{aligned} \sigma_{N_a}(\lambda) &= \frac{1}{2\pi} \int_{-\pi}^{\pi} K_1(\lambda - \sin k) \rho_{N_a}(k) dk - \frac{1}{2\pi} \int_{-\infty}^{\infty} K_2(\lambda - \mu) \sigma_{N_a}(\mu) d\mu \\ &+ \frac{1}{2\pi} \int_{-\pi}^{\pi} K_1(\lambda - \sin k) X_{N_a}^c(k) dk - \frac{1}{2\pi} \int_{-\infty}^{\infty} K_2(\lambda - \mu) X_{N_a}^s(\mu) d\mu, \end{aligned} \quad (27)$$

$$\begin{aligned} \rho_{N_a}(k) &= \frac{1}{2\pi} + \frac{1}{2\pi} \int_{-\infty}^{\infty} K_1(\sin k - \lambda) \cos k \sigma_{N_a}(\lambda) d\lambda \\ &+ \frac{1}{2\pi} \int_{-\infty}^{\infty} \cos k K_1(\sin k - \lambda) X_{N_a}^s(\lambda) d\lambda, \end{aligned} \quad (28)$$

with the quantities $X_{N_a}^s$ and $X_{N_a}^c$ defined as

$$X_{N_a}^c(k) = \frac{1}{N_a} \left[\sum_{j=1}^{N_a} \delta(k - k_j) \right] - \rho_{N_a}(k), \quad (29)$$

$$X_{N_a}^s(\lambda) = \frac{1}{N_a} \left[\sum_{\alpha=1}^{N_a/2} \delta(\lambda - \lambda_\alpha) \right] - \sigma_{N_a}(\lambda). \quad (30)$$

For $N = N_a - 2$ we do the same and obtain

$$\begin{aligned} \sigma_{N_a-2}(\lambda) &= \frac{1}{2\pi} \int_{-\pi}^{\pi} K_1(\lambda - \sin k) \rho_{N_a-2}(k) dk - \frac{1}{2\pi} \int_{-\infty}^{\infty} K_2(\lambda - \mu) \sigma_{N_a-2}(\mu) d\mu \\ &+ \frac{1}{2\pi} \int_{-\pi}^{\pi} K_1(\lambda - \sin k) X_{N_a-2}^c(k) dk - \frac{1}{2\pi} \int_{-\infty}^{\infty} K_2(\lambda - \mu) X_{N_a-2}^s(\mu) d\mu \\ &- \frac{1}{2\pi N_a} \sum_{k_h} K_1(\lambda - \sin k_h) + \frac{1}{2\pi N_a} \sum_{\lambda_h} K_2(\lambda - \lambda_h), \end{aligned} \quad (31)$$

$$\begin{aligned} \rho_{N_a-2}(k) &= \frac{1}{2\pi} + \frac{1}{2\pi} \int_{-\infty}^{\infty} K_1(\sin k - \lambda) \sigma_{N_a-2}(\lambda) \cos k d\lambda \\ &+ \frac{1}{2\pi} \int_{-\infty}^{\infty} K_1(\sin k - \lambda) X_{N_a-2}^s(\lambda) \cos k d\lambda - \frac{1}{2\pi N_a} \sum_{\lambda_h} K_1(\sin k - \lambda_h) \cos k. \end{aligned} \quad (32)$$

The functions $X_{N_a-2}^{s,c}$ are given by

$$X_{N_a-2}^s(\lambda) = \frac{1}{N_a} \sum_{\alpha=1}^{N_a/2} \delta(\lambda - \lambda_\alpha) - \sigma_{N_a-2}(\lambda), \quad (33)$$

$$X_{N_a-2}^c(k) = \frac{1}{N_a} \sum_{j=1}^{N_a} \delta(k - k_j) - \rho_{N_a-2}(k), \quad (34)$$

where the summation over k_j , λ_α includes the additional numbers k_h , λ_h .

We first consider the thermodynamic limit where the terms $X^{s,c}$ in Eqs. (27), (28), (31) and (32) vanish. For the half-filled case of $N = N_a$ the solution is known [8]:

$$\rho_{\infty, N_a}(k) = \frac{1}{2\pi} + \frac{\cos k}{2\pi} \int_0^{\infty} \frac{\cos(\omega \sin k) \exp(-\omega u)}{\cosh \omega u} J_0(\omega) d\omega, \quad (35)$$

$$\sigma_{\infty, N_a}(\lambda) = \frac{1}{2\pi} \int_0^{\infty} \frac{\cos \omega \lambda}{\cosh \omega u} J_0(\omega) d\omega. \quad (36)$$

Integrating the density of momenta over dk we obtain

$$Z_{\infty, N_a}^c(k) = \frac{k}{2\pi} + \frac{1}{2\pi} \int_0^{\infty} \frac{\sin(\omega \sin k) \exp(-\omega u)}{\omega \cosh \omega u} J_0(\omega) d\omega. \quad (37)$$

For the case of $N = N_a - 2$ the solution is easily found:

$$\begin{aligned}\sigma_{\infty, N_a-2} &= \sigma_{\infty, N_a} - \frac{1}{2\pi N_a} \sum_{k_h} \frac{\pi}{2u} \frac{1}{\cosh[\pi(\lambda - \sin k_h)/2u]} \\ &+ \frac{1}{2\pi N_a} \sum_{\lambda_h} \int_0^\infty \frac{\cos[\omega(\lambda - \lambda_h)]}{\cosh \omega u} \exp(-\omega u) d\omega,\end{aligned}\quad (38)$$

$$\begin{aligned}\rho_{\infty, N_a-2} &= \rho_{\infty, N_a} - \frac{\cos k}{2\pi N_a} \sum_{k_h} \int_0^\infty \frac{\cos[\omega(\sin k - \sin k_h)]}{\cosh \omega u} \exp(-\omega u) d\omega \\ &- \frac{\cos k}{2\pi N_a} \sum_{\lambda_h} \frac{\pi}{2u} \frac{1}{\cosh[\pi(\sin k - \lambda_h)/2u]},\end{aligned}\quad (39)$$

and

$$\begin{aligned}Z_{\infty, N_a-2}^c(k) &= Z_{\infty, N_a}^c(k) - \frac{1}{2\pi N_a} \sum_{k_h} \int_0^\infty \frac{\sin[\omega(\sin k - \sin k_h)] \exp(-\omega u)}{\omega \cosh \omega u} d\omega \\ &- \frac{1}{2\pi N_a} \sum_{\lambda_h} 2 \arctan \left[\tanh \left(\frac{\pi}{4u} (\sin k - \lambda_h) \right) \right].\end{aligned}\quad (40)$$

For calculating the finite size corrections we should find the differences between the densities $\rho_{N_a}(k)$, $\sigma_{N_a}(\lambda)$, $\sigma_{N_a-2}(\lambda)$, $\rho_{N_a-2}(k)$ and their thermodynamic limits values. Subtracting equations of the thermodynamic limit from Eqs. (27), (28) for the case $N = N_a$ we obtain:

$$\begin{aligned}\delta\sigma_{N_a}(\lambda) &= \sigma_{N_a}(\lambda) - \sigma_{\infty, N_a}(\lambda) = \frac{1}{2\pi} \int_{-\pi}^{\pi} K_1(\lambda - \sin k) \delta\rho_{N_a}(k) dk - \frac{1}{2\pi} \int_{-\infty}^{\infty} K_2(\lambda - \mu) \delta\sigma_{N_a}(\mu) d\mu \\ &+ \frac{1}{2\pi} \int_{-\pi}^{\pi} K_1(\lambda - \sin k) X_{N_a}^c(k) dk - \frac{1}{2\pi} \int_{-\infty}^{\infty} K_2(\lambda - \mu) X_{N_a}^s(\mu) d\mu,\end{aligned}\quad (41)$$

$$\begin{aligned}\delta\rho_{N_a}(k) &= \rho_{N_a}(k) - \rho_{\infty, N_a}(k) = \frac{1}{2\pi} \int_{-\infty}^{\infty} K_1(\sin k - \lambda) \cos k \delta\sigma_{N_a}(\lambda) d\lambda \\ &+ \frac{1}{2\pi} \int_{-\infty}^{\infty} \cos k K_1(\sin k - \lambda) X_{N_a}^s(\lambda) d\lambda.\end{aligned}\quad (42)$$

Equations (41) and (42) lead to algebraic equations for the Fourier transforms of $\delta\sigma_{N_a}(\lambda)$ and $\delta\rho_{N_a}(k)$, which yields:

$$\begin{aligned}\delta\sigma_{N_a}(\lambda) &= \int_{-\pi}^{\pi} \frac{\pi}{2u} \frac{1}{\cosh[\pi(\lambda - \sin k)/2u]} X_{N_a}^c(k) \frac{dk}{2\pi} \\ &- \int_{-\infty}^{\infty} d\mu \int_0^\infty \frac{\cos[\omega(\lambda - \mu)]}{\cosh(\omega u)} \exp(-\omega u) X_{N_a}^s(\mu) \frac{d\omega}{2\pi}.\end{aligned}\quad (43)$$

$$\begin{aligned}\delta\rho_{N_a}(k) &= \cos k \int_0^\infty \frac{d\omega}{2\pi} \int_{-\pi}^{\pi} \frac{\exp(-\omega u)}{\cosh(\omega u)} \cos[\omega(\sin k - \sin q)] X_{N_a}^c(q) dq \\ &+ \frac{\cos k}{2\pi} \int_{-\infty}^{\infty} \frac{\pi}{2u} \frac{1}{\cosh[\pi(\sin k - \lambda)/2u]} X_{N_a}^s(\lambda) d\lambda.\end{aligned}\quad (44)$$

Equations for $\delta\sigma_{N_a-2}(\lambda)$ and $\delta\rho_{N_a-2}(k)$ are obtained in a similar way taking into account that in the thermodynamic limit $k_h = \pm\pi$ and $\lambda_h = \pm\infty$. This gives Eqs. (41), (42) and (43), (44) where

N_a is replaced by $N_a - 2$ and $X^{s,c}$ by $\tilde{X}^{s,c}$. The quantity $\tilde{X}_{N_a-2}^s$ is given by Eq.(33) in which the values of λ_h are put equal to $+\infty$ and $-\infty$:

$$\tilde{X}_{N_a-2}^s(\lambda) = \frac{1}{N_a} \sum_{\alpha=1}^{N_a/2-2} \delta(\lambda - \lambda_\alpha) - \sigma_{N_a-2}(\lambda), \quad (45)$$

and $\tilde{X}_{N_a-2}^c$ by Eq. (34) where the values k_h are put equal to $+\pi$ and $-\pi$:

$$\tilde{X}_{N_a-2}^c(k) = \frac{1}{N_a} \left[\sum_{j=1}^{N_a-2} \delta(k - k_j) + \delta(k - \pi) + \delta(k + \pi) \right] - \rho_{N_a-2}(k). \quad (46)$$

The ground state energies (4) for the considered states can be rewritten as

$$\frac{E_{N_a}}{N_a} = -2 \int_{-\pi}^{\pi} \rho_{N_a}(k) \cos k \, dk - 2 \int_{-\pi}^{\pi} X_{N_a}^c(k) \cos k \, dk, \quad (47)$$

$$\begin{aligned} \frac{E_{N_a-2}}{N_a} &= -2 \int_{-\pi}^{\pi} \rho_{N_a-2}(k) \cos k \, dk - 2 \int_{-\pi}^{\pi} X_{N_a-2}^c(k) \cos k \, dk + \frac{2}{N_a} \sum_{k_h} \cos k_h \\ &= -2 \int_{-\pi}^{\pi} \rho_{N_a-2}(k) \cos k \, dk - 2 \int_{-\pi}^{\pi} \tilde{X}_{N_a-2}^c(k) \cos k \, dk - \frac{4}{N_a}. \end{aligned} \quad (48)$$

The finite size corrections to the ground state energy are given by

$$\delta E_N = E_N - E_{\infty, N}, \quad (49)$$

where the energies $E_{\infty, N}$ in the thermodynamic limit are

$$\frac{E_{\infty, N_a}}{N_a} = -2 \int_{-\pi}^{\pi} \rho_{\infty, N_a}(k) \cos k \, dk, \quad (50)$$

$$\frac{E_{\infty, N_a-2}}{N_a} = -2 \int_{-\pi}^{\pi} \rho_{\infty, N_a-2}(k) \cos k \, dk - \frac{4}{N_a}, \quad (51)$$

with $\rho_{\infty, N_a}(k)$ given by Eq. (35) and $\rho_{\infty, N_a-2}(k)$ by Eq. (39) with $k_h = \pm\pi$ and $\lambda_h = \pm\infty$.

Using Eqs. (47), (48), (46), (44), and Eq. (51) with ρ_{∞, N_a} and ρ_{∞, N_a-2} following from Eqs. (35) and Eq. (39), we obtain

$$\frac{\delta E_{N_a}}{N_a} = - \int_{-\pi}^{\pi} \epsilon_c(k) X_{N_a}^c(k) \, dk - \int_{-\infty}^{\infty} \epsilon_s(\lambda) X_{N_a}^s(\lambda) \, d\lambda, \quad (52)$$

$$\frac{\delta E_{N_a-2}}{N_a} = - \int_{-\pi}^{\pi} \epsilon_c(k) \tilde{X}_{N_a-2}^c(k) \, dk - \int_{-\infty}^{\infty} \epsilon_s(\lambda) X_{N_a-2}^s(\lambda) \, d\lambda, \quad (53)$$

where

$$\epsilon_c(k) = 2 \cos k + 2 \int_0^{\infty} \frac{J_1(\omega) \exp(-\omega u)}{\omega \cosh(\omega u)} \cos(\omega \sin k) \, d\omega, \quad (54)$$

$$\epsilon_s(\lambda) = 2 \int_0^{\infty} \frac{J_1(\omega)}{\omega \cosh(\omega u)} \cos(\omega \lambda) \, d\omega, \quad (55)$$

and we used the fact that $\epsilon_s(\pm\infty) = 0$, so that $\tilde{X}_{N_a-2}^s(\lambda)$ can be replaced by $X_{N_a-2}^s(\lambda)$ in Eq. (53).

For the repulsive Hubbard model at half filling, the gap is in the charge sector and the spin sector is gapless. Accordingly, the first term in the right-hand side of Eq. (52) describes the contribution of gapped charge excitations, and the second term is due to the contribution of gapless spin excitations.

We now return to Eq. (14) and using Eqs. (52), (53) write the finite size corrections to the gap of the attractive case in the form

$$\delta\Delta = \delta E_{N_a-2} - \delta E_{N_a} = \delta\Delta_{ng} + \delta\Delta_g + \delta, \quad (56)$$

where

$$\delta\Delta_{ng}/N_a = \int_{-\infty}^{\infty} \epsilon_s(\lambda) X_{N_a}^s d\lambda - \int_{-\infty}^{\infty} \epsilon_s(\lambda) X_{N_a-2}^s d\lambda, \quad (57)$$

is the contribution of the gapless sector,

$$\delta\Delta_g/N_a = \int_{-\pi}^{\pi} \epsilon_c(k) X_{N_a}^c dk - \int_{-\pi}^{\pi} \epsilon_c(k) X_{N_a-2}^c dk, \quad (58)$$

is due to gapped excitations, and we used a relation

$$\tilde{X}_{N_a-2}^c = X_{N_a-2}^c - \frac{1}{N_a} \sum_{h=1}^2 \delta(k - k_h) + \frac{1}{N_a} [\delta(k - \pi) + \delta(k + \pi)].$$

The term δ which is present in the gapped sector is given by

$$\delta = \sum_h \epsilon(k_h) - \epsilon(\pi) - \epsilon(-\pi) \quad (59)$$

and for $N_a\Delta_\infty \gtrsim 1$ it reduces to

$$\delta \approx \epsilon_c''(\pi)(k_h^+ - \pi)^2,$$

where k_h^+ is the value of k_h which is close to π at large N_a . This term behaves as $1/N_a^2$ for large N_a . In the limit of $u \ll 1$ the value of k_h^+ can be found from Eqs. (37) and (40) using the condition $Z_{N_a-2} = 1/2 - 1/2N_a$. We then obtain

$$\frac{\delta}{\Delta_\infty} = \frac{8\pi^2}{[\Delta_\infty N_a + (16/\pi) \ln 2]^2}, \quad (60)$$

where at half filling the gap of the thermodynamic limit Δ_∞ is given by

$$\Delta_\infty = 4u - 4 + 4 \int_0^\infty \frac{\exp(-\omega u) J_1(\omega)}{\omega \cosh \omega u} d\omega \quad (61)$$

at any interaction strength. The origin of the term δ is related to the definition of the gap. The state with two additional or two missing spin up (or spin down) particles contains unpaired fermions with energies above the gap. gapped excitations of our model are $S = 1/2$ -solitons which appear only in pairs. We thus calculate the exact energy of the states with 2 solitons which have different nonzero momenta and energies near the bottom of the excitation band. The contribution (59) takes into account these nonzero kinetic energies of the solitons and is proportional to the curvature of the excitation spectrum.

Note that the contribution of the gapped charge excitations for the repulsive Hubbard model corresponds to the contribution of gapped spin excitations for the attractive model, and the contribution of gapless spin excitations corresponds to the contribution of gapless charge excitations in the attractive case.

IV. EXPONENTIAL CORRECTIONS FOR HALF FILLING

In this Section we calculate the finite size corrections to the gap, originating from the gapped sector and given by Eq. (58). Using Eqs. (16), (29), (23) and the Poisson relation $\sum_m \delta(x - mn) = \sum_m \exp[2\pi xmi]$ we obtain

$$\int_{-\pi}^{\pi} \epsilon_c(k) X_N^c(k) dk = - \int_{-\pi}^{\pi} \epsilon_c(k) \left[\frac{\rho_N(k)}{\exp[-2\pi i N_a Z_N^c(k + i0)] + 1} + \frac{\rho_N(k)}{\exp[2\pi i N_a Z_N^c(k - i0)] + 1} \right] dk. \quad (62)$$

To lowest order, we take the functions $\rho_N(k)$ and $Z_N^c(k)$ equal to their values in the thermodynamic limit. So, they are given by Eq. (35) and Eq. (37) for the case $N = N_a$, and by Eq. (39) and Eq. (40) for $N = N_a - 2$, with $k_h = \pm\pi$ and $\lambda_h = \pm\infty$. One can show that the integral from $-\pi$ to π in the right-hand side of Eq. (62) is equal to the integral from $-\pi + i\text{arcsinh}u$ to $\pi + i\text{arcsinh}u$ for the first term of the integrand plus the integral from $-\pi - i\text{arcsinh}u$ to $\pi - i\text{arcsinh}u$ for the second term.

Equation (58) then reduces to

$$\begin{aligned} \frac{\delta\Delta_g}{N_a} &= \int_{-\pi+i\gamma}^{\pi+i\gamma} \left[\frac{\rho_{N_a}(k)\epsilon_c(k)}{1 + \exp[-2\pi i N_a Z_{N_a}^c(k)]} - \frac{\rho_{N_a-2}(k)\epsilon_c(k)}{1 + \exp[-2\pi i N_a Z_{N_a-2}^c(k)]} \right] dk \\ &+ c.c. = \frac{1}{2\pi i N_a} \int_{-\pi+i\gamma}^{\pi+i\gamma} \epsilon'_c(k) \ln \left(\frac{1 + \exp[2\pi i N_a Z_{N_a}^c(k)]}{1 + \exp[2\pi i N_a Z_{N_a-2}^c(k)]} \right) dk + c.c., \end{aligned} \quad (63)$$

where $\gamma = \text{arcsinh}u$.

It is convenient to present the ratio $\delta\Delta_g/\Delta_\infty$ as a function of $N_a\Delta_\infty$ and u . In Fig. 1 we show $\delta\Delta_g/\Delta_\infty$ versus $N_a\Delta_\infty$ for several values of u , and one clearly sees that for not very large $N_a\Delta_\infty$




FIG. 1: The correction $\delta\Delta_g/\Delta_\infty$ versus $N_a\Delta_\infty$. The solid curve is the result of exact calculations from Eq. (58), and the dashed curve represents the result of Eq. (63). In a) $u = 1$, in b) $u = 0.5$, and in c) $u = 0.25$ with the dotted curve showing the result of Eq. (72)

this correction becomes significant.

The edge points of the integration $k_0 = \pm\pi \pm i\text{arcsinh}u$ are the saddle points at which $\rho_{N_a}(k) = dZ_{N_a}/dk = 0$. For sufficiently large N_a we may use the saddle point approximation, and the expression for $\delta\Delta_g$ becomes

$$\delta\Delta_g \approx C \frac{|\epsilon'_c(k_0)|}{\pi \sqrt{N_a} |\rho'(k_0)|} \exp[-S_0], \quad (64)$$

where

$$\epsilon'_c(k_0) = i \left[2u - 2\sqrt{u^2 + 1} \int_0^\infty J_1(\omega) \tanh(\omega u) \exp(-\omega u) d\omega \right], \quad (65)$$

$$\rho'(k_0) = \frac{i}{2\pi} \left[\frac{u}{\sqrt{u^2 + 1}} + (u^2 + 1) \int_0^\infty \omega \tanh(\omega u) J_0(\omega) \exp(-\omega u) d\omega \right], \quad (66)$$

$$S_0 = -2\pi i (Z_{N_a}^c(k_0) - 1/2) = N_a \left[\gamma - \int_0^\infty \frac{\tanh(\omega u) \exp(-\omega u)}{\omega} J_0(\omega) d\omega \right], \quad (67)$$

$$C = \left[1 - \left(\frac{\Gamma(3/4)}{2\Gamma(5/4)} \right)^4 \right], \quad (68)$$

and we used the relation

$$Z_{N_a-2}^c(k_0) = Z_{N_a}^c(k_0) + \frac{i}{\pi N_a} \int_0^\infty \frac{\tanh(\omega u)}{\omega} \exp(-\omega u) d\omega = Z_{N_a}^c(k_0) + \frac{2i}{\pi N_a} \ln \left[\frac{2\Gamma(5/4)}{\Gamma(3/4)} \right]. \quad (69)$$

The saddle point approximation assumes that the exponent in Eq. (64) is large:

$$S_0 \gg 1. \quad (70)$$

In the case of strong interaction, $u \gg 1$, Eq. (64) gives

$$\delta\Delta_g \approx \frac{1}{\sqrt{N_a} u^{N_a-1}}, \quad (71)$$

and one sees that in this limit the correction $\delta\Delta_c$ is negligible.

The situation changes for $u < 1$. In the limit of $u \ll 1$, from Eq. (64) we obtain:

$$\delta\Delta_g \approx C \sqrt{\frac{2}{\pi}} \frac{\Delta_\infty \exp[-\Delta_\infty N_a/4]}{\sqrt{\Delta_\infty N_a}} \approx 0.6 \frac{\Delta_\infty \exp[-\Delta_\infty N_a/4]}{\sqrt{\Delta_\infty N_a}}, \quad (72)$$

where the gap in the thermodynamic limit, Δ_∞ , is given by Eq. (11). The criterion (70) then becomes $N_a \Delta_\infty \gg 1$. The obtained relation (72) is in accordance with the universal scaling behavior of the gap in massive quantum field theories [18], which is expected for the Hubbard model at $u \ll 1$.

V. POWER LAW CORRECTIONS

The correction to the gap, $\delta\Delta_{ng}$, provided by the gapless sector we calculate using the conformal field theory $1/N$ expansion for the energy [10, 11, 12]:

$$E_N - E_0 = \frac{2\pi}{N_a} v_s (\Delta_+ + \Delta_-), \quad E_0 = \epsilon N_a - \frac{\pi}{6N_a} v_s, \quad (73)$$

where ϵN_a is the energy in the thermodynamic limit, and the velocity of spin excitations is given by

$$v_s = \frac{d\epsilon}{dp} = \frac{\epsilon'_s(B)}{2\pi\sigma(B)}, \quad (74)$$

where B is defined by Eq. (9). The conformal dimensions of primary operators are equal to

$$\Delta_{\pm} = \frac{1}{2} \left(\xi_s(B) D_s \pm \frac{\Delta N_s}{2\xi_s(B)} \right)^2. \quad (75)$$

Integer or half-odd integer numbers ΔN_s , D_s are characterizing the excitation states. The component of the charge dressed matrix ξ_s is determined from the equation

$$\xi_s(\lambda) = 1 - \frac{1}{2\pi} \int_{-B}^B K_2(\lambda - \eta) \xi_s(\eta) d\eta. \quad (76)$$

For the half-filled case ($B = \infty$) the solution is $\xi_s(B = \infty) = 1/\sqrt{2}$. The energy and momentum of a spin excitation are given by

$$\epsilon_s(\lambda) = 2 \int_0^{\infty} \frac{\cos \omega \lambda}{\cosh \omega \lambda} J_1(\omega) \frac{d\omega}{\omega}, \quad (77)$$

$$p_s(\lambda) = \frac{\pi}{2} - \int_0^{\infty} \frac{\sin \omega \lambda}{\cosh \omega \lambda} J_0(\omega) \frac{d\omega}{\omega}, \quad (78)$$

$$\frac{dp_s}{d\lambda} = -2\pi\sigma(\lambda). \quad (79)$$

The excitation state with the energy $E(N_a/2, N_a/2 - 2)$ is characterized by the numbers $D_s = 0$, $\Delta N_s = 1$. According to Eq. (14), this leads to the correction

$$\delta\Delta_{ng} = \frac{\pi}{2N_a} \frac{v_s}{\xi_s^2}. \quad (80)$$

For the half-filled case we obtain

$$\delta\Delta_{ng} = \frac{2\pi}{N_a} \frac{I_1(\pi/2u)}{I_0(\pi/2u)}. \quad (81)$$

A more accurate result following from the calculations in Refs. [10, 12] reads:

$$\delta\Delta_{ng} = \frac{2\pi}{N_a} \frac{I_1(\pi/2u)}{I_0(\pi/2u)} \left(1 - \frac{1}{2 \ln[N_a I_0(\pi/2u)]} \right). \quad (82)$$

The comparison of $\delta\Delta_{ng}/\Delta_{\infty}$ (82) for $u = 1$ with the result of exact calculations from Eq. (57) shows the validity of conformal results even for not very large N_a (see Fig. 2). For example, at $u = 1$ even for $N_a = 10$, ($N_a\Delta_{\infty} \approx 13$) the relative difference is $\sim 15\%$.

In the limit of strong coupling Eq. (81) yields:

$$\delta\Delta_{ng} \simeq \frac{\pi^2}{2N_a u}; \quad u \gg 1, \quad (83)$$

and the power law correction (83) always dominates over the negligible exponential correction (71). The situation changes for $u \lesssim 1$. For not very large $\Delta_{\infty} N_a$, the non-conformal correction $\delta\Delta_g$ originating from gapped excitations becomes comparable with $\delta\Delta_{ng}$. This is seen from Fig. 3, where we present the two corrections for $u = 1$, $u = 0.5$ and $u = 0.25$. We also display the correction δ/Δ_{∞} which turns out to be significant for $N_a\Delta_{\infty}$ not greatly exceeding unity.




FIG. 2: The correction $\delta\Delta_{ng}/\Delta_\infty$ versus $N_a\Delta_\infty$ for $u = 1$. The solid curve shows the result of exact calculation from Eq. (57) and the dashed curve shows the result of Eq. (82).

VI. DISCUSSION OF THE RESULTS

In Fig. 4 we present the results of exact calculations for the gap Δ at half filling from Eqs. (2), (3), Eqs. (4) and (14) for the same u as in Fig. 3. A direct comparison of Δ with the gap of the thermodynamic limit Δ_∞ (61) shows that finite size corrections can be safely omitted for $N_a\Delta_\infty > 40$. On the other hand, already at $N_a\Delta_\infty < 10$, they dominate the gap. For $u = 0.25$ where $\Delta_\infty = 5 \cdot 10^{-3}$ this occurs already at $N_a < 1.5 \cdot 10^3$. For $u = 1$ we have $\Delta_\infty = 1.28$ and finite size corrections are important only for $N_a < 30$.

It is important that for $N_a\Delta_\infty \sim 10$ or even somewhat large the non-conformal corrections originating from the gapped sector become comparable with conformal corrections. This is seen in Fig. 1. For $u = 1$ even the exponential correction $\delta\Delta_g$ is comparable with $\delta\Delta_{ng}$. For $u = 0.5$ and $u = 0.25$ this is the correction δ that remains comparable with $\delta\Delta_{ng}$, whereas the exponential correction $\delta\Delta_g$ is about 10% of $\delta\Delta_{ng}$.

Qualitatively, the dependence $\Delta(N_a)$ remains the same for smaller filling factors. This is seen from Fig. 5, where we present our numerical results for $n = 0.2$. For $u = 1.25$ finite size effects become important only at a very small number of lattice sites $N_a < 20$. For $u = 0.25$ the thermodynamic-limit gap is $\Delta_\infty \approx 6 \cdot 10^{-2}$ and finite size effects are already important for $N_a \approx 500$.




FIG. 3: The result of exact calculations from Eqs. (57), (58) and (59) for $\delta\Delta_{ng}/\Delta_\infty$ (solid curve), $\delta\Delta_g/\Delta_\infty$ (dashed curve), and δ/Δ_∞ (dotted curve) versus $N_a\Delta_\infty$. The sum $\delta\Delta_{ng} + \delta\Delta_g + \delta + \Delta_\infty$ exactly coincides with the gap Δ calculated directly from the Bethe Ansatz equations (2) and (3) using Eqs. (4) and (14). In a) $u = 1$, in b) $u = 0.5$, and in c) $u = 0.25$.

VII. LIMIT OF $N_a u \ll 1$

The discussed results were related to the case where $N_a u \gg 1$. For example, the inequality (70) automatically requires the condition $N_a u \gg 1$, irrespective of whether u is large or small. In the opposite limit of $N_a u \ll 1$, which can be realized for $u \ll 1$, the energy spectrum of




FIG. 4: The gap Δ in units of t versus N_a at half filling for $u = 1$ in a), $u = 0.5$ in b), and $u = 0.25$ in c).

the attractive Hubbard Model shows no exponential gap and both charge and spin sectors are conformal. The analysis of Lieb-Wu equations for this case has been done in [19, 20], and found corrections $\sim u/N_a$. This can be understood from the conformal $1/N_a$ expansion as a consequence of the linear dependence of the velocities of elementary excitations on the interaction constant u [19, 21].

Here we consider the limit of $uN_a \ll 1$ for completeness and present first order corrections




FIG. 5: The gap Δ in units of t versus N_a , calculated numerically for $u = 0.25$ in a), and $u = 1.25$ in b) for the filling factor $n = 0.2$.

in u to the ground state energy and to the gap in the excitation spectrum. As in the previous sections, we calculate the energy for the Hubbard model in the repulsive case, where the Bethe Ansatz equations are easily solved, and then restore the energy for the attractive case using the particle-hole symmetry.

So, consider a system of N particles (N_\downarrow spin-down and $N - N_\downarrow$ spin-up) with repulsive interaction. From the Lieb-Wu equation (2) we obtain the momenta k_j to first order in u :

$$\exp(ik_j N_a) = \prod_{\alpha=1}^{N_\downarrow} \frac{\sin k_j - \lambda_\alpha + iu}{\sin k_j - \lambda_\alpha - iu} \Rightarrow \delta k_j = k_j - k_j^0 = \frac{1}{N_a} \sum_{\alpha=1}^{N_\downarrow} \frac{2u}{\sin(k_j^0) - \lambda_\alpha^0}, \quad (84)$$

where k^0 and λ^0 are the momenta and rapidities for $u \rightarrow 0$. The energy itself and the interaction-induced change of the energy are given by

$$E = -2 \sum_{j=1}^N \cos k_j, \quad \delta E = E - E^0 = \frac{4u}{N_a} \sum_{j=1}^N \sum_{\alpha=1}^{N_\downarrow} \frac{\sin k_j^0}{\sin(k_j^0) - \lambda_\alpha^0}, \quad (85)$$

where E_0 is the ground state energy for $u \rightarrow 0$. We now calculate the densities of momenta k and

rapidities λ in the thermodynamic limit from Eqs. (7) and (8). To the lowest order in u we obtain:

$$2\sigma(\lambda) = \int_{-Q}^Q \delta(\lambda - \sin k) \rho(k) dk, \quad (86)$$

$$\rho(k) = \frac{1}{2\pi} + \cos k \int_{-B}^B \delta(\lambda - \sin k) \sigma(\lambda) d\lambda. \quad (87)$$

The solution of Eqs. (86) and (87) is ($n_{\downarrow} \leq n_{\uparrow}$)

$$\rho(k) = \begin{cases} 1/\pi; & k \leq \pi n_{\downarrow}, \\ 1/2\pi; & \pi n_{\downarrow} < k \leq \pi(n - n_{\downarrow}). \end{cases} \quad (88)$$

$$\sigma(\lambda) = \frac{1}{2\pi} \frac{1}{\sqrt{1 - \lambda^2}}. \quad (89)$$

Then, using Eq. (9) for the total number of particles and the number of spin-down particles, we find an expression for the integration limits Q and B :

$$B = \sin(\pi n_{\downarrow}), \quad Q = \pi(n - n_{\downarrow}). \quad (90)$$

Using Eqs. (88) and (89) we obtain the interaction-induced change of the energy to first order in u :

$$\begin{aligned} \frac{\delta E}{N_a} &= 4u \int_{-Q}^Q dk \int_{-B}^B \frac{\sin k}{\sin k - \lambda} \sigma(\lambda) \rho(k) d\lambda = \\ &= \frac{4u}{\pi^2} (\pi n_{\downarrow})^2 + \frac{4u}{\pi^2} \int_{\pi n_{\downarrow}}^{\pi(n - n_{\downarrow})} dk \int_0^{\pi n_{\downarrow}} \frac{\sin^2 k}{\sin^2 k - \sin^2 q} dq, \end{aligned} \quad (91)$$

where we turned to the variable $q = \arcsin \lambda$. For the case of attraction we should substitute $n_{\uparrow} \rightarrow 1 - n_{\uparrow}$, $n_{\downarrow} \rightarrow n_{\downarrow}$, in accordance with the symmetry properties (12). Integrating over dq we find:

$$\frac{\delta E(-u)}{N_a} = -4un_{\downarrow} + 4un_{\downarrow}^2 + \frac{2u}{\pi^2} \left(2 \int_{\pi n_{\downarrow}}^{\pi/2} - \int_{\pi n_{\downarrow}}^{\pi n_{\uparrow}} \right) \tan k \ln \left(\frac{\tan k + \tan \pi n_{\downarrow}}{\tan k - \tan \pi n_{\downarrow}} \right) dk, \quad (92)$$

where n_{\downarrow} and n_{\uparrow} are already occupation numbers for the attractive Hubbard model, and we assume that $n_{\downarrow} \leq n_{\uparrow}$. Eq. (92) leads to the following result for the interaction-induced change of the energy to first order in $N_a u$:

$$\begin{aligned} \frac{\delta E(-u)}{N_a} &= 4un_{\downarrow}^2 - 4un_{\downarrow} + \frac{4u}{\pi} \arctan \pi n_{\downarrow} - \frac{u}{\pi^2} \ln \left(\frac{1 + \tan^2 \pi n_{\uparrow}}{1 + \tan^2 \pi n_{\downarrow}} \right) \ln \left(\frac{\tan \pi n_{\uparrow} + \tan \pi n_{\downarrow}}{\tan \pi n_{\uparrow} - \tan \pi n_{\downarrow}} \right) \\ &+ \frac{2u}{\pi^2} \left(-\text{Re}Li_2 \left(\frac{2 \tan \pi n_{\downarrow}}{\tan \pi n_{\downarrow} - i} \right) - \text{Re}Li_2 \left(\frac{\tan \pi n_{\uparrow} + \tan \pi n_{\downarrow}}{\tan \pi n_{\downarrow} - i} \right) + \text{Re}Li_2 \left(\frac{\tan \pi n_{\uparrow} - \tan \pi n_{\downarrow}}{-\tan \pi n_{\downarrow} - i} \right) \right), \end{aligned}$$

where $Li_2(z) = \sum_{k=1}^{\infty} z^k/k^2$ is a polylogarithmic function.

In the limit of small filling factors, $N_\uparrow \ll N_a$ and $N_\downarrow \ll N_a$, after a straightforward algebra we obtain:

$$\frac{\delta E}{N_a} \approx u(-4n_\downarrow - 4n_\uparrow n_\downarrow + 2n_\downarrow^2) \quad (93)$$

The limit of small filling factors in the Hubbard model corresponds to the gas phase of spin-1/2 fermions. For this case the ground state energy at $Nu \ll 1$ has been calculated in Refs. [15, 16, 17], and the result of Eq. (93) coincides with that of Refs. [15, 16, 17] in the attractive case.

Using Eq. (6) we then find a small interaction-induced correction to the gap in the excitation spectrum ($n_\uparrow = n_\downarrow = n/2$) of the attractive model to the lowest order in uN_a . For small filling factors we have: $\delta\Delta \approx 4u/N_a$, and in the considered limit of $N_a u \ll 1$ this correction is small compared to the level spacing $\sim 1/N_a$ in our finite size system.

VIII. CONCLUSIONS

In conclusion, we have studied finite size effects for the gap in the excitation spectrum of the 1D Fermi Hubbard model with one-site attraction. For the situation in which the thermodynamic-limit gap Δ_∞ exceeds the level spacing (near the Fermi energy) of the finite size system, there are two types of finite size corrections. For large interactions ($u \gg 1$) the leading is a power law conformal correction to Δ_∞ , which behaves as $1/N_a$ and originates from the gapless sector of the excitation spectrum. We also find non-conformal corrections originating from the gapped branch of the spectrum. As found at half filling, in the weakly interacting regime ($u \lesssim 1$) the non-conformal corrections can become of the order of the conformal correction.

For sufficiently small number of lattice sites (particles) the gap Δ is dominated by finite size effects. From a general point of view, this happens when $\Delta_\infty \lesssim 1/N_a$, i. e. Δ_∞ is smaller than the level spacing of the finite size system at energies close to the Fermi energy. Accordingly, for large interactions ($u \gg 1$) the finite size effects are not important as long as $N_a \gg 1$. However, in the weakly interacting regime ($u \lesssim 1$) they become dominant already at significantly larger N_a than a simple dimensional estimate $1/\Delta_\infty$. This is clearly seen from our results in Fig. 3 and Fig. 4 for $\Delta(N_a)$ at half filling.

Our findings are especially important for the studies of the 1D regime with cold atoms, where the number of particles in a 1D tube ranges from several tens to several hundreds [6, 7]. For such a system in the weakly interacting regime one can not use the result of the thermodynamic limit for the gap. Consequently, one can not employ the local density approximation for Δ based on this

result, for finding the spectrum of isospin gapped excitations in an external harmonic potential.

Acknowledgements

We are grateful to A. M. Tsvelik for fruitful discussions and acknowledge hospitality and support of Institut Henri Poincaré during the workshop “Quantum Gases” where part of this work has been done. We also acknowledge D.L. Kovrizhin for his useful suggestions on numerical methods. The work was supported by the IFRAF Institute, by ANR (grants 05-BLAN-0205 and 06-NANO-014-01), by the QUEDDIS program of ESF, and by the Dutch Foundation FOM. LPTMS is a mixed research unit No. 8626 of CNRS and Université Paris Sud.

-
- [1] F. H. L. Essler, H. Frahm, F. Gohmann, A. Klumper and V. Korepin, *The One-Dimensional Hubbard Model* (Cambridge University Press, Cambridge, 2005).
 - [2] A. A. Belavin, A. M. Polyakov and A. B. Zamolodchikov, Nuclear Physics B **241**, 333 (1984).
 - [3] J. L. Cardy, J. Phys. A: Math. Gen. **17**, L385 (1984).
 - [4] H. W. J. Blöte, J. L. Cardy, M. P. Nightingale, Phys. Rev. Lett **56**, 742 (1986).
 - [5] I. Affleck, Phys. Rev. Lett. **56**, 746 (1986).
 - [6] See for review: I. Bloch, J. Dalibard and W. Zwerger, Rev. Mod. Phys. **10** 885 (2008).
 - [7] See for review: S. Giorgini, L. P. Pitaevskii, and S. Stringari, Rev. Mod. Phys. (2008).
 - [8] E. H. Lieb and F.Y. Wu, Phys. Rev. Lett. **20**, 1445 (1968).
 - [9] H. J. de Vega and F. Woynarovich, Nuclear Physics B **251** [FS13], 439 (1985).
 - [10] F. Woynarovich and H.-P. Ecker, J. Phys. A: Math. Gen. **20**, L443 (1987).
 - [11] F. Woynarovich, J. Phys. A: Math. Gen.. **22**, 4243 (1989).
 - [12] H. Frahm, V. E. Korepin, Phys. Rev. B **42** 10553 (1990).
 - [13] V. Ya. Krivnov and A. A. Ovchinnikov, Sov. Phys. JETP **40**, 781 (1974).
 - [14] A. I. Larkin and J. Sak, Phys. Rev. Lett. **39**, 1025 (1977).
 - [15] M. T. Batchelor, M. Bortz, X. W. Guan and N. Oelkers, J. Phys. Conf. Ser. **42**, 5 (2006).
 - [16] G. E. Astrakharchik, D. Blume, S. Giorgini and L. P. Pitaevskii, Phys. Rev. Lett. **93**, 050402 (2004).
 - [17] J. N. Fuchs, A. Recati and W. Zwerger, Phys. Rev. Lett. **93**, 090408 (2004).
 - [18] M. Luscher, Commun. Math. Phys. **104**, 177 (1986).
 - [19] H. Frahm and M.P. Pfafmuller, Phys. Lett. A **204**, 347 (1995).
 - [20] L. Amico and A. Mastellone, Phys. Rev. B **73**, 214513 (2006).
 - [21] A.V. Chubukov, D.L. Maslov and F.H.L. Essler, arXiv:0801.1837.