

Mass Distribution, Rotation Curves and Gravity Theories

Rahul Datta

Space Applications Centre, Indian Space Research Organization,
Ahmedabad-380015, India

Dilip G. Banhatti

School of Physics, Madurai-Kamaraj University, Madurai 625021, India
[Email: dilip.g.banhatti@gmail.com]

ABSTRACT

{Comparison of mass density profiles of galaxies of varying sizes from some gravity theories based on observed galaxy rotation curves and assessing the need for dark matter.}
We present an analysis of five galaxies of varying galactic radii: NGC6822 (4.8 kpc), Large Magellanic Cloud (9 kpc), Milky Way (17 kpc), NGC3198 (30 kpc) and UGC9133 (102.5 kpc). The masses and mass density profiles of these galaxies have been computed using the scientific computing s/w package MATLAB taking the already available velocity profiles of the galaxies as the input, and without considering any dark matter contribution. We have plotted these profiles after computing them according to three different theories of gravity (and dynamics) : Newtonian (black line), Modified Newtonian Dynamics (MoND) (green line) and Vacuum Modified Gravity (red line). We also consider how the profile due to Newtonian theory would modify if we take into account a small negative value of the cosmological constant ($-5 \times 10^{-56} \text{ cm}^{-2}$ from theory) (blue line). Comparing these masses and mass density profiles, we try to form an idea regarding what could be a realistic theory of gravity and whether we need dark matter to explain the results.

Keywords : disk galaxy rotation curves, galaxy mass, mass density profile, dark matter, Newtonian theory, MoND, Vacuum Modified Gravity, negative cosmological constant

MOTIVATION

“Missing mass” was first inferred via dynamics for clusters of mostly elliptical galaxies using virial theorem. It was soon found to be dynamically present, although it was the light that was missing, same as in clusters, on even single galaxy scale, especially for spirals, most naturally taken to be balanced between rotation and gravity. Assigning mass-to-light ratios on various scales via putative astrophysical light emission processes on those scales, *dark matter*, which gravitated (and hence participated in dynamics), but emitted little or no light, became firmly established in astrophysics and cosmology.

In this paper, we consider the single galaxy scale, and use five galaxies, either disk spirals or dynamically similar galaxies, to examine the connection between mass distribution and rotation curve in a few representative models, without assuming any dark matter, on scales from 5 kpc (size of the smallest of the five galaxies) to 100 kpc (size of the largest of the 5).

Calculation / Computation Scheme

We first define the geometry and the notation for our calculation (computed using MATLAB s/w package). The galaxy disk is assumed axisymmetric.




FIG.1: Top view of a galaxy. The semicircular arcs represent rings constituting the disk. The other half forming the full circular arc is implied.

Definitions & notation (referring to FIGs. 1 & 2):

pc = parsec, $3.08568E13$ kms, 3.26151 light years

pi = 3.14159265

dr = radial thickness of a ring, pc

dth = $(pi / 180)$ radians

th = angle at the galaxy center, between galaxy radial line to test mass and radial line to ring segment, radians

alpha = angle subtended at the test mass by the center of the fundamental segment mass and the center of the galaxy, radians

beta = angle subtended at the test mass by the center of the fundamental segment mass and the elemental thickness dz of the fundamental segment mass, radians

pytks = multiplier to change v (pc/yr) to v (km/sec) = $9.7778E5$ km/pc*yr/sec

msun = mass of our sun, $1.989E33$ gms

htype indicates type of the galaxy




FIG.2: Edge-on view of a fundamental segment of a ring in an axisymmetric disk galaxy. Cf FIG.1.

$r(i)$ = radius to centerline circle of i th ring = $r_m - dr/2$ pc

$rr(i)$ = radius to rod used to represent fundamental segment mass for the i th ring =
 $r_m - (dr / 2) * (r_m - 0.575 dr) / (r_m - dr / 2)$ pc

N_r = number of rings

G = gravitational constant, $4.498E(-15)$ pc³/(msuns*yr²)

r_m = radius to outer edge of a ring, pc

r_{max} = galaxy maximum/rim radius, pc

r_t = radius to test mass from galaxy center, pcs

the test mass is located at the outer edge of each ring

$h(i)$ = galaxy thickness at radius r corresponding to i th ring, pc

$m_r(i)$ = mass contained only inside the i th ring, msuns

$M_r(i)$ = galaxy mass contained within the outer edge of the i th ring, i.e., sum of masses inside each ring starting from the galaxy center upto the i th ring, msuns

M = galaxy total mass, msuns

$dv(i)$ = volume of a fundamental segment of the i th ring, pc³
 $= h(i)*r(i)*dth*dr$

$rv(i)$ = volume of the i th ring, pc³
 $= 360*dv(i)$

$tvol = \text{galaxy total volume, pc}^3$
 $\rho(i) = \text{density of the } i\text{th ring, msuns/pc}^3$
 $\rho_{av} = \text{average density of galaxy} = M / tvol, \text{ msuns/pc}^3$
 $SMD(i) = \text{surface mass density of } i\text{th ring, msuns / pc}^2$

$SMD_{av} = \text{average surface mass density of galaxy}$
 $= M / (\pi * r_{max}^2) \text{ msuns / pc}^2$

$rSMD(i) = rd(i) * SMD(i)$

$v(i) = \text{computed orbital speed at test mass radius } r_t \text{ at the outer edge of } i\text{th ring, kms/sec}$

$v_{krim} = \sqrt{GM / r_{max}}$, the Kepler speed at the rim, kms/sec

$v_m = \text{measured speed, km/sec}$

$v_{max} = \text{maximum measured speed, km/sec}$

$mr_{nor}(i) = mr(i) / M$

$Mr_{nor}(i) = Mr(i) / M$

$hd(i), rd(i) = (h(i), r(i)) / r_{max}$

$\rho_{nor}(i) = \rho(i) / \rho_{av}$

$SMD_{nor}(i) = SMD(i) / SMD_{av}$

$rSMD_{nor}(i) = rd(i) * SMD_{nor}(i)$

$v_{nor}(i) = v(i) / v_{krim}$

FORWARD PROBLEM: Computation of the rotation profile from a given mass distribution.

Referring to FIG.2, acceleration due to one fundamental segment mass of a ring:

From Newton's law, centripetal acceleration = GM/r^2

Here, for a fundamental segment, ρ is a constant.

Hence, M/r^2 is equivalent to $\rho \int (Sdz/b^2) \cos(\beta)$, taking the component of the acceleration due to the mass of the elemental thickness, dz of the fundamental segment of a ring along the direction of the plane of the galactic disc, where S = cross-sectional area of the fundamental segment cut along the plane of the galaxy disc. Therefore,

$$Acc = G (\rho) \int (S dz/b^2) \cos(\beta)$$

$$= G (\rho) \int (S dz/b^2)(c/b)$$

But, $z = c \tan(\beta)$ and $b = c \sec(\beta)$ or, $dz = c \sec^2(\beta) d(\beta)$. Therefore, Acc

$$= G (\rho) S \int (c \sec^2(\beta) d(\beta)/b^2)(c/b)$$

$$= G (\rho) S \int (c^2 \sec^2(\beta)/b^3) d(\beta)$$

$$= G (\rho) S \int (c^2 \sec^2(\beta)/c^3 \sec^3(\beta)) d(\beta)$$

$$= G (\rho)(S/c) \int d(\beta)/\sec(\beta)$$

$$\begin{aligned}
&= G (\rho)(S/c) \int_{z=-h/2}^{z=+h/2} \cos(\beta)d(\beta) \\
&= G (\rho)(S/c) [\sin(\beta)]_{z=-h/2}^{z=+h/2} \\
&= G (\rho)(S/c) [z/(c^2+z^2)^{0.5}]_{z=-h/2}^{z=+h/2} \\
&= G (\rho)(S/c) [(h/2)/(c^2+(h/2)^2)^{0.5} + (h/2)/(c^2+(h/2)^2)^{0.5}] \\
&= G (\rho)(S/c) [h/(c^2+(h/2)^2)^{0.5}]
\end{aligned}$$

Referring to FIG.1, each ring consists of 360 such fundamental segments of mass and angle θ varies from 0 to π radians or 0 to 180 degrees, both in the clockwise and the anticlockwise direction from the line joining the galaxy center to the test mass. The centers of the fundamental segments correspond to 0.5 degrees, 1.5 degrees, 2.5 degrees and so on until 179.5 degrees both in the clockwise and anticlockwise direction. Therefore the centripetal acceleration of the test mass due to each ring:

$$v^2/rt = 2 (\text{pytks})^2 \int_{\theta=0}^{\theta=\pi} G (\rho)(S/c) [h/(c^2+(h/2)^2)^{0.5}] \cos(\alpha) d\theta$$

the $\cos(\alpha)$ term being multiplied for taking the component of the acceleration of the test mass due to each fundamental segment of the ring along the radial direction towards the galactic center and the term 2 being multiplied to account for both the clockwise and anticlockwise directions.

$$\cos(\alpha) = (rt - rr \cos(\theta))/c.$$

Hence, the squared orbital velocity due to each ring:

$$v^2 = 2 rt (\text{pytks})^2 \int_{\theta=0}^{\theta=\pi} G (\rho) (S/c^2) [h/(c^2+(h/2)^2)^{0.5}] (rt - rr \cos(\theta)) d\theta$$

Discretizing the above formula by converting the integral to summation for the purpose of computation, we have:

$$v^2 = 2 rt (\text{pytks})^2 \sum_{i=0}^{i=180} G (\rho) (S/c^2) [h/(c^2+(h/2)^2)^{0.5}] (rt - rr \cos((i-0.5)d\theta))$$

where $d\theta = (\pi/180)$ radians and

$$rr = (j dr) - (dr/2) \{ (j dr) - (0.575 dr)/(j dr) - (0.5 dr) \}$$

dr = radial thickness of each ring

$$S = r(j) d\theta dr$$

Summing up over all the rings, we have

$$v^2 = rt \sum_{j=0}^{Nr} \sum_{i=0}^{180} G(\rho)r(j)d\theta dr/c^2 [h(j)/(c^2+(h(j)/2)^2)^{0.5}] [rt - rr \cos((i-0.5)d\theta)]$$

REVERSE PROBLEM: Finding the mass distribution for a given rotation profile

The reverse problem is done by repeated applications of the forward problem using the equivalent thickness distribution, an arbitrary density distribution is input, and the speed profile computed using the above formula. The speed errors (measured minus computed) at each of the ring outer radii are then used to correct the densities of the rings and the process repeated until all speed errors are small enough to be neglected.

$$errv = (vm-v) / v_{max} ; f = 0.75*errv$$

if all $errv < 1E-6$ then quit

else

$$\rho(i) = (1+f) \rho(i-1) \text{ for each cycle } i$$

where vm = measured speed, and v_{max} = maximum computed speed

To compute the mass and mass density profiles after taking into account the contribution of a cosmological constant, λ taken from theory to be equal to $-5 \times 10^{-56} \text{cm}^{-2}$ in magnitude can be found out by using a new velocity profile as input, which is obtained after subtracting a quantity equal to $c*r(i)*\sqrt{\lambda/3}$ from the observed velocity profile. Note that we have considered a negative cosmological constant scenario.

To compute the mass and mass density profiles as per the Modified Newtonian Dynamics (MoND), we use the following equation for the FORWARD Algorithm and the REVERSE Algorithm, which incorporates the results of the Forward Algorithm, remains otherwise the same as earlier.

$$v_{MoND}^2 = v^2 (1 + \sqrt{(M_0/Mr_1(j)) (1 - \exp(-rt/r_0) (1+rt/r_0))})$$

$$v_{MoND}^2 = rt \sum_{j=0}^{Nr} \sum_{i=0}^{180} G(\rho)r(j)d\theta dr/c^2 [h(j)/(c^2+(h(j)/2)^2)^{0.5}] * [rt - rr \cos((i - 0.5)d\theta)] (1 + \sqrt{(M_0/Mr_1(j)) (1 - \exp(-rt/r_0) (1+rt/r_0))})$$

where,

$M_0 = 9.6e+11$ msuns for $r_{max} > 12000$ pc
 $= 2.4e+11$ msuns for $r_{max} = < 12000$ pc
 $r_0 = 13920$ pc for $r_{max} > 12000$ pc
 $= 6960$ pc for $r_{max} = < 12000$ pc

$Mr_1(j)$ = the quantity of mass contained within the radius at which the velocity is being computed.

To compute the mass and mass density profiles as per Vacuum Modified Gravity (VMG), we use the following equation for the FORWARD Algorithm and the REVERSE Algorithm, which incorporates the results of the Forward Algorithm, remains otherwise the same as earlier.

$$\begin{aligned}
 v^2 = & \text{rt} (\text{pytks})^2 \left\{ \sum_{j=0}^{j=N_r} \sum_{i=0}^{i=180} (G \rho r(j) \text{dth} \text{dr}/c^2) [h(j)/(c^2 + (h(j)/2)^2)^{0.5}] * \right. \\
 & \left. * [\text{rt} - \text{rr} \cos((i - 0.5)\text{dth})] (\cos(\text{rt}/\text{rc}_1(x)) + \sin(\text{rt}/\text{rc}_1(x))) \right. \\
 & \left. - \sum_{j=0}^{j=N_r} \sum_{i=0}^{i=180} (G \rho r(j) \text{dth} \text{dr}) [\text{atan}(4 c h(j)/(4 c^2 - h(j)^2))/c] * \right. \\
 & \left. * [\text{rt} - \text{rr} \cos((i - 0.5)\text{dth})] (\cos(\text{rt}/\text{rc}_1(j)) - \sin(\text{rt}/\text{rc}_1(j)))/\text{rc}_1(j) \right\}
 \end{aligned}$$

where

$$\rho_{v1}(j) = (1/3) * (\text{vel})^4 / 4 / 3.14159265 / Mr_1(j) / G / \beta$$

$$\text{rc}_1(j) = (\text{vel})^2 * \sqrt{1 / (3 * 4 * 3.14159265 * \rho_{v1}(j) * G)}$$

$Mr_1(j)$ = the quantity of mass contained within the radius at which the velocity is being computed.

$$\beta = 1.22 * 1.22 * (1.989E+30) / (3.08568E+16)^2 \text{ pc}/\sqrt{\text{msuns}}$$

$$\text{vel} = 3 / 9.7778 \text{ pc}/\text{year}$$


Results

Nine figures with self-explanatory captions are presented for each galaxy, followed by a table summarizing integral features of the galaxy. Thus there are 45 figures and 5 tables in all. The galaxies, their sizes & pages with their results are as follows:


NGC 6822.....	4.8 kpc.....	pp 8 to 17
Large Magellanic Cloud (LMC).....	9 kpc.....	pp 28 to 37
Milky Way.....	17 kpc.....	pp 38 to 47
NGC 3198.....	30 kpc.....	pp 48 to 57
UGC 9133.....	102.5 kpc.....	pp 18 to 27

NGC 6822 (4.8 kpc)


- Newtonian Gravity
- Modified Newtonian Dynamics
- Vacuum Modified Gravity
- Negative Cosmological Constant



- Newtonian Gravity
- Modified Newtonian Dynamics
- Vacuum Modified Gravity
- Negative Cosmological Constant




Density of the ring normalized by average density of the galaxy




Radius of the centreline circle of the ring normalized by the galaxy radius

Surface mass density of the ring normalized
by the average SMD of the galaxy




Radius of the centreline circle of the
ring normalized by the galaxy radius

Product of the normalized SMD and
normalized Radius of the ring




Radius of the centreline circle of the
ring normalized by the galaxy radius

Mass of matter in the ring normalized
by the total galaxy mass




Radius of the centreline circle of the
ring normalized by the galaxy radius


Mass of the galaxy contained within the ring
normalized by the total galaxy mass



Radius of the centreline circle of the
ring normalized by the galaxy radius




Radius of the centreline circle of the ring normalized by the galaxy radius




NGC 6822 (table)

20 Rings 4800 pc	Total Volume (pc ³)	Total Mass (solar mass, msuns)	Average Density (msuns/pc ³)	Average Surface Mass Density (msuns/pc ²)	Keplerian Velocity at Galaxy Rim (Kms/sec)
Newtonian Dynamics	7.6436 e+009	1.0032 E+009	0.1312	13.8593	29.9796
Newtonian Dynamics with Negative Cosmological constant	7.6436 e+009	9.8781 E+008	0.1292	13.6472	29.7493
MoND	1.1309 e+010	3.2500 E+008	0.0287	4.4901	17.0641
Vacuum Modified Gravity	1.1309 e+010	9.0237 E+008	0.0798	12.4667	28.4335

UGC 9133 (102.5 kpc)




- Newtonian Gravity
- Modified Newtonian Dynamics
- Vacuum Modified Gravity
- Negative Cosmological Constant




Density of the ring normalized by average density of the galaxy

- Newtonian Gravity
- Modified Newtonian Dynamics
- Vacuum Modified Gravity
- Negative Cosmological Constant




Radius of the centreline circle of the ring normalized by the galaxy radius

Surface mass density of the ring normalized
by the average SMD of the galaxy




Radius of the centreline circle of the
ring normalized by the galaxy radius



- Newtonian Gravity
- Modified Newtonian Dynamics
- Vacuum Modified Gravity
- Negative Cosmological Constant


Mass of matter in the ring normalized by the total galaxy mass




Radius of the centreline circle of the ring normalized by the galaxy radius


Mass of the galaxy contained within the ring
normalized by the total galaxy mass

- Newtonian Gravity
- Modified Newtonian Dynamics
- Vacuum Modified Gravity
- Negative Cosmological Constant



Radius of the centreline circle of the
ring normalized by the galaxy radius







UGC 9133 (table)

20 Rings 102500 pc	Total Volume (pc ³)	Total Mass (solar mass, msuns)	Average Density (msuns/pc ³)	Average Surface Mass Density (msuns/pc ²)	Keplerian Velocity at Galaxy Rim (Kms/sec)
Newtonian Dynamics	7.4429 e+013	8.2719 e+011	0.0111	25.0616	186.2949
Newtonian Dynamics with Negative Cosmological constant	7.4429 e+013	7.9198 e+011	0.0106	23.9949	182.2871
MoND	7.4429 e+013	4.1810 e+011	0.0056	12.6673	132.4460
Vacuum Modified Gravity	7.4429 e+013	5.5632 e+011	0.0075	16.8548	152.7774


Large Magellanic Cloud (LMC) (9 kpc)



- Newtonian Gravity
- Modified Newtonian Dynamics
- Vacuum Modified Gravity
- Negative Cosmological Constant




Density of the ring normalized by average density of the galaxy




Radius of the centreline circle of the ring normalized by the galaxy radius

Surface mass density of the ring normalized
by the average SMD of the galaxy

- Newtonian Gravity
- Modified Newtonian Dynamics
- Vacuum Modified Gravity
- Negative Cosmological Constant




Radius of the centreline circle of the
ring normalized by the galaxy radius



Mass of matter in the ring normalized
by the total galaxy mass


- Newtonian Gravity
- Modified Newtonian Dynamics
- Vacuum Modified Gravity
- Negative Cosmological Constant




Radius of the centreline circle of the
ring normalized by the galaxy radius

Mass of the galaxy contained within the ring
normalized by the total galaxy mass


- Newtonian Gravity
- Modified Newtonian Dynamics
- Vacuum Modified Gravity
- Negative Cosmological Constant



Radius of the centreline circle of the
ring normalized by the galaxy radius




Radius of the centreline circle of the ring normalized by the galaxy radius





Large Magellanic Cloud (LMC) (table)

20 Rings 9000 pc	Total Volume (pc ³)	Total Mass (solar mass, msuns)	Average Density (msuns/pc ³)	Average Surface Mass Density (msuns/pc ²)	Keplerian Velocity at Galaxy Rim (Kms/sec)
Newtonian Dynamics	8.0439 e+010	5.3772 E+009	0.0668	21.1309	50.6891
Newtonian Dynamics with Negative Cosmological constant	8.0439 e+010	5.2885 E+009	0.0657	20.7824	50.2694
MoND	8.0439 e+010	1.9989 E+009	0.0248	7.8551	30.9052
Vacuum Modified Gravity	8.0439 e+010	3.5654 E+009	0.0443	14.0113	41.2758

Milky Way (17 kpc)









Radius of the centreline circle of the ring normalized by the galaxy radius

Surface mass density of the ring normalized
by the average SMD of the galaxy



Radius of the centreline circle of the
ring normalized by the galaxy radius


Product of the normalized SMD and
normalized Radius of the ring



Radius of the centreline circle of the
ring normalized by the galaxy radius


- Newtonian Gravity
- Modified Newtonian Dynamics
- Vacuum Modified Gravity
- Negative Cosmological Constant

Mass of matter in the ring normalized
by the total galaxy mass




Radius of the centreline circle of the
ring normalized by the galaxy radius


Mass of the galaxy contained within the ring
normalized by the total galaxy mass



Radius of the centreline circle of the
ring normalized by the galaxy radius




Radius of the centreline circle of the ring normalized by the galaxy radius




Milky Way (table)


20 Rings 17000 pc	Total Volume (pc ³)	Total Mass (solar mass, msuns)	Average Density (msuns/pc ³)	Average Surface Mass Density (msuns/pc ²)	Keplerian Velocity at Galaxy Rim (Kms/sec)
Newtonian Dynamics	5.4211 e+011	1.4955 E+011	0.2759	164.7222	194.5070
Newtonian Dynamics with Negative Cosmological constant	5.4211 e+011	1.4842 E+011	0.2738	163.4686	193.7654
MoND	5.4211 e+011	1.0694 E+011	0.1973	117.7889	164.4793
Vacuum Modified Gravity	5.4211 e+011	1.1782 E+011	0.2173	129.7735	172.6443

NGC 3198 (30 kpc)




- Newtonian Gravity
- Modified Newtonian Dynamics
- Vacuum Modified Gravity
- Negative Cosmological Constant






Radius of the centreline circle of the ring normalized by the galaxy radius


Surface mass density of the ring normalized
by the average SMD of the galaxy



Radius of the centreline circle of the
ring normalized by the galaxy radius




Mass of matter in the ring normalized
by the total galaxy mass




Radius of the centreline circle of the
ring normalized by the galaxy radius

Mass of the galaxy contained within the ring
normalized by the total galaxy mass


- Newtonian Gravity
- Modified Newtonian Dynamics
- Vacuum Modified Gravity
- Negative Cosmological Constant



Radius of the centreline circle of the
ring normalized by the galaxy radius



Radius of the centreline circle of the ring normalized by the galaxy radius



NGC 3198 (table)

20 Rings 30000 pc	Total Volume (pc ³)	Total Mass (solar mass, msuns)	Average Density (msuns/pc ³)	Average Surface Mass Density (msuns/pc ²)	Keplerian Velocity at Galaxy Rim (Kms/sec)
Newtonian Dynamics	2.9792 e+012	8.7450 E+010	0.0294	30.9290	111.9639
Newtonian Dynamics with Negative Cosmological constant	2.9792 e+012	8.5459 E+010	0.0287	30.2250	110.6824
MoND	2.9792 e+012	4.1890 E+010	0.0141	14.8154	77.4911
Vacuum Modified Gravity	2.9792 e+012	5.4961 e+010	0.0184	19.4384	88.7617

Conclusions

We can observe from the figures & tables, infer and conclude as follows:

1. As we move from smaller to larger galaxies, there is an increasing similarity between the mass and mass density profiles predicted as per different theories. For small galaxies, NGC6822 and LMC, there is a significant variation between what the different theories predict. As we move to Milky Way and NGC3198, we see much more similar looking curves and for UGC9133, the curves appear strikingly similar.
2. The well-studied luminosity profiles and mass-to-light ratios of galaxies suggest that there is an exponential decrease in the mass of visible matter as we move away from the centre of the galaxy. But, considering that ordinary matter in the outer reaches of a galaxy should be much cooler and may escape detection and the matter near the galactic centre should be very hot, we can probably explain why we see a very sharp fall in the mass and mass densities near the galactic centre and an almost constant profile as we move further outwards.
3. The mass and mass density profiles show an unexpected rising trend towards the outer edges of the galaxies and this trend is more prominent for smaller galaxies

and the extent of the rise decreases as we move towards larger galaxies and becomes almost unnoticeable for very large galaxies.

4. For the larger galaxies, all the existing and proposed theories of gravity show nearly identical predictions, which appear intuitively satisfying considering what we already know regarding structures of galaxies. But for smaller galaxies, the predictions from different theories appear to vary significantly and also the results appear to contradict our expectations.

5. For example, MoND seems to give predictions very close to what is expected for galaxies of every size that we have examined whereas Newtonian theory gives bizarre predictions for small galaxies, though as we move to larger galaxies the predictions are closer to our expectations. Similar is the situation with Vacuum Modified Gravity.

6. Hence, what we may conclude is that there is a need to study smaller galaxies extensively in order to arrive at a conclusion regarding whether at all we need dark matter and whether MoND can be considered as a valid theory of gravity.

7. From the trend that is observed it appears that either MoND is correct or else there is a need to invoke dark matter along within Newtonian theory in order to explain the observations. And this dark matter, if there, seems to be concentrated near the outer edges of the galaxies with its gravitational effect on visible matter increasing as we move towards the galactic rim. And also the effect of dark matter compared to ordinary matter is more pronounced for smaller galaxies than the larger ones and is almost negligible for the very large ones.

REFERENCES

1. astro-ph/0006140

Disk-Galaxy Density Distribution from Orbital Speeds using Newton's Law

Kenneth F. Nicholson

2. astro-ph/0006330 23 Jun 2000

Disk-Galaxy Density Distribution from Orbital Speeds using Newton's Law

Version 1.1

Kenneth F. Nicholson

3. astro-ph/0101401 24 Jan 2001

Galaxy Mass Distributions for Some Extreme Orbital-Speed Profiles

Kenneth F. Nicholson

- 4.** astro-ph/0309762 v2, revised 1 Mar 2007
Galactic mass distribution without dark matter or modified Newtonian mechanics
Kenneth F Nicholson
- 5.** astro-ph/0506370v4
Galaxy Rotation Curves Without Non-Baryonic Dark Matter
J. R. Brownstein and J. W. Moffat
- 6.** astro-ph/0806.1131
Newtonian mechanics & gravity fully model disk galaxy rotation curves without dark matter
Dilip G. Banhatti
[For IAU Symposium 254 on Galaxy Disk in Cosmological Context, 9-14 June 2008, Copenhagen, Denmark]
- 7.** astro-ph/0703430v7 [25 Apr 2008 *Current Science* **94** (8) 960+986-95]
Disk galaxy rotation curves and dark matter distribution
Dilip G. Banhatti
- 8.** astro-ph/0712.1110v1
Vacuum Modified Gravity as an explanation for flat galaxy rotation curves
R. Van Nieuwenhove
- 9.** RHCPP99-20T astro-ph/9911485 November 1999
A possible explanation of Galactic Velocity Rotation Curves in terms of a Cosmological Constant
Steven B Whitehouse and George V. Kraniotis