

THE GALACTIC POTENTIAL AND THE ASYMMETRIC DISTRIBUTION OF HYPERVELOCITY STARS

HAGAI B. PERETS¹, XUFEN WU², HONGSHENG ZHAO², BENOIT FAMAËY³, GIANFRANCO GENTILE⁴ AND TAL ALEXANDER^{1,5}

¹ Weizmann Institute of Science, POB 26, Rehovot 76100, Israel

² Scottish University Physics Alliances, University of St Andrews, KY16 9SS, U.K.

³ Institut d'Astronomie et d'Astrophysique, Université Libre de Bruxelles, Boulevard du Triomphe, B-1050 Bruxelles, Belgium

⁴ Sterrenkundig Observatorium, Ghent University, Krijgslaan 281, S9, B-9000 Ghent, Belgium

⁵ The William Z. and Eda Bess Novick career development chair

Draft version December 29, 2018

Abstract

In recent years several hypervelocity stars (HVSs) have been observed in the halo of our Galaxy. Such HVSs have possibly been ejected from the Galactic center and then propagated in the Galactic potential up to their current position. The recent survey for candidate HVSs show an asymmetry in the kinematics of candidate HVSs (position and velocity vectors), where more outgoing stars than ingoing stars (i.e. positive Galactocentric velocities vs. negative ones) are observed. We show that such kinematic asymmetry, which is likely due to the finite lifetime of the stars and Galactic potential structure, could be used in a novel method to probe and constrain the Galactic potential, identify the stellar type of the stars in the survey and estimate the number of HVSs. Kinematics-independent identification of the stellar types of the stars in such surveys (e.g. spectroscopic identification) could further improve these results. We find that the observed asymmetry between ingoing and outgoing stars favors specific Galactic potential models. It also implies a lower limit of $\sim 54 \pm 8$ main sequence HVSs in the survey sample ($\gtrsim 648 \pm 96$ in the Galaxy), assuming that all of the main sequence stars in the survey originate from the Galactic center. The other stars in the survey are likely to be hot blue horizontal branch stars born in the halo rather than stars ejected from the Galactic center.

Subject headings: black hole physics — galaxies: nuclei — stars: kinematics

1. INTRODUCTION

Hypervelocity stars (HVSs) are stars with extremely high peculiar velocities relative to the velocity distribution of their parent population. In recent years several HVSs have been observed in the Galactic halo, some of them unbound to the Galaxy (with velocities beyond the escape velocity; Brown et al. 2007b). From these a Galactic population of 96 ± 10 such HVSs was inferred (Brown et al. 2007b), up to the 100 kpc distance limit of the survey. Many similar bound HVSs (with velocities lower than the escape velocity) have been observed at larger numbers. Most of the observed HVSs are B-type stars (Brown et al. 2005, 2006a,b, 2007a,b; Edelmann et al. 2005; future observations of HVSs of other stellar types are discussed in Kollmeier & Gould 2007; Brown et al. 2008; Kenyon et al. 2008). Given the color selection of the targeted survey for these stars (Brown et al. 2006a), such stars could be either main sequence (MS; or blue straggler) B stars or hot blue horizontal branch (BHB) stars. Currently only three of the stars in the survey have specific unambiguous identification and were found to be MS stars (Fuentes et al. 2006; Lopez-Morales & Bonanos 2008; Przybilla et al. 2008a).

Extreme velocities as found for these stars most likely suggest a dynamical origin from an interaction with or close to the massive black hole (MBH) in the Galactic center (GC; Hills 1988; Yu & Tremaine 2003; Levin 2006; O'Leary & Loeb 2007; Perets et al. 2007). In the following we discuss only HVSs ejected from the GC¹ and observed in

the Galactic halo, > 10 kpc from the GC, which would require ejection velocities from the GC of $\gtrsim 800 \text{ km s}^{-1}$. Such HVSs could serve as probes of the GC environment, stellar population and dynamics (see e.g. Perets 2007; Sesana et al. 2007a; Kenyon et al. 2008) and serve as an independent evidence for the existence of a MBH in the Galactic center (Hills 1988). Most of the B-type stars observed through the HVSs survey have lower velocities and are either bound HVSs (Brown et al. 2007a) or are just halo stars. The survey shows an asymmetry in the kinematics of the stars, where more stars have positive radial velocities in Galactocentric coordinates (i.e. outgoing stars) than negative ones (ingoing or returning stars). As we show this asymmetry is dependent on both the absolute velocity and the distance of the stars from the GC, and could be used in a novel method to probe and constrain the Galactic potential, identify the stellar type of the stars in the survey and estimate the number of HVSs.

This paper is organized as follows. We first briefly describe the HVSs survey (§2), and then suggest a novel method to probe the Galactic potential using such surveys (§3). In §4 we use a similar analysis to infer the statistics of the stellar type of stars in the HVSs survey and estimate a lower limit to the number of HVSs in the Galaxy.

2. THE VELOCITY-DISTANCE DISTRIBUTION OF HVSS

HVSs of almost any stellar type could theoretically be observed, since the currently suggested scenarios for the origin of HVSs give rise to only limited number of constraints on their stellar characteristics (e.g. Hansen 2007; Perets 2007). However, given their relatively small numbers in the Galaxy, it is practically impossible to find HVSs close by. For this reason, following the discovery of the first HVSs in the Galactic halo (Brown et al. 2005; Hirsch et al. 2005; Edelmann et al.

high velocity stars ejected from the Galactic disk.

Electronic address: hagai.perets@weizmann.ac.il

¹ Note that recent observations possibly suggest a different origin for two of HVSs observed serendipitously (Bonanos et al. 2008; Przybilla et al. 2008b; Heber et al. 2008, but see Perets 2008). Throughout this paper we assume that most (if not all) of the MS stars in the HVSs survey sample have a GC origin. We note however that the methods we describe here to constrain the Galactic potential could similarly be used, in principle, for the analysis of

2005), Brown et al. (2006b) have issued a survey of HVSs extending to large distances. They have searched for HVSs among color selected B-type halo stars of limited magnitude (a more recent survey also searches for A type stars; Brown et al. 2008; not included in our analysis). Such stars are luminous enough to be observed at large distances in the Galactic halo where the relative frequency of HVSs is much higher, and are less likely to be contaminated by the disk and halo stellar population. These stars could be either MS B stars (or blue stragglers) with masses of $3 - 4 M_{\odot}$ and short lifetimes ($1 - 4 \times 10^8$ yrs) or hot blue horizontal branch (BHB) stars also with short lifetimes (few $\times 10^8$ yrs on the horizontal branch), but long progenitor lifetime. The absolute magnitudes of observed stars and hence their inferred distances depend on their stellar type. Any analysis of the distribution of HVSs should take both possibilities into account.

Fig. 2 in Brown et al. (2007b) shows a definite asymmetry in the distribution of ingoing and outgoing HVSs, where the velocities of ingoing stars do not extend beyond 300 km s^{-1} . Given that the escape velocity at these distances is much higher, one would expect to see bound HVSs returning at velocities up to the escape velocity, in striking contrast with observations. In order to study this behavior, we turn to the velocity-distance distribution of these stars. In fig. 1 we show the radial velocity-distance distribution (relative to the GC) for all of the observed B-type stars in the Brown et al. (2007b) survey, assuming they are either MS stars (and are therefore more luminous and more distant; fig. 1a), or hot BHB stars (and therefore closer; fig. 1b).

3. PROBING THE GALACTIC POTENTIAL USING HYPERVELOCITY STARS

Many studies have been done to constrain the Galactic potential at large distances through observations (see e.g. Fich & Tremaine 1991; Battaglia et al. 2005; Smith et al. 2007; Xue et al. 2008, and references therein). Some of these studies use the velocity dispersion of observed objects to constrain the Galactic potential (e.g. Battaglia et al. 2005; Xue et al. 2008), however these suffer from uncertainties regarding the velocity anisotropy and the behavior of the stellar halo density at very large distances, and require some apriori assumptions regarding these parameters, which may lead to large uncertainties (see, for example, discussion in Dehnen et al. 2006). In additions many objects are needed in order to obtain the velocity dispersion at a given distance from the GC. Other studies explore the local escape velocity from the Galaxy through observations of high velocity stars (Smith et al. 2007). However, such analysis contains degeneracies and depends on the unknown structure of the tail of the velocity distribution of the high velocity stars. Consequently specific assumptions must be taken for the velocity distribution, which could be strongly affected by the small number statistics of the observed highest velocity stars in the distribution tail. Moreover, the assumptions used for the stellar velocities depend on their being extended up to the escape velocity from the Galaxy. Although large surveys may help solve this problem, very high velocity stars are quite rare in the Galaxy, and would be difficult to find especially in surveys limited to relatively close environment of the solar neighborhood.

Gnedin et al. (2005) and Yu & Madau (2007) suggested to use the kinematics of HVSs in order to probe the Galactic potential using the position and velocity vectors of HVSs at large Galactocentric distances. Under the assumption that HVSs were ejected from the GC they suggest to measure the slight

departure from purely radial orbits of these HVSs, due to the (possible) triaxiality of the Galactic potential. These methods require the accurate distance and 3D velocity of HVSs, and focus on the triaxiality of the Galactic potential, which is important in the context of hierarchical, cold dark matter (CDM) models of structure formation (e.g. Hayashi et al. 2007), although we note this should be interesting also in respect to modified Newtonian dynamics (MOND) theories (Milgrom 1983). Recently, Kenyon et al. (2008) have studied the propagation of HVSs in the Galactic potential and showed that it could depend strongly on the Galactic potential at the central regions of the Galaxy (200pc). They also showed the dependence of the HVSs radial distributions on the stellar type, and their observational implications.

In the following we suggest a method which is more general in nature and more useful in constraining and distinguishing between different Galactic mass distributions which are required by the different Galactic potential models at large distances (although it could also be relevant to probing the triaxiality of the Galactic potential). This can also be used in order to discriminate between different CDM Galactic potential models and/or between Galactic potentials in MOND theories. This method makes use of the asymmetry in velocity distributions of ingoing and outgoing HVSs (see also related discussion in Kenyon et al. 2008). We begin with a naive description of the method, assuming one could observe even the oldest bound HVSs (i.e. those that could not leave the Galaxy, and could have gone through the Galaxy a few times). We then continue with a more realistic treatment which takes into account the finite lifetime of stars observable in the halo, given the limited observational capabilities.

3.1. Long lived, observable hypervelocity stars

Let us assume that HVSs have been continuously ejected from the GC with some distribution of velocities, which would produce both bound and unbound HVSs. Unbound stars eventually leave the galaxy. Bound stars reach the apocenter point of their orbit and then return back to the GC with negative radial velocity (in Galactocentric coordinates). The Galactocentric distance-velocity distribution of ingoing stars would then have a cut-off, which would correspond to the escape velocity of these stars at a given distance from the GC. Such a cut off is distance dependent and thus more distant HVSs would have lower absolute velocities. At the same time we should see that the distribution of outgoing HVSs extends to much higher absolute velocities, since this population includes the unbound stars, on their way out of the galaxy. Consequently, a clear asymmetry should be observed between the distribution of ingoing and outgoing stars. This asymmetry or cut-off in the distance-velocity distribution would map the escape velocity of stars from the galaxy at any given distance where HVSs are observed, and serve as a direct probe of the galactic potential. Note, however, that very different galactic potential may have escape velocities which are quite similar at a wide range of distances from the GC (Wu et al. 2008). In such cases observations of more distant HVSs (i.e. wider distance range) may be required to distinguish between such potentials. Since such stars would be fainter, this would be more difficult observationally.

The use of ingoing and outgoing halo HVSs has three advantages over methods used to probe the local escape velocity from the Galaxy. First, there is a clear natural separation between bound and unbound stars, and the latter can not contaminate the sample of ingoing high velocity bound stars which

are used to calculate the escape velocity. Second, there is no required assumption regarding the structure of the velocity distribution and its tail. Third, the HVSs are observed over a large distance range, and could thus map the Galactic potential in this full range, and not only at the local scale as have been done with high velocity stars in surveys such as the Rave survey (Smith et al. 2007).

3.2. Realistic short lived hypervelocity stars

In reality, observable stars in the HVSs targeted survey may not have an unlimited propagation time, as was our naive assumption. This may result either because of their short lifetimes (after which they evolve to a different stellar type, which can not be observed at such distances with current instruments) or due to their possible origin from a burst like event, which ejected HVSs only over a limited short time, and not as a continuous process occurring over the lifetime of the Galaxy (see Perets 2007)². Nevertheless, the general method prescribed above could still be applicable, with some modifications. In fact, as we the limited propagation time of HVSs may prove to be more advantageous in some respects).

Assuming some finite propagation time for HVSs, we would still expect an asymmetry in the ingoing and outgoing HVSs distance-velocity distributions. However, in this case the cutoff in the ingoing HVSs distribution would be at much lower velocities than the escape velocity. This cutoff corresponds to the maximal return velocity of HVSs which could still be observed coming back during their short propagation time. Assuming a maximal propagation time for the HVSs (its lifetime), this cutoff could thus be used as a probe of the Galactic potential in the same way as the naive method outlined above. Moreover, the short propagation time of stars can be advantageous for our purposes. Stars of different stellar types have different maximal lifetimes and would produce different distance-velocity cutoffs. Consequently these different populations can supply us with several independent probes of the Galactic potential, that, combined together, would further assist in constraining the Galactic potential. a higher velocity at a given distance implies a longer travel time (as the travel time depends on the potential out to the apo-apse of the orbit), ingoing HVSs provide information not only on the escape velocity at the point where they are observed today, but even further away. Furthermore, up to the distance-velocity cutoff the distribution of outgoing and ingoing stars (with lower velocities than the cutoff velocity at their position) could be compared^{2,3}, in terms of the number of stars; the two dimensional distance-(absolute) velocity distribution and the stellar types (if known) or color-color distributions of the samples of outgoing and ingoing stars. Any difference between these distributions is due to the further propagation in the Galactic potential of the ingoing HVSs to the apocenter of their orbit and back. Different Galactic potential models give different return times, that are also highly sensitive to the velocity of the HVSs. Consequently, the statistical correlation between the ingoing and outgoing distance-velocity distributions could serve as a quantitatively sensitive method for discriminating between models for the Galactic potential than just the popu-

lation of the highest velocity ingoing HVSs.

3.3. Discriminating between Galactic potential models

Currently, only three HVSs in the HVSs survey have an unambiguous stellar type identification, and the analysis suggested here for probing the Galactic potential can not be used directly. The clear cut-off in the ingoing stars distribution would not be observed, as it would be smeared by the existence of hot BHB stars contaminating the sample. Nevertheless we still expect a statistical asymmetry between the number of ingoing vs. outgoing stars in the sample, which should be observable beyond the theoretical cut-off. As an illustrative example we show the critical asymmetry lines for propagation of MS B stars HVSs up to a maximal lifetime of 4×10^8 . In fig. 1a we show the critical asymmetry lines for such time limited propagation in different models for the Galactic potential. The different models we use include five dark matter (CDM) potentials and one MOND potential. Beside the Paczynski (1990) model (hereafter PAC) all models are described in detail in Wu et al. (2008), where the same reference names are used (KZS, BSC, RAVE1-3 and MOND). The PAC, KZS, BSC and the MOND models are almost indistinguishable in this range of distances, where as the RAVE 1-2 (indistinguishable in this range) and RAVE 3 models show very different behavior. We look for asymmetry by counting the numbers of outgoing vs. ingoing stars in the sample, for the different potentials, i.e. counting the number of stars above and below the positive and negative velocity lines of the critical asymmetry curves shown in fig. 1a. We look for the best fit model, which should show the largest asymmetry. We find a total of 166 outgoing stars vs. 112 ingoing stars for the PAC, KZS, BSC and MOND models (the probability for getting such an asymmetric distribution from an a priori symmetric distribution is $p = 10^{-3}$) and 378 (452) outgoing stars vs. 319 (388) ingoing stars for the RAVE 3 (RAVE 1-2) models ($p = 0.025$ (0.027)). Although all models show asymmetric distributions, the stars counted for the RAVE models contain the contribution from the PAC/KZS/BSC/MOND models. When subtracting this contribution we find that the RAVE models do not show any additional asymmetry [212 (286) outgoing vs. 207 (276) ingoing stars; $p = 0.8$ (0.67)]. Therefore the HVSs sample favors the PAC, KZS, BSC and MOND models over the RAVE models.

As we have shown even the contaminated sample of HVSs could already constrain Galactic potential models. Future identification of the stellar types of the stars in the survey which could purify it could give even stronger constraints on these and other Galactic potential models. We also note that stars with longer MS lifetimes could probe larger distance range during their propagation and still return during their MS lifetimes to be observed as ingoing HVS⁴. Such stars could therefore serve as observable probes of the Galactic potential at even larger distances not accessible in any other ways (and could possibly discriminate between CDM and MOND models that differ only at these distance ranges). This would require the identification of later-type HVSs among halo stars (e.g. Brown et al. 2008, not analyzed here).

4. PROPAGATION IN THE GALACTIC POTENTIAL AND THE LIFETIMES OF OBSERVED HYPERVELOCITY STARS

² In such a case, for example, the highest velocity bound HVSs may never be observed as ingoing HVSs, since a longer propagation time is required for them to reach the apo-apse point of their orbit and become ingoing HVSs.

³ This, however, is true only under the assumption of a continuous and constant ejection rate, which is not the case for burst ejection of HVSs by an inspiralling IMBH (Levin 2006; Baumgardt et al. 2006; Lückmann & Baumgardt 2007; Sesana et al. 2007b)

⁴ In fact rejuvenated blue straggler stars could therefore contaminate the sample of the HVSs, however this population is not expected to be large (Perets 2008).




FIG. 1.— The distance-velocity distribution of Halo B-type stars from Brown et al. (2007b) observations (blue circles). (a) The observed stars are assumed to be main sequence stars. The lines represent the critical asymmetry lines (see text) for various potentials (see legend), for propagation up to the maximal lifetime of such stars ($\sim 4 \times 10^8$ yrs). (b) The same, but now assuming all the stars are hot BHB stars, and therefore less distant. The velocity-distance distribution of regular halo BHB stars (\times marks; taken from Xue et al. 2008), is shown for comparison. Dashed middle line in both panels separates between outgoing and ingoing stars. The escape velocity (in the KZS model) is also shown for comparison in both panels.

The method described above uses the finite lifetimes of stars and their kinematics to constrain the Galactic potential. A very similar approach could also be used to constrain the number of the HVSSs and their lifetimes, given a specific Galactic potential. Making use of this approach we show that most of the stars in the HVSSs survey, especially with ingoing velocities are likely to be halo hot BHB stars and have not been ejected from the GC (see also Kollmeier & Gould 2007; Yu & Madau 2007; Brown et al. 2007b; Kenyon et al. 2008, for related discussions), but nevertheless the number of HVSSs ejected from the GC could be much higher than previously thought.

In the previous section we described the critical distance-velocity asymmetry lines. For a given Galactic potential and a given propagation time of a HVS, one could find the critical line outside which no such ingoing stars should be observed. In other words any ingoing star beyond this line can not be

a HVS from the GC with such (or shorter) lifetime. We can therefore identify at least some of the HVSSs sample stars as stars that are not MS stars ejected from the GC using this criteria (see Svensson et al. 2007 for a related discussion). Moreover, since the distribution of such stars should be symmetric (as observed for other samples of halo objects, such as regular halo BHB stars) any asymmetry beyond the critical lines is due to the outgoing MS HVSSs from the GC (or from the Galactic disk¹, a possibility which we do not discuss here).

Assuming the PAC model (or the KZS and BSC model that give similar results) and a maximal propagation time of 4×10^8 yrs (lifetime of a $3 M_\odot$ MS B star) we find an overabundance of $166 - 112 = 54$ outgoing stars, where an asymmetry of 8 stars correspond to the 1σ probability level. We therefore give a lower limit estimate for the number of HVSSs beyond the critical line of $\gtrsim 54 \pm 8$ HVSSs, from which we infer, following the calculations by Brown et al. (2007b), that a total number of $\gtrsim 648 \pm 96$ stars (main sequence B stars of $3-4 M_\odot$ at distances of $10 \text{ kpc} \lesssim r \lesssim 100 \text{ kpc}$ ejected from the Galactic center) exist in the Galaxy. We expect that most if not all of the ingoing stars beyond the critical line are hot BHB stars, where outgoing stars beyond the line could be both hot BHB stars or MS stars, with a ratio of 2 to 1 ($166 - 54 = 112$ vs. 54). These large estimated number of HVSSs may suggest a different contamination from high velocity stars ejected from the Galactic disc, that could also produce an asymmetric distribution due to finite MS lifetimes of the stars. Here we do not address this possibility, which requires a more detailed study and would be addressed in a another paper (Perets et al., in preparation).

In this calculation we assumed a specific maximal propagation time for the GC HVSSs. Instead, we can look at the asymmetric distributions for stars that propagated in the Galactic potential for shorter propagation times. Such stars could be either more massive MS B stars with shorter MS lifetimes, or MS stars ejected from the GC only after evolving for some time in the GC (or ejected more recently from the GC). Since the frequency of more massive stars is small, the latter possibilities are more likely to apply. One would therefore expect to see an asymmetric distribution even at times shorter than the MS lifetime of $3 M_\odot$ B MS star (4×10^8 yrs), likely of the order of half this time ($\sim 2 \times 10^8$ yrs), assuming a continuous ejection rate of HVSSs. In fig. 2 we show the number of outgoing vs. ingoing stars for a range of propagation times. For the PAC potential model we used, the distribution begins to become asymmetric at propagation times of 2.5×10^8 yrs, in good agreement with our expectations.

For completeness we discuss the possibility that the HVSSs are not MS stars but hot BHB stars ejected from the GC. In this case their absolute magnitude is different from that of MS stars, and the inferred distances change accordingly (see fig. 1b). Again we can look for the typical propagation time at which we see an asymmetric distribution of these stars assuming they were ejected from the GC. We find this to be at a few 10^8 yrs. This time is comparable, but longer, than the lifetime of hot BHB stars at this phase. It is also much shorter than the lifetime of hot BHB progenitors, which could extend up to a few Gyrs. Both of these inconsistencies, together with the dynamical constraints against the ejection of hot BHB HVSSs (Perets 2007) suggest a different identification of these stars. Furthermore, comparing their velocity-distance distribution to that of regular BHB stars in the halo (Xue et al. 2008) show very similar distribution (see fig. 1b), i.e. the observed cut-off in the distribution is not related to the critical lines we de-




FIG. 2.— The asymmetric distribution of outgoing vs. ingoing stars. The curves show the integrated number of outgoing vs. ingoing stars below (or above, respectively) the critical asymmetry lines (PAC model), corresponding to the range of propagation times in the Galactic potential. The asymmetry begins at propagation times of $\sim 2.5 \times 10^8$ yrs (see left dotted vertical line). No ingoing stars are observed to correspond to propagation times beyond $\sim 1.3 \times 10^9$ yrs.

scribed, but due to the limit of the symmetric distribution of halo objects.

Another possibility is that asymmetries in the HVSs candidates sample are not due to the population of HVSs, but due to

stellar streams located at specific regions of the Galactic halo, which could have correlated velocities, and not an isotropic distribution as we assumed. To check this we repeated the asymmetry calculations described above, but this time for several different and distinct regions in the Galactic halo (different Galactic longitudes). Although some differences in the strength of the asymmetry are observed, all regions showed a clear asymmetry bias towards outgoing stars.

5. SUMMARY

In this paper we studied the characteristics, the origins and the use of stars observed in the survey for HVSs in the Galactic halo. The kinematics of currently observed HVSs ejected from the Galactic center depend strongly on their lifetimes and their propagation in the Galactic potential. We suggest a novel method to probe the Galactic potential up to large distances using the kinematics and the spectral identification of HVSs. We also use a reverse method, where a specific Galactic potential model is assumed, to give lower limit estimates on the number of HVSs ejected from the Galactic center. Future observations of HVSs in M31 (Sherwin et al. 2007) and in other Galaxies (Perets et al., in preparation) could have a similar use for studies of Galactic structures and potentials.

REFERENCES

- Battaglia, G. et al. 2005, *MNRAS*, 364, 433
 Baumgardt, H., Gualandris, A., & Portegies Zwart, S. 2006, *MNRAS*, 372, 174
 Bonanos, A. Z. et al. 2008, *ApJ*, 675, L77
 Brown, W. R., Geller, M. J., & Kenyon, S. J. 2008, *ArXiv e-prints*, 808
 Brown, W. R. et al. 2005, *ApJ*, 622, L33
 —. 2006a, *ApJ*, 640, L35
 —. 2006b, *ApJ*, 647, 303
 —. 2007a, *ApJ*, 660, 311
 —. 2007b, *ApJ*, 671, 1708
 Dehnen, W., McLaughlin, D. E., & Sachania, J. 2006, *MNRAS*, 369, 1688
 Edelmann, H. et al. 2005, *ApJ*, 634, L181
 Fich, M. & Tremaine, S. 1991, *ARA&A*, 29, 409
 Fuentes, C. I. et al. 2006, *ApJ*, 636, L37
 Gnedin, O. Y. et al. 2005, *ApJ*, 634, 344
 Hansen, B. 2007, *arXiv:0711.1164*
 Hayashi, E., Navarro, J. F., & Springel, V. 2007, *MNRAS*, 377, 50
 Heber, U. et al. 2008, *A&A*, 483, L21
 Hills, J. G. 1988, *Nature*, 331, 687
 Hirsch, H. A. et al. 2005, *A&A*, 444, L61
 Kenyon, S. J. et al. 2008, *ApJ*, 680, 312
 Kollmeier, J. A. & Gould, A. 2007, *ApJ*, 664, 343
 Levin, Y. 2006, *ApJ*, 653, 1203
 Löckmann, U. & Baumgardt, H. 2007, *arXiv:0711.1326*
 Lopez-Morales, M. & Bonanos, A. Z. 2008, *ArXiv: 0802.2945*
 Milgrom, M. 1983, *ApJ*, 270, 365
 O’Leary, R. M. & Loeb, A. 2007, *MNRAS*, 1076
 Paczynski, B. 1990, *ApJ*, 348, 485
 Perets, H. B. 2007, *ArXiv: 0712.1888*
 —. 2008, *ArXiv: 0802.1004*
 Perets, H. B., Hopman, C., & Alexander, T. 2007, *ApJ*, 656, 709
 Przybilla, N., Nieva, M. F., Tillich, A., Heber, U., Butler, K., & Brown, W. R. 2008a, *A&A*, 488, L51
 Przybilla, N. et al. 2008b, *A&A*, 480, L37
 Sesana, A., Haardt, F., & Madau, P. 2007a, *MNRAS*, 379, L45
 —. 2007b, *arXiv:0710.4301*
 Sherwin, B. D., Loeb, A., & O’Leary, R. M. 2007, *ArXiv: 0709.1156*
 Smith, M. C. et al. 2007, *MNRAS*, 379, 755
 Svensson, K. M., Church, R. P., & Davies, M. B. 2007, *arXiv:0709.3748*
 Wu, X. et al. 2008, *MNRAS*, 386, 2199
 Xue, X. et al. 2008, *ArXiv: 0801.1232*
 Yu, Q. & Madau, P. 2007, *MNRAS*, 379, 1293
 Yu, Q. & Tremaine, S. 2003, *ApJ*, 599, 1129

We would like to thank Warren Brown for helpful discussions and for supplying us with the data on the kinematics of the stars in the HVSs sample. HBP would also like to thank the Israeli Commercial & Industrial Club for their support through the Ramon scholarship. BF is a postdoctoral researcher of the FNRS (Belgium). GG is a postdoctoral fellow with the National Science Fund (FWO-Vlaanderen). TA is supported by ISF grant 928/06 and ERC Starting Grant 202996 and a new Faculty grant by sir H. Djangoly, CBE, of London UK.