COHOMOLOGICAL NON-RIGIDITY OF GENERALIZED REAL BOTT MANIFOLDS OF HEIGHT 2

MIKIYA MASUDA

Abstract. We investigate when two generalized real Bott manifolds of height 2 have isomorphic cohomology rings with $\mathbb{Z}/2$ coefficients and also when they are diffeomorphic. It turns out that cohomology rings with $\mathbb{Z}/2$ coefficients do not distinguish those manifolds up to diffeomorphism in general. This gives a counterexample to the cohomological rigidity problem for real toric manifolds posed in [\[5\]](#page-6-0). We also prove that generalized real Bott manifolds of height 2 are diffeomorphic if they are homotopy equivalent.

1. INTRODUCTION

A toric manifold is a compact smooth toric variety and a real toric manifold is the set of real points of a toric manifold. In [\[7\]](#page-6-1) we asked whether toric manifolds are diffeomorphic if their cohomology rings with $\mathbb Z$ coefficients are isomorphic as graded rings, which is now called cohomological rigidity problem for toric manifolds. No counterexample and some partial affirmative solutions are known to the problem (see [\[3\]](#page-6-2), [\[7\]](#page-6-1)). If X is a toric manifold and $X(\mathbb{R})$ is the real toric manifold associated to X, then $H^*(X(\mathbb{R}); \mathbb{Z}/2)$ is isomorphic to $H^{2*}(X;\mathbb{Z})\otimes \mathbb{Z}/2$ as graded rings. Motivated by this, we posed in [\[5\]](#page-6-0) the following analogous problem.

Cohomological rigidity problem for real toric manifolds. Are two real toric manifolds are diffeomorphic if their cohomology rings with $\mathbb{Z}/2$ coefficients are isomorphic as graded rings?

We say that *cohomological rigidity over* $\mathbb{Z}/2$ holds for a family of closed smooth manifolds if the manifolds in the family are distinguished up to diffeomorphism by their cohomology rings with $\mathbb{Z}/2$ coefficients. A real Bott manifold is the total space of an iterated $\mathbb{R}P¹$ bundles where each $\mathbb{R}P¹$ bundle is the projectivization of a Whitney sum of two real line bundles. A real Bott manifold is not only a real toric manifold but also a compact flat riemannian manifold. We proved in [\[5\]](#page-6-0) (and [\[6\]](#page-6-3)) that cohomological rigidity over $\mathbb{Z}/2$ holds for the family of real Bott manifolds.

Date: October 24, 2018.

²⁰⁰⁰ Mathematics Subject Classification. Primary 57R91; Secondary 14M25.

Key words and phrases. real toric manifold, cohomological rigidity, generalized real Bott manifold of height 2.

The author was partially supported by Grant-in-Aid for Scientific Research 19204007.

2 M. MASUDA

In this paper we consider real toric manifolds obtained as the total spaces of projectivization of Whitney sums of real line bundles over a real projective space. We call such a real toric manifold a generalized real Bott manifold of height 2. In this paper we will investigate when those two manifolds have isomorphic cohomology rings with $\mathbb{Z}/2$ coefficients and also when they are diffeomorphic. As a result, we will see that cohomological rigidity over $\mathbb{Z}/2$ fails to hold for some family of generalized real Bott manifolds of height 2, which gives a negative answer to the cohomological rigidity problem for real toric manifolds above. We also prove that generalized real Bott manifolds of height 2 are diffeomorphic if they are homotopy equivalent.

The author thanks Y. Nishimura for pointing out a mistake in an earlier version of this paper and T. Panov for his comments.

2. Cohomological condition

Let a, b be positive integers and we fix them. Let γ be the tautological line bundle over $\mathbb{R}P^{a-1}$ and let 1 denote a trivial real line bundle over an appropriate space. For a real vector bundle E , we denote by $P(E)$ the total space of the projectivization of E. For an integer q such that $0 \le q \le b$, we set

$$
M(q) := P(q\gamma \oplus (b-q)\mathbf{1}).
$$

Note that

(2.1)
$$
M(q)
$$
 is diffeomorphic to $M(b-q)$

because $P(E \otimes L)$ and $P(E)$ are diffeomorphic for any smooth vector bundle E and line bundle L over a smooth manifold.

A simple computation shows that

(2.2)
$$
H^*(M(q); \mathbb{Z}/2) = \mathbb{Z}/2[x, y]/(x^a, (x+y)^q y^{b-q})
$$

where x is the pullback of the first Stiefel-Whitney class of γ to $M(q)$ and y is the first Stiefel-Whitney class of the tautological line bundle over $M(q)$. One easily sees that a set $\{x^i y^j \mid 0 \le i < a, 0 \le j < b\}$ is an additive basis of $H^*(M(q); \mathbb{Z}/2)$.

Lemma 2.1. If $0 < q < b$, then both y^a and $(x + y)^a$ are non-zero.

Proof. Suppose $y^a = 0$. Then it follows from [\(2.2\)](#page-1-0) that there are constants $c, d \in \mathbb{Z}/2$ and a homogeneous polynomial $f(x, y)$ in x, y over $\mathbb{Z}/2$ such that

$$
y^{a} = \begin{cases} cx^{a} & \text{if } a < b \\ dx^{a} + f(x, y)(x + y)^{q}y^{b-q} & \text{if } a \geq b \end{cases}
$$

as polynomials in x, y . Clearly the former does not occur and the latter also does not occur because $q > 0$ by assumption. This is a contradiction, so $y^a \neq 0$.

If we set $X = x$ and $Y = x + y$, then $x + y = Y$ and $y = X + Y$, so that the role of x and $x + y$ will be interchanged. Since $b - q > 0$ by assumption, the above argument applied to Y instead of y proves that $(x+y)^a \neq 0$. \Box

Definition. $h(a) := \min\{n \in \mathbb{N} \cup \{0\} \mid 2^n \geq a\}.$

For example,

$$
h(1) = 0, h(2) = 1, h(3) = h(4) = 2, h(5) = h(6) = h(7) = h(8) = 3,
$$

$$
h(9) = \cdots = h(16) = 4, \dots.
$$

Lemma 2.2. Let q and q' be non-negative integers. Then $\binom{q'}{q}$ $\binom{q'}{i} \equiv \binom{q}{i}$ $\binom{q}{i}$ (mod 2) for $0 \leq \forall i < a$ if and only if $q' \equiv q \pmod{2^{h(a)}}$, where $\binom{n}{m}$ $\binom{n}{m}$ is understood to be 0 when $n < m$ as usual.

Proof. When $q' = q$, the lemma is trivial. We may assume that $q' > q$ without loss of generality. We note that the former congruence relations in the lemma are equivalent to the following congruence relation of polynomials in t with $\mathbb{Z}/2$ coefficients:

(2.3)
$$
(1+t)^{q'-q} \equiv 1 \pmod{t^a}.
$$

We shall prove the "if" part first. Suppose $q' \equiv q \pmod{2^{h(a)}}$. Then $q' - q = 2^{h(a)}R$ with some positive integer R and the left hand side of [\(2.3\)](#page-2-0) turns into

$$
(1+t)^{q'-q} = (1+t^{2^{h(a)}})^R \equiv 1 \pmod{t^a}
$$

where the last congruence relation holds because $2^{h(a)} \geq a$. This verifies $(2.3).$ $(2.3).$

We shall prove the "only if" part by induction on a. When $a = 1$, $2^{h(a)} = 1$ and hence the congruence relation $q' \equiv q \pmod{2^{h(a)}}$ trivially holds. Suppose that the induction assumption is satisfied for $a - 1$ with $a \geq 2$ and that [\(2.3\)](#page-2-0) holds for a. Then (2.3) holds for $a-1$, so the induction assumption implies $q' \equiv q \pmod{2^{h(a-1)}}$. When $a-1$ is not a power of 2, $h(a-1) = h(a)$; so the congruence relation $q' \equiv q \pmod{2^{h(a)}}$ holds for a. When $a - 1$ is a power of 2, say 2^s ,

$$
h(a-1) = s
$$
, $h(a) = s+1$

and $q'-q = 2^s Q$ with some positive integer Q because $q' \equiv q \pmod{2^{h(a-1)}}$. Therefore

$$
(1+t)^{q'-q} = (1+t^{2^s})^Q = 1 + Qt^{2^s} + \text{higher degree terms.}
$$

Since this is congruent to 1 modulo t^a and $a > 2^s = a - 1$, Q must be even. This shows that $q' \equiv q \pmod{2^{s+1}}$, proving the induction assumption for a because $s + 1 = h(a)$. This completes the induction step and proves the "only if" part. \Box

Theorem 2.3. Let $0 \le q, q' \le b$. Then $H^*(M(q); \mathbb{Z}/2)$ and $H^*(M(q'); \mathbb{Z}/2)$ are isomorphic as graded rings if and only if $q' \equiv q$ or $b - q \pmod{2^{h(a)}}$.

Proof. If both q and q' are in $\{0, b\}$, then the theorem is trivial. So we may assume $0 < q < b$ without loss of generality. We denote by x' and y' the generators in $H^*(M(q'); \mathbb{Z}/2)$ corresponding to x and y.

4 M. MASUDA

The "if" part easily follows from [\(2.2\)](#page-1-0) and Lemma [2.2.](#page-2-1) We shall prove the "only if" part. Suppose that there is an isomorphism

$$
\varphi \colon H^*(M(q'); \mathbb{Z}/2) \to H^*(M(q); \mathbb{Z}/2)
$$

as graded rings. Since $\varphi(x')^a = \varphi(x'^a) = 0$, $\varphi(x')$ is neither y nor $x +$ y by Lemma [2.1.](#page-1-1) Therefore $\varphi(x') = x$ and hence $\varphi(y') = y$ or $x + y$. Suppose $\varphi(y') = y$. (When $\varphi(y') = x + y$, the role of q and $b - q$ will be interchanged.) Then $(x'+y')^{q'}y'^{b-q'}$ maps to $(x+y)^{q'}y^{b-q'}$ by φ and it is zero in $H^*(M(q); \mathbb{Z}/2)$, so there are constants $c, d \in \mathbb{Z}/2$ and a homogeneous polynomial $f(x, y)$ in x, y over $\mathbb{Z}/2$ such that

$$
(x+y)^{q'}y^{b-q'} = \begin{cases} c(x+y)^q y^{b-q} & \text{in case } a > b \\ d(x+y)^q y^{b-q} + f(x,y)x^a & \text{in case } a \le b \end{cases}
$$

as polynomials in x, y. Clearly c is non-zero, so $c = 1$. Therefore $q' = q$ in case $a > b$. If $d = 0$, then the right-hand side of the identity above in case $a \leq b$ is divisible by x as $a \geq 1$ while the left-hand side is not. Therefore $d = 1$ and the identity above in case $a \leq b$ implies the former congruence relations in Lemma [2.2](#page-2-1) by comparing the coefficients of $x^i y^{b-i}$ for $i < a$; so $q' \equiv q \pmod{2^{h(a)}}$ by Lemma [2.2.](#page-2-1)

Corollary 2.4. Cohomological rigidity over $\mathbb{Z}/2$ holds for $M(q)$'s if $b \leq$ $2^{h(a)}$.

Proof. Suppose that $M(q)$ and $M(q')$ have isomorphic cohomology rings with $\mathbb{Z}/2$ coefficients. Then q and q' must satisfy the congruence relation in Theorem [2.3.](#page-2-2) But since $b \leq 2^{h(a)}$, the congruence implies that $q' = q$ or $b - q$. This together with [\(2.1\)](#page-1-2) shows that $M(q')$ is diffeomorphic to $M(q)$.

3. KO theoretical condition

In this section, we use KO theory to deduce a necessary and sufficient condition on q and q' for $M(q)$ and $M(q')$ to be diffeomorphic. We begin with a general lemma.

Lemma 3.1. Let $E \rightarrow X$ be a real smooth vector bundle over a smooth manifold X. Let $\pi: P(E) \to X$ be the associated real projective bundle and let η be the tautological real line bundle over $P(E)$. Then the tangent bundle $\tau P(E)$ of $P(E)$ with 1 added is isomorphic to $\text{Hom}(\eta, \pi^*(E)) \oplus \pi^*(\tau X)$.

Proof. A point ℓ of $P(E)$ is a line in E and the fibers of η over ℓ are vectors in the line ℓ , so η is a subbundle of $\pi^*(E)$. We give a fiber metric on E. It induces a fiber metric on $\pi^*(E)$ and we denote by η^{\perp} the orthogonal complement of η in $\pi^*(E)$. Then $\tau_f P(E)$ the tangent bundle along the fiber of $\pi: P(E) \to X$ is isomorphic to $\text{Hom}(\eta, \eta^{\perp})$. This is proved in [\[8,](#page-6-4) Lemma 4.4] when X is a point and the same argument works for any X . In fact, the argument is as follows. We note that the unit S^0 bundle $S(\eta)$ of η can naturally be identified with the unit sphere bundle $S(E)$ of E. Let $v \in S(\eta)$ be in the fiber over $\ell \in P(E)$, that is, v is a vector in the line ℓ with unit length. To an element $\psi \in \text{Hom}(\eta, \eta^{\perp})$ over $\ell \in P(E)$, we assign $\psi(v)$. It is tangent to the fiber of $S(E)$ over $\pi(\ell) \in X$ at $v \in S(E) = S(\eta)$ and $\psi(-v) = -\psi(v)$, so $\psi(v)$ defines an element of $\tau_f P(E)$ over ℓ . This correspondence gives an isomorphism from $\text{Hom}(\eta, \eta^{\perp})$ to $\tau_f P(E)$.

Thus we obtain

$$
\tau_f P(E) \oplus \mathbf{1} \cong \text{Hom}(\eta, \eta^{\perp}) \oplus \text{Hom}(\eta, \eta) \cong \text{Hom}(\eta, \pi^*(E)).
$$

This implies the lemma because $\tau P(E) \cong \tau_f P(E) \oplus \pi^*(\tau X)$.

Definition. $k(a) := \#\{n \in \mathbb{N} \mid 0 \le n \le a \text{ and } n \equiv 0, 1, 2, 4 \pmod{8}\}.$

For example,

$$
k(1) = 0, \ k(2) = 1, \ k(3) = k(4) = 2, \ k(5) = k(6) = k(7) = k(8) = 3, \ k(9) = 4, \ k(10) = 5, \ k(11) = k(12) = 6, \dots
$$

It is known that $\widetilde{KO}(\mathbb{R}P^{a-1})$ is a cyclic group of order $2^{k(a)}$ generated by $\gamma-1$ ([\[1,](#page-6-5) Theorem 7.4]). This implies that $2^{k(a)}\gamma$ is trivial because the fiber dimension (that is $2^{k(a)}$) is strictly larger than the dimension of the base space (that is $a - 1$).

Theorem 3.2. Let $0 \le q, q' \le b$. Then $M(q)$ and $M(q')$ are diffeomorphic if and only if $q' \equiv q$ or $b - q \pmod{2^{k(a)}}$.

Proof. We shall prove the "if" part first. If $2^{k(a)} \ge b$ (this is the case when $a \geq b$, then $q' = q$ or $b - q$ and hence $M(q) \cong M(q')$ by [\(2.1\)](#page-1-2). Suppose $2^{k(a)} < b$. Then $a < b$ so that the bundles $q' \gamma \oplus (b-q')$ and $q \gamma \oplus (b-q)$ are in the stable range and these bundles are isomorphic because $\widetilde{KO}(\mathbb{R}P^{a-1})$ is a cyclic group of order $2^{k(a)}$ generated by $\gamma - 1$ and $q' \equiv q \pmod{2^{k(a)}}$. Hence $M(q) \cong M(q')$.

We shall prove the "only if" part. Suppose $M(q) \cong M(q')$ and let $f: M(q) \to M(q')$ be a diffeomorphism. Then

$$
f^*(\tau M(q')) = \tau M(q) \quad \text{in } \widetilde{KO}(M(q)).
$$

Since $\tau(\mathbb{R}P^{a-1})\oplus \mathbf{1}\cong a\gamma$, it follows from Lemma [3.1](#page-3-0) that the identity above implies

(3.1)
$$
f^* \big(\operatorname{Hom}(\eta', q' \gamma \oplus (b - q')\mathbf{1}) \oplus a\gamma \big) = \operatorname{Hom}(\eta, q\gamma \oplus (b - q)\mathbf{1}) \oplus a\gamma \quad \text{in } \widetilde{KO}(M(q))
$$

where η and η' denote the tautological line bundles over $M(q)$ and $M(q')$ respectively and γ is regarded as a line bundle over $M(q)$ and $M(q')$ through the projections onto $\mathbb{R}P^{a-1}$.

If both q and q' are in $\{0, b\}$, then the "only if" part is obviously satisfied. Therefore we may assume that $0 < q < b$. Then $f^*(x') = x$ and $f^*(y') = y$ or $x+y$ by Lemma [2.1.](#page-1-1) Therefore $f^*(\gamma) = \gamma$ and $f^*(\eta') = \eta$ or $\gamma\eta$. Suppose

 $f^*(\eta') = \eta$ occurs. (When $f^*(\eta') = \gamma\eta$ occurs, the role of q and $b - q$ will be interchanged.) Then [\(3.1\)](#page-4-0) reduces to

$$
\operatorname{Hom}(\eta, q' \gamma \oplus (b - q')\mathbf{1}) = \operatorname{Hom}(\eta, q \gamma \oplus (b - q)\mathbf{1}) \quad \text{in } \widetilde{KO}(M(q)).
$$

The fibration $M(q) \to \mathbb{R}P^{a-1}$ has a cross-section and we send the identity above to $\widetilde{KO}(\mathbb{R}P^{a-1})$ through the cross-section. Then η becomes trivial or γ because a line bundle over $\mathbb{R}P^{a-1}$ is either trivial or γ . In any case, the identity above reduces to

(3.2)
$$
(q'-q)(\gamma-1) = 0 \quad \text{in } \widetilde{KO}(\mathbb{R}P^{a-1})
$$

and this implies $q' \equiv q \pmod{2^{k(a)}}$

One easily sees that $h(a) \leq k(a)$ for any a and the equality holds if and only if $a \leq 9$. Corollary [2.4](#page-3-1) can be improved as follows.

Theorem 3.3. Cohomological rigidity over $\mathbb{Z}/2$ holds for $M(q)$'s if and only if $a \leq 9$ or $b \leq 2^{h(a)}$.

Proof. If $a \leq 9$, then $h(a) = k(a)$. So the "if" part follows from Theo-rems [2.3](#page-2-2) and [3.2](#page-4-1) when $a \le 9$ and from Corollary [2.4](#page-3-1) when $b \le 2^{h(a)}$.

Suppose $a \ge 10$ (so $k(a) > h(a) \ge 4$) and $b > 2^{h(a)}$. Then we take

$$
(q, q') = \begin{cases} (1, 2^{h(a)} + 1) & \text{when } b \text{ is a multiple of } 2^{h(a)}, \\ (0, 2^{h(a)}) & \text{when } b \text{ is not a multiple of } 2^{h(a)}.\end{cases}
$$

In both cases above, $q' \equiv q \pmod{2^{h(a)}}$ but q' is not congruent to neither q nor $b-q$ modulo $2^{k(a)}$ since $k(a) > h(a) \geq 4$. Therefore $M(q)$ and $M(q')$ are not diffeomorphic by Theorem [3.2](#page-4-1) while they have isomorphic cohomology rings with $\mathbb{Z}/2$ coefficients by Theorem [2.3.](#page-2-2)

4. HOMOTOPICAL RIGIDITY

Cohomological rigidity over $\mathbb{Z}/2$ does not hold for $M(q)$'s in general, but the following holds.

Theorem 4.1. If $M(q)$ and $M(q')$ are homotopy equivalent, then they are diffeomorphic.

Proof. For a finite CW complex X, $J(X)$ denotes the J group of X and $J: \widetilde{KO}(X) \to J(X)$ denotes the J homomorphism. Let $f: M(q) \to M(q')$ be a homotopy equivalence. Then

$$
J(f^*(\tau M(q'))) = J(\tau M(q)) \text{ in } J(M(q))
$$

by a theorem of Atiyah ([\[2,](#page-6-6) Theorem 3.6]). The same argument as in the latter part of the proof of Theorem [3.2](#page-4-1) shows that we may assume that $0 < q < b$ and $f^*(\eta') = \eta$, and then

$$
J((q'-q)(\gamma-1)) = 0 \quad \text{in } J(\mathbb{R}P^{a-1}).
$$

This implies [\(3.2\)](#page-5-0) because $J: \widetilde{KO}(\mathbb{R}P^{a-1}) \to J(\mathbb{R}P^{a-1})$ is an isomorphism (see [\[4,](#page-6-7) Theorem 13.9]). Hence $M(q)$ and $M(q')$ are diffeomorphic.

 \Box

Theorem [4.1](#page-5-1) motivates us to ask whether two real toric manifolds are diffeomorphic (or homeomorphic) if they are homotopy equivalent, which we may call homotopical rigidity problem for real toric manifolds.

REFERENCES

- [1] J. F. Adams, Vector fields on spheres, Ann. of Math. 75 (1962), 603–632.
- [2] M. F. Atiyah, Thom complexes, Proc. London Math. Soc. (3) 11 (1961), 291–310.
- [3] S. Choi, M. Masuda and D. Y. Suh, Topological classification of generalized Bott towers, preprint, [arXiv:0807.4334.](http://arxiv.org/abs/0807.4334)
- [4] D. Husemoller, Fiber Bundles, Third Edition, Graduate Texts in Math. 20, Springer-Verlag 1993.
- [5] Y. Kamishima and M. Masuda, Cohomological rigidity of real Bott manifolds, preprint, [arXiv:0807.4263.](http://arxiv.org/abs/0807.4263)
- [6] M. Masuda, Classification of real Bott manifolds, preprint, [arXiv:0809.2178.](http://arxiv.org/abs/0809.2178)
- [7] M. Masuda and D. Y. Suh, Classification problems of toric manifolds via topology, Proc. of Toric Topology, Contemp. Math. 460 (2008), 273–286, [arXiv:0709.4579.](http://arxiv.org/abs/0709.4579)
- [8] J. W. Milnor and J. D. Stasheff, Characteristic Classes, Ann. of Math. Studies 76, Princeton Univ. Press 1974.

Department of Mathematics, Osaka City University, Sugimoto, Sumiyoshiku, Osaka 558-8585, Japan

E-mail address: masuda@sci.osaka-cu.ac.jp