

# Induced nilpotent orbits and birational geometry

Yoshinori Namikawa

To the memory of Professor Masayoshi Nagata

## Introduction

Let  $G$  be a complex simple algebraic group and let  $\mathfrak{g}$  be its Lie algebra. A nilpotent orbit  $\mathcal{O}$  in  $\mathfrak{g}$  is an orbit of a nilpotent element of  $\mathfrak{g}$  by the adjoint action of  $G$  on  $\mathfrak{g}$ . Then  $\mathcal{O}$  admits a natural symplectic 2-form  $\omega$  and the nilpotent orbit closure  $\bar{\mathcal{O}}$  has symplectic singularities in the sense of [Be] and [Na3] (cf. [Pa], [Hi]). In [Ri], Richardson introduced the notion of so-called the *Richardson orbit*. A nilpotent orbit  $\mathcal{O}$  is called Richardson if there is a parabolic subgroup  $Q$  of  $G$  such that  $\mathcal{O} \cap n(\mathfrak{q})$  is an open dense subset of  $n(\mathfrak{q})$ , where  $n(\mathfrak{q})$  is the nil-radical of  $\mathfrak{q}$ . Later, Lusztig and Spaltenstein [L-S] generalized this notion to the *induced orbit*. A nilpotent orbit  $\mathcal{O}$  is an induced orbit if there are a parabolic subgroup  $Q$  of  $G$  and a nilpotent orbit  $\mathcal{O}'$  in the Levi subalgebra  $\mathfrak{l}(\mathfrak{q})$  of  $\mathfrak{q} := \text{Lie}(Q)$  such that  $\mathcal{O}$  meets  $n(\mathfrak{q}) + \mathcal{O}'$  in an open dense subset. If  $\mathcal{O}$  is an induced orbit, one has a natural map (cf. (1.2))

$$\nu : G \times^Q (n(\mathfrak{q}) + \bar{\mathcal{O}}') \rightarrow \bar{\mathcal{O}}.$$

The map  $\nu$  is a generically finite, projective, surjective map. This map is called the *generalized Springer map*. In this paper, we shall study the induced orbits from the view point of *birational geometry*. For a Richardson orbit  $\mathcal{O}$ , the Springer map  $\nu$  is a map from the cotangent bundle  $T^*(G/Q)$  of the flag variety  $G/Q$  to  $\bar{\mathcal{O}}$ . In [Fu], Fu proved that, if  $\bar{\mathcal{O}}$  has a crepant (projective) resolution, it is a Springer map. Note that  $Q$  is not unique (even up to the conjugate) for a Richardson orbit  $\mathcal{O}$ . This means that  $\bar{\mathcal{O}}$  has many different crepant resolutions. In [Na], the author has given a description of all crepant resolutions of  $\bar{\mathcal{O}}$  and proved that any two different crepant resolutions are connected by *Mukai flops*. The purpose of this paper is to generalize

these to *all* nilpotent orbits  $\mathcal{O}$ . If  $\mathcal{O}$  is not Richardson,  $\bar{\mathcal{O}}$  has no crepant resolution. The substitute of a crepant resolution, is a **Q-factorial terminalization**. Let  $X$  be a complex algebraic variety with rational Gorenstein singularities. A partial resolution  $f : Y \rightarrow X$  of  $X$  is said to be a **Q-factorial terminalization** of  $X$  if  $Y$  has only **Q-factorial** terminal singularities and  $f$  is a birational projective morphism such that  $K_Y = f^*K_X$ . A **Q-factorial terminalization** is a crepant resolution exactly when  $Y$  is smooth. Recently, Birkar-Cascini-Hacon-McKernan [B-C-H-M] have established the existence of minimal models of complex algebraic varieties of general type. As a corollary of this, we know that  $X$  always has a **Q-factorial terminalization**. In particular,  $\bar{\mathcal{O}}$  should have a **Q-factorial terminalization**. The author would like to pose the following conjecture.

**Conjecture.** *Let  $\mathcal{O}$  be a nilpotent orbit of a complex simple Lie algebra  $\mathfrak{g}$ . Let  $\tilde{\mathcal{O}}$  be the normalization of  $\bar{\mathcal{O}}$ . Then one of the following holds:*

- (1)  $\tilde{\mathcal{O}}$  has **Q-factorial terminal singularities**.
- (2) *There are a parabolic subalgebra  $\mathfrak{q}$  of  $\mathfrak{g}$  with Levi decomposition  $\mathfrak{q} = \mathfrak{l} \oplus \mathfrak{n}$  and a nilpotent orbit  $\mathcal{O}'$  of  $\mathfrak{l}$  such that (a):  $\mathcal{O} = \text{Ind}_{\mathfrak{l}}^{\mathfrak{g}}(\mathcal{O}')$  and (b): the normalization of  $G \times^{\mathbb{Q}} (n(\mathfrak{q}) + \tilde{\mathcal{O}}')$  is a **Q-factorial terminalization** of  $\tilde{\mathcal{O}}$  via the generalized Springer map.*

*Moreover, if  $\tilde{\mathcal{O}}$  does not have **Q-factorial terminal singularities**, then every **Q-factorial terminalization** of  $\tilde{\mathcal{O}}$  is of the form (2). Two **Q-factorial terminalizations** are connected by Mukai flops (cf. [Na], p.91).*

In this paper, we shall prove that Conjecture is true when  $\mathfrak{g}$  is classical.

*Acknowledgement:* The author received from B. Fu, in 2006, a handwritten manuscript on explicit calculations of the **Q-factorial terminalization** for certain nilpotent orbit. The author would like to thank him.

**Notations and convention.** Let  $X$  be a normal algebraic variety. Then,  $X$  is called **Q-factorial** if, for any Weil divisor  $D$  on  $X$ , its suitable multiple  $mD$  ( $m > 0$ ) is a Cartier divisor. Assume that the canonical divisor  $K_X$  is Cartier. We say that  $X$  has only *terminal singularities* if, for a resolution  $\pi : Y \rightarrow X$ ,  $K_Y = \pi^*K_X + \sum a_i E_i$  with  $a_i > 0$  for all  $i$ . Here  $E_i$  run through all  $\pi$ -exceptional prime divisors. For details in birational geometry, see the first part of [Ka] (or one can find a quick guide in [Na 4], Preliminaries). On the other hand, for details on the basic properties on nilpotent orbits, see [C-M] and [K-P].

## §1. Preliminaries

(1.1) *Nilpotent orbits and resolutions:* Let  $G$  be a complex simple algebraic group and let  $\mathfrak{g}$  be its Lie algebra.  $G$  has the adjoint action on  $\mathfrak{g}$ . The orbit  $\mathcal{O}_x$  of a nilpotent element  $x \in \mathfrak{g}$  for this action is called a nilpotent orbit. By the Jacobson-Morozov theorem, one can find a semi-simple element  $h \in \mathfrak{g}$ , and a nilpotent element  $y \in \mathfrak{g}$  in such a way that  $[h, x] = 2x$ ,  $[h, y] = -2y$  and  $[x, y] = h$ . For  $i \in \mathbf{Z}$ , let

$$\mathfrak{g}_i := \{z \in \mathfrak{g} \mid [h, z] = iz\}.$$

Then one can write

$$\mathfrak{g} = \bigoplus_{i \in \mathbf{Z}} \mathfrak{g}_i.$$

Let  $\mathfrak{h}$  be a Cartan subalgebra of  $\mathfrak{g}$  with  $h \in \mathfrak{h}$ . Let  $\Phi$  be the corresponding root system and let  $\Delta$  be a base of simple roots such that  $h$  is  $\Delta$ -dominant, i.e.  $\alpha(h) \geq 0$  for all  $\alpha \in \Delta$ . In this situation,

$$\alpha(h) \in \{0, 1, 2\}.$$

The weighted Dynkin diagram of  $\mathcal{O}_x$  is the Dynkin diagram of  $\mathfrak{g}$  where each vertex  $\alpha$  is labeled with  $\alpha(h)$ . A nilpotent orbit  $\mathcal{O}_x$  is completely determined by its weighted Dynkin diagram. A Jacobson-Morozov parabolic subalgebra for  $x$  is the parabolic subalgebra  $\mathfrak{p}$  defined by

$$\mathfrak{p} := \bigoplus_{i \geq 0} \mathfrak{g}_i.$$

Let  $P$  be the parabolic subgroup of  $G$  determined by  $\mathfrak{p}$ . We put

$$\mathfrak{n}_2 := \bigoplus_{i \geq 2} \mathfrak{g}_i.$$

Then  $\mathfrak{n}_2$  is an ideal of  $\mathfrak{p}$ ; hence,  $P$  has the adjoint action on  $\mathfrak{n}_2$ . Let us consider the vector bundle  $G \times^P \mathfrak{n}_2$  over  $G/P$  and the map

$$\mu : G \times^P \mathfrak{n}_2 \rightarrow \mathfrak{g}$$

defined by  $\mu([g, z]) := Ad_g(z)$ . Then the image of  $\mu$  coincides with the closure  $\bar{\mathcal{O}}_x$  of  $\mathcal{O}_x$  and  $\mu$  gives a resolution of  $\bar{\mathcal{O}}_x$  (cf. [K-P], Proposition 7.4). We call  $\mu$  the *Jacobson-Morozov resolution* of  $\bar{\mathcal{O}}_x$ . The orbit  $\mathcal{O}_x$  has a natural closed non-degenerate 2-form  $\omega$  (cf. [C-G], Prop. 1.1.5., [C-M], 1.3). By  $\mu$ ,  $\omega$  is regarded as a 2-form on a Zariski open subset of  $G \times^P \mathfrak{n}_2$ . By [Pa], [Hi], it

extends to a 2-form on  $G \times^P \mathfrak{n}_2$ . In other words,  $\bar{\mathcal{O}}_x$  has symplectic singularity. Let  $\tilde{\mathcal{O}}_x$  be the normalization of  $\bar{\mathcal{O}}_x$ . In many cases, one can check the  $\mathbf{Q}$ -factoriality of  $\tilde{\mathcal{O}}_x$  by applying the following lemma to the Jacobson-Morozov resolution:

**Lemma (1.1.1).** *Let  $\pi : Y \rightarrow X$  be a projective resolution of an affine variety  $X$  with rational singularities. Let  $\rho$  be the relative Picard number for  $\pi$ . If  $\text{Exc}(\pi)$  contains  $\rho$  different prime divisors, then  $X$  is  $\mathbf{Q}$ -factorial.*

*Proof.* Recall that two line bundles  $L, L'$  on  $Y$  are called  $\pi$ -numerically equivalent if  $(L.C) = (L'.C)$  for every proper curve  $C$  on  $Y$  such that  $\pi(C)$  is a point. Let  $N^1(\pi)$  be the group of  $\pi$ -numerical classes of line bundles on  $Y$ , and put  $N^1(\pi)_{\mathbf{Q}} := N^1(\pi) \otimes \mathbf{Q}$ . By definition, the Picard number  $\rho$  is the dimension of the  $\mathbf{Q}$ -vector space  $N^1(\pi)_{\mathbf{Q}}$ . Let  $E_i$ , ( $1 \leq i \leq \rho$ ) be the prime divisors contained in  $\text{Exc}(\pi)$ . We shall prove that  $N^1(\pi)_{\mathbf{Q}} = \oplus \mathbf{Q}[E_i]$ . It suffices to show that  $[E_i]$ 's are linearly independent. Put  $d := \dim Y$ . Assume  $\sum a_i [E_i] = 0$ . Put  $m$  to be the largest number among  $\{\dim \pi(E_i)\}$ . We take  $m$  very ample divisors  $H_1, \dots, H_m$  on  $X$  in such a way that  $\cap H_j$  intersects with  $\pi(E_i)$ , in finite points, for each  $i$  with  $\dim \pi(E_i) = m$ , but  $\cap H_j$  does not intersect with  $\pi(E_i)$  for each  $i$  with  $\dim \pi(E_i) < m$ . By the Bertini theorem, we may assume that  $\cap \pi^*(H_j)$  is non-singular. We cut out  $\cap \pi^*(H_j)$  further by  $d - m - 2$  very ample divisors  $L_1, \dots, L_{d-m-2}$  on  $Y$ . Then the resulting variety is a smooth surface  $S$ . The restriction of  $\pi$  to  $S$  gives a birational contraction map  $\pi|_S : S \rightarrow \bar{S}$ . For  $i$  with  $\dim \pi(E_i) = m$ , we put  $C_i := \pi^*(H_1) \cap \dots \cap \pi^*(H_m) \cap L_1 \cap \dots \cap L_{d-m-2} \cap E_i$ . Then  $C_i$  is a (possibly reducible) curve contained in  $\text{Exc}(\pi|_S)$ . Moreover, two different such  $C_i$ 's do not have no common irreducible components. By a theorem of Grauert, the intersection matrix of  $\pi|_S$ -exceptional curves, is negative definite, which implies that  $[C_i]$  are linearly independent in  $N^1(\pi|_S)_{\mathbf{Q}}$ . Let us consider the restriction map

$$\iota : N^1(\pi)_{\mathbf{Q}} \rightarrow N^1(\pi|_S)_{\mathbf{Q}}.$$

Note that  $\iota([E_i]) = 0$  for  $i$  with  $\dim \pi(E_i) < m$ , and  $\iota([E_i]) = [C_i]$  for  $i$  with  $\dim \pi(E_i) = m$ . Since  $\sum a_i [E_i] = 0$  in  $N^1(\pi)_{\mathbf{Q}}$ , we have  $\sum a_i [C_i] = 0$  in  $N^1(\pi|_S)_{\mathbf{Q}}$ . This implies that  $a_i = 0$  for all  $i$  with  $\dim \pi(E_i) = m$ . Next, replace  $m$  by the second largest number among  $\{\dim \pi(E_i)\}$  and repeat the same procedure; then we finally conclude that  $a_i = 0$  for all  $i$ . Now let us prove that  $X$  is  $\mathbf{Q}$ -factorial. Let  $D$  be a prime Weil divisor of  $X$  and let  $D'$  be the prime divisor of  $Y$  obtained as the proper transform of  $D$ . There are rational numbers  $b_i$  such that  $[D'] + \sum b_i [E_i] = 0$  in  $N^1(\pi)_{\mathbf{Q}}$ . Since  $X$  has only

rational singularities,  $l(D' + \sum b_i[E_i])$  is the pull-back of a Cartier divisor  $M$  on  $X$  for some integer  $l > 0$ . This implies that  $lD$  is linearly equivalent to the Cartier divisor  $M$ .

(1.2) *Induced orbits*

(1.2.1). Let  $G$  and  $\mathfrak{g}$  be the same as in (1.1). Let  $Q$  be a parabolic subgroup of  $G$  and let  $\mathfrak{q}$  be its Lie algebra with Levi decomposition  $\mathfrak{q} = \mathfrak{l} \oplus \mathfrak{n}$ . Here  $\mathfrak{n}$  is the nil-radical of  $\mathfrak{q}$  and  $\mathfrak{l}$  is a Levi-part of  $\mathfrak{q}$ . Fix a nilpotent orbit  $\mathcal{O}'$  in  $\mathfrak{l}$ . Then there is a unique nilpotent orbit  $\mathcal{O}$  in  $\mathfrak{g}$  meeting  $\mathfrak{n} + \mathcal{O}'$  in an open dense subset ([L-S]). Such an orbit  $\mathcal{O}$  is called the nilpotent orbit induced from  $\mathcal{O}'$  and we write

$$\mathcal{O} = \text{Ind}_{\mathfrak{l}}^{\mathfrak{g}}(\mathcal{O}').$$

Note that when  $\mathcal{O}' = 0$ ,  $\mathcal{O}$  is the Richardson orbit for  $Q$ . Since the adjoint action of  $Q$  on  $\mathfrak{q}$  stabilizes  $\mathfrak{n} + \mathcal{O}'$ , one can consider the variety  $G \times^Q (\mathfrak{n} + \mathcal{O}')$ . There is a map

$$\nu : G \times^Q (\mathfrak{n} + \mathcal{O}') \rightarrow \bar{\mathcal{O}}$$

defined by  $\nu([g, z]) := Ad_g(z)$ . Since  $\text{Codim}_{\mathfrak{l}}(\mathcal{O}') = \text{Codim}_{\mathfrak{g}}(\mathcal{O})$  (cf. [C-M], Prop. 7.1.4),  $\nu$  is a generically finite dominating map. Moreover,  $\nu$  is factorized as

$$G \times^Q (\mathfrak{n} + \mathcal{O}') \rightarrow G/Q \times \bar{\mathcal{O}} \rightarrow \bar{\mathcal{O}}$$

where the first map is a closed embedding and the second map is the 2-nd projection; this implies that  $\nu$  is a projective map. In the remainder, we call  $\nu$  the generalized Springer map for  $(Q, \mathcal{O}')$ .

(1.2.2). Assume that  $Q$  is contained in another parabolic subgroup  $\bar{Q}$  of  $G$ . Let  $\bar{L}$  be the Levi part of  $\bar{Q}$  which contains the Levi part  $L$  of  $Q$ . Let  $\bar{\mathfrak{q}} = \bar{\mathfrak{l}} \oplus \bar{\mathfrak{n}}$  be the Levi decomposition. Note that  $\bar{L} \cap Q$  is a parabolic subgroup of  $\bar{L}$  and  $\mathfrak{l}(\bar{L} \cap Q) = \mathfrak{l}$ . Let  $\mathcal{O}_1 \subset \bar{\mathfrak{l}}$  be the nilpotent orbit induced from  $(\bar{L} \cap Q, \mathcal{O}')$ . Then there is a natural map

$$\pi : G \times^Q (\mathfrak{n} + \mathcal{O}') \rightarrow G \times^{\bar{Q}} (\bar{\mathfrak{n}} + \mathcal{O}_1)$$

which factorizes  $\nu$  as  $\bar{\nu} \circ \pi = \nu$ . Here  $\bar{\nu}$  is the generalized Springer map for  $(\bar{Q}, \mathcal{O}_1)$ .

(1.2.3). Assume that there are a parabolic subgroup  $Q_L$  of  $L$  and a nilpotent orbit  $\mathcal{O}_2$  in the Levi subalgebra  $\mathfrak{l}(Q_L)$  such that  $\mathcal{O}'$  is the nilpotent orbit induced from  $(Q_L, \mathcal{O}_2)$ . Then there is a parabolic subgroup  $Q'$  of  $G$

such that  $Q' \subset Q$ ,  $\mathfrak{l}(Q') = \mathfrak{l}(Q_L)$  and  $\mathcal{O}$  is the nilpotent orbit induced from  $(Q', \mathcal{O}_2)$ . The generalized Springer map  $\nu'$  for  $(Q', \mathcal{O}_2)$  is factorized as

$$G \times^{Q'} (\mathfrak{n}' + \bar{\mathcal{O}}_2) \rightarrow G \times^Q (\mathfrak{n} + \bar{\mathcal{O}}') \rightarrow \bar{\mathcal{O}}.$$

**Lemma (1.2.4).** *Let*

$$\nu : G \times^Q (\mathfrak{n} + \bar{\mathcal{O}}') \rightarrow \bar{\mathcal{O}}$$

*be a generalized Springer map defined in (1.2.1). Then the normalization of  $G \times^Q (\mathfrak{n} + \bar{\mathcal{O}}')$  is a symplectic variety.*

*Proof.* We shall prove that the pull-back of the Kostant-Kirillov form  $\omega$  on  $\bar{\mathcal{O}}$  gives a non-degenerate 2-form on  $G \times^Q (\mathfrak{n} + \bar{\mathcal{O}}')$ . This is enough for proving (1.2.4). In fact,  $G \times^Q (\mathfrak{n} + \bar{\mathcal{O}}')$  is locally a product of  $\bar{\mathcal{O}}'$  and a non-singular variety; hence the normalization of  $G \times^Q (\mathfrak{n} + \bar{\mathcal{O}}')$  has only rational Gorenstein singularities. The Kostant-Kirillov form extends to a regular 2-form on any resolution of  $G \times^Q (\mathfrak{n} + \bar{\mathcal{O}}')$  as explained in (1.1). Therefore, the normalization of  $G \times^Q (\mathfrak{n} + \bar{\mathcal{O}}')$  is a symplectic variety. The following argument is analogous to [Pa]. Let  $\mathfrak{h}$  be a Cartan subalgebra of  $\mathfrak{g}$  such that  $\mathfrak{h} \subset \mathfrak{l}$ . There is an involution  $\phi_{\mathfrak{g}}$  of  $\mathfrak{g}$  which stabilizes  $\mathfrak{h}$  and which acts on the root system  $\Phi$  via  $-1$ . Put  $\mathfrak{n}_- := \phi_{\mathfrak{g}}(\mathfrak{n})$ . Take a point  $[1, y + y'] \in G \times^Q (\mathfrak{n} + \bar{\mathcal{O}}')$  so that  $y \in \mathfrak{n}$ ,  $y' \in \bar{\mathcal{O}}'$  and  $y + y' \in \mathcal{O}$ . The tangent space of  $G \times^Q (\mathfrak{n} + \bar{\mathcal{O}}')$  at  $[1, y + y']$  is decomposed as

$$T_{[1, y + y']} = \mathfrak{n}_- \oplus T_{y + y'}(\mathfrak{n} + \bar{\mathcal{O}}').$$

Since  $Q \cdot (y + y')$  coincides with the Zariski open dense subset  $\mathcal{O} \cap (\mathfrak{n} + \bar{\mathcal{O}}') \subset \mathfrak{n} + \bar{\mathcal{O}}'$ , an element  $v \in T_{[1, y + y']}$  can be written as

$$v = v_1 + [v_2, y + y'], \quad v_1 \in \mathfrak{n}_-, \quad v_2 \in \mathfrak{q}.$$

Let  $d\nu_* : T_{[1, y + y']} \rightarrow T_{\nu([1, y + y'])}\bar{\mathcal{O}}$  be the tangential map for  $\nu$ . Then

$$d\nu_*(v) = [v_1 + v_2, y + y'].$$

Take one more element  $w \in T_{[1, y + y']}$  in such a way that

$$w = w_1 + [w_2, y + y'], \quad w_1 \in \mathfrak{n}_-, \quad w_2 \in \mathfrak{q}.$$

Denote by  $\langle \cdot, \cdot \rangle$  the Killing form of  $\mathfrak{g}$ . By the definition of the Kostant-Kirillov form, one has


$$\omega(d\nu_*(v), d\nu_*(w)) := \langle y + y', [v_1 + v_2, w_1 + w_2] \rangle.$$

Note that  $\langle y + y', [v_1, w_1] \rangle = \langle y, [v_1, w_1] \rangle$ , and  $\langle y + y', [v_2, w_2] \rangle = \langle y', [v_2, w_2] \rangle$ . Therefore,

$$\begin{aligned} \omega(d\nu_*(v), d\nu_*(w)) = & \\ & \langle y, [v_1, w_1] \rangle + \langle y + y', [v_1, w_2] \rangle + \langle y + y', [v_2, w_1] \rangle + \langle y', [v_2, w_2] \rangle = \\ & \langle y, [v_1, w_1] \rangle + \langle v_1, [w_2, y + y']_n \rangle - \langle [v_2, y + y']_n, w_1 \rangle + \omega([v_2, y'], [w_2, y']), \end{aligned}$$

where  $[w_2, y + y']_n$  (resp.  $[v_2, y + y']_n$ ) is the nil-radical part of  $[w_2, y + y']$  (resp.  $[v_2, y + y']$ ) in the decomposition  $T_{y+y'}(\mathfrak{n} + \bar{\mathcal{O}}') = \mathfrak{n} + T_{y'}\mathcal{O}'$ . Let  $\mathcal{O}_r \subset \mathfrak{g}$  be the Richardson orbit for  $Q$ , and let  $\pi : T^*(G/Q) \rightarrow \bar{\mathcal{O}}_r$  be the Springer map. The first part  $\langle y, [v_1, w_1] \rangle + \langle v_1, [w_2, y + y']_n \rangle - \langle [v_2, y + y']_n, w_1 \rangle$  corresponds to the 2-form on  $T^*(G/Q)$  obtained by the pull-back of the Kostant-Kirillov 2-form on  $\mathcal{O}_r$  by  $\pi$  (cf. [Pa]), which is non-degenerate on  $T^*(G/Q)$ . Let us consider the second part  $\omega([v_2, y'], [w_2, y'])$ . Denote by  $[v_2, y + y']_l$  (resp.  $[w_2, y + y']_l$ ) the  $T_{y'}\mathcal{O}'$ -part of  $[v_2, y + y']$  (resp.  $[w_2, y + y']$ ) in the decomposition  $T_{y+y'}(\mathfrak{n} + \bar{\mathcal{O}}') = \mathfrak{n} + T_{y'}\mathcal{O}'$ . Then  $[v_2, y'] = [v_2, y + y']_l$  and  $[w_2, y'] = [w_2, y + y']_l$ ; hence, the second part is the Kostant-Kirillov form on  $\mathcal{O}'$ . The arguments above show that  $\nu^*\omega$  is non-degenerate at  $[1, y + y']$  for an arbitrary  $y \in \mathfrak{n}$  and for an arbitrary  $y' \in \mathcal{O}'$ . By the  $G$ -equivariance of  $\nu$ , we have the lemma.

(1.3) *Nilpotent orbits in classical Lie algebras:* When  $\mathfrak{g}$  is a classical Lie algebra,  $\mathfrak{g}$  is naturally a Lie subalgebra of  $\text{End}(V)$  for a complex vector space  $V$ . Then we can attach a partition  $\mathbf{d}$  of  $n := \dim V$  to each orbit as the Jordan type of an element contained in the orbit. Here a partition  $\mathbf{d} := [d_1, d_2, \dots, d_k]$  of  $n$  is a set of positive integers with  $\sum d_i = n$  and  $d_1 \geq d_2 \geq \dots \geq d_k$ . Another way of writing  $\mathbf{d}$  is  $[d_1^{s_1}, \dots, d_k^{s_k}]$  with  $d_1 > d_2 > \dots > d_k > 0$ . Here  $d_i^{s_i}$  is an  $s_i$  times  $d_i$ 's:  $d_i, d_i, \dots, d_i$ . The partition  $\mathbf{d}$  corresponds to a Young diagram. For example,  $[5, 4^2, 1]$  corresponds to



When an integer  $e$  appears in the partition  $\mathbf{d}$ , we say that  $e$  is a *member* of  $\mathbf{d}$ . We call  $\mathbf{d}$  *very even* when  $\mathbf{d}$  consists with only even members, each having even multiplicity.

Let us denote by  $\epsilon$  the number 1 or  $-1$ . Then a partition  $\mathbf{d}$  is  $\epsilon$ -admissible if all even (resp. odd) members of  $\mathbf{d}$  have even multiplicities when  $\epsilon = 1$  (resp.  $\epsilon = -1$ ). The following result can be found, for example, in [C-M, §5].

**Proposition (1.3.1)** *Let  $\mathcal{N}o(\mathfrak{g})$  be the set of nilpotent orbits of  $\mathfrak{g}$ .*

(1)( $A_{n-1}$ ): *When  $\mathfrak{g} = \mathfrak{sl}(n)$ , there is a bijection between  $\mathcal{N}o(\mathfrak{g})$  and the set of partitions  $\mathbf{d}$  of  $n$ .*

(2)( $B_n$ ): *When  $\mathfrak{g} = \mathfrak{so}(2n+1)$ , there is a bijection between  $\mathcal{N}o(\mathfrak{g})$  and the set of  $\epsilon$ -admissible partitions  $\mathbf{d}$  of  $2n+1$  with  $\epsilon = 1$ .*

(3)( $C_n$ ): *When  $\mathfrak{g} = \mathfrak{sp}(2n)$ , there is a bijection between  $\mathcal{N}o(\mathfrak{g})$  and the set of  $\epsilon$ -admissible partitions  $\mathbf{d}$  of  $2n$  with  $\epsilon = -1$ .*

(4)( $D_n$ ): *When  $\mathfrak{g} = \mathfrak{so}(2n)$ , there is a surjection  $f$  from  $\mathcal{N}o(\mathfrak{g})$  to the set of  $\epsilon$ -admissible partitions  $\mathbf{d}$  of  $2n$  with  $\epsilon = 1$ . For a partition  $\mathbf{d}$  which is not very even,  $f^{-1}(\mathbf{d})$  consists of exactly one orbit, but, for very even  $\mathbf{d}$ ,  $f^{-1}(\mathbf{d})$  consists of exactly two different orbits.*

Take an  $\epsilon$ -admissible partition  $\mathbf{d}$  of a positive integer  $m$ . If  $\epsilon = 1$ , we put  $\mathfrak{g} = \mathfrak{so}(m)$  and if  $\epsilon = -1$ , we put  $\mathfrak{g} = \mathfrak{sp}(m)$ . We denote by  $\mathcal{O}_{\mathbf{d}}$  a nilpotent orbit in  $\mathfrak{g}$  with Jordan type  $\mathbf{d}$ . Note that, except when  $\epsilon = 1$  and  $\mathbf{d}$  is very even,  $\mathcal{O}_{\mathbf{d}}$  is uniquely determined. When  $\epsilon = 1$  and  $\mathbf{d}$  is very even, there are two possibilities for  $\mathcal{O}_{\mathbf{d}}$ . If necessary, we distinguish the two orbits by the labelling:  $\mathcal{O}_{\mathbf{d}}^I$  and  $\mathcal{O}_{\mathbf{d}}^{II}$ . Let us fix a classical Lie algebra  $\mathfrak{g}$  and study the relationship among nilpotent orbits in  $\mathfrak{g}$ . When  $\mathfrak{g}$  is of type  $B$  or  $D$  (resp.  $C$ ), we only consider the  $\epsilon$ -admissible partitions with  $\epsilon = 1$  (resp.  $\epsilon = -1$ ). We introduce a partial order in the set of the partitions of (the same number): for two partitions  $\mathbf{d}$  and  $\mathbf{f}$ ,  $\mathbf{d} \geq \mathbf{f}$  if  $\sum_{i \leq k} d_i \geq \sum_{i \leq k} f_i$  for all  $k \geq 1$ . On the other hand, for two nilpotent orbits  $\mathcal{O}$  and  $\mathcal{O}'$  in  $\mathfrak{g}$ , we write  $\mathcal{O} \geq \mathcal{O}'$  if  $\mathcal{O}' \subset \bar{\mathcal{O}}$ . Then,  $\mathcal{O}_{\mathbf{d}} \geq \mathcal{O}_{\mathbf{f}}$  if and only if  $\mathbf{d} \geq \mathbf{f}$ . When  $\mathbf{d}$  and  $\mathbf{f}$  are  $\epsilon$ -admissible partitions with  $\mathbf{f} \geq \mathbf{g}$ , we call this pair an  $\epsilon$ -degeneration or simply a *degeneration*.

Now let us consider the case  $\mathfrak{g}$  is of type  $B$ ,  $C$  or  $D$ .

Assume that an  $\epsilon$ -degeneration  $\mathbf{d} \geq \mathbf{f}$  is *minimal* in the sense that there is no  $\epsilon$ -admissible partition  $\mathbf{d}'$  (except  $\mathbf{d}$  and  $\mathbf{f}$ ) such that  $\mathbf{d} \geq \mathbf{d}' \geq \mathbf{f}$ . Kraft and Procesi [K-P] have studied the *normal slice*  $N_{\mathbf{d}, \mathbf{f}}$  of  $\mathcal{O}_{\mathbf{f}} \subset \bar{\mathcal{O}}_{\mathbf{d}}$  in such cases. If, for two integers  $r$  and  $s$ , the first  $r$  rows and the first  $s$  columns of  $\mathbf{d}$  and  $\mathbf{f}$  coincide and the partition  $(d_1, \dots, d_r)$  is  $\epsilon$ -admissible, then one can erase these rows and columns from  $\mathbf{d}$  and  $\mathbf{f}$  respectively to get new partitions  $\mathbf{d}'$  and  $\mathbf{f}'$  with  $\mathbf{d}' \geq \mathbf{f}'$ . If we put  $\epsilon' := (-1)^s \epsilon$ , then  $\mathbf{d}'$  and  $\mathbf{f}'$  are both



$\epsilon'$ -admissible. The pair  $(\mathbf{d}', \mathbf{f}')$  is also minimal. Repeating such process, one can reach a degeneration  $\mathbf{d}_{irr} \geq \mathbf{f}_{irr}$  which is *irreducible* in the sense that there are no rows and columns to be erased. By [K-P], Theorem 2,  $N_{\mathbf{d}, \mathbf{f}}$  is analytically isomorphic to  $N_{\mathbf{d}_{irr}, \mathbf{f}_{irr}}$  around the origin. According to [K-P], a minimal and irreducible degeneration  $\mathbf{d} \geq \mathbf{f}$  is one of the following:

- a:  $\mathfrak{g} = sp(2)$ ,  $\mathbf{d} = (2)$ , and  $\mathbf{f} = (1^2)$ .
- b:  $\mathfrak{g} = sp(2n)$  ( $n > 1$ ),  $\mathbf{d} = (2n)$ , and  $\mathbf{f} = (2n - 2, 2)$ .
- c:  $\mathfrak{g} = so(2n + 1)$  ( $n > 0$ ),  $\mathbf{d} = (2n + 1)$ , and  $\mathbf{f} = (2n - 1, 1^2)$ .
- d:  $\mathfrak{g} = sp(4n + 2)$  ( $n > 0$ ),  $\mathbf{d} = (2n + 1, 2n + 1)$ , and  $\mathbf{f} = (2n, 2n, 2)$ .
- e:  $\mathfrak{g} = so(4n)$  ( $n > 0$ ),  $\mathbf{d} = (2n, 2n)$ , and  $\mathbf{f} = (2n - 1, 2n - 1, 1^2)$ .
- f:  $\mathfrak{g} = so(2n + 1)$  ( $n > 1$ ),  $\mathbf{d} = (2^2, 1^{2n-3})$ , and  $\mathbf{f} = (1^{2n+1})$ .
- g:  $\mathfrak{g} = sp(2n)$  ( $n > 1$ ),  $\mathbf{d} = (2, 1^{2n-2})$ , and  $\mathbf{f} = (1^{2n})$ .
- h:  $\mathfrak{g} = so(2n)$  ( $n > 2$ ),  $\mathbf{d} = (2^2, 1^{2n-4})$ , and  $\mathbf{f} = (1^{2n})$ .

In the first 4 cases (a,b,c,d,e),  $\mathcal{O}_{\mathbf{f}}$  have codimension 2 in  $\bar{\mathcal{O}}_{\mathbf{d}}$ . In the last 3 cases (f,g,h),  $\mathcal{O}_{\mathbf{f}}$  have codimension  $\geq 4$  in  $\bar{\mathcal{O}}_{\mathbf{d}}$ .

**Proposition (1.3.2)** *Let  $\mathcal{O}$  be a nilpotent orbit in a classical Lie algebra  $\mathfrak{g}$  of type B, C or D with Jordan type  $\mathbf{d} := [(d_1)^{s_1}, \dots, (d_k)^{s_k}]$  ( $d_1 > d_2 > \dots > d_k$ ). Let  $\Sigma$  be the singular locus of  $\bar{\mathcal{O}}$ . Then  $\text{Codim}_{\bar{\mathcal{O}}}(\Sigma) \geq 4$  if and only if the partition  $\mathbf{d}$  has full members, that is, any integer  $j$  with  $1 \leq j \leq d_1$  is a member of  $\mathbf{d}$ . Otherwise,  $\text{Codim}_{\bar{\mathcal{O}}}(\Sigma) = 2$ .*

*Proof.* Assume that  $\text{Codim}_{\bar{\mathcal{O}}}(\Sigma) \geq 4$ . We shall prove that  $\mathbf{d}$  has full members. Suppose, to the contrary, that there is some  $i$  with  $d_i \geq d_{i+1} + 2$ . Then one can find a minimal degeneration  $\mathbf{d} \geq \mathbf{f}$  where  $\mathbf{f}$  is one of the following:

$$\begin{aligned} & [\dots, d_i^{s_i-1}, (d_{i+1} + 1)^2, d_{i+1}^{s_{i+1}-1}, \dots], \quad d_i = d_{i+1} + 2 \\ & \quad [\dots, d_i^{s_i-1}, d_i - 2, d_{i+1} + 2, d_{i+1}^{s_{i+1}-1}, \dots] \\ & \quad [\dots, d_i^{s_i-1}, d_i - 2, (d_{i+1} + 1)^2, d_{i+1}^{s_{i+1}-2}, \dots] \\ & \quad [\dots, d_i^{s_i-2}, (d_i - 1)^2, d_{i+1} + 2, d_{i+1}^{s_{i+1}-1}, \dots] \\ & \quad [\dots, d_i^{s_i-2}, (d_i - 1)^2, (d_{i+1} + 1)^2, d_{i+1}^{s_{i+1}-2}, \dots]. \end{aligned}$$

By the row-column erasing one gets an irreducible, minimal degeneration  $\mathbf{d}_{irr} \geq \mathbf{f}_{irr}$  of type a,b,c,d or e. This is a contradiction; hence  $\mathbf{d}$  has full members. Conversely, assume that  $\mathbf{d}$  has full members, i.e.  $d_j = k - j + 1$

for all  $1 \leq j \leq k$ . Then, for every minimal degeneration  $\mathbf{d} \geq \mathbf{f}$ ,  $\mathbf{f}$  is one of the following:

$$[\dots, (k-i+1)^{s_i-2}, (k-i)^{s_{i-1}+4}, (k-i-1)^{s_{i-2}-2}, \dots]$$

$$[\dots, (k-i+1)^{s_i-1}, (k-i)^{s_{i-1}+2}, (k-i-1)^{s_{i-2}-1}, \dots].$$

By the row-column erasing one gets an irreducible, minimal degeneration  $\mathbf{d}_{irr} \geq \mathbf{f}_{irr}$  of type  $f, g$  or  $h$ . Therefore,  $\text{Codim}_{\bar{\mathcal{O}}}(\Sigma) \geq 4$ .

(1.4) *Jacobson-Morozov resolutions in the case of classical Lie algebras*(cf. [CM], 5.3): Let  $V$  be a complex vector space of dimension  $m$  with a non-degenerate symmetric (or skew-symmetric) form  $\langle \cdot, \cdot \rangle$ . In the symmetric case, we take a basis  $\{e_i\}_{1 \leq i \leq m}$  of  $V$  in such a way that  $\langle e_j, e_k \rangle = 1$  if  $j+k = m+1$  and otherwise  $\langle e_j, e_k \rangle = 0$ . In the skew-symmetric case, we take a basis  $\{e_i\}_{1 \leq i \leq m}$  of  $V$  in such a way that  $\langle e_j, e_k \rangle = 1$  if  $j < k$  and  $j+k = m+1$ , and  $\langle e_j, e_k \rangle = 0$  if  $j+k \neq m+1$ . When  $(V, \langle \cdot, \cdot \rangle)$  is a symmetric vector space,  $\mathfrak{g} := \mathfrak{so}(V)$  is the Lie algebra of type  $B_{(m-1)/2}$  (resp.  $D_{m/2}$ ) if  $m$  is odd (resp. even). When  $(V, \langle \cdot, \cdot \rangle)$  is a skew-symmetric vector space,  $\mathfrak{g} := \mathfrak{sp}(V)$  is the Lie algebra of type  $C_{m/2}$ . In the remainder of this paragraph,  $\mathfrak{g}$  is one of these Lie algebra contained in  $\text{End}(V)$ . Let  $\mathfrak{h} \subset \mathfrak{g}$  be the Cartan subalgebra consisting of all diagonal matrices, and let  $\Delta$  be the standard base of simple roots. Let  $x \in \mathfrak{g}$  be a nilpotent element. As in (1.1), one can choose  $h, y \in \mathfrak{g}$  in such a way that  $\{x, y, h\}$  is a  $sl(2)$ -triple. If necessary, by replacing  $x$  by its conjugate element, one may assume that  $h \in \mathfrak{h}$  and  $h$  is  $\Delta$ -dominant. Assume that  $x$  has Jordan type  $\mathbf{d} = [d_1, \dots, d_k]$ . The diagonal matrix  $h$  is described as follows. Let us consider the sequence of integers of length  $m$ :

$$d_1-1, d_1-3, \dots, -d_1+3, -d_1+1, d_2-1, d_2-3, \dots, -d_2+3, -d_2+1, \dots, d_k-1, d_k-3, \dots, -d_k+3, -d_k+1.$$

Rearrange this sequence in the non-increasing order and get a new sequence  $p_1^{t_1}, \dots, p_l^{t_l}$  with  $p_1 > p_2 > \dots > p_l$  and  $\sum t_i = m$ . Then

$$h = \text{diag}(p_1^{t_1}, \dots, p_l^{t_l}).$$

Here  $p_i^{t_i}$  means the  $t_i$  times of  $p_i$ 's:  $p_i, p_i, \dots, p_i$ . It is then easy to describe explicitly the Jacobson-Morozov parabolic subalgebra  $\mathfrak{p}$  of  $x$  and its ideal  $\mathfrak{n}_2$  (cf. (1.1)). The Jacobson-Morozov parabolic subgroup  $P$  is the stabilizer group of certain isotropic flag  $\{F_i\}_{1 \leq i \leq r}$  of  $V$ . Here, an isotropic flag of  $V$

(of length  $r$ ) is an increasing filtration  $0 \subset F_1 \subset F_2 \subset \dots \subset F_r \subset V$  such that  $F_{r+1-i} = F_i^\perp$  for all  $i$ . The flag type of  $P$  is  $(t_1, \dots, t_l)$ . The nilradical  $\mathfrak{n} := \bigoplus_{i>0} \mathfrak{g}_i$  of  $\mathfrak{p}$  consists of the elements  $z$  of  $\mathfrak{g}$  such that  $z(F_i) \subset F_{i-1}$  for all  $i$ . On the other hand, it depends on the weighted Dynkin diagram for  $x$  how  $\mathfrak{n}_2$  takes its place in  $\mathfrak{n}$ .

**Example (1.4.1).** Assume that  $\mathfrak{d}$  has full members (cf. (1.3.2)). Then, it can be checked that

$$\mathfrak{n}_2 = \{z \in \mathfrak{g}; z(F_i) \subset F_{i-2} \text{ for all } i\}.$$

Let us consider the Jacobson-Morozov resolution

$$\mu : G \times^P \mathfrak{n}_2 \rightarrow \bar{\mathcal{O}}_{\mathfrak{d}}.$$

Take  $z \in \mathcal{O}_{\mathfrak{d}}$ . One can look at the fiber  $\mu^{-1}(z)$  by using the characterization of  $\mathfrak{n}_2$  above. Assume that  $G = Sp(V)$ . We prepare two kinds of skew-symmetric vector space  $V_d$  ( $d$ : even), and  $W_{2d}$  ( $d$ : odd) as follows. The  $V_d$  is a  $d$  dimensional vector space with a basis  $e_{d-1}, e_{d-3}, \dots, e_{-d+3}, e_{-d+1}$ . The skew-symmetric form  $\langle \cdot, \cdot \rangle$  is defined in such a way that  $\langle e_i, e_j \rangle = 1$  (resp.  $\langle e_i, e_j \rangle = -1$ ) when  $i + j = 0$  and  $i > j$  (resp.  $i + j = 0$  and  $i < j$ ), and  $\langle e_i, e_j \rangle = 0$  when  $i + j \neq 0$ . Let  $z_d$  be the endomorphism of  $V_d$  defined by the  $d \times d$  matrix  $J_d$  with  $J_d(i, i+1) = 1$  ( $1 \leq i \leq d/2$ ),  $J_d(i, i-1) = -1$  ( $d/2+1 \leq i \leq d-1$ ) and otherwise  $J_d(i, j) = 0$ . The  $W_{2d}$  is a  $2d$  dimensional vector space with a basis  $f_{d-1}, f_{d-3}, \dots, f_{-d+3}, f_{-d+1}, f'_{d-1}, f'_{d-3}, \dots, f'_{-d+3}, f'_{-d+1}$ . The skew-symmetric form  $\langle \cdot, \cdot \rangle$  is defined in such a way that  $\langle f_i, f'_j \rangle = 1$  ( $\langle f'_i, f_j \rangle = -1$ ) when  $i + j = 0$ , and otherwise  $\langle f_i, f'_j \rangle = 0$ ,  $\langle f'_i, f_j \rangle = 0$ . Let  $z_{2d}$  be the endomorphism defined by the matrix  $J'_{2d}$  with  $J'_{2d}(i, i+1) = 1$  ( $1 \leq i \leq d-1$ ),  $J'_{2d}(d, d+1) = 0$ ,  $J'_{2d}(i, i+1) = -1$  ( $d+1 \leq i \leq 2d-1$ ) and otherwise  $J'_{2d}(i, j) = 0$ . Write  $\mathfrak{d}$  as  $[k^{s_k}, (k-1)^{s_{k-1}}, \dots, 2^{s_2}, 1^{s_1}]$  with  $s_i > 0$ . Note that when  $i$  is odd,  $s_i$  is even. In the notation above,  $d_1 = \dots = d_{s_k} = k$ ,  $d_{s_k+1} = \dots = d_{s_k+s_{k-1}} = k-1$  and so on. Then,  $t_1 = s_k$ ,  $t_2 = s_{k-1}$ ,  $t_3 = s_k + s_{k-2}$ ,  $t_4 = s_{k-1} + s_{k-3}$ , ... We may assume that

$$V = \left( \bigoplus_{d:\text{even}} (V_d)^{\oplus s_d} \right) \oplus \left( \bigoplus_{d:\text{odd}} (W_{2d})^{\oplus s_{d/2}} \right)$$

and  $z|_{V_d} = z_d$ ,  $z|_{W_{2d}} = z_{2d}$ . An element of  $\mu^{-1}(z)$  is the isotropic flag  $\{F_i\}$  of  $V$  with flag type  $(t_1, \dots, t_l)$  which satisfies  $z(F_i) \subset F_{i-2}$  for all  $i$ . One can find such a flag as follows. Put  $F_1 := z^{k-1}(V)$  ( $= \text{Im}(z^{k-1})$ ) and define  $F_{2k-2} :=$

$F_1^\perp$ . We next put  $F_2 := z^{k-2}(F_{2k-2})$  and  $F_{2k-3} := F_2^\perp$ . The subsequent step is similar; we put inductively  $F_i := z^{k-i}(F_{2k-i})$  and  $F_{2k-i-1} := F_i^\perp$ .

When  $G = SO(V)$ ,  $\mathbf{d}$  can be written as  $[k^{s_k}, (k-1)^{s_{k-1}}, \dots, 2^{s_2}, 1^{s_1}]$  where  $s_i$  is even when  $i$  is even. We prepare two kinds of symmetric vector spaces  $V_d$  ( $d$ : odd) and  $W_{2d}$  ( $d$ : even). Then

$$V = \left( \bigoplus_{d:\text{odd}} (V_d)^{\oplus s_d} \right) \oplus \left( \bigoplus_{d:\text{even}} (W_{2d})^{\oplus s_{d/2}} \right).$$

The description of the flag corresponding to  $\mu^{-1}(z)$  is quite similar.

**Lemma (1.4.2)** *Assume that  $\mathbf{d}$  has full members. For each minimal  $\epsilon$ -degeneration  $\mathbf{d} \geq \mathbf{f}$ , the fiber  $\mu^{-1}(\mathcal{O}_{\mathbf{f}})$  has codimension 1 in  $G \times^P \mathfrak{n}_2$ .*

*Proof. (1):* By the proof of (1.3.2), if  $\mathbf{d}$  has full members, then, for every minimal degeneration  $\mathbf{d} \geq \mathbf{f}$ , its reduction  $\mathbf{d}_{irr} \geq \mathbf{f}_{irr}$  is of type  $f$ ,  $g$  or  $h$ . If it is of type  $f$ , the normal slice  $N_{\mathbf{d},\mathbf{f}}$  of  $\mathcal{O}_{\mathbf{f}} \subset \bar{\mathcal{O}}_{\mathbf{d}}$  is isomorphic to the germ of  $\bar{\mathcal{O}}_{[2^2, 1^{2n-3}]}(\subset so(2n+1))$  ( $n > 1$ ) at 0. Similarly, if it is of type  $g$  (resp.  $h$ ),  $N_{\mathbf{d},\mathbf{f}}$  is isomorphic to the germ of  $\bar{\mathcal{O}}_{[2, 1^{2n-2}]}(\subset sp(2n))$  ( $n > 1$ ) (resp.  $\bar{\mathcal{O}}_{[2^2, 1^{2n-4}]}(\subset so(2n))$  ( $n > 2$ )) at 0. Note that they are all isolated singularities. Except when  $\mathbf{d}_{irr} \geq \mathbf{f}_{irr}$  is of type  $h$  with  $n = 3$ , these germs have  $\mathbf{Q}$ -factorial terminal singularities. Indeed, they are isolated symplectic singularities of  $\dim \geq 4$ ; hence they have only terminal singularities. The  $\mathbf{Q}$ -factoriality of them is checked by using the Jacobson-Morozov resolutions of them. The exceptional locus of each Jacobson-Morozov resolution consists of a flag variety with  $b_2 = 1$ ; this implies the  $\mathbf{Q}$ -factoriality. If  $\text{Codim } \mu^{-1}(\mathcal{O}_{\mathbf{f}}) \geq 2$ , then  $N_{\mathbf{d},\mathbf{f}}$  is not  $\mathbf{Q}$ -factorial. Therefore, the lemma follows in these cases.

**(2):** The only exception is when  $\mathbf{d}_{irr} \geq \mathbf{f}_{irr}$  is of type  $h$  with  $n = 3$ . In this case, the germ of  $\bar{\mathcal{O}}_{[2^2, 1^2]}(\subset so(6))$  at 0 has only terminal singularity because it is an isolated symplectic singularity with  $\dim 6$ . But its Jacobson-Morozov resolution has  $\text{Gr}_{iso}(2, 6)$  as its exceptional divisor and  $b_2(\text{Gr}_{iso}(2, 6)) = 2$ . By this observation, we know that  $\bar{\mathcal{O}}_{[2^2, 1^2]}$  is not  $\mathbf{Q}$ -factorial. Let  $\mathbf{d} = [k^{s_k}, (k-1)^{s_{k-1}}, \dots, 2^{s_2}, 1^{s_1}]$  be an  $\epsilon$ -admissible partition of  $m$ . Our exceptional case only occurs when  $s_j = 2$  for some  $j$  with  $(-1)^{j+1} = \epsilon$ . Then

$$\mathbf{f} = [\dots, (j+1)^{s_{j+1}-2}, j^6, (j-1)^{s_{j-1}-2}, \dots].$$

Let

$$\mu : G \times^P \mathfrak{n}_2 \rightarrow \bar{\mathcal{O}}_{\mathbf{d}}$$

be the Jacobson-Morozov resolution. Take  $z \in \mathcal{O}_{\mathbf{f}}$ . We shall prove that the isotropic Grassmann variety  $\text{Gr}_{iso}(2, 6)$  is contained in  $\mu^{-1}(z)$  by a similar

argument to (1.4.1). We only discuss here the case when  $G = Sp(V)$ , but the case of  $G = SO(V)$  is quite similar. As in (1.4.1), we take two skew-symmetric spaces  $V_d$  ( $d$ : even) and  $W_{2d}$  ( $d$ : odd). We may assume that  $V$  is isomorphic to

$$\bigoplus_{d:\text{even}, d \neq j-1, j, j+1} (V_d)^{\oplus s_d} \oplus W_{2j-2}^{\oplus s_{j-1}/2-1} \oplus V_j^{\oplus 6} \oplus W_{2j+2}^{\oplus s_{j+1}/2-1} \oplus \bigoplus_{d:\text{odd}, d \neq j-1, j, j+1} (W_{2d})^{\oplus s_d/2}$$

and  $z|_{V_d} = z_d$ ,  $z|_{W_{2d}} = z_{2d}$ . We must find isotropic flags  $\{F_i\}$  of  $V$  with flag type  $(t_1, \dots, t_l)$  which satisfies  $z(F_i) \subset F_{i-2}$  for all  $i$ . Here  $(t_1, \dots, t_l)$  is the same one as in (1.4.1). As in (1.4.1), we define  $F_i$  with  $i \leq k - j - 1$  by  $F_i := z^{k-i}(F_{2k-i})$  and  $F_{2k-i-1} := F_i^\perp$ . But, the situation is different from (1.4.1) when  $i = k - j$ . We cannot put  $F_{k-j} := z^j(F_{k+j})$ . In fact,  $\dim z^j(F_{k+j})$  is exactly 2 less than the dimension  $F_{k-j}$  should have because the exponent  $s_{j+1} - 2$  of  $j + 1$  in  $\mathbf{f}$  is exactly 2 less than that of  $j + 1$  in  $\mathbf{d}$ . Let us consider the direct summand  $V_j^{\oplus 6}$  of  $V$ . The kernel of the endomorphism  $z_j : V_j \rightarrow V_j$  has dimension 1 and is spanned by  $e_{j-1}$ . Take 6 copies  $e_{j-1}^{(1)}, \dots, e_{j-1}^{(6)}$  of  $e_{j-1}$ . Then,  $\text{Ker}(z_j^{\oplus 6})$  is a 6 dimensional vector space spanned by  $e_{j-1}^{(1)}, \dots, e_{j-1}^{(6)}$ . We want to choose two dimensional subspace  $L \subset \text{Ker}(z_j^{\oplus 6})$  and want to define  $F_{k-j}$  as  $z^j(F_{k+j}) + L$ . Since  $F_{k-j}$  should be contained in  $z^{j-1}((F_{k-j})^\perp)$ , we must choose  $L$  in such a way that

$$z^j(F_{k+j}) + L \subset z^{j-1}((z^j(F_{k+j}) + L)^\perp).$$

Let  $v = \sum a_i e_{j-1}^{(i)}$  and  $w = \sum b_i e_{j-1}^{(i)}$  be a basis of  $L$ . Then the condition above is equivalent to

$$\sum a_i^2 = \sum b_i^2 = \sum a_i b_i = 0.$$

This means that  $[L] \in Gr_{iso}(2, 6)$ . For such an  $L$ , we put  $F_{k-j} := z^j(F_{k+j}) + L$ . Once  $F_{k-j}$  is fixed, we define, for  $i \geq k - j + 1$ ,  $F_i := z^{k-i}(F_{2k-i})$  and  $F_{2k-i-1} := F_i^\perp$ . One can check that  $\{F_i\}$  is a desired flag. Therefore,  $Gr_{iso}(2, 6) \subset \mu^{-1}(z)$ . Since  $\dim Gr_{iso}(2, 6) = 5$  and  $\dim N_{\mathbf{d}, \mathbf{f}} = 6$ , this implies that  $\mu^{-1}(\mathcal{O}_{\mathbf{f}})$  has codimension 1 in  $G \times^P \mathfrak{n}_2$ .

**Corollary (1.4.3)** *Assume that  $\mathbf{d}$  is an  $\epsilon$ -admissible partition and it has full members. Let  $\tilde{\mathcal{O}}_{\mathbf{d}}$  be the normalization of  $\bar{\mathcal{O}}_{\mathbf{d}}$ . Then,  $\tilde{\mathcal{O}}_{\mathbf{d}}$  has only  $\mathbf{Q}$ -factorial terminal singularities except when  $\mathfrak{g} = so(4n + 2)$ ,  $n \geq 1$  and  $\mathbf{d} = [2^{2n}, 1^2]$ .*

*Proof.* Let  $k$  be the maximal member of  $\mathbf{d}$ . Then there are  $k - 1$  minimal degenerations  $\mathbf{d} \geq \mathbf{f}$ . By Lemma (1.4.2),  $\text{Exc}(\mu)$  contains at least  $k - 1$

irreducible divisors. When  $\epsilon = 1$  (i.e.  $\mathfrak{g} = \mathfrak{so}(V)$ ) and there is a minimal degeneration  $\mathbf{d} \geq \mathbf{f}$  with  $\mathbf{f}$  very even, there are two nilpotent orbits with Jordan type  $\mathbf{f}$ . Thus, in this case,  $\text{Exc}(\mu)$  contains at least  $k$  irreducible divisors. On the other hand, for the Jacobson-Morozov parabolic subgroup  $P$ ,  $b_2(G/P) = k - 1$  when  $\mathfrak{g} = \mathfrak{sp}(V)$ , or  $\mathfrak{g} = \mathfrak{so}(V)$  with  $\dim V$  odd. When  $\mathfrak{g} = \mathfrak{so}(V)$  and  $\dim V$  is even, we must be careful; if the flag type of  $P$  is of the form  $(p_1, \dots, p_{k-1}; 2; p_{k-1}, \dots, p_1)$ ,  $b_2(G/P) = k$ . This happens when  $\dim V = 4n + 2$  and  $\mathbf{d} = [2^{2n}, 1^2]$  or when  $\dim V = 8m + 4n + 4$  and  $\mathbf{d} = [4^{2m}, 3, 2^{2n}, 1]$ . In the latter case,  $\mathbf{d}$  has a minimal degeneration  $\mathbf{d} \geq \mathbf{f}$  with  $\mathbf{f} = [4^{2m}, 2^{2n+2}]$ , which is very even. Note that  $b_2(G/P)$  coincides with the relative Picard number  $\rho$  of the Jacobson-Morozov resolution. By these observations, we know that  $\mu$  has at least  $\rho$  exceptional divisors except when  $\mathfrak{g} = \mathfrak{so}(4n + 2)$ ,  $n \geq 1$  and  $\mathbf{d} = [2^{2n}, 1^2]$ . Therefore,  $\tilde{\mathcal{O}}_{\mathbf{d}}$  are  $\mathbf{Q}$ -factorial in these cases. By (1.3.2) they have terminal singularities. When  $\mathfrak{g} = \mathfrak{so}(4n+2)$ ,  $n \geq 1$  and  $\mathbf{d} = [2^{2n}, 1^2]$ ,  $\mathcal{O}_{\mathbf{d}}$  is a Richardson orbit and the Springer map gives a small resolution of  $\tilde{\mathcal{O}}_{\mathbf{d}}$ . Therefore,  $\tilde{\mathcal{O}}_{\mathbf{d}}$  has non- $\mathbf{Q}$ -factorial terminal singularities.

(1.5) *Induced orbits in classical Lie algebras:* Let  $\mathbf{d} = [d_1^{s_1}, \dots, d_k^{s_k}]$  be an  $\epsilon$ -admissible partition of  $m$ . According as  $\epsilon = 1$  or  $\epsilon = -1$ , we put  $G = SO(m)$  or  $G = Sp(m)$  respectively. Assume that  $\mathbf{d}$  does not have full members. In other words, for some  $p$ ,  $d_p \geq d_{p+1} + 2$  or  $d_k \geq 2$ . We put  $r = \sum_{1 \leq j \leq p} s_j$ . Then  $\mathcal{O}_{\mathbf{d}}$  is an induced orbit (cf. [C-M], 7.3). More explicitly, there are a parabolic subgroup  $Q$  of  $G$  with (isotropic) flag type  $(r, m - 2r, r)$  with Levi decomposition  $\mathfrak{q} = \mathfrak{l} \oplus \mathfrak{n}$ , and a nilpotent orbit  $\mathcal{O}'$  of  $\mathfrak{l}$  such that  $\mathcal{O}_{\mathbf{d}} = \text{Ind}_{\mathfrak{l}}^{\mathfrak{g}}(\mathcal{O}')$ . Here,  $\mathfrak{l}$  has a direct sum decomposition  $\mathfrak{l} = \mathfrak{gl}(r) \oplus \mathfrak{g}'$ , where  $\mathfrak{g}'$  is a simple Lie algebra of type  $B_{(m-2r-1)/2}$  (resp.  $D_{(m-2r)/2}$ , resp.  $C_{(m-2r)/2}$ ) when  $\epsilon = 1$  and  $m$  is odd (resp.  $\epsilon = 1$  and  $m$  is even, resp.  $\epsilon = -1$ ). Moreover,  $\mathcal{O}'$  is a nilpotent orbit of  $\mathfrak{g}'$  with Jordan type  $[(d_1 - 2)^{s_1}, \dots, (d_p - 2)^{s_p}, d_{p+1}^{s_{p+1}}, \dots, d_k^{s_k}]$ . Let us consider the generalized Springer map

$$\nu : G \times^Q (n(\mathfrak{q}) + \tilde{\mathcal{O}}') \rightarrow \tilde{\mathcal{O}}_{\mathbf{d}}$$

(cf. (1.2)).

**Lemma (1.5.1).** *The map  $\nu$  is birational. In other words,  $\deg(\nu) = 1$ .*

*Proof.* We only discuss the case  $G = Sp(m)$ . It is enough to prove that  $\nu^{-1}(z)$  is a point for  $z \in \mathcal{O}_{\mathbf{d}}$ . We prepare two kinds of skew-symmetric vector space  $V_d$  ( $d$ : even), and  $W_{2d}$  ( $d$ : odd) as follows. The  $V_d$  is a  $d$

dimensional vector space with a basis  $e_1, e_2, \dots, e_d$ . The skew-symmetric form  $\langle \cdot, \cdot \rangle$  is defined in such a way that  $\langle e_i, e_j \rangle = 1$  (resp.  $\langle e_i, e_j \rangle = -1$ ) when  $i + j = d + 1$  and  $i > j$  (resp.  $i + j = d + 1$  and  $i < j$ ), and  $\langle e_i, e_j \rangle = 0$  when  $i + j \neq d + 1$ . Let  $z_d$  be the endomorphism of  $V_d$  defined by the  $d \times d$  matrix  $J_d$  with  $J_d(i, i + 1) = 1$  ( $1 \leq i \leq d/2$ ),  $J_d(i, i + 1) = -1$  ( $d/2 + 1 \leq i \leq d - 1$ ) and otherwise  $J_d(i, j) = 0$ . The  $W_{2d}$  is a  $2d$  dimensional vector space with a basis  $f_1, \dots, f_d, f'_1, \dots, f'_d$ . The skew-symmetric form  $\langle \cdot, \cdot \rangle$  is defined in such a way that  $\langle f_i, f'_j \rangle = 1$  ( $\langle f'_i, f_j \rangle = -1$ ) when  $i + j = d + 1$ , and otherwise  $\langle f_i, f'_j \rangle = 0$ ,  $\langle f'_i, f_j \rangle = 0$ . Let  $z_{2d}$  be the endomorphism defined by the matrix  $J'_{2d}$  with  $J'_{2d}(i, i + 1) = 1$  ( $1 \leq i \leq d - 1$ ),  $J'_{2d}(d, d + 1) = 0$ ,  $J'_{2d}(i, i + 1) = -1$  ( $d + 1 \leq i \leq 2d - 1$ ) and otherwise  $J'_{2d}(i, j) = 0$ . Note that, in the partition  $\mathbf{d}$ ,  $s_i$  is even if  $d_i$  is odd. When  $d_i$  is even, we put  $U_i := V_{d_i}^{\oplus s_i}$  and define  $z_i \in \text{End}(U_i)$  by  $z_i = z_{d_i}^{\oplus s_i}$ . When  $d_i$  is odd, we put  $U_i := W_{2d_i}^{\oplus s_i/2}$  and define  $z_i \in \text{End}(U_i)$  by  $z_i = z_{2d_i}^{\oplus s_i/2}$ . Let us consider the skew-symmetric vector space  $V$  defined by

$$V := \bigoplus_{1 \leq i \leq k} U_i.$$

Then we may assume that  $z$  is an endomorphism of  $V$  defined by  $z = \bigoplus z_i$ . Each  $U_i$  has a filtration  $0 \subset U_{i,1} \subset U_{i,2} \subset \dots \subset U_{i,d_i} = U_i$  defined by  $U_{i,j} := \text{Im}(z_i^{d_i-j})$ . By definition,  $U_{i,d_i-1} = (U_{i,1})^\perp$ . The problem is to find an  $r$  dimensional isotropic subspace  $F$  of  $V$  in such a way that  $z(F) = 0$  and  $z|_{(F^\perp/F)}$  has Jordan type  $[(d_1 - 2)^{s_1}, \dots, (d_p - 2)^{s_p}, d_{p+1}^{s_{p+1}}, \dots, d_k^{s_k}]$ . We shall prove that  $F = \bigoplus_{1 \leq i \leq p} U_{i,1}$ . First note that  $F \subset \bigoplus_{1 \leq i \leq k} U_{i,1}$  since  $z(F) = 0$ . Next, one can check that  $F^\perp \subset \bigoplus_{i \geq 2} U_i \oplus U_{1,d_1-1}$ . In fact, if not, then one can find some  $v \in F^\perp/F$  with  $z^{d_1-2}(v) \neq 0$ , which contradicts that  $z|_{F^\perp/F}$  has Jordan type  $[(d_1 - 2)^{s_1}, \dots, (d_p - 2)^{s_p}, d_{p+1}^{s_{p+1}}, \dots, d_k^{s_k}]$ . By taking the dual with respect to the skew-symmetric form, one has  $U_{1,1} \subset F$ . We put  $F_2 := F \cap \bigoplus_{2 \leq i \leq k} U_{i,1}$ . Let  $(F_2)^\perp$  be the orthogonal complement of  $F_2$  in  $\bigoplus_{2 \leq i \leq k} U_i$  (not in  $V$ ). Then

$$F^\perp/F = \left( \bigoplus_{2 \leq j \leq d_1-1} U_{1,j} \right) \oplus (F_2)^\perp/F_2.$$

Then  $x|_{(F_2)^\perp/F_2}$  has Jordan type  $[(d_2 - 2)^{s_1}, \dots, (d_p - 2)^{s_p}, d_{p+1}^{s_{p+1}}, \dots, d_k^{s_k}]$ . We apply the same argument to  $F_2$  to conclude that  $U_{2,1} \subset F_2$ . In particular,  $U_{2,1} \subset F$ . In this way, we can prove inductively that  $U_{i,1} \subset F$  for  $i \leq p$ . Since  $\dim(\bigoplus_{1 \leq i \leq p} U_{i,1}) = r$ , we have  $F = \bigoplus_{1 \leq i \leq p} U_{i,1}$ .

**Remark (1.5.2).** A nilpotent orbit is called *rigid* if it is not induced from any other nilpotent orbit in a proper Levi subalgebra of  $\mathfrak{g}$ . In a simple Lie

algebra  $\mathfrak{g}$  of type B, C or D,  $\mathcal{O}_{\mathbf{d}}$  is rigid if and only if  $\mathbf{d}$  has full members and any odd (resp. even) member  $d_i$  does not have multiplicity  $s_i = 2$  when  $\epsilon = 1$  (resp.  $\epsilon = -1$ ) (cf. [CM]). By Corollary (1.4.3), for a rigid orbit  $\mathcal{O}_{\mathbf{d}}$ ,  $\tilde{\mathcal{O}}_{\mathbf{d}}$  has only  $\mathbf{Q}$ -factorial terminal singularities. But, even if  $\tilde{\mathcal{O}}_{\mathbf{d}}$  has only  $\mathbf{Q}$ -factorial terminal singularities,  $\mathcal{O}_{\mathbf{d}}$  is not necessarily rigid. Assume that an odd (resp. even) member  $d_p$  has multiplicity 2 when  $\epsilon = 1$  (resp.  $\epsilon = -1$ ). In this case, we have an induction of *another type*. Namely, put  $r = \sum_{1 \leq j \leq p-1} s_j + 1$ . Then, there are a parabolic subgroup  $Q$  of  $G$  with (isotropic) flag type  $(r, m - 2r, r)$  with Levi decomposition  $\mathfrak{q} = \mathfrak{l} \oplus \mathfrak{n}$ , and a nilpotent orbit  $\mathcal{O}'$  in  $\mathfrak{l}$  such that  $\mathcal{O}_{\mathbf{d}} = \text{Ind}_{\mathfrak{l}}^{\mathfrak{g}}(\mathcal{O}')$ . The orbit  $\mathcal{O}'$  is contained in  $\mathfrak{g}'$ , and its Jordan type is  $[(d_1 - 2)^{s_1}, \dots, (d_{p-1} - 2)^{s_{p-1}}, (d_p - 1)^2, d_{p+1}^{s_{p+1}}, \dots, d_k^{s_k}]$ . As explained above, if  $\mathbf{d}$  has full members, but  $\mathcal{O}_{\mathbf{d}}$  is not rigid, then  $\mathcal{O}_{\mathbf{d}}$  has an induction of this type. But, for such an induction, the generalized Springer map  $\nu$  is not birational.

## §2. Main Results

(2.1) Let  $X$  be a complex algebraic variety with rational Gorenstein singularities. A partial resolution  $f : Y \rightarrow X$  of  $X$  is said to be a  $\mathbf{Q}$ -factorial terminalization of  $X$  if  $Y$  has only  $\mathbf{Q}$ -factorial terminal singularities and  $f$  is a birational projective morphism such that  $K_Y = f^*K_X$ . In particular, when  $Y$  is smooth,  $f$  is called a crepant resolution. In general,  $X$  has no crepant resolution; however, by [B-C-H-M],  $X$  always has a  $\mathbf{Q}$ -factorial terminalization. But, in our case, the  $\mathbf{Q}$ -factorial terminalization can be constructed very explicitly without using the general theory in [B-C-H-M].

**Proposition (2.1.1).** *Let  $\mathcal{O}$  be a nilpotent orbit of a classical simple Lie algebra  $\mathfrak{g}$ . Let  $\tilde{\mathcal{O}}$  be the normalization of  $\tilde{\mathcal{O}}$ . Then one of the following holds:*

(1)  $\tilde{\mathcal{O}}$  has  $\mathbf{Q}$ -factorial terminal singularities.

(2) *There are a parabolic subalgebra  $\mathfrak{q}$  of  $\mathfrak{g}$  with Levi decomposition  $\mathfrak{q} = \mathfrak{l} \oplus \mathfrak{n}$  and a nilpotent orbit  $\mathcal{O}'$  of  $\mathfrak{l}$  such that (a):  $\mathcal{O} = \text{Ind}_{\mathfrak{l}}^{\mathfrak{g}}(\mathcal{O}')$  and (b): the normalization of  $G \times^{\mathbf{Q}} (n(\mathfrak{q}) + \tilde{\mathcal{O}}')$  is a  $\mathbf{Q}$ -factorial terminalization of  $\tilde{\mathcal{O}}$  via the generalized Springer map.*

*Proof.* When  $\mathfrak{g}$  is of type A, every  $\tilde{\mathcal{O}}$  has a Springer resolution; hence (2) always holds. Let us consider the case  $\mathfrak{g}$  is of B, C or D. Assume that (1) does not hold. Then, by (1.4.3), the Jordan type  $\mathbf{d}$  of  $\mathcal{O}$  does not have full members except when  $\mathfrak{g} = \mathfrak{so}(4n + 2)$ ,  $n \geq 1$  and  $\mathbf{d} = [2^{2n}, 1^2]$ . In the exceptional case,  $\mathcal{O}$  is a Richardson orbit and the Springer map gives a crepant resolution of  $\tilde{\mathcal{O}}$ ; hence (2) holds. Now assume that  $\mathbf{d}$  does not have



full members. Then, by (1.5),  $\mathcal{O}$  is an induced nilpotent orbit and there is a generalized Springer map

$$\nu : G \times^Q (n(\mathfrak{q}) + \bar{\mathcal{O}}') \rightarrow \bar{\mathcal{O}}.$$

This map is birational by (1.5.1). Let us consider the orbit  $\mathcal{O}'$  instead of  $\mathcal{O}$ . If (1) holds for  $\mathcal{O}'$ , then  $\nu$  induces a  $\mathbf{Q}$ -factorial terminalization of  $\tilde{\mathcal{O}}$ . If (1) does not hold for  $\mathcal{O}'$ , then  $\mathcal{O}'$  is an induced orbit. By (1.2.3), one can replace  $Q$  with a smaller parabolic subgroup  $Q'$  in such a way that  $\mathcal{O}$  is induced from  $(Q', \mathcal{O}_2)$  for some nilpotent orbit  $\mathcal{O}_2 \subset \mathfrak{l}(Q')$ . The generalized Springer map  $\nu'$  for  $(Q', \mathcal{O}_2)$  is factorized as

$$G \times^{Q'} (\mathfrak{n}' + \bar{\mathcal{O}}_2) \rightarrow G \times^Q (\mathfrak{n} + \bar{\mathcal{O}}') \rightarrow \bar{\mathcal{O}}.$$


The second map is birational as explained above. The first map is locally obtained by a base change of the generalized Springer map


$$L(Q) \times^{L(Q) \cap Q'} (\mathfrak{n}(L(Q) \cap Q') + \bar{\mathcal{O}}_2) \rightarrow \bar{\mathcal{O}}'.$$

This map is birational by (1.5.1). Therefore, the first map is also birational, and  $\nu'$  is birational. This induction step terminates and (2) finally holds.

(2.2) We shall next show that *every*  $\mathbf{Q}$ -factorial terminalization of  $\tilde{\mathcal{O}}$  is of the form in Proposition (2.1.1) except when  $\tilde{\mathcal{O}}$  itself has  $\mathbf{Q}$ -factorial terminal singularities. In order to do that, we need the following Proposition.

**Proposition (2.2.1).** *Let  $\mathcal{O}$  be a nilpotent orbit of a classical simple Lie algebra  $\mathfrak{g}$ . Assume that a  $\mathbf{Q}$ -factorial terminalization of  $\tilde{\mathcal{O}}$  is given by the normalization of  $G \times^Q (n(\mathfrak{q}) + \bar{\mathcal{O}}')$  for some  $(Q, \mathcal{O}')$  as in (2.1.1). Assume that  $Q$  is a maximal parabolic subgroup of  $G$  (i.e.  $b_2(G/Q) = 1$ ), and this  $\mathbf{Q}$ -factorial terminalization is small. Then  $Q$  is a parabolic subgroup corresponding to one of the following marked Dynkin diagrams and  $\mathcal{O}' = 0$ :*





*Proof.* Assume that  $\mathfrak{g}$  is of type  $A$ . Then every nilpotent orbit closure has a crepant resolution via the Springer map. Once  $\tilde{\mathcal{O}}$  has a crepant resolution, every  $\mathbf{Q}$ -factorial terminalization is a crepant resolution (cf. [Na 2]). Then the claim follows from [Na], Proposition 5.1. Assume that  $\mathfrak{g}$  is of type  $B$ ,  $C$  or  $D$ . Since the  $\mathbf{Q}$ -factorial terminalization is small,  $\tilde{\mathcal{O}}$  has terminal singularities. By (1.3.2) and (1.4.3),  $\mathcal{O}$  is the nilpotent orbit in  $so(4n+2)$ ,  $n \geq 1$  with Jordan type  $[2^{2n}, 1^2]$ . In this case,  $\mathcal{O}$  is a Richardson orbit and  $\tilde{\mathcal{O}}$  has exactly two crepant resolutions via the Springer maps  $G \times^Q n(\mathfrak{q}) \rightarrow \tilde{\mathcal{O}}$  corresponding to two marked Dynkin diagrams of type  $D$  listed above (cf. [Na], p.92). Therefore,  $\tilde{\mathcal{O}}$  has no other  $\mathbf{Q}$ -factorial terminalizations.

The following is the main theorem:

**Theorem (2.2.2).** *Let  $\mathcal{O}$  be a nilpotent orbit of a classical simple Lie algebra  $\mathfrak{g}$ . Then  $\tilde{\mathcal{O}}$  always has a  $\mathbf{Q}$ -factorial terminalization. If  $\tilde{\mathcal{O}}$  itself does not have  $\mathbf{Q}$ -factorial terminal singularities, then every  $\mathbf{Q}$ -factorial terminalization is given by the normalization of  $G \times^Q (n(\mathfrak{q}) + \tilde{\mathcal{O}}')$  in (2.1.1). Moreover, any two such  $\mathbf{Q}$ -factorial terminalizations are connected by a sequence of Mukai flops of type  $A$  or  $D$  defined in [Na], pp. 91, 92.*

*Proof.* The first statement is nothing but (2.1.1). The proof of the second statement is quite similar to that of [Na], Theorem 6.1. Assume that  $\tilde{\mathcal{O}}$  does not have  $\mathbf{Q}$ -factorial terminal singularities. Then, by (2.1.1), one can find a generalized Springer (birational) map

$$\nu : G \times^Q (n(\mathfrak{q}) + \tilde{\mathcal{O}}') \rightarrow \tilde{\mathcal{O}}.$$

Let  $X_Q$  be the normalization of  $G \times^Q (n(\mathfrak{q}) + \tilde{\mathcal{O}}')$ . Then  $\nu$  induces a  $\mathbf{Q}$ -factorial terminalization  $f : X_Q \rightarrow \tilde{\mathcal{O}}$ . The relative nef cone  $\overline{\text{Amp}}(f)$  is a rational, simplicial, polyhedral cone of dimension  $b_2(G/Q)$  (cf. (1.2.2) and [Na], Lemma 6.3). Each codimension one face  $F$  of  $\overline{\text{Amp}}(f)$  corresponds to a birational contraction map  $\phi_F : X_Q \rightarrow Y_Q$ . The construction of  $\phi_F$  is as follows. The parabolic subgroup  $Q$  corresponds to a marked Dynkin diagram  $D$ . In this diagram, there are exactly  $b_2(G/Q)$  marked vertexes. Choose a marked vertex  $v$  from  $D$ . The choice of  $v$  determines a codimension one face  $F$  of  $\overline{\text{Amp}}(f)$ . Let  $D_v$  be the maximal, connected, single marked Dynkin subdiagram of  $D$  which contains  $v$ . Let  $\bar{D}$  be the marked Dynkin diagram obtained from  $D$  by erasing the marking of  $v$ . Let  $\bar{Q}$  be the parabolic

subgroup of  $G$  corresponding to  $\bar{D}$ . Then, as in (1.2.2), we have a map

$$\pi : G \times^Q (\mathfrak{n} + \bar{\mathcal{O}}') \rightarrow G \times^{\bar{Q}} (\bar{\mathfrak{n}} + \bar{\mathcal{O}}_1).$$

Let  $Y_Q$  be the normalization of  $G \times^{\bar{Q}} (\bar{\mathfrak{n}} + \bar{\mathcal{O}}_1)$ . Then  $\pi$  induces a birational map  $X_Q \rightarrow Y_Q$ . This is the map  $\phi_F$ . Note that  $\pi$  is locally obtained by a base change of the generalized Springer map

$$L(\bar{Q}) \times^{L(\bar{Q}) \cap Q} (\mathfrak{n}(L(\bar{Q}) \cap Q) + \bar{\mathcal{O}}') \rightarrow \bar{\mathcal{O}}_1.$$

Let  $Z(\mathfrak{l}(\mathfrak{q}))$  (resp.  $Z(\mathfrak{l}(\bar{\mathfrak{q}}))$ ) be the center of  $\mathfrak{l}(\mathfrak{q})$  (resp.  $\mathfrak{l}(\bar{\mathfrak{q}})$ ). By the definition of  $\bar{Q}$ , the simple factors of  $\mathfrak{l}(\bar{\mathfrak{q}})/Z(\mathfrak{l}(\bar{\mathfrak{q}}))$  are common to those of  $\mathfrak{l}(\mathfrak{q})/Z(\mathfrak{l}(\mathfrak{q}))$  except one factor, say  $\mathfrak{m}$ . Put  $\mathcal{O}'' := \mathcal{O}' \cap \mathfrak{m}$ . By (2.2.1),  $\pi$  (or  $\phi_F$ ) is a small birational map if and only if  $\mathcal{O}'' = 0$  and  $D_v$  is one of the single Dynkin diagrams listed in (2.2.1). In this case, one can make a new marked Dynkin diagram  $D'$  from  $D$  by replacing  $D_v$  by its dual  $D_v^*$  (cf. [Na], Definition 1). Let  $Q'$  be the parabolic subgroup of  $G$  corresponding to  $D'$ . We may assume that  $Q$  and  $Q'$  are both contained in  $\bar{Q}$ . The Levi part of  $Q'$  is conjugate to that of  $Q$ ; hence there is a nilpotent orbit in  $\mathfrak{l}(\mathfrak{q}')$  corresponding to  $\mathcal{O}'$ . We denote this orbit by the same  $\mathcal{O}'$ . Then  $\mathcal{O}$  is induced from  $(Q', \mathcal{O}')$ . As above, let  $X_{Q'}$  be the normalization of  $G \times^{Q'} (\mathfrak{n}(\mathfrak{q}') + \mathcal{O}')$ . Then we have a birational map  $\phi'_F : X_{Q'} \rightarrow Y_Q$ . The diagram

$$X_Q \rightarrow Y_Q \leftarrow X_{Q'}$$

is a flop. Assume that  $g : X \rightarrow \tilde{\mathcal{O}}$  is a  $\mathbf{Q}$ -factorial terminalization. Then, the natural birational map  $X \dashrightarrow X_Q$  is an isomorphism in codimension one. Let  $L$  be a  $g$ -ample line bundle on  $X$  and let  $L_0 \in \text{Pic}(X_Q)$  be its proper transform of  $L$  by this birational map. If  $L_0$  is  $f$ -nef, then  $X = X_Q$  and  $f = g$ . Assume that  $L_0$  is not  $f$ -nef. Then one can find a codimension one face  $F$  of  $\overline{\text{Amp}}(f)$  which is negative with respect to  $L_0$ . Since  $L_0$  is  $f$ -movable, the birational map  $\phi_F : X_Q \rightarrow Y_Q$  is small. Then, as seen above, there is a new (small) birational map  $\phi'_F : X_{Q'} \rightarrow Y_Q$ . Let  $f' : X_{Q'} \rightarrow \tilde{\mathcal{O}}$  be the composition of  $\phi'_F$  with the map  $Y_Q \rightarrow \tilde{\mathcal{O}}$ . Then  $f'$  is a  $\mathbf{Q}$ -factorial terminalization of  $\tilde{\mathcal{O}}$ . Replace  $f$  by this  $f'$  and repeat the same procedure; but this procedure ends in finite times (cf. [Na], Proof of Theorem 6.1 on pp. 104, 105). More explicitly, there is a finite sequence of  $\mathbf{Q}$ -factorial terminalizations of  $\tilde{\mathcal{O}}$ :

$$X_0(= X_Q) \dashrightarrow X_1(= X_{Q'}) \dashrightarrow \dots \dashrightarrow X_k(= X_{Q_k})$$

such that  $L_k \in \text{Pic}(X_k)$  is  $f_k$ -nef. This means that  $X = X_{Q_k}$ .

**Example (2.3).** We put  $G = SP(12)$ . Let  $\mathcal{O}$  be the nilpotent orbit in  $sp(12)$  with Jordan type  $[6, 3^2]$ . Let  $Q_1 \subset G$  be a parabolic subgroup with flag type  $(3, 6, 3)$ . The Levi part  $\mathfrak{l}_1$  of  $\mathfrak{q}_1$  has a direct sum decomposition

$$\mathfrak{l}_1 = \mathfrak{gl}(3) \oplus C_3,$$

where  $C_3$  is isomorphic to  $sp(6)$  up to center. Let  $\mathcal{O}'$  be the nilpotent orbit in  $sp(6)$  with Jordan type  $[4, 1^2]$ . Then  $\mathcal{O} = \text{Ind}_{\mathfrak{l}_1}^{sp(12)}(\mathcal{O}')$ . Next consider the parabolic subgroup  $Q_2 \subset SP(6)$  with flag type  $(1, 4, 1)$ . The Levi part  $\mathfrak{l}_2$  of  $\mathfrak{q}_2$  has a direct sum decomposition

$$\mathfrak{l}_2 = \mathfrak{gl}(1) \oplus C_2$$

where  $C_2$  is isomorphic to  $sp(4)$  up to center. Let  $\mathcal{O}''$  be the nilpotent orbit in  $sp(4)$  with Jordan type  $[2, 1^2]$ . Then  $\mathcal{O}' = \text{Ind}_{\mathfrak{l}_2}^{sp(6)}(\mathcal{O}'')$ . One can take a parabolic subgroup  $Q$  of  $SP(12)$  with flag type  $(3, 1, 4, 1, 3)$  in such a way that the Levi part  $\mathfrak{l}$  of  $\mathfrak{q}$  contains the nilpotent orbit  $\mathcal{O}''$ . Then  $\mathcal{O}$  is the nilpotent orbit induced from  $\mathcal{O}''$ . We shall illustrate the induction step above by

$$([2, 1^2], sp(4)) \rightarrow ([4, 1^2], sp(6)) \rightarrow ([6, 3^2], sp(12)).$$

Since  $\tilde{\mathcal{O}}''$  has only  $\mathbf{Q}$ -factorial terminal singularities, the  $\mathbf{Q}$ -factorial terminalization of  $\tilde{\mathcal{O}}$  is given by the generalized Springer map

$$\nu : G \times^{\mathbf{Q}} (n(\mathfrak{q}) + \tilde{\mathcal{O}}'') \rightarrow \tilde{\mathcal{O}}.$$

The induction step is not unique; we have another induction step

$$([2, 1^2], sp(4)) \rightarrow ([4, 3^2], sp(10)) \rightarrow ([6, 3^2], sp(12)).$$

By these inductions, we get another generalized Springer map

$$\nu' : G \times^{Q'} (n(\mathfrak{q}') + \tilde{\mathcal{O}}'') \rightarrow \tilde{\mathcal{O}},$$

where  $Q'$  is a parabolic subgroup of  $G$  with flag type  $(1, 3, 4, 3, 1)$ . This gives another  $\mathbf{Q}$ -factorial terminalization of  $\tilde{\mathcal{O}}$ . The two  $\mathbf{Q}$ -factorial terminalizations of  $\tilde{\mathcal{O}}$  are connected by a Mukai flop of type  $A_3$ .

## References

- [BCHM] Birkar, C., Cascini, P., Hacon, C., McKernan, J.: Existence of minimal models for varieties of general type, [math.AG/0610203](#)
- [Be] Beauville, A. : Symplectic singularities, *Invent. Math.* **139** (2000), 541-549
- [C-G] Chriss, M. , Ginzburg, V. : Representation theory and complex geometry, *Progress in Math.* , Birkhauser, 1997
- [C-M] Collingwood, D. , McGovern, W. : Nilpotent orbits in semi-simple Lie algebras, van Nostrand Reinhold, *Math. Series*, 1993
- [Fu] Fu, B. : Symplectic resolutions for nilpotent orbits, *Invent. Math.* **151**. (2003), 167-186
- [Hi] Hinich, V. : On the singularities of nilpotent orbits, *Israel J. Math.* **73** (1991), 297-308
- [Ka] Kawamata, Y.: Crepant blowing-up of 3-dimensional canonical singularities and its application to degenerations of algebraic surfaces, *Ann. Math.* **127** (1988), 93-163
- [K-P] Kraft, H., Procesi, C.: On the geometry of conjugacy classes in classical groups, *Comment. Math. Helv.* **57**. (1982), 539-602
- [L-S] Lusztig, G., Spaltenstein, N.: Induced unipotent classes, *J. London Math. Soc.* **19**. (1979), 41-52
- [Na] Namikawa, Y. : Birational geometry of symplectic resolutions of nilpotent orbits, *Advances Studies in Pure Mathematics* **45**, (2006), *Moduli Spaces and Arithmetic Geometry* (Kyoto, 2004), pp. 75-116
- [Na 2] Namikawa, Y.: Flops and Poisson deformations of symplectic varieties, *Publ. RIMS* **44**. No 2. (2008), 259-314
- [Na 3] Namikawa, Y.: Extensions of 2-forms and symplectic varieties, *J. Reine Angew. Math.* **539** (2001), 123-147
- [Na 4] Namikawa, Y.: Birational geometry and deformations of nilpotent orbits, *Duke Math. J.* **143** (2008), 375-405

- [Pa] Panyushev, D. I. : Rationality of singularities and the Gorenstein property of nilpotent orbits, *Funct. Anal. Appl.* **25** (1991), 225-226
- [Ri] Richardson, R. : Conjugacy classes in parabolic subgroups of semi-simple algebraic groups, *Bull. London Math. Soc.* **6** (1974), 21-24

Yoshinori Namikawa  
Department of Mathematics, Graduate School of Science, Kyoto University,  
JAPAN  
namikawa@math.kyoto-u.ac.jp