

Two new Probability inequalities and Concentration Results

Ravindran Kannan
Microsoft Research Labs., India

1 Introduction

The study of stochastic combinatorial problems as well as Probabilistic Analysis of Algorithms are among the many subjects which use concentration inequalities. A central concentration inequality is the Höfdding-Azuma (H-A) inequality: For real-valued random variables X_1, X_2, \dots, X_n satisfying respectively absolute bounds and the Martingale (difference) condition:

$$|X_i| \leq 1 \quad ; \quad E(X_i | X_1, X_2, \dots, X_{i-1}) = 0,$$

the H-A inequality asserts the following tail bound: $\Pr(|\sum_{i=1}^n X_i| \geq t) \leq c_1 e^{-c_2 t^2/n}$, for some constants c_1, c_2 (which are the tails of $N(0, n)$, the standard normal density with variance n , but for constants.) Here, we present two theorems both of which considerably weaken the assumption of an absolute bound, as well as the Martingale condition, while retaining the strength of the conclusion. As consequences of our theorems, we derive new concentration results for many combinatorial problems.

Our Theorem 1 is simply stated. It weakens the absolute bound of 1 on $|X_i|$ to a weaker condition than a bound of 1 on some moments (upto the m th moment) of X_i . It weakens the Martingale difference assumption to requiring that certain correlations be non-positive. The conclusion upper bounds $E(\sum_{i=1}^n X_i)^m$ (essentially) by the m th moment of $N(0, n)$; it will be easy to get tail bounds from these moment bounds. Note that both the hypotheses and the conclusion involve bounds on moments upto the same m ; so finite moments are sufficient to get some conclusions, unlike in H-A as well as Chernoff bounds in both of which, one uses the absolute bound to get a bound on $E(e^{X_i})$. Note that if X_i have power law tails (with only finite moments bounded), no automatic bound on $E(e^{X_i})$ is available. But, both H-A inequality and Chernoff bounds follow as very special cases of our Theorem 1.

The study of the minimum length of a Hamilton tour through n random points chosen in i.i.d. trials from the uniform density in the unit square, was started by the seminal work of Bearwood, Halton and Hammersley [10]. The algorithmic question - of finding an approximately optimal Hamilton tour in this i.i.d. setting was tackled by Karp [32] - and his work not only pioneered the field of Probabilistic Analysis of Algorithms, but also inspired later TSP algorithms for deterministic inputs, like Arora's [7]. Earlier hard concentration results for the minimal length of a Hamilton tour in the i.i.d. case were made easy by Talagrand's inequality [43]. But all these concentration results for the Hamilton tour problem as well as many other combinatorial problems [41] make crucial use of the fact that the points are i.i.d. and so random variables like the number of points in a region in the unit square are very concentrated - have exponential tails. In the modern setting, heavier tailed distributions are of interest. There are many models of what "heavy-tailed" distribution should mean; this is not the subject of this paper. But as we will see, our theorems are amenable to "bursts in space", where each region of space chooses (independently) the number of points that fall in it, but then may choose that many points possibly adversarially; further, the number of points may have power-law tails instead of exponential tails. In other problems, one may have "bursts in time", where, each time unit may choose from a power-law tailed distribution the number of arrivals/new items/jobs.

Using Theorem 1, we are able to prove as strong concentration as was known for the i.i.d. case of TSP (but for constants), but, now allowing bursts in space. We do the same for the minimum

weight spanning tree problem as well. We then consider random graphs where edge probabilities are not equal. We show a concentration result for the chromatic number (which has been well-studied under the traditional model with equal edge probabilities.) In these cases, we use the traditional Doob Martingale construction to first cast the problem as a Martingale problem. The moment conditions needed for the hypotheses of our theorems follow naturally.

But an application where we do not have a Martingale, but do have the weaker hypothesis of Theorem 1 is when we pick a random vector(s) of unit length as in the well-known Johnson-Lindenstrauss (JL) Theorem on Random Projections. Using Theorem 1, we prove a more general theorem than JL where heavier-tailed distributions are allowed.

A further weakening of the hypotheses of H-A is obtained in our Main Theorem - Theorem (7) whose proof is more complicated. In Theorem (7), we use information on conditional moments of X_i conditioned on “typical values” of $X_1 + X_2 + \dots + X_{i-1}$ as well as the “worst-case” values. This is very useful in many contexts as we show. Using Theorem 2, we settle the concentration question for (the discrete case of) the well studied stochastic bin-packing problem [17] proving concentration results which we show are best possible. Here, we prove a bound on the variance of X_i using Linear Programming duality; we then exploit a feature of Theorem (7) (which is also present in Theorem (1)): higher moments have lower weight in our bounds, so for bin-packing, it turns out that higher moments don’t need to be carefully bounded. This feature is also used for the next application which is the well-studied problem of proving concentration for the number X of triangles in the standard random graph $G(n, p)$. While many papers have proved good tail bounds for large deviations, we prove here the first sub-Gaussian tail bounds for all values of p - namely that X has $N(0, \mathbf{Var}X)$ tails for deviations upto $(np)^{7/4}$ (see Definition (1)). [Such sub-Gaussian bounds were partially known for the easy case when $p \geq 1/\sqrt{n}$, but not for the harder case of smaller p .] We also give a proof of concentration for the longest increasing subsequence problem. It is hoped that the theorems here will provide a tool to deal with heavy-tailed distributions and inhomogeneity in other situations as well.

There have been many sophisticated probability inequalities. Besides H-A (see McDiarmid [35] for many useful extensions) and Chernoff, Talagrand’s inequality already referred to ([43]) has numerous applications. Burkholder’s inequality for Martingales [15] and many later developments give bounds based on finite moments. A crucial point here is that unlike the other inequalities, different moments have different weights in the bounds (the second moment has the highest) and this helps get better tail bounds. We will discuss comparisons of our results with these earlier results in section 14. But one more note is timely here: many previous inequalities have also used Martingale bounds after excluding “atypical” cases. But usually, they insist on an absolute bound in the typical case, whereas, here we only insist on moment bounds. It is important to note that many (probably all) individual pieces of our approach have been used before; the contribution here is in carrying out the particular combination of them which is then able to prove results for a wide range applications.

2 Theorem 1

In theorem (1) below, we weaken the absolute bound $|X_i| \leq 1$ of H-A to (2). Since this will be usually applied with $n \geq m$, (2) will be weaker than $E(X_i^l | X_1 + X_2 + \dots + X_{i-1}) \leq 1$ which is in turn weaker than the absolute bound - $|X_i| \leq 1$. We replace the Martingale difference condition $E(X_i | X_1, X_2, \dots, X_{i-1}) = 0$ by the obviously weaker condition (1) which we will call *strong negative correlation*; it is only required for odd l which we see later relates to negative correlation. Also, we only require these conditions for all l upto a certain even m . We prove a bound on the (same) m

(which is even) th moment of $\sum_{i=1}^n X_i$. Thus, the higher the moment bounded by the hypothesis, the higher the moment bounded by the conclusion. This in particular will allow us to handle things like “power-law” tails. The following definition will be useful to describe tail bounds.

Definition 1. Let a, σ be positive reals. We say that a random variable X has $N(0, \sigma^2)$ tails upto a if there exist constants c, c' such that for all $t \in [0, a]$, we have

$$\Pr(|X - EX| \geq t) \leq c'e^{-ct^2/\sigma^2}.$$

Here there is a hidden parameter n (which will be clear from the context) and the constants c, c' are independent of n , whereas a, σ could depend on n .

Theorem 1. Let X_1, X_2, \dots, X_n be real valued random variables and m an even positive integer satisfying the following for $i = 1, 2, \dots, n$ ¹:

$$EX_i(X_1 + X_2 + \dots + X_{i-1})^l \leq 0, l < m, \text{ odd.} \quad (1)$$

$$E(X_i^l | X_1 + X_2 + \dots + X_{i-1}) \leq \left(\frac{n}{m}\right)^{(l-2)/2} l!, l \leq m, \text{ even.} \quad (2)$$

Then, we have

$$E \left(\sum_{i=1}^n X_i \right)^m \leq (48nm)^{m/2}.$$

$$\sum_{i=1}^n X_i \text{ has } N(0, n) \text{ tails upto } \sqrt{nm}.$$

Remark 1. Since under the hypothesis of (H-A), (1) and (2) hold for all m , (H-A) follows from the last statement of the theorem. We will also show that Chernoff bounds follow as a simple corollary.

Remark 2. Note that for the upper bound in (2), we have

$$\left[\left(\frac{n}{m}\right)^{\frac{l}{2}-1} l! \right]^{1/l} \approx \left(\frac{n}{m}\right)^{\frac{1}{2}-\frac{1}{l}} l.$$

The last quantity is an increasing function of l when $n \geq m$, which will hold in most applications. Thus the requirements on $(E(X_i^l | X_1 + X_2 + \dots + X_{i-1}))^{1/l}$ are the “strongest” for $l = 2$ and the requirements get progressively “weaker” for higher moments. This will be useful, since, in applications, it will be easier to bound the second moment than the higher ones. The same qualitative aspect also holds for the Main Theorem.

Remark 3. Here, we give one comparison of Theorem (1) with perhaps the closest result to it in the literature, namely a result proved by de la Peña ((1.7) of [18] - slightly specialized to our situation) which asserts: If X_1, X_2, \dots, X_n is a Martingale difference sequence with $E(X_i^2 | X_1, X_2, \dots, X_{i-1}) \leq 2$ for all i and $E(X_i^l | X_1, X_2, \dots, X_{i-1}) \leq (l/2)\alpha^{(l/2)-1}$, for all positive even integers l , where α is some fixed real, then

$$\Pr \left(\sum_{i=1}^n X_i \geq t \right) \leq \exp \left(-\frac{ct^2}{n + \sqrt{\alpha t}} \right).$$

¹ E will denote the expectation of the entire expression which follows.

It is easy to see that this implies $N(0, n)$ tails upto $n/\sqrt{\alpha}$.

Setting $\alpha = \frac{n}{m}$, the hypothesis of Theorem (1) implies [18]’s hypothesis **upto** $l \leq m$, not for all l as required there. Were we to be given this hypothesis for all l and furthermore assume X_i are Martingale differences (rather than the more general (1) condition), then since $n/\sqrt{\alpha} = \sqrt{nm}$, we would get the same conclusion as Theorem (1). [18]’s result is stronger in other directions (which we won’t discuss here), but, a main point of our theorem is to assume only finite moments since we would like to deal with long-tailed distributions. Further, note that we can apply our theorem with $m = O(\sqrt{n})$, whence, 2 allows moment bounds to grow with n unlike [18].

Proof Let $M_l = \text{MAX}_{i=1}^n E(X_i^l | X_1 + X_2 + \dots + X_{i-1})$ for even $l \leq m$. For $1 \leq i \leq n$ and $q \in \{0, 2, 4, \dots, m-2, m\}$, define

$$f(i, q) = E \left(\sum_{j=1}^i X_j \right)^q.$$

Using the two assumptions, we derive the following recursive inequality for $f(n, m)$, which we will later solve (much as one does in a Dynamic Programming algorithm):

$$f(n, m) \leq f(n-1, m) + \frac{11}{5} \sum_{t \in \{2, 4, 6, \dots, m\}} \frac{m^t}{t!} M_t f(n-1, m-t), \quad (3)$$

Proof of (3): Let $A = X_1 + X_2 + \dots + X_{n-1}$. Let $a_l = \frac{m^l}{l!} E|X_n|^l |A|^{m-l}$. Expanding $(A + X_n)^m$, we get

$$E(A + X_n)^m \leq EA^m + mEX_nA^{m-1} + \sum_{l=2}^m a_l. \quad (4)$$

Now, we note that $EX_nA^{m-1} \leq 0$ by hypothesis (1) and so the second term may be dropped. [In fact, this would be the only use of the Martingale difference condition if we had assumed it; we use SNC instead, since it clearly suffices.] We will next bound the “odd terms” in terms of the two even terms on the two sides using a simple “log-convexity” of moments argument. For odd $l \geq 3$, we have

$$E|X_n|^l |A|^{m-l} \leq E \left(X_n^{l+1} A^{m-l-1} X_n^{l-1} A^{m-l+1} \right)^{1/2} \leq (E(X_n^{l+1} A^{m-l-1}))^{1/2} (E(X_n^{l-1} A^{m-l+1}))^{1/2}$$

$$\text{Also, } \frac{1}{l!} \leq \frac{6}{5} \frac{1}{\sqrt{(l+1)!}} \frac{1}{\sqrt{(l-1)!}}$$

So, a_l is at most $6/5$ times the geometric mean of a_{l+1} and a_{l-1} and hence is at most $6/5$ times their arithmetic mean. Plugging this into (4), we get

$$E \left(\sum_{i=1}^n X_i \right)^m \leq EA^m + \frac{11}{5} (a_2 + a_4 + \dots + a_m) \quad (5)$$

Now, we use the standard trick of “integrating over” X_n first and then over A (which is also crucial for proving H-A) to get for even l : $EX_n^l A^{m-l} = E_A (A^{m-l} E_{X_n}(X_n^l | A)) \leq M_l EA^{m-l}$ which yields (3).

We view (3) as a recursive inequality for $f(n, m)$. We will use this same inequality for the proof of the Main theorem, but there we use an inductive proof; here, instead, we will now “unravel” the recursion to solve it. [Note that we cannot use induction since we only know the upper bound




Figure 1: Recursion Tree

involving $(n/m)^{(l/2)-1}$ on the moments (as in the hypothesis of the theorem) and as n decreases for induction, this bound gets tighter.] Note that the dropping the $EX_n A^{m-1}$ ensured that the coefficient of EA^m is 1 instead of the $11/5$ we have in front of the other terms. This is important: if we had $11/5$ instead, since the term does not reduce m , but only n , we would get a $(11/5)^n$ when we unwind the recursion. This is no good; we can get m terms in the exponent in the final result, but not n .

Imagine a directed graph (see figure Recursion Tree) constructed as follows: The graph has a root marked $f(n, m)$. The root has $(m/2) + 1$ directed edges out of it going to $(m/2) + 1$ nodes marked (respectively) $f(n-1, m), f(n-1, m-2), \dots, f(n-1, 0)$. The edges have weights associated with them which are (respectively) $1, \frac{11}{5} \frac{m^2}{2!} M_2, \frac{11}{5} \frac{m^4}{4!} M_4, \dots, \frac{11}{5} \frac{m^m}{m!} M_m$. In general, a node of the directed graph marked $f(i, q)$ (for $i \geq 2, 0 \leq q \leq m$, even) has $(q/2) + 1$ edges going from it to nodes marked $f(i-1, q), f(i-1, q-2), \dots, f(i-1, 0)$; these edges have “weights” respectively $1, \frac{11}{5} \frac{q^2}{2!} M_2, \frac{11}{5} \frac{q^4}{4!} M_4, \dots, \frac{11}{5} \frac{q^q}{q!} M_q$ which are respectively at most

$$1, \frac{11}{5} \frac{m^2}{2!} M_2, \frac{11}{5} \frac{m^4}{4!} M_4, \dots, \frac{11}{5} \frac{m^q}{q!} M_q.$$

A node marked $f(1, q)$ has one child - a leaf marked $f(0, 0)$ connected by an edge of weight M_q . Define the weight of a path from a node to a leaf as the product of the weights of the edges along the path. It is easy to show by induction on the depth of a node that $f(i, q)$ is the sum of weights of all paths from node marked $f(i, q)$ to a leaf. [For example, if the assertion holds for all $i \leq n$, then (3) implies that it holds for the root.] We do not formally prove this here. A similar (slightly more complicated) Lemma - Lemma (1)- will be proved during the proof of the Main Theorem.

Now, there is a 1-1 correspondence between paths from $f(n, m)$ to a leaf and elements of the following set : $L = \{(l_1, l_2, \dots, l_n) : l_i \geq 0, \text{ even} ; \sum_{i=1}^n l_i = m\}$; l_i indicates that at level i we take the l_i th edge - i.e., we go from node $f(i, m - l_n - l_{n-1} - \dots - l_{i+1})$ to $f(i-1, m - l_n - l_{n-1} - \dots - l_i)$ on this path. For an $l = (l_1, l_2, \dots, l_n) \in L$ and $t \in \{0, 2, 4, \dots, m\}$, define

$$g_t(l) = \text{number of } i \text{ with } l_i = t .$$

Clearly, the vector $g(l) = (g_0(l), g_2(l), \dots, g_m(l))$ belongs to the set

$$H = \{h = (h_0, h_2, h_4, \dots, h_m) : \sum_t th_t = m; h_t \geq 0; \sum_t h_t = n\}.$$

Since the weight of an edge corresponding to l_i at any level is at most $(\frac{11}{5})^z M_{l_i} \frac{m^{l_i}}{l_i!}$, where $z = 1$ iff $l_i \geq 2$, and the number of non-zero l_i along any path is at most $m/2$, we have

$$f(n, m) \leq \sum_{l \in L} \left(\frac{11}{5}\right)^{m/2} \prod_t M_t^{g_t(l)} \frac{m^{tg_t(l)}}{(t!)^{g_t(l)}}$$

For an $h \in H$, the number of $l \in L$ with $g_t(l) = h_t \forall t$ is the number of ways of picking subsets of the n variables of cardinalities $h_0, h_2, h_4, \dots, h_m$, namely,

$$\binom{n}{h_0, h_2, h_4, \dots, h_m} = \frac{n!}{h_0! h_2! h_4! \dots h_m!} \leq \frac{n^{h_2+h_4+\dots+h_m}}{h_2! h_4! \dots h_m!}.$$

Thus, we have (using the assumed upper bound on conditional moments)

$$\begin{aligned} f(n, m) &\leq \left(\frac{11}{5}\right)^{m/2} \sum_{h \in H} \frac{n^{h_2+h_4+\dots+h_m}}{h_2! h_4! \dots h_m!} \prod_t m^{th_t} \frac{n^{h_t((t/2)-1)}}{m^{h_t((t/2)-1)}} \\ &\leq \left(\frac{11}{5}\right)^{m/2} \sum_h (nm)^{\sum_t th_t/2} \frac{m^{h_2+h_4+\dots+h_m}}{h_2! h_4! \dots h_m!} \\ &\leq \left(\frac{11}{5} nm\right)^{m/2} |H| \text{MAX}_{h \in H} \frac{(em)^{h_2+h_4+\dots+h_m}}{h_2^{h_2} h_4^{h_4} \dots h_m^{h_m}}, \end{aligned} \quad (6)$$

using Stirling inequality for factorial. Now we will show that the maximum is attained when $h_2 = m/2$ and the other h_t are all zero. In what follows t only ranges over values ≥ 2 for which $h_t \neq 0$.

$$\prod_t \frac{e^{h_t} m^{h_t}}{h_t^{h_t}} = \prod_t e^{h_t} \left(\frac{m}{th_t}\right)^{h_t} t^{h_t} \leq \prod_t e^{h_t} \left(1 + \left(\frac{m}{th_t} - 1\right)\right)^{h_t} t^{h_t} \leq \exp\left(\sum_t h_t + \frac{m}{t} - h_t + h_t \ln t\right),$$

using $1 + x \leq e^x$ for all real x . Now, the function $\sum_t \left(\frac{m}{t} + h_t \ln t\right)$ (considered as a function of the h_t) is linear and so its maximum over the simplex - $h \geq 0; \sum_t th_t = m$ - is attained at an extreme point. Hence $\sum_t \left(\frac{m}{t} + h_t \ln t\right) \leq \text{MAX}_t \left(\frac{m}{t} + \frac{m \ln t}{t}\right)$. Now considered as a function of t , $\frac{m}{t} + \frac{m \ln t}{t}$ is decreasing, so the maximum of this over our range is at $t = 2$. Thus, we have

$$\prod_t \frac{e^{h_t} m^{h_t}}{h_t^{h_t}} \leq (2e)^{m/2}. \quad (7)$$

Now, we bound $|H|$: each element of H corresponds to a unique $\frac{m}{2}$ -vector $(h_2, 2h_4, 4h_8, \dots)$ with coordinates summing to $m/2$. Thus $|H|$ is at most the number of partitions of $m/2$ into $m/2$ parts which is $\binom{m}{m/2} \leq 2^m$. Plugging this and (7) into (6), we get the moment bound in the theorem.

The bound on m th moment of $\sum_i X_i$ in the theorem will be used in a standard fashion to get tail bounds. For any t , by Markov inequality, we get from the theorem $\Pr(|\sum_i X_i| \geq t) \leq \frac{(24nm)^{m/2}}{t^m}$. The right hand side is minimized at $m = t^2/(cn)$. So since the hypothesis of the theorem holds for this m , we get the claimed tail bounds. \square

The following Corollary is a strengthening of Chernoff Bounds.

Corollary 2. Suppose X_1, X_2, \dots, X_n are real valued random variables, σ a positive real and m an even positive integer such that

$$\begin{aligned} E(X_i^k | X_1 + X_2 + \dots + X_{i-1}) &\leq \sigma^2 && \text{for } k \text{ even, } \quad k \leq m \\ EX_i(X_1 + X_2 + \dots + X_{i-1})^k &\leq 0 && \text{for } k \text{ odd, } \quad k \leq m. \end{aligned}$$

Then, $E(\sum_{i=1}^n X_i)^m \leq (c_2 n m \sigma^2)^{m/2}$ and $\sum_{i=1}^n X_i$ has $N(0, n\sigma^2)$ tails upto $\text{Min}(n\sigma^2, \sqrt{m n} \sigma)$.

Proof Let $t \in [0, \text{Min}(n\sigma^2, \sqrt{m n} \sigma)]$. We will apply the theorem with m equal to the even integer nearest to $t^2/(c_1 n \sigma^2)$ for a suitable $c_1 > 2$. Since $t \leq n\sigma^2$, it is easy to see that $\sigma^2 \leq \sigma^k (n/m)^{(k/2)-1}$ for any even k , so the hypothesis of the theorem applies to the set of random variables - $(X_1/\sigma), (X_2/\sigma), \dots, (X_n/\sigma)$. So from the theorem, we get that

$$E\left(\sum_{i=1}^n X_i\right)^m \leq (c_2 n m \sigma^2)^{m/2}$$

and so by Markov, we get

$$\Pr\left(\left|\sum_{i=1}^n X_i\right| \geq t\right) \leq \left(\frac{c_2 n m \sigma^2}{t^2}\right)^{m/2}.$$

Now choose c suitably so that $\frac{c_2 n m \sigma^2}{t^2} \leq \frac{1}{2}$ and we get the Corollary.

Remark 4. The set-up for Chernoff bounds is: X_1, X_2, \dots, X_n are i.i.d. Bernoulli random variables with $EX_i = \nu$. For any $t \leq n\nu$ Chernoff bounds assert: $\Pr(|\sum_{i=1}^n (X_i - \nu)| > t) \leq e^{-ct^2/(n\nu)}$. We get this from the Corollary applied to $X_i - \nu$, since $E(X_i - \nu)^2 \leq \nu$ and since $|X_i - \nu| \leq 1$, higher even moments of $X_i - \nu$ are at most the second moment. So, the hypothesis of the Corollary hold with $\sigma^2 = \nu$ and we can apply it.

The general Chernoff bounds deal with the case when the Bernoulli trials are independent, but not identical - EX_i may be different for different i . This unfortunately is one of the points this simple theorem cannot deal with. However, the Main Theorem does deal with it and we can derive the general Chernoff bounds as a simple corollary of that theorem - see Remark (8).

3 Functions of independent random variables

Theorem 1 and the Main Theorem (7) will often be applied to a real-valued function $f(Y_1, Y_2, \dots, Y_n)$ of independent (not necessarily real-valued) random variables Y_1, Y_2, \dots to show concentration of f . This is usually done using the Doob's Martingale construction which we recall in this section. While there is no new stuff in this section, we will introduce notation used throughout the paper.

Let Y_1, Y_2, \dots, Y_n be independent random variables. Denote $Y = (Y_1, Y_2, \dots, Y_n)$. Let $f(Y)$ be a real-valued function of Y . One defines the classical Doob's Martingale:

$$X_i = E(f | Y_1, Y_2, \dots, Y_i) - E(f | Y_1, Y_2, \dots, Y_{i-1}).$$

It is a standard fact that the X_i form a Martingale difference sequence and so (1) is satisfied. We will use the short-hand $E^i f$ to denote $E(f | Y_1, Y_2, \dots, Y_i)$, so

$$X_i = E^i f - E^{i-1} f.$$

Let $Y^{(i)}$ denote the $n - 1$ -tuple of random variables $Y_1, Y_2, \dots, Y_{i-1}, Y_{i+1}, \dots, Y_n$ and suppose $f(Y^{(i)})$ is also defined. Let

$$\Delta_i = f(Y) - f(Y^{(i)}).$$

$$\text{Then, } X_i = E^i \Delta_i - E_{Y_i} (E^i \Delta_i), \quad (8)$$

since $Y^{(i)}$ does not involve Y_i . f, Y_i, X_i, Δ_i will all be reserved for these quantities throughout the paper. We use c to denote a generic constant which can have different values.

4 Random TSP with Inhomogeneous, heavy-tailed distributions

One of the earliest problems to be studied under Probabilistic Analysis [41] is the concentration of the length f of the shortest Hamilton cycle through a set of n points picked uniformly independently at random from a unit square. Similarly, Karp's algorithm for the problem [32] was one of the earliest polynomial time algorithms for the random variant of a problem which is NP-hard in the worst-case; see also [42]. It is known that $Ef \in \Theta(\sqrt{n})$ and that f has $N(0, 1)$ tails. This was proved after many earlier steps by Rhee and Talagrand [38] and Talagrand's inequality yielded a simpler proof of this. But Talagrand's method works only for independent points; under independence, the number of points in any sub-region of the unit square follows Poisson distribution which has exponentially falling tails. Here, we will give a simple self-contained proof of the concentration result for more general distributions (of number of points in sub-regions) than the Poisson. Two important points of our more general distribution are

- Inhomogeneity (some areas of the unit square having greater probability than others) is allowed.
- heavier tails (for example with power-law distributions) than the Poisson are allowed.

We divide the unit square into n small squares, each of side $1/\sqrt{n}$. We will generate at random a set Y_i of points in the i th small square, for $i = 1, 2, \dots, n$. We assume that the $|Y_i|$ are independent, but not necessarily identical random variables. Once the $|Y_i|$ are chosen, the actual sets Y_i can be chosen in any (possibly dependent) manner (subject to the cardinalities being what was already chosen.) This thus allows for collusion where points in a small square can choose to bunch together or be spread out in any way.

Theorem 3. *Suppose there is a fixed $c_1 \in (0, 1)$, an even positive integer $m \leq n$, and an $\epsilon > 0$, such that for $1 \leq i \leq n$ and $1 \leq l \leq m/2$,*

$$\Pr(|Y_i| = 0) \leq c_1 \quad ; \quad E|Y_i|^l \leq (O(l))^{(2-\epsilon)l}.$$

Suppose $f = f(Y_1, Y_2, \dots, Y_n)$ is the length of the shortest Hamilton tour through $Y_1 \cup Y_2 \cup \dots \cup Y_n$. Then, f has $N(0, 1)$ tails upto \sqrt{m} .

Remark 5. *If each Y_i is generated according to a Poisson of intensity 1 (=Area of small square times n), then $E|Y_i|^l \leq l^l$ and so the conditions of the theorem are satisfied for all m (with room to spare).*

Remark 6. *Note that if the hypothesis hold only upto a certain m , we get normal tails upto \sqrt{m} . So for example $|Y_i|$ can have power law tails and we still get a result, whereas the older results require exponential tails.*

Proof Order the small squares in \sqrt{n} layers - the first layer consists of all squares touching the bottom or left boundary; the second layer consists of all squares which are 1 square away from the bottom and left boundary etc. until the last layer is the top right square (order within each layer is arbitrary.) Fix an i . Let S_i be the i th square. Let $\tau = \tau(Y_{i+1}, \dots, Y_n)$ be the minimum distance

from a point of S_i to a point in $Y_{i+1} \cup \dots \cup Y_n$ and $\tau_0 = \text{Min}(\tau, 2\sqrt{2})$. τ_0 depends only on Y_{i+1}, \dots, Y_n . (So, $E^i \tau_0 = E\tau_0$.) We wish to bound $\Delta_i = f(Y) - f(Y^{(i)})$ (see notation in section (3)). For this, suppose we had a tour \mathcal{T} through $Y^{(i)}$. We can break this tour at a point in $Y_{i+1} \cup Y_{i+2} \cup Y_n$ (if it is not empty) closest to S_i , detour to S_i , do a tour of Y_i and then return to \mathcal{T} . If $Y_{i+1} \cup Y_{i+2} \cup Y_n$ is empty, we just break \mathcal{T} at any point and do a detour through Y_i . So, we have

$$\begin{aligned} \Delta_i &\leq \tau_0 + \text{dist. from a point in } S_i \text{ to a point in } Y_i + \text{length of tour thro' } Y_i + \tau_0 \\ &\leq 2\tau_0 + O(1/\sqrt{n}) + f(Y_i). \end{aligned}$$

Since $\Delta_i \geq 0$, we get using (8) for any even l :

$$-E_{Y_i}(E^i \Delta_i) \leq X_i \leq E^i \Delta_i \implies |X_i| \leq 2E\tau_0 + O(1/\sqrt{n}) + f(Y_i) \implies E^{i-1} X_i^l \leq c^l (E\tau_0)^l + \frac{c^l}{n^{l/2}} + \frac{c^l}{n^{l/2}} E|Y_i|^{l/2} \quad (9)$$

where the last step uses the following well-known fact [41].

Claim 1. *For any square B of side α in the plane and any set of s points in B , there is a Hamiltonian tour through the points of length at most $c\alpha\sqrt{s}$.*

First focus on $i \leq n - 100 \ln n$. We will see that we can get a good bound on $E\tau_0$ for these i . For any $\lambda \in [0, 5\sqrt{\ln n}/\sqrt{n}]$, there is a square region T_λ of side λ inside $S_{i+1} \dots S_n$ (indeed, inside the later layers) which touches S_i . So, $\Pr(\tau \geq \sqrt{2}\lambda) \leq \Pr(T_\lambda \cap (Y_{i+1} \cup \dots \cup Y_n) = \emptyset) \leq e^{-c\lambda^2}$ by the hypothesis that $\Pr(|Y_j| = 0) < c_1 < 1$. This implies that

$$\begin{aligned} E\tau_0 &\leq \Pr\left(\tau \geq 5\sqrt{\ln n}/\sqrt{n}\right) (2\sqrt{2}) + E(\tau | \tau \leq 5\sqrt{\ln n}/\sqrt{n}) \\ &\leq \frac{c}{\sqrt{n}} + \left(E(\tau^2 | \tau \leq 5\sqrt{\ln n}/\sqrt{n})\right)^{1/2} \leq \frac{c}{\sqrt{n}} + \left(\int_0^\infty \lambda e^{-c\lambda^2}\right)^{1/2} \leq \frac{c}{\sqrt{n}}. \end{aligned}$$

Plugging this and the fact that $E|Y_i|^{l/2} \leq (O(l))^{(2-\epsilon)(l/2)} \leq (O(l))^l$ into (9), we get $E^{i-1} X_i^l \leq \frac{l^l}{n^{l/2}}$. We now apply theorem (1) to $c_6\sqrt{n}X_i$, for $i = 1, 2, \dots, n - 100 \ln n$ to get

$$E \left(\sum_{i=1}^{n-100 \ln n} X_i \right)^m \leq (cm)^{m/2}. \quad (10)$$

Now, we consider $i \geq n - 100 \ln n + 1$. All of these squares are inside a square of side $\sqrt{\ln n}/\sqrt{n}$. So, we have $|\sum_{i=n-100 \ln n+1}^n X_i| \leq 2\sqrt{2} + \frac{c\sqrt{\ln n} \sqrt{\sum_{i=n-100 \ln n+1}^n |Y_i|}}{n^{1/2}}$. Now using $E \left(\sum_{i=n-100 \ln n+1}^n |Y_i| \right)^{m/2} \leq c(\ln n)^{m/2} m^{-cm}$, we get $E \left(\sum_{i=1}^n X_i \right)^m \leq (cm)^{m/2}$ which by the usual argument via Markov inequality, yields the tail bounds asserted. \square

5 Minimum Weight Spanning tree

This problem is tackled similarly to the TSP in the previous section. We will get the same result as Talagrand's inequality is able to derive, the proof is more or less the same as our proof for the TSP, except that there is an added complication because adding points does not necessarily increase the weight of the minimum spanning tree. The standard example is when we already have the vertices of an equilateral triangle and add the center to it.

Theorem 4. *Under the same hypotheses and notation as in Theorem (3), suppose $f = f(Y_1, Y_2, \dots, Y_n)$ is the length of the minimum weight spanning tree on $Y_1 \cup Y_2 \cup \dots \cup Y_n$. f has $N(0, 1)$ tails upto \sqrt{m} .*

Proof If we already have a MWST for $Y \setminus Y_i$, we can again connect the point in Y_{i+1}, \dots, Y_n closest to S_i to S_i , then add on a MWST on Y_i to get a spanning tree on Y . This implies again that $\Delta_i \leq \tau_0 + \frac{c\sqrt{|Y_i|}}{\sqrt{n}}$. But now, we could have $f(Y) < f(\hat{Y})$. We show that

Claim 2. $\Delta_i \geq -c_{10}\tau_0 - \frac{c\sqrt{|Y_i|}}{\sqrt{n}}$.

Proof We may assume that $Y_i \neq \emptyset$. Consider the MWST T of Y . We call an edge of the form $(x, y) \in T : x \in Y_i, y \in Y \setminus Y_i$, with $|x - y| \geq c_9/\sqrt{n}$, a long edge and an edge $(x, y) \in T : x \in Y_i, y \in Y \setminus Y_i$, with $|x - y| < c_9/\sqrt{n}$ a short edge. It is well-known that the degree of each vertex in T is $O(1)$ (we prove a more complicated result in the next para), so there are at most $6|Y_i|$ short edges; we remove all of them and add a MWST on the non- Y_i ends of them. Since the edges are short, the non- Y_i ends all lie in a square of side $O(1/\sqrt{n})$, so a MWST on them is of length at most $O(\sqrt{|Y_i|}/\sqrt{n})$ by Claim (1).

We claim that there are at most $O(1)$ long edges - indeed if $(x, y), (w, z)$ are any two long edges with $x, w \in Y_i$, we have $|y - z| \geq |x - y| - \frac{\sqrt{2}}{\sqrt{n}}$, since otherwise, $(T \setminus (x, y)) \cup (y, z) \cup (x, w)$ would contain a better spanning tree than T . Similarly, $|y - z| \geq |w - z| - \frac{\sqrt{2}}{\sqrt{n}}$. Let x_0 be the center of square S_i . The above implies that in the triangle x_0, y, z , we have $|y - z| \geq |x_0 - y| - \frac{6}{\sqrt{n}}, |x_0 - z| - \frac{6}{\sqrt{n}}$. But $|y - z|^2 = |y - x_0|^2 + |z - x_0|^2 - 2|y - x_0||z - x_0| \cos(\angle y, x_0, z)$. Assume without loss of generality that $|y - x_0| \geq |z - x_0|$. If the angle $\angle y, x_0, z$ were less than 10 degrees, then we would have $|y - z|^2 \leq |y - x_0|^2 + |z - x_0|^2 - 1.8|y - x_0||z - x_0| < (|y - x_0| - 0.4|z - x_0|)^2$ a contradiction. So, we must have that the angle is at least 10 degrees which implies that there are at most 36 long edges.

Let a be the point in Y_{i+1}, \dots, Y_n closest to S_i if $Y_{i+1} \cup \dots \cup Y_n$ is non-empty; otherwise, let a be the point in $Y_1 \cup Y_2 \cup \dots \cup Y_{i-1}$ closest to S_i . We finally replace each long edge $(x, y), x \in Y_i$ by edge (a, y) . This clearly only costs us $O(\tau_0)$ extra, proving the claim.

Now the proof of the theorem is completed analogously to the TSP. □

6 Chromatic Number of inhomogeneous random graphs

Martingale inequalities have been used in different (beautiful) ways on the chromatic number χ of an (ordinary) random graph $G(n, p)$, where each edge is chosen independently to be in with probability p (see for example [39],[11], [12],[22], [34], [6], [2]).

Here we study chromatic number in a more general model. An inhomogeneous random graph - denoted $G(n, P)$ - has vertex set $[n]$ and a $n \times n$ matrix $P = \{p_{ij}\}$ where p_{ij} is the probability that edge (i, j) is in the graph. Edges are in/out independently. Let

$$p = \frac{\sum_{i,j} p_{ij}}{\binom{n}{2}}$$

be the average edge probability. Let $\chi = \chi(G(n, P))$ be the chromatic number. Since each node can change the chromatic number by at most 1, it is trivial to see that $\Pr(|\chi - E\chi| \geq t) \leq c_1 e^{-c_2 t^2/n}$ by H-A. Here we prove the first non-trivial result, which is stronger than the trivial one when the graph is sparse, i.e., when $p \in o(1)$.

Theorem 5. χ of $G(n, P)$ has $N(0, n \ln n \sqrt{p})$ tails upto $n\sqrt{p}$.

Remark 7. Given only p , note that χ could be as high as $\Omega(n\sqrt{p})$: for example, p_{ij} could be $\Omega(1)$ for $i, j \in T$ for some T with $|T| = O(n\sqrt{p})$ and zero elsewhere.

Proof Let $p_i = \sum_j p_{ij}$ be the expected degree of i . Let

$$S = \{i : p_i \geq n\sqrt{p}\}.$$

$|S| \leq 2n\sqrt{p}$. Split the $n - |S|$ vertices of $[n] \setminus S$ into $k = (n - |S|)\sqrt{p}$ groups G_1, G_2, \dots, G_k by picking for each vertex a group uniformly at random independent of other vertices. It follows by routine application of Chernoff bounds that with probability at least $1/2$, we have : (i) for each i , the sum of $p_{ij}, j \in$ (same group as i) $\leq O(\ln n)$ and (ii) $|G_t| \in O(\ln n/\sqrt{p})$ for all t . We choose any partition of $[n] \setminus S$ into G_1, G_2, \dots, G_k satisfying (i) and (ii) at the outset and fix this partition. Then we make the random choices to choose $G(n, P)$. We put the vertices of S into singleton groups - $G_{k+1}, \dots, G_{k+|S|}$.

Define Y_i for $i = 1, 2, \dots, k + |S|$ as the set of edges (of $G(n, P)$) in $G_i \times (G_1 \cup G_2 \cup \dots \cup G_{i-1})$. We can define the Doob's Martingale $X_i = E(\chi | Y_1, Y_2, \dots, Y_i) - E(\chi | Y_1, Y_2, \dots, Y_{i-1})$. First consider $i = 1, 2, \dots, k$. Define Δ_i as in section 3. Let d_j be the degree of vertex j in G_i in the graph induced on G_i alone. Δ_i is at most $\max_{j \in G_i} d_j + 1$, since we can always color G_i with this many additional colors. d_j is the sum of independent Bernoulli random variables with $Ed_j = \sum_{l \in G_i} p_{jl} \leq O(\ln n)$. By Remark (8), we have that $E(d_j - Ed_j)^l \leq \text{MAX}((cl \ln n)^{l/2}, (cl)^l)$. Hence, $E^{i-1}(\Delta_i^l) \leq (cl)^l + (cl \ln n)^{l/2}$.

We will apply Theorem (1) to the sum

$$\frac{c_7 X_1}{\ln n} + \frac{c_7 X_2}{\ln n} + \dots + \frac{c_7 X_k}{\ln n}.$$

It follows from the above that these satisfy the hypothesis of the Theorem provided $m \leq k$. From this, we get that

$$E \left(\sum_{i=1}^k X_i \right)^m \leq (cmk \ln n)^{m/2}.$$

For $i = k + 1, \dots, k + |S|$, Δ_i are absolutely bounded by 1, so by the Theorem $E(X_{k+1} + X_{k+2} + \dots + X_{k+|S|})^m \leq (c|S|m)^{m/2}$. Thus,

$$E \left(\sum_{i=1}^{k+|S|} X_i \right)^m \leq (cmk \ln n)^{m/2}.$$

Let $t \in (0, n\sqrt{p})$. We take $m =$ the even integer nearest to $t^2/(c_4 n\sqrt{p} \ln n)$ to get the theorem. \square

7 Random Projections

A famous theorem of Johnson-Lindenstrauss [44] asserts that if v is picked uniformly at random from the surface of the unit ball in \mathbf{R}^n , then for $k \leq n$, and $\epsilon \in (0, 1)$, $\sum_{i=1}^k v_i^2$ has $N(0, \frac{k}{n^2})$ tails upto $\frac{k}{n}$.

The original proof exploits the details of the uniform density and simpler later proofs ([8], [20], [25]) use the Gaussian in the equivalent way of picking v . Here, we will prove the same conclusion

²A clearly equivalent statement talks about the length of the projection of a fixed unit length vector onto a random k - dimensional sub-space.

under weaker hypotheses which allows again longer tails (and so does not use any special property of the uniform or the Gaussian). This is the first application which uses the Strong Negative Correlation condition rather than the Martingale Difference condition.

Theorem 6. *Suppose $Y = (Y_1, Y_2, \dots, Y_n)$ is a random vector picked from a distribution such that (for a $k \leq n$) (i) $E(Y_i^2 | Y_1^2 + Y_2^2 + \dots + Y_{i-1}^2)$ is a non-increasing function of $Y_1^2 + Y_2^2 + \dots + Y_{i-1}^2$ for $i = 1, 2, \dots, k$ and (ii) for even $l \leq k$, $E(Y_i^l | Y_1^2 + Y_2^2 + \dots + Y_{i-1}^2) \leq (cl)^{l/2} / n^{l/2}$. Then, $\sum_{i=1}^k Y_i^2$ has $N(0, \frac{k}{n^2})$ tails upto $\frac{k}{n}$.*

Proof The theorem will be applied with $X_i = Y_i^2 - EY_i^2$. First, (i) implies for odd l : $EX_i(X_1 + X_2 + \dots + X_{i-1})^l \leq 0$, by (an elementary version) of say, the FKG inequality. [If $X_1 + X_2 + \dots + X_{i-1} = W$, then since W^l is an increasing function of W for odd l and $E(X_i | W)$ a non-increasing function of W , we have $EX_i W^l = E_W(E(X_i | W) W^l) \leq E_W(E(X_i | W)) E W^l = EX_i E W^l = 0$.] Now, for even l , $E^{i-1}(X_i^l) \leq 2^l E Y_i^{2l} + 2^l (E Y_i^2)^l \leq (cl)^l / n^l$. So we may apply the theorem to the scaled variables $c_7 n X_i$, for $i = 1, 2, \dots, k$ for $m \leq k$ to get that $n \sum_{i=1}^k X_i$ has $N(0, k)$ tails upto $O(\sqrt{kk}) = O(k)$. So, $\sum_{i=1}^k X_i$ has $N(0, \frac{k}{n^2})$ tails upto $O(k/n^2)$ as claimed. \square

Question A common use of J-L is the following: suppose we have N vectors v_1, v_2, \dots, v_N in \mathbf{R}^n , where n, N are high. We wish to project the v_i to a space of dimension $k \ll n$ and still preserve all distances $|v_i - v_j|$. Clearly, J-L guarantees that for one $v_i - v_j$, if we pick a random k dimensional space, its length is more or less preserved (within a scaling factor). Since the tail probabilities fall off exponentially in k , it suffices to take k a polynomial in $\log N$ to ensure all distances are preserved. In this setting, it is useful to find more general choices of random subspaces (instead of picking them uniformly at random from all subspaces) and there has been some work on this ([8], [1], [3]). The question is whether Theorem 1 here or the Main Theorem can be used to derive more general results.

8 Main Probability Inequality

Now, we come to the main theorem. We will again assume Strong Negative Correlation (1) of the real-valued random variables X_1, X_2, \dots, X_n . The first main point of departure from Theorem (1) is that we allow different variables to have different bounds on conditional moments. A more important point will be that we will use information on conditional moments conditioned on “typical” values of previous variables as well as the pessimistic “worst-case” values. More specifically, we assume the following bounds on moments for $i = 1, 2, \dots, n$ (m again is an even positive integer):

$$E(X_i^l | X_1 + X_2 + \dots + X_{i-1}) \leq M_{il} \quad \text{for } l = 2, 4, 6, 8 \dots m. \quad (11)$$

In some cases, the bound M_{il} may be very high for the “worst-case” $X_1 + X_2 + \dots + X_{i-1}$. We will exploit the fact that for a “typical” $X_1 + X_2 + \dots + X_{i-1}$, $E(X_i^l | X_1 + X_2 + \dots + X_{i-1})$ may be much smaller. To this end, suppose

$$\mathcal{E}_{i,l}, \quad l = 2, 4, 6, \dots, m; i = 1, 2, \dots, n$$

are events. $\mathcal{E}_{i,l}$ is to represent the “typical” case. \mathcal{E}_{1l} will be the whole sample space. In addition to (11), we assume that

$$E(X_i^l | X_1 + X_2 + \dots + X_{i-1}, \mathcal{E}_{i,l}) \leq L_{il} \quad (12)$$

$$\Pr(\mathcal{E}_{i,l}) = 1 - \delta_{i,l} \text{ for } l = 2, 4, 6, 8 \dots m \quad (13)$$

Two quantities play a role in the theorem. The first is the “average typical l th moment” L_l which we define as

$$L_l = \frac{1}{n} \sum_{i=1}^n L_{i,l} \quad \text{for } l = 2, 4, 6, 8 \dots m.$$

The second has to do with worst-case moments, but modulated by $\delta_{i,l}$. Let

$$\hat{M}_{i,l} = M_{i,l} \delta_{i,l}^{2/(m-l+2)}.$$

Note that while $M_{i,l}$ may be very large, one can make $\hat{M}_{i,l}$ smaller by controlling $\delta_{i,l}$.

Theorem 7 (Main Theorem). *Let X_1, X_2, \dots, X_n be real valued random variables satisfying Strong Negative Correlation (1) and m be a positive even integer and $L_l, \hat{M}_{i,l}, \delta_{i,l}$ be as above. Then for $X = \sum_{i=1}^n X_i$,*

$$EX^m \leq (cm)^m \left(\sum_{l=1}^{m/2} \frac{1}{l^2} \left(\frac{nL_{2l}}{m} \right)^{\frac{1}{l}} \right)^{m/2} + (cm)^m \sum_{l=1}^{m/2} \frac{1}{nl^2} \sum_{i=1}^n \left(n\hat{M}_{i,2l} \right)^{m/2l}.$$

Besides the distinction between typical case and worst-case conditional moments which we already mentioned, a second feature of the Theorem is similar to Theorem (1) in that the second moment term will often be the important one. The L term on the right hand side of the theorem is at most

$$(cm)^{m/2} \left(nL_2 + \sqrt{nm}L_4^{1/2} + \dots \right)^{m/2},$$

where we note that for $m \ll n$, (which is the usual parameter setting with which the theorem will be applied) the coefficients of higher moments decline fast, so that under reasonable conditions, the nL_2 term is what matters. In this case, it will not be difficult to see that we get $N(0, nL_2)$ tails, as we would in the ideal case when X_i are independent and in the limit $X_1 + X_2 + \dots + X_n$ behaves like the normal (with variance equal to sum of the variances of the X_i , namely nL_2).

Remark 8. *The general Chernoff bounds are a very special case: suppose $X_i, i = 1, 2, \dots, n$ are independent Bernoulli trials with $EX_i = \nu_i$. We will apply the theorem to bound the m th moment of $X = \sum_i (X_i - \nu_i)$ and from that the tail probability. It is easy to see that $E(X_i - \nu_i)^l \leq \nu_i$ for all even l , so we may take $L_{i,2l} = \nu_i$ to satisfy the hypothesis of the Theorem for every m . Let $\sum_i \nu_i = \nu$. We get*

$$EX^m \leq (cm)^{m/2} \left(m \sum_l (1/l^2) \frac{\nu^{1/l}}{m^{1/l}} \right)^{m/2}.$$

The maximum of $(\nu/m)^{1/l}$ occurs at $l = 1$ if $\nu \geq m$ and at $l = m/2$ otherwise; in any case, it is at most $1 + (\nu/m)$ and so we get (using $\sum_l (1/l^2) \leq 4$) for any $t > 0$,

$$EX^m \leq (cm(\nu + m))^{m/2} \implies Pr(|X| \geq t) \leq \left(\frac{cm(\nu + m)}{t^2} \right)^{m/2}.$$

Now putting $m = \frac{t^2}{2(\nu+t)}$, we get $Pr(|X| \geq t) \leq e^{-ct^2/(2(\nu+t))}$, which are Chernoff bounds.

9 Proof of the Main Theorem

[The proof is complicated, not for lack of efforts on the part of the author. While certainly some of the intricate use of inequalities to get things to the final form which is usable may be necessary, it is possible that the reader may be luckier in simplifying the proof.]

We will use induction on n, m . At a general step of the argument, we will need to bound $E(\sum_{i=1}^r X_i)^q$, where, $r \leq n$ and $q \leq m$, even. To bound this, let $A = X_1 + X_2 + \dots + X_{r-1}$. Binomial expansion gives us

$$E\left(\sum_{i=1}^r X_i\right)^q = E(A + X_r)^q = EA^q + qEX_r A^{q-1} + \sum_{l=2}^q \binom{q}{l} EX_r^l A^{q-l}.$$

The second term is non-positive by hypothesis. Also arguing exactly as in the proof of theorem (1), for odd $l \geq 3$,

$$EX_r^l A^{q-l} \leq \frac{3}{5}(EX_r^{l+1} A^{q-l-1} + EX_r^{l-1} A^{q-l+1}),$$

and so we get

$$E\left(\sum_{i=1}^r X_i\right)^q \leq EA^q + 3 \sum_{\substack{l \geq 2 \\ l \text{ even}}} \binom{q}{l} EX_r^l A^{q-l}. \quad (14)$$

Without confusion, we will use \mathcal{E}_{rl} to mean the 0-1 indicator variable of the event (defined earlier) \mathcal{E}_{rl} . Then, for even $l \geq 2$, we get

$$\begin{aligned} EX_r^l A^{q-l} &= EX_r^l A^{q-l} \mathcal{E}_{rl} + EX_r^l A^{q-l} (1 - \mathcal{E}_{rl}) \leq L_{rl} EA^{q-l} + M_{rl} EA^{q-l} (1 - \mathcal{E}_{rl}) \\ &\leq L_{rl} EA^{q-l} + M_{rl} \left(EA^{q-l+2} \right)^{\frac{q-l}{q-l+2}} (E(1 - \mathcal{E}_{rl}))^{\frac{2}{q-l+2}} \quad \text{H\"older} \\ &\leq L_{rl} EA^{q-l} + M_{rl} \delta_{rl}^{\frac{2}{m-l+2}} \left(EA^{q-l+2} \right)^{\frac{q-l}{q-l+2}} \quad \text{since } m \geq q \\ &\leq L_{rl} EA^{q-l} + \left(\hat{M}_{rl}^{\frac{2q}{l(q-l+2)}} (3m^2 n^{2/l})^{\frac{q-l}{q-l+2}} \right) \left(\frac{\hat{M}_{rl}^{\frac{(q-l)(l-2)}{l(q-l+2)}} (EA^{q-l+2})^{\frac{q-l}{q-l+2}}}{(3m^2 n^{2/l})^{\frac{(q-l)}{q-l+2}}} \right) \\ &\leq L_{rl} EA^{q-l} + \hat{M}_{rl}^{q/l} (3m^2 n^{2/l})^{\frac{q-l}{2}} + \frac{\hat{M}_{rl}^{\frac{l-2}{l}} EA^{q-l+2}}{3m^2 n^{2/l}}, \end{aligned}$$

where, in the last step, we have used Young's inequality which says that for any $a, b > 0$ real and $s, r > 0$ with $\frac{1}{s} + \frac{1}{r} = 1$, we have $ab \leq a^s + b^r$; we have applied this with $s = (q-l+2)/2$ and $r = (q-l+2)/(q-l)$.

Plugging this into (14), we get:

$$E\left(\sum_{i=1}^r X_i\right)^q \leq \sum_{\substack{l \geq 0 \\ \text{even}}}^q a_{rl}^{(q)} E\left(\sum_{i=1}^{r-1} X_i\right)^{q-l},$$

$$\begin{aligned}
a_{rl}^{(q)} &= 1 + 3 \frac{1}{3m^2n} \binom{q}{2}, & l = 0 \\
a_{rl}^{(q)} &= 3 \binom{q}{l} L_{rl} + 3 \binom{q}{l+2} \hat{M}_{r,l+2}^{l/(l+2)} \frac{1}{3m^2n^{2/(l+2)}}, & 2 \leq l \leq q-2 \\
a_{rl}^{(q)} &= 3L_{rq} + 3 \sum_{\substack{l_1 \geq 2 \\ \text{even}}}^q \binom{q}{l_1} \hat{M}_{rl_1}^{q/l_1} (3m^2)^{(q-l_1)/2} n^{(q-l_1)/l_1}, & l = q.
\end{aligned}$$

It is easy to see that

$$\begin{aligned}
a_{rl}^{(q)} &\leq a_{rl} = 1 + \frac{1}{n}, \quad l = 0 \\
a_{rl}^{(q)} &\leq a_{rl} = 3 \binom{m}{l} \left(L_{rl} + \hat{M}_{r,l+2}^{l/(l+2)} n^{-2/(l+2)} \right), \quad 2 \leq l \leq q-2 \\
a_{rq}^{(q)} &\leq \hat{a}_{rq} = 3L_{rq} + 3 \sum_{\substack{l_1 \geq 2 \\ \text{even}}}^q \binom{q}{l_1} \hat{M}_{rl_1}^{q/l_1} (3m^2)^{(q-l_1)/2} n^{(q-l_1)/l_1}.
\end{aligned}$$

It is important to make $a_{r0}^{(q)}$ not be much greater than 1 because in this case only n is reduced and so in the recurrence, this could happen n times. Note that except for $l = q$, the other a_{rl} do not depend upon q ; we have used \hat{a}_{rq} to indicate that this extra dependence. With this, we have

$$E \left(\sum_{i=1}^r X_i \right)^q \leq \hat{a}_{rq} + \sum_{\substack{l \geq 0 \\ \text{even}}}^{q-2} a_{rl} E \left(\sum_{i=1}^{r-1} X_i \right)^{q-l}.$$

We wish to solve these recurrences by induction on r, q . Intuitively, we can imagine a directed graph with root marked (n, m) . The root has $\frac{m}{2} + 1$ children which are marked $(n-1, m-l)$ for $l = 0, 2, \dots, m$; the node marked (r, q) is trying to bound $E(\sum_{i=1}^r X_i)^q$. There are also weights on the edges of a_{rl} . The graph keeps going until we reach the leaves - which are marked $(1, *)$ or $(r, 0)$. This is very similar to the recursion tree picture accompanying the proof of Theorem (1). It is intuitively easy to argue that the bound we are seeking at the root is the sum over all paths from the root to the leaves of the product of the edge weights on the path. We formalize this in a lemma.

For doing that, for $1 \leq r \leq n; 2 \leq q \leq m, q$ even and $1 \leq i \leq r$ define $S(r, q, i)$ as the set of $s = (s_i, s_{i+1}, s_{i+2}, \dots, s_r)$ with $s_i > 0; s_{i+1}, s_{i+2}, \dots, s_r \geq 0$ and $\sum_{j=i}^r s_j = q; s_j$ even.

Lemma 1. *For any $1 \leq r \leq n$ and any $q \leq m$ even, we have*

$$E \left(\sum_{i=1}^r X_i \right)^q \leq \sum_{i=1}^r \sum_{s \in S(r, q, i)} \hat{a}_{i, s_i} \prod_{j=i+1}^r a_{j, s_j}.$$

Proof Indeed, the statement is easy to prove for the base case of the induction - $r = 1$ since \mathcal{E}_{1l} is the whole sample space and $EX_1^q \leq L_{1q}$. For the inductive step, we proceed as follows.

$$\begin{aligned}
E \left(\sum_{i=1}^r X_i \right)^q &\leq \sum_{\substack{s_r \geq 0 \\ \text{even}}}^{q-2} a_{r, s_r} E \left(\sum_{i=1}^{r-1} X_i \right)^{q-s_r} + \hat{a}_{r, q} \\
&\leq \hat{a}_{r, q} + \sum_{i=1}^{r-1} \sum_{\substack{s_r \geq 0 \\ \text{even}}}^{q-2} a_{r, s_r} \sum_{s \in S(r-1, q-s_r, i)} \hat{a}_{i, s_i} \prod_{j=i+1}^{r-1} a_{j, s_j}.
\end{aligned}$$

We clearly have $S(m, q, m) = \{q\}$ and for each fixed $i, 1 \leq i \leq r-1$, there is a 1-1 map $S(r-1, q, i) \cup S(r-1, q-2, i) \cup \dots \cup S(r-1, 2, i) \rightarrow S(r, q, i)$ given by $s = (s_i, s_{i+1}, \dots, s_{r-1}) \rightarrow s' = (s_i, \dots, s_{r-1}, q - \sum_{j=i}^{r-1} s_j)$ and it is easy to see from this that we have the inductive step, finishing the proof of the Lemma. \square

The “sum of products” form in the lemma is not so convenient to work with. We will now get this to the “sum of moments” form stated in the Theorem. This will require a series of (mainly algebraic) manipulations with ample use of Young’s inequality, the inequality asserting $(a_1 + a_2 + \dots + a_r)^q \leq r^{q-1}(a_1^q + a_2^q + \dots + a_r^q)$ for positive reals a_1, a_2, \dots and $q \geq 1$ and others.

So far, we have (moving the $l = 0$ terms separately in the first step)

$$\begin{aligned}
E \left(\sum_{i=1}^n X_i \right)^m &\leq \left(\prod_{i=1}^n a_{i0} \right) \sum_{i=1}^n \sum_{s \in S(n, m, i)} \hat{a}_{i, s_i} \prod_{\substack{j=i+1 \\ s_j \neq 0}}^n a_{j, s_j} \\
&\leq 3 \sum_{i=1}^n \sum_{s \in S(n, m, i)} \hat{a}_{i, s_i} \prod_{\substack{j=i+1 \\ s_j \neq 0}}^n a_{j, s_j} \\
&\leq 3 \sum_{t \geq 1}^{m/2} \left(\sum_{i=1}^n \hat{a}_{i, 2t} \right) \sum_{s \in Q(m-2t)} \prod_{\substack{j=1 \\ s_j \neq 0}}^n a_{j, s_j}
\end{aligned} \tag{15}$$

$$\text{where, } Q(q) = \left\{ s = (s_1, s_2, \dots, s_n) : s_i \geq 0 \text{ even ; } \sum_j s_j = q \right\}$$

Fix q for now. For $s \in Q(q), l = 0, 1, 2, \dots, q/2$, let $T_l(s) = \{j : s_j = 2l\}$ and $t_l(s) = |T_l(s)|$. Note that $\sum_{l=0}^{q/2} l t_l(s) = q/2$. Call $t(s) = (t_0(s), t_1(s), t_2(s), \dots, t_{q/2}(s))$ the “signature” of s . In the special case when a_{il} is independent of i , the signature clearly determines the “ s term” in the sum (15). For the general case too, it will be useful to group terms by their signature. Let (the set of possible signatures) be T . [T consists of all $t = (t_0, t_1, t_2, \dots, t_{q/2})$ with $t_l \geq 0$ $\sum_{l=1}^{q/2} l t_l = q/2$; $\sum_{l=0}^{q/2} t_l = n$.

$$\begin{aligned}
\text{Now, } \sum_{s \in Q(q)} \prod_{\substack{j=1 \\ s_j \neq 0}}^n a_{j, s_j} &= \sum_{t \in T} \sum_{T_0, T_1, T_2, \dots, T_{q/2} : |T_l| = t_l} \prod_{l=1}^{q/2} \prod_{i \in T_l} a_{i, 2l} \\
&\leq \sum_{t \in T} \prod_{l=1}^{q/2} \frac{1}{t_l!} \left(\sum_{i=1}^n a_{i, 2l} \right)^{t_l},
\end{aligned}$$

since the expansion of $(\sum_{i=1}^n a_{i, 2l})^{t_l}$ contains $t_l!$ copies of $\prod_{i \in T_l} a_{i, 2l}$ (as well other terms we do not need.) Now define $R = \{r = (r_1, r_2, \dots, r_{q/2}) : r_l \geq 0; \sum_l r_l = q/2\}$. We have

$$\begin{aligned}
\sum_{t \in T} \prod_{l=1}^{q/2} \frac{1}{t_l!} \left(\sum_{i=1}^n a_{i, 2l} \right)^{t_l} &\leq \sum_{r \in R} \prod_l \frac{1}{(r_l/l)!} \left(\sum_{i=1}^n a_{i, 2l} \right)^{r_l/l} \\
&\leq \frac{1}{(q/2)!} \left(\sum_{l=1}^{q/2} m^{1-(1/l)} \left(\sum_i a_{i, 2l} \right)^{1/l} \right)^{q/2},
\end{aligned} \tag{16}$$

where the first inequality is seen by substituting $r_l = t_l l$ and noting that the terms corresponding to the r such that $l|r_l \forall l$ are sufficient to cover the previous expression and the other terms are

non-negative. To see the second inequality, we just expand the last expression and note that the expansion contains $\prod_l (\sum_i a_{i,2l})^{r_l/l}$ with coefficient $\binom{q/2}{r_1, r_2, \dots, r_{q/2}}$ for each $r \in R$. Now, it only remains to see that $m^{r_l(1-(1/l))} \geq \frac{r_l!}{(r_l/l)!}$, which is obvious. Thus, we have plugging in (16) into (15), (for some constant $c > 0$; recall c may stand for different constants at different points):

$$EX^m \leq c^m \sum_{t=1}^{\frac{m}{2}} \frac{3}{\left(\frac{m}{2} - t\right)!} \left(\sum_{i=1}^n \hat{a}_{i,2t} \right) \left(\sum_{l=1}^{\frac{m}{2}-t} m^{1-\frac{1}{l}} \left(\sum_i a_{i,2l} \right)^{\frac{1}{l}} \right)^{\frac{m}{2}-t}.$$

$$\text{Now, } \left(\frac{m}{2} - t\right)! \geq \left(\frac{m}{2} - t\right)^{\frac{m}{2}-t} e^{-\frac{m}{2}} e^t \geq m^{\frac{m}{2}-t} e^{-\frac{m}{2}} \text{Min}_t \left[\left(\frac{\frac{m}{2} - t}{m}\right)^{\frac{m}{2}-t} e^t \right] \geq m^{\frac{m}{2}-t} (2e)^{-\frac{m}{2}},$$

the last using Calculus to differentiate the log of the expression with respect to t to see that the min is at $t = 0$. Thus,

$$EX^m \leq c^m \sum_t \left[\left(\frac{3}{m} \sum_{l=1}^{\frac{m}{2}-t} m^{1-\frac{1}{l}} \left(\sum_i a_{i,2l} \right)^{\frac{1}{l}} \right)^{\frac{m}{2}-t} \right] \left[\sum_{i=1}^n \hat{a}_{i,2t} \right].$$

Let α, β denote the quantities in the 2 square brackets respectively. Young's inequality gives us: $\alpha\beta \leq \alpha^{m/(m-2t)} + \beta^{m/2t}$. Thus,

$$EX^m \leq \sum_{t=1}^{\frac{m}{2}} \left(\sum_i \hat{a}_{i,2t} \right)^{\frac{m}{2t}} + \sum_t \left(\sum_{l=1}^{\frac{m}{2}-t} m^{-\frac{1}{l}} \left(\sum_i a_{i,2l} \right)^{\frac{1}{l}} \right)^{\frac{m}{2}} \quad (17)$$

In what follows, let l_1 run over even values to m and i run from 1 to n .

$$\begin{aligned} \sum_{t=1}^{\frac{m}{2}} \left(\sum_i \hat{a}_{i,2t} \right)^{\frac{m}{2t}} &\leq c^m \sum_t \left(\sum_i L_{i,2t} \right)^{\frac{m}{2t}} \\ &+ c^m m^m \sum_t \left(\frac{1}{n} \sum_i \sum_{l_1 \leq 2t} \left(\frac{2t}{l_1 m} \right)^{l_1} (n \hat{M}_{i,l_1})^{2t/l_1} \right)^{\frac{m}{2t}} \leq \\ &c^m \sum_t \left(\sum_i L_{i,2t} \right)^{\frac{m}{2t}} + m^m \sum_{t, l_1} \frac{t^{\frac{m}{2t}}}{l_1^{l_1} n^{\frac{m}{2t}}} \left(\sum_i (n \hat{M}_{i,l_1})^{2t/l_1} \right)^{\frac{m}{2t}} \\ &\leq c^m \sum_t \left(\sum_i L_{i,2t} \right)^{\frac{m}{2t}} + c^m m^m \sum_{l_1} \frac{1}{n l_1^{l_1}} \sum_i (n \hat{M}_{i,l_1})^{m/l_1}, \end{aligned} \quad (18)$$

(using $t^{m/2t} \leq c^m$.)

$$\begin{aligned}
& \sum_{l=1}^{\frac{m}{2}-1} m^{-(1/l)} \left(\sum_i a_{i,2l} \right)^{\frac{1}{l}} \leq \\
& \sum_{l=1}^{\frac{m}{2}-1} m^{-\frac{1}{l}} \left(\frac{m^{2l}}{(2l)!} \right)^{\frac{1}{l}} \left(\sum_i L_{i,2l} + \frac{\hat{M}_{i,2l+2}^{\frac{l}{l+1}}}{n^{1/(l+1)}} \right)^{\frac{1}{l}} \leq \\
& m^2 \sum_{l=1}^{\frac{m}{2}-1} \frac{m^{-\frac{1}{l}}}{l^2} \left(\left(\sum_i L_{i,2l} \right)^{\frac{1}{l}} + \left(\sum_i \hat{M}_{i,2l+2} \right)^{\frac{1}{l+1}} \right) \leq \\
& m^2 \sum_{l=1}^{\frac{m}{2}} \frac{m^{-\frac{1}{l}}}{l^2} \left(\sum_i L_{i,2l} \right)^{1/l} + m^2 \sum_{l=2}^{\frac{m}{2}} \frac{1}{(l-1)^2} \left(\sum_i \hat{M}_{i,2l} \right)^{\frac{1}{l}}. \tag{19}
\end{aligned}$$

We will further bound the last term using Hölder's inequality:

$$\begin{aligned}
& \left(\sum_{l=2}^{\frac{m}{2}} \frac{\left(\sum_i \hat{M}_{i,2l} \right)^{1/l}}{(l-1)^2} \right)^{\frac{m}{2}} \leq \\
& \left(\sum_{l=1}^{\infty} \frac{1}{l^2} \right)^{(m-2)/2} \left(\sum_l \frac{1}{(l-1)^2} \left(\sum_i \hat{M}_{i,2l} \right)^{\frac{m}{2l}} \right) \\
& \leq 2^m \sum_{l=1}^{\frac{m}{2}} \frac{1}{nl^2} \sum_i (n \hat{M}_{i,2l})^{\frac{m}{2l}}. \tag{20}
\end{aligned}$$

Now plugging (19,18,20) into (17) and noting that $cm^{2-(1/l)}/l^2 \geq 1$, we get the Theorem. \square

10 Bin Packing

Now we tackle bin packing. The input consists of n i.i.d. items - $Y_1, Y_2, \dots, Y_n \in (0, 1)$. Suppose $EY_1 = \mu$ and $\mathbf{Var}Y_1 = \sigma$. Let $f = f(Y_1, Y_2, \dots, Y_n)$ be the minimum number of capacity 1 bins into which the items Y_1, Y_2, \dots, Y_n can be packed. It was shown (after many successive developments) using non-trivial bin-packing theory ([37]) that f has $N(0, n(\mu^2 + \sigma^2))$ tails upto $O(n(\mu^2 + \sigma^2))$. Talagrand [43] gives a simple proof of this from his inequality (this is the first of the six or so examples in his paper.) [We can also give a simple proof of this from our theorem.]

Talagrand [43] says (in our notation) “especially when μ is small, one expects that the behavior of f resembles the behavior of $\sum_{i=1}^n Y_i$. Thereby, one should expect that f should have tails of $N(0, n\sigma^2)$ or, at least, less ambitiously, $N(0, n(\mu^2 + \sigma^2))$ ”.

However, $N(0, n\sigma^2)$ (as for sums of independent random variables) is easily seen to be impossible. An example is when items are of size $1/k$ or $(1/k) + \epsilon$ (k a positive integer and $\epsilon \ll 1/k$ is a positive real) with probability $1/2$ each. σ is $O(\epsilon)$. It is clear that the number n_1 of $1/k$ items can be in $\frac{n}{2} \pm \Theta(\sqrt{n})$. Now, a bin can have at most $k-1$ items if it has any $(1/k) + \epsilon$ item; it can have k items if they are all $1/k$. Thus if n_1 number of $1/k$ items, we get

$$f = \frac{n_1}{k} + \frac{n - n_1}{k-1} + O(1) = \frac{n}{2} \left(\frac{1}{k} + \frac{1}{k-1} \right) \pm \frac{\sqrt{n}}{k^2}.$$

From this it can be seen that the standard deviation of f is $\Omega(\sqrt{n}\mu^2) \gg \sqrt{n}\sigma$, establishing what we want.

Here we prove the best possible interval of concentration when the items take on only one of a fixed finite set of values (discrete distributions - a case which has received much attention in the literature for example [19] and references therein). [While our proof of the upper bound here is only for problems with a fixed finite number of types, it would be nice to extend this to continuous distributions.]

Theorem 8. *Suppose Y_1, Y_2, \dots, Y_n are i.i.d. drawn from a discrete distribution with r atoms, $r \in O(1)$, each with probability at least $\frac{1}{\log n}$. Let $EY_1 = \mu \leq \frac{1}{r^2 \log n}$ and $\mathbf{Var}Y_i = \sigma^2$. Then for any $t \in (0, n(\mu^3 + \sigma^2))$, we have*

$$\Pr(|f - Ef| \geq t + r) \leq c_1 e^{-ct^2/(n(\mu^3 + \sigma^2))}.$$

Further, there is distribution for Y_i in which $\mathbf{Var}(f) \in \Omega(n(\mu^3 + \sigma^2))$.

Proof Let item sizes be $\zeta_1, \zeta_2, \dots, \zeta_j \dots \zeta_r$ and the probability of picking type j be p_j . [We will reserve j to denote the j th item size.] We have : mean $\mu = \sum_j p_j \zeta_j$ and standard deviation $\sigma = (\sum_j p_j (\zeta_j - \mu)^2)^{1/2}$.

Note that if $\mu \leq r/\sqrt{n}$, then earlier results already give concentration in an interval of length $O(\sqrt{n}(\mu + \sigma))$ which is then $O(r + \sigma)$, so there is nothing to prove. So assume that $\mu \geq r/\sqrt{n}$.

Define a ‘‘bin Type’’ as an r - vector of non-negative integers specifying number of items of each type which are together packable into one bin. If bin type i packs a_{ij} items of type j for $j = 1, 2, \dots, r$ we have $\sum_j a_{ij} \zeta_j \leq 1$. Note that s , the number of bin types depends only on ζ_j , not on n .

For any set of given items, we may write a Linear Programming relaxation of the bin packing problem whose answers are within additive error r of the integer solution. If there are n_j items of size ζ_j in the set, the Linear program, which we call ‘‘Primal’’ (since later we will take its dual) is :

Primal : (x_i number of bins of type i .)

$$\text{Min} \sum_{i=1}^s x_i \quad \text{subject to} \quad \sum_{i=1}^s x_i a_{ij} \geq n_j \forall j ; x_i \geq 0.$$

Since an optimal basic feasible solution has at most r non-zero variables, we may just round these r up to integers to get an integer solution; thus the additive error is at most r as claimed. In what follows, we prove concentration not for the integer program’s value, but for the value of the Linear Program. The Linear Program has the following dual :

$$\text{MAX} \sum_{j=1}^r n_j y_j \quad \text{s.t.} \quad \sum_j a_{ij} y_j \leq 1 \text{ for } i = 1, 2, \dots, s; y_j \geq 0.$$

(y_j may be interpreted as the ‘‘imputed’’ size of item j) Let $Y = (Y_1, Y_2, \dots, Y_n)$ and (for an i we fix attention on) $Y' = (Y_1, Y_2, \dots, Y_{i-1}, Y_{i+1}, \dots, Y_n)$. We denote by $f(Y)$ the value of the Linear Program for the set of items Y . Let $\Delta_i = f(Y) - f(Y')$. The typical events \mathcal{E}_i will just be that the number of copies of each ζ_j among Y_1, Y_2, \dots, Y_{i-1} is close to its expectation:

$$\mathcal{E}_i : \forall j, |\text{no.of copies of } \zeta_j \text{ in } Y_1, Y_2, \dots, Y_{i-1} - (i-1)p_j| \leq 100 \sqrt{m \ln \frac{10m}{\mu} p_j (i-1)},$$

where m is to be specified later, but will satisfy $\frac{1}{10}n(\mu^3 + \sigma^2)$. We will use the Theorem with this parameter m .

We will make crucial use of the fact that second moments count highly for the bound in the theorem. So the main technical part of the proof is the following Lemma bounding typical conditional second moments.

Lemma 2. *Under \mathcal{E}_i , $\mathbf{Var}(\Delta_i|Y_1, Y_2, \dots, Y_{i-1}) \in O(\mu^3 + \sigma^2)$.*

Proof Suppose now, we have already chosen all but Y_i . Now, we pick Y_i at random; say $Y_i = \zeta_k$. Let $Y = (Y_1, Y_2, \dots, Y_n)$ and $Y' = (Y_1, Y_2, \dots, Y_{i-1}, Y_{i+1}, \dots, Y_n)$

Let

$$\Delta_i = f(Y) - f(Y').$$

Suppose we have the optimal solution of the LP for Y' . There is a bin type which packs $\lfloor 1/\zeta_k \rfloor$ copies of item of type k ; let i_0 be the index of this bin type. Clearly if we increase x_{i_0} by $\frac{1}{\lfloor 1/\zeta_k \rfloor}$, we get a feasible solution to the new primal LP for Y . So $0 \leq \Delta_i \leq \frac{1}{\lfloor 1/\zeta_k \rfloor} \leq \zeta_k + 2\zeta_k^2$, which implies

$$\begin{aligned} E(\Delta_i^2|Y') &\leq \sum_j p_j (\zeta_j + 2\zeta_j^2)^2 \leq \sum_j p_j \zeta_j^2 + 8 \sum_j p_j \zeta_j^3 \\ &\leq \mu^2 + \sigma^2 + 8 \times 8 \sum_j p_j |\zeta_j - \mu|^3 + 8 \times 8 \sum_j p_j \mu^3 \leq \mu^2 + 65\sigma^2 + 64\mu^3. \end{aligned} \quad (21)$$

Now, we lower bound Δ_i by looking at the dual. For this, let y be the dual optimal solution for Y' . (Note : Thus, $y = y(Y')$ is a function of Y' .) y is feasible to the new dual LP too (after adding in Y_i), since the dual constraints do not change. So, we get: $\Delta_i \geq y_k$.

$$E(\Delta_i|Y') \geq \sum_j p_j y_j(Y'). \quad (22)$$

Also, recalling the bin type i_0 defined earlier, we see that $y_k \leq 1/\lfloor (1/\zeta_k) \rfloor \leq \zeta_k + 2\zeta_k^2$. Say the number of items of type j in Y' is $(n-1)p_j + \gamma_j$. It is easy to see that ζ is a feasible dual solution. Since y is an optimal solution, we have

$$\sum_j ((n-1)p_j + \gamma_j)y_j \geq \sum_j ((n-1)p_j + \gamma_j)\zeta_j.$$

$$\begin{aligned} \mu - \sum_j p_j y_j &= \sum_j p_j (\zeta_j - y_j) = \frac{1}{n-1} \left(\sum_j ((n-1)p_j + \gamma_j)(\zeta_j - y_j) \right) + \frac{1}{n-1} \sum_j \gamma_j (y_j - \zeta_j) \\ &\leq \frac{1}{n-1} \sum_j \gamma_j (y_j - \zeta_j) \\ &\leq \frac{1}{n-1} \left(\sum_j (\gamma_j^2/p_j) \right)^{1/2} \left(\sum_j p_j (y_j - \zeta_j)^2 \right)^{1/2} \\ &\leq \frac{32(\mu + \sigma)r}{n} \text{MAX}_j |\gamma_j/\sqrt{p_j}|, \end{aligned} \quad (23)$$

where we have used the fact that $-\zeta_j \leq y_j - \zeta_j \leq 2\zeta_j^2 \leq 2\zeta_j$. Let $(i-1)p_j + \gamma'_j$ and $(n-i)p_j + \gamma''_j$ respectively be the number of items of size ζ_j among Y_1, Y_2, \dots, Y_{i-1} and Y_{i+1}, \dots, Y_n . Since γ''_j is the

sum of $n - i$ i.i.d. random variables, each taking on value $-p_j$ with probability $1 - p_j$ and $1 - p_j$ with probability p_j , we have $E(\gamma_j'')^2 = \mathbf{Var}(\gamma_j'') \leq np_j$. Now, we wish to bound the conditional moment of γ_j' conditioned on Y_1, Y_2, \dots, Y_{i-1} . But under the worst-conditioning, this can be very high. [For example, all fractions upto $i - 1$ could be of the same type.] Here we exploit the typical case conditioning. The expected number of “successes” in the $i - 1$ Bernoulli trials is $p_j(i - 1)$. By using Chernoff, we get (recall the definition of \mathcal{E}_i) $\Pr(\neg\mathcal{E}_i) = (\text{say}) \delta_i \leq \mu^{4m} m^{-4m}$. Using (22) and (23), we get

$$\begin{aligned} & E(\Delta_i | Y_1, Y_2, \dots, Y_{i-1}; \mathcal{E}_i) \\ & \geq \mu - \frac{32(\mu + \sigma)r}{n} E\left(\max_j \frac{1}{\sqrt{p_j}} (100\sqrt{m \ln(10m/\mu)p_j(i-1)} + (E(\gamma_j'')^2)^{1/2})\right) \\ & \geq \mu - c(\mu + \sigma)^{5/2} r \sqrt{\ln(10m/\mu)} - \frac{c\mu r}{\sqrt{n}} \geq \mu - O(\mu^2), \end{aligned}$$

using $m \leq \frac{1}{10}n(\mu^3 + \sigma^2)$. So, we get recalling (21),

$$\mathbf{Var}(\Delta_i | Y_1, Y_2, \dots, Y_{i-1}; \mathcal{E}_i) = E(\Delta_i^2 | Y_1, Y_2, \dots, Y_{i-1}; \mathcal{E}_i) - (E(\Delta_i | Y_1, Y_2, \dots, Y_{i-1}; \mathcal{E}_i))^2 \leq c(\mu^3 + \sigma^2),$$

using $\frac{r}{\sqrt{n}} \leq \mu \leq \frac{1}{r^2 \log n}$. This completes the proof of the Lemma. \square

Now, we have for the worst-case conditioning,

$$\mathbf{Var}(\Delta_i | Y_1, Y_2, \dots, Y_{i-1}) \leq E(\Delta_i^2 | Y_1, Y_2, \dots, Y_{i-1}) \leq c\mu^2.$$

We now appeal to (8) to see that these also give upper bounds on $\mathbf{Var}(X_i)$. As promised, dealing with higher moments is easy: note that $|\Delta_i| \leq 1$ implies that $L_{i,2l} \leq L_{i,2}$. Now to apply the Theorem, we have $L_{i,2l} \leq c(\mu^3 + \sigma^2)$. So the “ L terms” are bounded as follows :

$$\sum_{l=1}^{m/2} \frac{m^{1-(1/l)}}{l^2} \left(\sum_{i=1}^n L_{i,2l} \right)^{1/l} \leq \sum_{l=1}^{m/2} \frac{m}{l^2} \left(\frac{cn(\mu^3 + \sigma^2)}{m} \right)^{1/l} \leq cn(\mu^3 + \sigma^2)$$

noting that $m \leq n(\mu^3 + \sigma^2)$ implies that the maximum of $((n/m)(\mu^3 + \sigma^2))^{1/l}$ is attained at $l = 1$ and also that $\sum_l (1/l^2) \leq 2$. Now, we work on the M terms in the Theorem. $\max_i \delta_i \leq \mu^{4m} m^{-4m} = \delta^*$ (say).

$$\sum_{l=1}^{m/2} (1/n) \sum_{i=1}^n (n\hat{M}_{i,2l})^{m/2l} = \sum_{l=1}^{m/2} e^{h(l)},$$

where $h(l) = \frac{m}{2l} \log n + \frac{m}{l(m-2l+2)} \log \delta^*$. We have $h'(l) = -\frac{m}{2l^2} \log n - \log \delta^* \frac{m(m-4l+2)}{l^2(m-2l+2)^2}$. Thus for $l \geq (m/4) + (1/2)$, $h'(l) \leq 0$ and so $h(l)$ is decreasing. Now for $l < (m/4) + (1/2)$, we have $\frac{m}{2l^2} \log n \geq -(\log \delta^*) \frac{m(m-4l+2)}{l^2(m-2l+2)^2}$, so again $h'(l) \leq 0$. Thus, $h(l)$ attains its maximum at $l = 1$, so $(36m)^{m+2} \sum_{l=1}^{m/2} e^{h(l)} \leq m(36m)^{m+3} n^{m/2} \delta^*$ giving us $(36m)^{m+2} \sum_{l=1}^{m/2} (n\hat{M}_{i,2l}^*)^{m/2l} \leq (cnm(\mu^3 + \sigma^2))^{m/2}$. Thus we get from the Main Theorem that $E(f - Ef)^m \leq (cmn(\mu^3 + \sigma^2))^{\frac{m}{2}}$, from which Theorem (8) follows by the choice of $m = \lfloor \frac{t^2}{c_5 n(\mu^3 + \sigma^2)} \rfloor$.

10.1 Lower Bound on Spread for Bin Packing

This section proves the last statement in the theorem. Suppose the distribution is :

$$\Pr\left(Y_1 = \frac{k-1}{k(k-2)}\right) = \frac{k-2}{k-1}; \quad \Pr\left(Y_1 = \frac{1}{k}\right) = \frac{1}{k-1}.$$

This is a “perfectly packable distribution” (well-studied class of special distributions) ($k - 2$ of the large items and 1 of the small one pack.) Also, σ is small. But we can have number of $1/k$ items equal to $\frac{n}{k-1} - c\sqrt{\frac{n}{k}}$. Number of bins required $\geq \sum_i X_i = \frac{n}{k} + \frac{n}{k(k-1)} + c\sqrt{\frac{n}{k}} \left(\frac{1}{k} \left(\frac{k-1}{k-2} - 1 \right) \right) \geq \frac{n}{k-1}$. So at least $c\sqrt{\frac{n}{k}}$ bins contain only $(k - 1)/k(k - 2)$ sized items (the big items). The gap in each such bin is at least $1/k$ for a total gap of $\Omega(\sqrt{n}/k^{3/2})$. On the other hand, if the number of small items is at least $n/(k - 1)$, then each bin except two is perfectly fillable.

11 Longest Increasing Subsequence

Let Y_1, Y_2, \dots, Y_n be i.i.d., each distributed uniformly in $[0, 1]$. We consider here $f(Y) =$ the length of the longest increasing subsequence (LIS) of Y . This is a well-studied problem. It is known that $Ef = (2 + o(1))\sqrt{n}$ (see for example [5]). Since changing one Y_i changes f by at most 1, traditional H-A yields $N(0, n)$ tails which is not so interesting. Frieze [23] gave a clever argument (using a technique Steele [41] calls “flipping”) to show concentration in intervals of length $n^{1/3}$. Talagrand [43] gave the first (very simple) proof of $N(0, \sqrt{n})$ tails. Here, we also supply a (fairly simple) proof from Theorem (7) of $N(0, \sqrt{n})$ tails. [But by now better intervals of concentration, namely $O(n^{1/6})$ are known, using detailed arguments specific to this problem [9].] Our argument follows from two claims below. Call Y_i essential for Y if Y_i belongs to every LIS of Y (equivalently, $f(Y \setminus Y_i) = f(Y) - 1$.) Fix Y_1, Y_2, \dots, Y_{i-1} and for $j \geq i$, let $a_j = \Pr(Y_j \text{ is essential for } Y | Y_1, Y_2, \dots, Y_{i-1})$

Claim 3. a_i, a_{i+1}, \dots, a_n form a non-decreasing sequence.

Proof Let $j \geq i$. Consider a point ω in the sample space where Y_j is essential, but Y_{j+1} is not. Map ω onto ω' by swapping the values of Y_j and Y_{j+1} ; this is clearly a 1-1 measure preserving map. If θ is a LIS of ω with $j \in \theta, j + 1 \notin \theta$, then $\theta \setminus j \cup j + 1$ is an increasing sequence in ω' ; so $f(\omega') \geq f(\omega)$. If $f(\omega') = f(\omega) + 1$, then an LIS α of ω' must contain both j and $j + 1$ and so contains no k such that Y_k is between Y_j, Y_{j+1} . Now $\alpha \setminus j$ is an LIS of ω contradicting the assumption that j is essential for ω . So $f(\omega') = f(\omega)$. So, $j + 1$ is essential for ω' and j is not. So, $a_j \leq a_{j+1}$. \square

Claim 4. $a_i \leq c/\sqrt{n - i + 1}$.

Proof $a_i \leq \frac{1}{n-i+1} \sum_{j \geq i} a_j$. Now $\sum_{j \geq i} a_j = a$ (say) is the expected number of essential elements among Y_i, \dots, Y_n which is clearly at most $Ef(Y_i, Y_{i+1}, \dots, Y_n) \leq c\sqrt{n - i + 1}$, so the claim follows. \square

Δ_i is a 0-1 random variable with $E(\Delta_i | Y_1, Y_2, \dots, Y_{i-1}) \leq c/\sqrt{n - i + 1}$. Thus it follows (using (8) of section (3)) that

$$E(X_i^2 | Y_1, Y_2, \dots, Y_{i-1}) \leq c/\sqrt{n - i + 1}.$$

Clearly, $E(X_i^l | Y_1, Y_2, \dots, Y_{i-1}) \leq E(X_i^2 | Y_1, Y_2, \dots, Y_{i-1})$ for $l \geq 2$, even. Thus we may apply the main Theorem with \mathcal{E}_{il} equal to the whole sample space. Assuming $p \leq \sqrt{n}$, we see that (using $\sum_l (1/l^2) = O(1)$)

$$E(f - Ef)^p \leq (c_1 p)^{(p/2)+2} n^{p/4},$$

from which one can derive the asserted sub-Gaussian bounds.

12 Number of Triangles in a random graph

Let $f = f(G(n, p))$ be the number of triangles in the random graph $G(n, p)$, where each edge is independently put in with probability p . There has been much work on the concentration of f .

[33], [45] discuss in detail why Talagrand's inequality cannot prove good concentration when p the edge probability is $o(1)$. [But we assume that $np \geq 1$, so that $Ef = O(n^3p^3)$ is $\Omega(1)$.] It is known (by a simple calculation - see [26]) that

$$\mathbf{Var} f = O(\text{MAX}(n^3p^3, n^4p^5)).$$

Our main result here is that f has $N(0, \mathbf{Var} f)$ tails upto $O^*((np)^{7/4})$, where, as usual, the $*$ hides log factors. By a simple example, we see that f does not have $N(0, \mathbf{Var} f)$ tails beyond $(np)^{9/4}$. We note that our result is the first sub-Gaussian tail bound (with the correct variance) for the case when $p \leq 1/\sqrt{n}$. [For the easier case when $p = n^{-\alpha}$, $\alpha < 1/2$, such a tail bound was known [45], but only upto $(np)^\epsilon$ for a small $\epsilon > 0$.]

The most popular question about concentration of f has been to prove upper bounds on $\Pr(f \geq (1 + \epsilon)Ef)$ for essentially $\epsilon \in \Omega(1)$ (see [33], [27]), i.e., for deviations as large as $\Omega(Ef)$. In a culmination of this line of work, [28] have proved that

$$\Pr(f \geq (1 + \epsilon)Ef) \leq ce^{-c\epsilon^2 n^2 p^2}.$$

This is a special case of their theorem on the number of copies of any fixed graph in $G_{n,p}$. Their main focus is large deviations, but for general t , putting $\epsilon = t/n^3p^3$ would only give us $e^{-t^2/(n^4p^4)}$. Also, [33] develops a concentration inequality specially for polynomial functions of independent bounded random variables and [45] develops and surveys many applications of this inequalities; [45] discusses the concentration of the number of triangles as the ‘‘principal example’’.

Theorem 9. f has $N(0, \mathbf{Var} f)$ tails upto $O^*((np)^{7/4})$.

Proof Let Y_i be the set of neighbors of vertex i among $[i - 1]$ and imagine adding the Y_i in order. [This is often called the vertex-exposure Martingale.] We will also let Y_{ij} be the 0-1 variable denoting whether there is an edge between i and j for $j < i$. The number of triangles f can be written as $f = \sum_{i>j>k} Y_{ij}Y_{jk}Y_{ik}$.

As usual consider the Doob Martingale difference sequence

$$X_i = E(f|Y_1, Y_2, \dots, Y_i) - E(f|Y_1, Y_2, \dots, Y_{i-1}).$$

It is easy to see that

$$X_i = \sum_{j<k \in [i-1]} Y_{jk}(Y_{ij}Y_{ik} - p^2) + (n-i)p^2 \sum_{j<i} (Y_{ij} - p) = X_{i,1} + X_{i,2} \text{ (say).}$$

Let E^i denote $E(\cdot|Y_1, Y_2, \dots, Y_{i-1})$. We will be applying our main concentration inequality Theorem (7) with $m = O(t^2/\mathbf{Var} f)$. Let q be any even integer between 2 and m . $E^i(X_i^q) \leq 2^q E^i(X_{i,1}^q) + 2^q E^i(X_{i,2}^q)$. Of the two, it is much easier to deal with $X_{i,2}$. Indeed we have using Corollary (2).

$$E^i(X_{i,2}^q) \leq c^q n^q p^{2q} (npq)^{q/2} \leq (cn^3 p^5 q)^{q/2}. \tag{24}$$

Let \mathcal{E}_i be the event: (recall, as always, c stands for $\text{poly}(\log n)$ and may have different values in different places)

$$\begin{aligned} \mathcal{E}_i : & |Y_j| \leq cnp \text{ for } j \leq i \\ \forall S \subseteq [i-1], \text{ with } |S| \leq cnp, \text{ we have } & \sum_{j,k \in S} Y_{jk} \leq \max(cn^2 p^3, cnp) \end{aligned}$$

Now,

$$E^i(X_{i,1}^2) = \sum_{j_1 < k_1 < i} \sum_{j_2 < k_2 < i} Y_{j_1 k_1} Y_{j_2 k_2} E(Y_{i j_1} Y_{i k_1} - p^2)(Y_{i j_2} Y_{i k_2} - p^2).$$

Only terms where there are 2 or 3 distinct vertices among j_1, j_2, k_1, k_2 contribute to the expectation. The number of terms with 2 distinct vertices (and thus only one edge in $[i-1]$) is at most $n^2 p$ under \mathcal{E}_i and $E(Y_{i j_1} Y_{i k_1} - p^2)^2 \leq p^2$, so the contribution of these terms is $O(n^2 p^3)$. If there are 3 distinct vertices, we have a path of length 2 in $[i-1]$; there are $n^2 p$ choices for the first edge of the path and np choices of second edge under \mathcal{E}_i ; finally, we have $|E(Y_{i j_1} Y_{i k_1} - p^2)(Y_{i j_2} Y_{i k_2} - p^2)| = O(p^3)$; so the total of these terms is $O(n^3 p^5)$. Thus, we have

$$E^i(X_{i,1}^2 | \mathcal{E}_i) \leq \frac{c \mathbf{Var} f}{n}.$$

Further, under \mathcal{E}_i , $|X_{i,1}| \leq a$, where, $a = \max(cn^2 p^3, cnp)$ so we have for any even $l \geq 2$, $E^i X_{i,1}^l \leq \mathbf{Var} f a^{l-2}/n$. We note also that $E^i(X_{i,2}^q) \leq \mathbf{Var} f a^{l-2}/n$, since $q \leq m \leq O^*(\sqrt{np})$ as is easy to see. Plugging these bounds into the “ L terms” of theorem (7), we get

$$\sum_{l=1}^{m/2} \frac{1}{l^2} \left(\frac{\sum_i L_{i,2l}}{m} \right)^{1/l} \leq ca^2 \sum_l \frac{1}{l^2} \left(\frac{\mathbf{Var} f}{a^2 m} \right)^{1/l}.$$

Since by the choice of m , we have $ma^2 < \mathbf{Var} f$, the maximum of $\left(\frac{\mathbf{Var} f}{a^2 m} \right)^{1/l}$ is attained at $l = 1$. Also $\sum_l (1/l^2) \in O(1)$. So, we have

$$m^m \left(\sum_{l=1}^{m/2} \frac{1}{l^2} \left(\frac{\sum_i L_{i,2l}}{m} \right)^{1/l} \right)^{m/2} \leq (m \mathbf{Var} f)^{m/2}. \quad (25)$$

Now, we bound the M terms. Since the expected number of edges within a particular $S \subseteq [i-1]$ with $|S| \leq cnp$ is $O(n^2 p^3)$, the probability that there are more than $\max(cn^2 p^3, cnp)$ edges is most e^{-cnp} for a particular S . Since there are at most $np \binom{n}{np}$ S ’s to consider, union bound gives us:

$$\delta_i = \Pr(-\mathcal{E}_i) \leq e^{-cnp}.$$

We use a crude bound of $|X_i| \leq n^2$ to get $M_{i,l} \leq n^{2l}$. So,

$$(cm)^m \sum_{l=1}^{m/2} \frac{1}{n l^2} \sum_{i=1}^n \left(n \hat{M}_{i,2l} \right)^{m/2l} \leq (cm)^m \sum_l (1/l^2) n^{m/2l} n^{2m} e^{-cnp/m}.$$

Again, it is easy to see that $m \leq O^*(\sqrt{np})$; so the above is at most $(cm \mathbf{Var} f)^{m/2}$. Together with the bound on the L - terms, we now have

$$E(f - Ef)^m \leq (cm \mathbf{Var} f)^{m/2},$$

from which the tail bound follows by using Markov as before.

Remark 9. *It is easy to see that we do not have $N(0, \mathbf{Var} f)$ tails beyond $(np)^{9/4}$: just take a random $G(n, p)$. Now add all the $(np)^{3/2}$ edges among the first $(np)^{3/4}$ vertices; the probability of all these edges being present is $e^{-c(np)^{3/2}}$ which is $e^{-t^2/n^3 p^3}$, where the deviation t from Ef is $(np)^{9/4}$, namely the triangles among the first $(np)^{3/4}$ vertices.*

Remark 10. *The inequalities in [33] and [45] bound tails of polynomial functions of independent variables; the papers give many applications of them. Since most of the situations considered here are not polynomial functions, these are not applicable. But number of triangles is a polynomial of degree 3 in the underlying variables Y_{ij} and so the main theorem of [45] (Theorem (4.2)) and Corollaries do apply. In that theorem, we have to choose $\tilde{k} = 2$ or 3 and it is easy to see that with the conditions, we only get a tail bound which falls as $e^{-\dots t}$ and not t^2 as required for sub-Gaussian bounds.*

13 Questions

Many interesting open questions remain. Since the TSP is a classic problem, it would be interesting to strengthen/generalize results for the TSP. The first is to assume more limited independence: if one divides the unit square into l pieces which have Y_1, Y_2, \dots, Y_l as the set of points inside each respectively, can we prove concentration when $l \in o(n)$ and $E|Y_i| = n/l$ and assuming some moment conditions. Then, we have the question of extending concentration results under “bursts in space” to 3 and higher dimensions and finally, there are many other combinatorial problems [41] for which it would be interesting to prove such results.

We have not dealt much with “bursts in time”, but the theorems here would seem to be applicable to such situations. In the bin-packing problem, it would be natural to assume that at each time i , one first picks the number of items which would arrive at that time and then have the items pick either adversarially or stochastically their sizes and prove concentration for the minimum number of bins. On-line versions of this problem are of interest. Queueing Theory has many examples of handling bursts and it remains to be seen how the results here may help in that area.

The count of the number of not only triangles, but also other fixed graphs has been well-studied, but only for large deviations of the order of the expectation. It would be interesting to establish sub-Gaussian bounds as done here for triangles. This has some relation to the study of clustering coefficients and local communities in large (web-like) graphs.

14 Comparisons with other inequalities

The main purpose of this paper was to formulate and prove general probability inequalities which can be used to tackle the complicated combinatorial and other examples discussed. Here, we will compare our inequality to some others in the literature. For this we consider basic situations rather than complex ones to illustrate things better.

The “sub-Gaussian” behavior - $e^{-t^2 \dots}$ with the “correct” variance (for example in Theorem (1) and Corollary (2)) needs that the exponent of m in the upper bound in Theorem (1) be $\frac{m}{2}$. Moment inequalities are of course well-studied and there are many sophisticated developments. One type of inequality is the Rosenthal type inequalities [16] which assert for Martingale difference sequence X_1, X_2, \dots, X_n and even integer m :

$$E \left(\sum_{i=1}^n X_i \right)^m \leq f(m) \left(E \left(\sum_i E(X_i^2 | X_1, X_2, \dots, X_{i-1}) \right)^{m/2} + E \max_i X_i^m \right).$$

Here, $f(m)$ has to be at least $cm/\ln m$ as shown by a simple example of [30], which means that we cannot get sub-Gaussian bounds from these inequalities. The example is: The X_i are i.i.d. Bernoulli random variables with $X_i = 1 - (1/n)$ with probability $1/n$ and $-1/n$ with probability $1 - (1/n)$ and $n = cm/\ln m$. For this, we have $E(\sum_{i=1}^n X_i)^m \geq n^m (1 - (1/n))^m \Pr(X_i = (1 - (1/n)) \forall i) \geq n^{m-o(m)}$.

Our Theorem (1) can tackle the example: Note that for $l \geq \sqrt{\ln m}$, we have $(n/m)^{(l/2)-1}l! \geq 1$ and since $EX_i^l \leq 1/n$, the hypothesis of our Theorem (1) is satisfied. For $l \leq \sqrt{\ln m}$, we see that $(n/m)^{(l/2)-1}l! \geq 1/n$ and this also suffices. So, our Theorem yields $E(\sum_i X_i)^m \leq (nm)^{m/2}$. But the example proves that $f(m) \geq (cm/\ln m)^m$.

Another class of inequalities are the Burkholder [15] type inequalities which assert

$$E(X_1 + X_2 + \dots + X_n)^m \leq g(m)E(X_1^2 + X_2^2 + \dots + X_n^2)^{m/2},$$

for even integers m when X_i are Martingale differences. Here, since the right hand side involves taking the expectation of a power of the sum of n quantities, we only gain if we could argue (in essence) that not many of them can be simultaneously high. Indeed, if we do not have any such information, then the best we might say is $X_1^2 + X_2^2 + \dots + X_n^2 \leq n \max_i |X_i|^2$, which only bounds the r.h.s. by $g(m)n^{m/2}E \max_i X_i^m$ and since it is known that $g(m)$ has to be at least $(cm)^{m/2}$, this does not give as strong results as Theorem (1). [The fact that $g(m) \geq (cm)^{m/2}$ follows from the simple example when X_i are i.i.d., each equal to ± 1 with probability $1/2$ each.] But, here is a simple natural example where Burkholder inequality provably cannot derive something as strong as Theorem (1): let Z_i be i.i.d., each Poisson with mean 1 and let $X_i = \pm Z_i, i = 1, 2, \dots, n$, with probability $1/2$ each, so $EX_i = 0$. It is well-known that for even l , $EX_i^l = EZ_i^l = (cl)^l$, where c here (and the rest of this section) involves constant and logarithmic (in l) factors. Theorem (1) directly yields $N(0, n)$ tails for $X = \sum_{i=1}^n X_i$ upto n . But to apply Burkholder, we must deal with $E(\sum_i X_i^2)^{m/2}$ for even m . We have

$$E\left(\sum_i X_i^2\right)^{m/2} \geq \binom{n}{m/2} (m/2)! (EX_1^2)^{m/2} + nEX_1^m \geq (cn)^{m/2} + (cm)^m.$$

So, the best one can ever prove is $EX^m \leq (cnm)^{m/2} + (cm)^{3m/2}$. Consider a tail probability $\Pr(|X| \geq t)$; the best we could get for this from Burkholder type inequalities is

$$\Pr(|X| \geq t) \leq \frac{(cm)^{3m/2}}{t^m} + \frac{(cnm)^{m/2}}{t^m}.$$

The minimum value of $(cm)^{3m/2}/t^m$ is easily seen by Calculus to be $e^{-ct^{2/3}}$ and when $n^{3/4} \in o(t)$, we have $t^{2/3} \in o(t^2/n)$, so we do not get $N(0, n)$ tails beyond $n^{3/4}$. One can ask if this is a cooked up example. But it occurs naturally - in many geometric probability results for example, where, n i.i.d. points are picked uniformly from the unit square, it turns out that the ‘‘Poisson approximation’’ where instead one runs a Poisson process of intensity n to get the points is more useful since, then, points in non-intersecting regions of the square are independent [4]. In this process, clearly the number of points in any region of area $1/n$ is Poisson with mean 1 and indeed, in our TSP and minimum weight spanning tree analysis, we used a generalization of this, allowing longer tails and dependence for the generation process and were still able to use Theorem (1).

The author has received many queries about how particular inequalities (the literature is clearly rich in this area with a number of clever papers, a majority appearing in the venerable journal: *Annals of Probability*) compares to the theorems here. An exhaustive comparison with each inequality in the literature would not be possible. But some more comparisons are given here. We consider three particular corollaries of our theorems - Generalized Chernoff bounds (GC) (Corollary (2), Remarks (4 and 8)), H-A and the Poisson example above. Our theorems can derive tail bounds for all of these.

A recent result on the line of Burkholder inequalities is for example, one in [36], which asserts that

$$E(X_1 + X_2 + \dots + X_n)^m \leq (cmn)^{m/2} EX_1^m,$$

where m is again even and X_i are now stationary Martingale differences. The $m^{m/2}$ is promising for getting sub-Gaussian bounds, but the high moment EX_1^m on the right hand side means that Chernoff bounds don't follow from this. On the other hand, for stationary martingale differences, this is a strengthening of H-A. Talagrand's inequality can of course derive Chernoff bounds, but it only applies to independent random variables and so cannot derive H-A or GC. There are also inequalities based on the beautiful technique of Decoupling, for example Theorems 1.2A to 1.5B of [18]. This works only for Martingale differences, requires bounds similar to our theorem (1), but for all moments, not just up to an m precluding our Corollary (2) and all other applications assuming only finite moments. But, we note that this does tackle the Poisson example and indeed, our theorem (1) is close in spirit to this, as discussed in Remark (3). [Needless to add all the inequalities mentioned have their virtues which for want of space, we do not describe.] Here is a little table summarizing these comparisons.

	Poiss	H-A	GC
THM 1	Yes	Yes	Yes
Rosent	X	X	X
Burkh	X	Yes	Yes
Decoup	Yes	Yes	X
Talag	???	X	X

Legend: Poiss - the Poisson example above. GC - Generalized Chernoff.

Our crucial advantage is that while earlier moment inequalities generally do not focus on differentiating between the coefficients of different moments, the current paper pays particular attention to the terms involving different moments. We are able to get a smaller coefficient on the higher moments which thus matter less; this is helpful, since lower moments are easier to bound tightly. This enables us to get the sub-Gaussian tails in the combinatorial situations discussed, whereas traditional inequalities do not get such bounds. It is worth noting that if we settle for an extra factor of $m^{m/2}$ in the bounds of our Theorem (7) (thus abandoning correct Gaussian tails) and also restrict only to Martingale differences instead of (1), then Burkholder's inequality would imply the theorem.

Another family of inequalities are the Efron-Stein type inequalities. A recent result of Boucheron, Bousquet, Lugosi and Massart [13] proves concentration for a real-valued function F of independent random variables Y_1, Y_2, \dots, Y_n . Let $Z = F(Y_1, Y_2, \dots, Y_n)$ and suppose functions $Z_i = Z_i(Y_1, Y_2, \dots, Y_{i-1}, Y_{i+1}, \dots, Y_n)$ are arbitrary functions. Their main theorem is that

$$E((Z - EZ)_+)^m \leq (cm)^{m/2} E \left(\sum_{i=1}^n (Z - Z_i)^2 \right)^{m/2}. \quad (26)$$

[In the setting of independent random variables, this is in a way similar to Burkholder.]

Here, again, we sum up the variations in Z caused by all the n variables and then take a high moment of it. The advantage of this would be in situations where one can show that not too many of individual Y_i cause large changes for typical Y_1, Y_2, \dots, Y_n . [See [13].] This general line of approach is also reminiscent of Talagrand's inequality; but Talagrand allows simultaneous change of variables. Note that (26) has an exponent of $m/2$ on the m which can lead to the ideal sub-Gaussian behavior. In contrast, our inequality (like Rosenthal's) only considers variations of one individual variable at a time which is in many cases easier to bound. We saw this in the case of Bin-Packing, coloring and other examples. Even for the classical Longest Increasing Subsequence (LIS) problem, where for example, Talagrand's crucial argument is that only a small number $O(\sqrt{n})$ of elements (namely those in the current LIS) cause a decrease in the length of the LIS by their deletion, we

are able to bound individual variations (in essence arguing that EACH variable has roughly only a $O(1/\sqrt{n})$ probability of changing the length of the LIS sufficiently to get a concentration result.

Note that if one can only handle individual variations, then (26) again essentially yields only

$$E((Z - EZ)_+)^m \leq (cmn)^{m/2} \max_i E(Z - Z_i)^m.$$

In this case, arguments as in Theorem (1) as well as what we do for Bin-Packing and LIS which is based mainly on the second moment, do not work, since the above involves a high moment. There are many other specialized ingenious probability inequalities in the literature; we have only touched upon general ones.

Besides the situation like JL theorem, the Strong Negative correlation condition is also satisfied by the so-called “negatively associated” random variables ([29],[21], [14] for example). Variables in occupancy (balls and bins) problems, 0-1 variables produced by a randomized rounding algorithm of Srinivasan [40] etc. are negatively associated.

Acknowledgements Thanks to David Aldous, Alesandro Arlotto, Alan Frieze, Svante Janson, Manjunath Krishnapur, Claire Mathieu, Assaf Naor, Yuval Peres and Mike Steele, for helpful discussions.

References

- [1] D. Achlioptas, Database friendly random projections, Proc. Principles of Database systems (PODS) 274-281 (2001).
- [2] D. Achlioptas and A. Naor, The two possible values of the chromatic number of a random graph. Dimitris Achlioptas , Assaf Naor . Ann. of Math. (2) 162 (2005), no. 3, 1335–1351.
- [3] N. Ailon, B. Chazelle, The Fast Johnson-Lindenstrauss Transform and Approximate Nearest Neighbors, , SIAM J. Comput. 39 (2009), 302-322. Prelim. version in STOC 2006.
- [4] D. Aldous, Probability Approximations via the Poisson clumping heuristic, Springer-Verlag, 1989, New York.
- [5] D. Aldous and P. Diaconis, Hammersely’s interacting particle process and longest increasing subsequences, Probability Theory and related fields, 103, 1995, pp199-213.
- [6] N. Alon, M. Krivelevich, The concentration of the chromatic number of random graphs, Combinatorica, **17**, 1997, 303-313.
- [7] S. Arora, Polynomial-time Approximation Schemes for Euclidean TSP and other Geometric Problems. Journal of the ACM 45(5) 753-782, 1998..
- [8] R. Arriaga and S. Vempala, An algorithmic theory of learning: Robust concepts and random projections, Proceedings of Foundations of Computer Science, 1999, 616-623
- [9] J. Baik, P. Deift and K. Johansson, On the Distribution of the length of the longest increasing subsequence of random permutations, Journal of the American Mathematical Society 12 (1999), no. 4, 1119–1178.
- [10] J. Bearwood, J. H. Halton and J. M. Hammersley, “The shortest path through many points”, Proceedings of the Cambridge Philosophical Society, 55, pp.299-327 (1959).

- [11] B. Bollobás, Random Graphs, Cambridge Studies in advanced mathematics, 73 (2001)
- [12] B. Bollobás, Martingales, isoperimetric inequalities and random graphs, in *Combinatorics*, Proceedings Eger., (1987) (Hajnal, et. al Eds) Colloq. Math. Soc. Janós Bolyai, **52** North-Holland, Amsterdam, pp 113-139.
- [13] S. Boucheron, O. Bousquet, G. Lugosi and P. Massart, Moment inequalities for functions of independent random variables, The Annals of Probability, 2005 Vol 33, No. 2, p 514-560.
- [14] M. Boutsikas and M. V. Koutras, A bound for the distribution of the sum of discrete associated or negatively associated random variables, The annals of Applied Probability, Vol 10, No. 4, 1137-1150 (2000).
- [15] D. L. Burkholder, Distribution function inequalities for Martingales, Annals of Probability, **1**, 1973, 19-42.
- [16] D. L. Burkholder, Inequalities for operators on Martingales, Proc. Intl. Congress Math., (Nice 1970) **2** pp 551-557, Gauthier-Villars, Paris.
- [17] E. G. Coffman, Jr., and G. S. Lueker, Probabilistic Analysis of Packing and Partitioning Algorithms, Wiley & Sons, 1991.
- [18] V. de la Peña, “A general class of exponential inequalities for Martingales and ratios”, The Annals of Probability, Vol. 27, No. 1, (1999)
- [19] J. Csirik, D. S. Johnson, C. Kenyon, J. B. Orlin, P. W. Shor, and R. R. Weber On the Sum-of-Squares Algorithm for Bin Packing, . Thirty-Second Annual ACM Symposium on Theory of Computing (STOC), 208-217, 2000. Journal version in JACM, 53(1), 1-65, 2006.
- [20] S. Dasgupta and A. Gupta, An elementary proof of the Johnson-Lindenstrauss Lemma, International Computer Science Institute, TR-99-006, (1999).
- [21] D. Dubhashi and D. Ranjan, Balls and bins: A study in negative dependence, Random Structures and Algorithms, 13, 99-124 (1998).
- [22] A. M. Frieze, On the independence number of random graphs, Discrete Mathematics, **81** pp171-175 (1983).
- [23] A. M. Frieze, On the length of the longest monotone increasing subsequence in a random permutation, Annals of Applied Probability, 1, 1991, 301-305.
- [24] P. Hitczenko, Best constants in martingale version of Rosenthal’s inequality, The Annals of Probability, (1990) Vol. 18, No. 4, p. 1656-1668.
- [25] P. Indyk and R. Motwani, Approximate nearest neighbors: Towards removing the curse of dimensionality, Proceedings of Symposium on Theory of Computing, 1998, 604-613.
- [26] S. Janson, T. Luczak and A. Rucinski, Random Graphs, Wiley- Interscience Series in Discrete Mathematics and Optimization (2000).
- [27] S. Janson, A. Rucinski, The deletion method for upper tail estimates, Combinatorica, **24** (4) p. 615-640.

- [28] S. Janson, K. Oleszkiewicz and A. Rucinski, Upper tails for subgraph counts in random graphs, *Israeli Journal of Mathematics*, (2004).
- [29] K. Joag-Dev and F. Proschan, Negative-association of random variables with applications, *Annals of Statistics*, **11** 286-295 (1983).
- [30] W. B. Johnson, G. Schechtman and J. Zinn, Best constants in moment inequalities for linear combinations of independent random variables, *Annals of Probability*, Vol. 13, 1985, No. 1, 234-253.
- [31] R. M. Karp, The probabilistic analysis of some combinatorial search algorithms, in *Algorithms and Complexity: New Directions and recent results*, J. F. Traub, ed., Academic Press, New York, 1976, pp 1-19.
- [32] R. M. Karp, Probabilistic Analysis of partitioning algorithms for the Traveling Salesman problem in the plane, *Mathematics of Operations research* 2, 1977, pp 209-224.
- [33] J.H. Kim and V. Vu, Concentration of multivariate polynomials and its applications, *Combinatorica*, **20**, (2000) p 417-434.
- [34] T. Luczak, The chromatic number of random graphs, *Combinatorica*, **11** 1991, 45-54.
- [35] C. McDiarmid. Concentration. In *Probabilistic Methods for Algorithmic Discrete Mathematics*, edited by M. Habib, C. McDiarmid, J. Ramirez-Alfonsin, and B. Reed, pp. 195-248, *Algorithms and Combinatorics* 16. Berlin: Springer, 1998.
- [36] M. Peligrad, S. Utev and W. B. Wu, A maximal L_p -inequality for stationary sequences and its applications, *Proc. American Math. Soc.*, (2005)
- [37] W. Rhee, Inequalities for the bin packing problem III, *Optimization*, **29** (1994) p 381-385
- [38] W. Rhee and M. Talagrand, A sharp deviation for the stochastic Traveling salesman problem, *Annals of Probability*, **17**, pp 1-8 (1989).
- [39] E. Shamir and J. Spencer, Sharp concentration of the chromatic number of random graphs, *Combinatorica* **7**, p 121-129.
- [40] A. Srinivasan, Distributions on level sets with applications to approximation algorithms, in the *Proc. of the 42 nd IEEE Symposium on Foundations of Computer Science*, (FOCS) 2001.
- [41] J. M. Steele, *Probability Theory and Combinatorial Optimization*, CBMS-NSF Regional Conference Series in Applied Mathematics, SIAM (1997)
- [42] J. M. Steele, Probabilistic Algorithm for the directed traveling salesman problem, *Mathematics of Operations Research*, **11**, 1986, 343-350.
- [43] M. Talagrand, Concentration of measure and isoperimetric inequalities in product spaces, *Publications mathématiques de l'I.H.É.S.*, tome 81 (1995) p. 73-205.
- [44] S. Vempala, The random projection method, *DIMACS Series in Discrete Mathematics and TCS*, volume 65 (2000)
- [45] V. Vu, Concentration of non-Lipschitz functions and applications, *Random Structures and Algorithms*, **20** (3) (2002), 262- 316