

Dynamical symmetry of Dirac hydrogen atom with spin symmetry and its connection with Ginocchio's oscillator

Fu-Lin Zhang, Bo Fu, and Jing-Ling Chen*

*Theoretical Physics Division, Chern Institute of Mathematics,
Nankai University, Tianjin 300071, People's Republic of China*

(Dated: September 15, 2021)

The Dirac hydrogen atom with spin symmetry is shown has a $SO(4)$ symmetry. The generators are derived, and the corresponding Casimir operator leads to the energy spectrum naturally. This type hydrogen atom is connected to a four-dimensional Dirac system with equal scalar and vector harmonic oscillator potential, by the Kustaanheimo-Stiefel transformation with a constraint.

PACS numbers: 03.65.Pm; 03.65.-w; 02.20.-a; 21.10.Sf

Introduction- Hydrogen atom and harmonic oscillator are two simple, solvable and real models. A common feature of them is that, their orbits of motion are closed in classical mechanics [1]. This indicates there are more constants of motion in these systems than the orbit angular momentum. They have been shown as the Rung-Lenz vector [2, 3] in the hydrogen atom and the second order tensors [4] in the harmonic oscillator. These conserved quantities as well as angular momentum generate the $SO(4)$ and $SU(3)$ Lie groups respectively. They are not geometrical but the symmetries in the phase space, and are called dynamical symmetries. These symmetries lead to an algebraic approach to determine the energy levels. Generally, the N -dimensional (ND) hydrogen atom has the $SO(N+1)$ and the isotropic oscillator has the $SU(N)$ symmetry. The $SO(4)$ generators in the 3D hydrogen atom can be written as two sets decoupled $SU(2)$ operators, whose Casimir operators are equal [5]. On the other hand, one can introduce a constraint condition on the 4D oscillator, and separate it into two 2D oscillators with the same energy. These two systems have the same algebraic structure, and can be connected by the famous Kustaanheimo-Stiefel (KS) transformation [6]. The same transformation has been applied to the path integral treatment of the hydrogen atom [7] and the radial coherent state for the Coulomb problem [8].

In the relativistic quantum mechanics, the motion of spin-1/2 particle satisfies the Dirac equation. Neither the Dirac hydrogen atom [9] nor the Dirac oscillator [10] has a dynamical symmetry. The main reason of the breaking of dynamical symmetries is the spin-orbit coupling. There wasn't any transformation reported to connect these two Dirac systems.

In recent years, the Dirac equation with scalar and vector potentials of equal magnitude (SVPEM) are studied widely [11, 12, 13, 14, 15, 16, 17, 18]. When the potentials are spherical, the Dirac equation is said to have the spin or pseudospin symmetry corresponding to the same or opposite sign. Then, the total angular momentum can be divided into conserved orbital and spin parts (see Eq. (2) for the spin symmetry case), which form the $SU(2)$ algebra separately. These symmetries, which

have been observed in the hadron and nuclear spectroscopies for a long time [19, 20], are derived from the investigation of the dynamics between a quark and an antiquark [21, 22, 23]. The very lately studies [17, 18] have revealed that, the motion of a spin-1/2 particle with SVPEM satisfies the same differential equation and has the same energy spectrum as a scalar particle. For the spin or pseudospin symmetry systems, Alberto *et. al.* [17] have indicated that the spin-orbit and Darwin terms of either the upper component or the lower component of the Dirac spinor vanish, which made it equivalent, as far as energy is concerned, to a spin-0 state. These results suggest that, one can image the spin-1/2 particle with SVPEM as a relativistic scalar particle with an additional spin but without the spin-orbit coupling. From this point of view, we speculate, the dynamical symmetries in non-relativistic hydrogen atom and harmonic oscillator would be holden in the Dirac systems with spin or pseudospin symmetry.

Actually, Ginocchio [24, 25] has found the $U(3)$ and pseudo- $U(3)$ symmetry in the Dirac equation with spin or pseudospin symmetry when the potential takes the harmonic oscillator form. The aim of this work is to answer the following questions: Does the Dirac equation with spin or pseudospin symmetry has a $SO(4)$ dynamical symmetry, when the potential takes the Coulomb form? And what is the relation of it and the Ginocchio's oscillator? For the sake of brevity, we only given the details of the spin symmetry case. The results of the pseudospin symmetry case will be easily obtained by making some corrections.

$SO(4)$ symmetry- The Dirac Hamiltonian with spin symmetry, in the relativistic units, $\hbar = c = 1$, is given by

$$H = \vec{\alpha} \cdot \vec{p} + \beta M + (1 + \beta) \frac{V(r)}{2}, \quad (1)$$

where $\vec{\alpha}$ and β are the Dirac matrices, and M is the mass. It commutes with the deformed orbital and spin angular momentum [22]

$$\vec{L} = \begin{bmatrix} \vec{l} & 0 \\ 0 & U_p \vec{l} U_p^\dagger \end{bmatrix}, \quad \vec{S} = \begin{bmatrix} \vec{s} & 0 \\ 0 & U_p \vec{s} U_p^\dagger \end{bmatrix}, \quad (2)$$

where $\vec{l} = \vec{r} \times \vec{p}$, $\vec{s} = \frac{\vec{\sigma}}{2}$ are the usual spin generators, $\vec{\sigma}$ are the Pauli matrices, and $U_p = U_p^\dagger = \frac{\vec{\sigma} \cdot \vec{p}}{p}$ is the helicity unitary operator [26]. The components of them form the $SU(2)$ Lie algebra separately, and the sum of them equals to the total angular momentum on account of $U_p \vec{l} U_p^\dagger + U_p \vec{s} U_p^\dagger = \vec{l} + \vec{s}$. Ginocchio has shown \vec{L} in Eq. (2) to be three of the generators of the $U(3)$ symmetry group. The other conserved quantities are supposed to take the form as

$$Q = \begin{bmatrix} Q_{11} & Q_{12} \vec{\sigma} \cdot \vec{p} \\ \vec{\sigma} \cdot \vec{p} Q_{21} & \vec{\sigma} \cdot \vec{p} Q_{22} \vec{\sigma} \cdot \vec{p} \end{bmatrix}.$$

If it commutes with the spin symmetry Hamiltonian (1), the matrix elements should satisfy the equations

$$\begin{aligned} Q_{12} &= Q_{21}, \\ [Q_{11}, V(r)] + [Q_{12}, p^2] &= 0, \\ [Q_{12}, V(r)] + [Q_{22}, p^2] &= 0, \\ Q_{11} &= Q_{12}(2M + V(r)) + Q_{22}p^2. \end{aligned} \quad (3)$$

In non-relativistic hydrogen atom, the constants of motion are the orbital angular momentum \vec{l} and the Runge-Lenz vector [2, 3, 5], $\vec{R} = \frac{\vec{f}}{2Mk} - \frac{\vec{r}}{r}$, where $\vec{f} = \vec{p} \times \vec{l} - \vec{l} \times \vec{p}$, and k is the parameter in the Coulomb potential $V(r)^h = -\frac{k}{r}$. One can obtain the following relations easily

$$\begin{aligned} [\vec{f}, p^2] &= 0, \quad [-\frac{\vec{r}}{r}, V^h(r)] = 0, \\ \frac{1}{2Mk} [\vec{f}, V^h(r)] + \frac{1}{2M} [-\frac{\vec{r}}{r}, p^2] &= 0. \end{aligned}$$

Insert them into Eq. (3), we can find the solutions of the Coulomb potential $V(r) = V^h(r) = -\frac{k}{r}$

$$\vec{Q} = \begin{bmatrix} 2M\vec{R} + \frac{k\vec{r}}{r^2} & (-\frac{\vec{r}}{r})\vec{\sigma} \cdot \vec{p} \\ \vec{\sigma} \cdot \vec{p}(-\frac{\vec{r}}{r}) & \vec{\sigma} \cdot \vec{p}(\frac{1}{k} \frac{\vec{f}}{p^2})\vec{\sigma} \cdot \vec{p} \end{bmatrix}.$$

The commutation relations of conserved \vec{L} and \vec{Q} are

$$\begin{aligned} [L_i, L_j] &= i\epsilon_{ijk}L_k, \\ [L_i, Q_j] &= i\epsilon_{ijk}Q_k, \\ [Q_i, Q_j] &= i\frac{-4}{k^2}(H^2 - M^2)\epsilon_{ijk}L_k, \end{aligned} \quad (4)$$

where $i, j, k = 1, 2, 3$. And,

$$\begin{aligned} \vec{Q} \cdot \vec{L} &= 0, \\ \vec{Q}^2 &= \frac{4}{k^2}(H^2 - M^2)(L^2 + 1) + (H + M)^2. \end{aligned} \quad (5)$$

For a fixed energy level, H is a constant. We can define the normalized generators $\vec{A} = [\frac{-4}{k^2}(H^2 - M^2)]^{-\frac{1}{2}}\vec{Q}$. Evidently, the two later relations in Eq. (4) are transformed into $[L_i, A_j] = i\epsilon_{ijk}A_k$ and $[A_i, A_j] = i\epsilon_{ijk}L_k$. These results show that the Dirac hydrogen atom with spin symmetry has a $SO(4)$ symmetry.

By the standard process of the non-relativistic hydrogen atom [5], the relations of the generators lead to the energy spectrum. Set $\vec{I} = (\vec{L} + \vec{A})/2$ and $\vec{K} = (\vec{L} - \vec{A})/2$, which satisfy the commutation relations, $[I_i, I_j] = i\epsilon_{ijk}I_k$, $[K_i, K_j] = i\epsilon_{ijk}K_k$, $[I_i, K_j] = 0$. \vec{I} and \vec{K} constitute two commutative $SU(2)$ algebra. Considering Eq. (5), it is immediate to obtain the Casimir operators

$$\vec{I}^2 = \vec{K}^2 = \frac{1}{4} \left[-\frac{k^2}{4} \frac{H + M}{H - M} - 1 \right] = j(j + 1),$$

where $j = 0, 1/2, 1, 3/2, \dots$. Hence, the eigenvalues of the Hamiltonian are given by

$$E^\pm = \frac{\pm 4n^2 - k^2}{4n^2 + k^2} M, \quad n = 2j + 1 = 1, 2, 3, \dots \quad (6)$$

which come up to [11, 12].

When $M \rightarrow \infty$, $H \rightarrow M$, the non-relativistic limit of the energy levels is $E^+ \rightarrow M - \frac{k^2}{2n^2}M$, the second term of which agrees with the non-relativistic results [5]. We can also get the non-relativistic limits of the conserved quantities

$$H - M \rightarrow \begin{bmatrix} \frac{p^2}{2M} - \frac{k}{r} & 0 \\ 0 & \frac{p^2}{2M} \end{bmatrix}, \quad \frac{\vec{Q}}{2M} \rightarrow \begin{bmatrix} \vec{R} & 0 \\ 0 & \frac{1}{2Mk} \vec{f} \end{bmatrix}.$$

The upper-left elements of the above matrices are nothing but the non-relativistic hydrogen atom Hamiltonian and the Runge-Lenz vector, and the lower-right ones are their limits when $k \rightarrow 0$.

KS transformation- The treatment of dynamical symmetry in the above section can be generalized to ND Dirac systems with SVPEM. This will be discussed in a forthcoming paper. Here we give the spectrum of the 4D Dirac system with equal scalar and vector harmonic oscillator potentials (4D Ginocchio's oscillator), and show its relation with the Dirac hydrogen atom with spin symmetry. Denoting the 4D spatial coordinates and momentums as u_μ and P_μ ($\mu = 1, 2, 3, 4$), the Dirac Hamiltonian with SVPEM is given by

$$\mathcal{H} = \alpha^\mu P_\mu + \beta m + (1 + \beta) \frac{\mathcal{V}(u)}{2}.$$

Here, $\alpha^{1,2,3} = \vec{\sigma}$ and β are the same as the 3D case, and $\alpha^4 = -\sigma_2 \otimes I$. Then, the Dirac equation can be written as

$$\begin{bmatrix} m + \mathcal{V}(u) - \varepsilon & B \\ B^\dagger & -m - \varepsilon \end{bmatrix} \begin{bmatrix} \phi_1 \\ \phi_2 \end{bmatrix} = 0, \quad (7)$$

where $B = \sigma^\mu P_\mu$ with $\sigma^{1,2,3} = \vec{\sigma}$ and $\sigma^4 = i$. It equals to the second order differential equations

$$\begin{aligned} [P^2 + (\varepsilon + m)\mathcal{V}(u) - (\varepsilon^2 - m^2)]\phi_1 &= 0, \\ (\varepsilon + m)\phi_2 + B^\dagger \phi_1 &= 0. \end{aligned}$$

Considering ε is a constant for a given energy level, the first equation takes the same form as the Schrödinger equation with the mass $\tilde{m} = (\varepsilon + m)/2$ and eigenvalue $\tilde{\varepsilon} = \varepsilon - m$. When $\mathcal{V}(u) = \frac{1}{2}m\omega^2 u^2 = \frac{1}{2}\tilde{m}\Omega^2 u^2$, the energy spectrum of the 4D Schrödinger equation is $\tilde{\varepsilon} = (N+2)\Omega$, where $N = n_1 + n_2 + n_3 + n_4$ is the total number operator [8] of the 4D non-relativistic harmonic oscillator. Hence, the eigenvalues in Eq. (7) satisfy the equation

$$(\varepsilon + m)^2(\varepsilon - m)^2 - 2m\omega^2(\varepsilon + m)(N + 2)^2 = 0. \quad (8)$$

The so-called KS transformation, between the spatial coordinates $\vec{r} = (x_1, x_2, x_3)$ and $u_\mu = (u_1, u_2, u_3, u_4)$, is given by [6]

$$\begin{aligned} x_1 &= 2(u_1 u_3 - u_2 u_4), \\ x_2 &= 2(u_1 u_4 + u_2 u_3), \\ x_3 &= u_1^2 + u_2^2 - u_3^2 - u_4^2, \end{aligned}$$

which connects the 3D hydrogen atom and 4D harmonic oscillator with the constraint condition

$$u_4 P_1 - u_1 P_4 + u_2 P_3 - u_3 P_2 = 0. \quad (9)$$

Under this transformation and constraint, it is easy to obtain $B = 2\Gamma\vec{\sigma}\cdot\vec{p}$, where $\Gamma = u_1 + iu_2\sigma_3 - iu_3\sigma_2 + iu_4\sigma_1$, $\Gamma\Gamma^\dagger = u^2 = r$. Then, the Dirac equation (7) can be written as

$$\begin{bmatrix} 2\Gamma & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} \frac{m-\varepsilon}{4r} + \frac{1}{8}m\omega^2 & \vec{\sigma}\cdot\vec{p} \\ \vec{\sigma}\cdot\vec{p} & -m-\varepsilon \end{bmatrix} \begin{bmatrix} 2\Gamma^\dagger & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} \phi_1 \\ \phi_2 \end{bmatrix} = 0. \quad (10)$$

Setting

$$M + E = m + \varepsilon, \quad M - E = \frac{1}{8}m\omega^2, \quad k = \frac{1}{4}(\varepsilon - m), \quad (11)$$

Eq. (10) is equivalent to nothing but the 3D Dirac equation of the hydrogen atom with spin symmetry

$$\begin{bmatrix} M - \frac{k}{r} - E & \vec{\sigma}\cdot\vec{p} \\ \vec{\sigma}\cdot\vec{p} & -M - E \end{bmatrix} \begin{bmatrix} \psi_1 \\ \psi_2 \end{bmatrix} = 0, \quad (12)$$

where $\psi_1 = 2\Gamma^\dagger\phi_1$ and $\psi_2 = \phi_2$. The constraint in Eq. (9) requires the number operators in 4D Schrödinger harmonic oscillator satisfy [8] $n_1 + n_2 = n_3 + n_4$. Therefore, the quantum number in Eq. (8) is confined to be even, $N+2 = 2n = 2, 4, 6, \dots$. Substituting the relations from Eq. (11) into Eq. (8), one can obtain the energy spectrum in Eq. (6).

Conclusion- In summary, we have shown that the 3D Dirac hydrogen atom with spin symmetry has the $SO(4)$ symmetry. The nature of this symmetry is not geometrical but dynamical. The relation of its Hamiltonian and the Casimir operators of the symmetry group leads to an algebraic solution of the relativistic energy spectrum. The non-relativistic limits of the conserved quantities coincide with their non-relativistic counterpart accurately. As a result of the symmetries of the Coulomb

and harmonic oscillator problem, we prove the Dirac hydrogen atom with spin symmetry can be connected to a 4D Ginocchio's oscillator by the KS transformation with a constraint condition. Dynamical symmetry and the KS transformation are two important concepts in non-relativistic quantum mechanics. The results in this paper and [25] show they also exist in Dirac systems with SVPEM. Since the spin symmetry and pseudospin symmetry exist frequently in antinucleon and nucleon spectrum [11]. We can foretell there are more properties of the Schrödinger equations presence in these systems.

This work is supported in part by NSF of China (Grants No. 10575053 and No. 10605013) and Program for New Century Excellent Talents in University. The Project-sponsored by SRF for ROCS, SEM.

* Email:chenjl@nankai.edu.cn

- [1] J. Bertrand, C. R. Acad. Sci. Paris. **77** (1873).
- [2] W. Lenz, Z. Phys. **24**, 197 (1925).
- [3] W. Pauli, Z. Phys. **36**, 336 (1926).
- [4] D. M. Fradkin, American Journal of Physics **33**, 207 (1965).
- [5] W. Greiner and B. Müller, *Quantum Mechanics (Symmetries)* (Springer, 1994), p. 453.
- [6] F. H. J. Cornish, J. Phy. A: Math. Gen. **17**, 323 (1984).
- [7] R. Ho and A. Inomata, Phys. Rev. Lett. **48**, 231 (1982).
- [8] C. C. Gerry, Phys. Rev. A **33**, 6 (1986).
- [9] W. Greiner, *Relativistic Quantum Mechanics: wave equation* (Reprinted in China by Beijing World Publishing Corporation, 2003), p. 225.
- [10] M. Moshinsky and A. Szczepaniak, J. Phys. A: Math. Gen. **22**, L817 (1989).
- [11] J. N. Ginocchio, Phys. Rep. **414**, 165 (2005).
- [12] A. Leviatan, Phys. Rev. Lett. **92**, 202501 (2004).
- [13] A. L. Blokhin, C. Bahri, and J. P. Draayer, Phys. Rev. Lett. **74**, 4149 (1995).
- [14] S.-G. Zhou, J. Meng, and P. Ring, Phys. Rev. Lett. **91**, 262501 (2003).
- [15] K. Sugawara-Tanabe and A. Arima, Phys. Rev. C **58**, R3065 (1998).
- [16] J. N. Ginocchio, Phys. Rev. Lett. **78**, 436 (1997).
- [17] P. Alberto, A. S. de Castro, and M. Malheiro, Phys. Rev. C **75**, 047303 (2007).
- [18] A. D. Alhaidari, H. Bahlouli, and A. Al-Hasan, Phys. Lett. A **349**, 87 (2006).
- [19] A. Arima, M. Harvey, and K. Shimizu, Phys. Lett. B **30**, 517 (1969).
- [20] K. Hecht and A. Adler, Nucl. Phys. A **137**, 129 (1969).
- [21] G. Smith and L. Tassie, Ann. Phys. **65**, 352 (1971).
- [22] J. S. Bell and H. Ruegg, Nucl. Phys. B **98**, 151 (1975).
- [23] H. W. Crater and P. V. Alstine, Phys. Rev. D **37**, 1982 (1988).
- [24] J. N. Ginocchio, Phys. Rev. C **69**, 034318 (2004).
- [25] J. N. Ginocchio, Phys. Rev. Lett. **95**, 252501 (2005).
- [26] A. L. Blokhin, C. Bahri, and J. P. Draayer, Phys. Rev. Lett. **74**, 4149 (1995).