Aharonov-Bohm effect on AdS_2 and nonlinear supersymmetry of reflectionless Pöschl-Teller system

Francisco Correa, Vít Jakubský and Mikhail S. Plyushchay

Departamento de Física, Universidad de Santiago de Chile, Casilla 307, Santiago 2, Chile E-mails: fco.correa.s@gmail.com, v.jakubsky@gmail.com, mplyushc@lauca.usach.cl

Abstract

We explain the origin and the nature of a special nonlinear supersymmetry of a reflectionless Pöschl-Teller system by the Aharonov-Bohm effect for a nonrelativistic particle on the AdS_2 . A key role in the supersymmetric structure appearing after reduction by a compact generator of the AdS_2 isometry is shown to be played by the discrete symmetries related to the space and time reflections in the ambient Minkowski space. We also observe that a correspondence between the two quantum non-relativistic systems is somewhat of the AdS/CFT holography nature.

1 Introduction

Exactly solvable quantum hyperbolic Pöschl-Teller system [1]–[6] finds diverse applications in a variety of physical problems. It appears, for instance, in the form of a stability equation for soliton (kink) solutions in 1+1-dimensional φ^4 and sine-Gordon field theories [7]. It describes static solutions in the Gross-Neveu model [8, 9, 10], and was used in investigation of the tachyon condensation phenomenon in gauge field string dynamics [11, 12]. This system, on the other hand, plays a fundamental role in the inverse scattering problem method for nonlinear evolution equations [13, 14].

All the mentioned applications are related to a peculiar property of the Pöschl-Teller system, whose potential $U(x) = -\lambda \cosh^{-2} x$ becomes reflectionless at special values $\lambda = m(m+1), m = 1, 2, ...,$ of the coupling constant. This property finds a simple explanation in terms of almost isospectral transformations underlying supersymmetric quantum mechanics [15]. Namely, the reflectionless Pöschl-Teller (RPT) system with $\lambda = m(m+1)$ can be related to a free particle model (m = 0) by a Crum-Darboux transformation of order m. The RPT system possesses also a shape-invariance property, due to which the energies of its m bound states are found algebraically [15, 16, 17].

The RPT system with m bound states is characterized by an integral of motion that is a differential operator of order 2m + 1. This operator is known as a nontrivial operator of the Lax pair of a non-periodic problem for the m-th order Korteweg-de Vries equation [14]. It was rediscovered recently in the context of nonlinear supersymmetric structure associated with the RPT system, see Refs. [18, 19], and [20, 21]. More specifically, it was observed in [18] that the RPT system is characterized by a hidden bosonized nonlinear N = 2 supersymmetry

[22]. The nontrivial Lax operator plays the role of one of its two supercharges, and the parity (reflection) operator R, $R\psi(x) = \psi(-x)$, is identified as the \mathbb{Z}_2 -grading operator Γ . As a consequence, the extended system composed from the two RPT systems with parameters mand l < m (including the case l = 0 that corresponds to a free particle system), is described by a nonlinear tri-supersymmetry [23, 24]. The tri-supersymmetric structure admits three alternative choices of the grading operator, $\Gamma = \sigma_3$, R, or $\sigma_3 R$, where σ_3 is a diagonal Pauli matrix [25]. For $\Gamma = \sigma_3$, the local part of this unusual supersymmetric structure corresponds to a nonlinear N = 4 supersymmetry, in which the higher order integral associated with the nontrivial Lax operator plays the role of a bosonic central charge [24].

In the present paper we show that the hidden bosonized nonlinear supersymmetry of the RPT system and the related nonlinear tri-supersymmetric structure of the extended case originate from a quantum problem of a charged non-relativistic particle on the AdS_2 surface in the presence of a singular magnetic-like vortex. Classically, the particle performs a free, geodesic motion on the AdS₂. Aharonov-Bohm effect [26] influences essentially on the quantum properties of the system. For half-integer values of the flux, the spectrum is characterized by a specific additional double degeneration related to an involutive automorphism of the AdS_2 isometry. The system reveals a reflectionless quantum dynamics in this case. These two related properties underlie the peculiar nonlinear supersymmetric structure of the RPT system obtainable by an appropriate angular momentum reduction. At the same time for a generic magnetic flux value, the Aharonov-Bohm effect on the AdS_2 provides a new vision on the nature of the Crum-Darboux transformations for the quantum hyperbolic Pöschl-Teller system in the context of the AdS/CFT correspondence [27, 28]. We also discuss this aspect in the background of non-relativistic AdS/CFT. The latter is based on non-relativistic conformal symmetry [29]–[35], and became recently a hot topic due to the diverse range of applications, see [36, 37].

The paper is organized as follows. In Section 2 we summarize briefly the results on the nonlinear hidden bosonized supersymmetric structure of the RPT system and the trisupersymmetric structure of the pair of such systems in the light of the Crum-Darboux transformations, discussed in detail in [18, 24]. In Section 3 we investigate the classical and quantum theory of a non-relativistic particle on the AdS_2 , minimally coupled to a U(1) gauge field given by the Aharonov-Bohm vector potential. In Section 4 we show how the nonlinear supersymmetric structure of the RPT system emerges from such a non-relativistic system at half-integer values of the magnetic flux. In Section 5 we discuss a realization of the conformal and super-conformal dynamical symmetries in the RPT system. We conclude with a discussion of the results and open research questions in Section 6.

2 Crum-Darboux transformations and tri-supersymmetry

A hyperbolic reflectionless Pöschl-Teller system is described by the Hamiltonian

$$H_m = -\frac{d^2}{dx^2} - \frac{m(m+1)}{\cosh^2 x}.$$
 (2.1)

Here and in what follows we put $\hbar = 1$. It is convenient to treat m as a parameter that can take any integer value, $m \in \mathbb{Z}$. Then m = 0 corresponds to a free particle case, and we have

the identity

$$H_m = H_{-(m+1)}.$$
 (2.2)

In terms of the first order differential operators

$$\mathcal{D}_m = \frac{d}{dx} + m \tanh x, \qquad \mathcal{D}_{-m} = -\mathcal{D}_m^{\dagger}, \qquad (2.3)$$

the second order operator (2.1) can be presented in a form

$$H_m = -\mathcal{D}_{-m}\mathcal{D}_m - m^2, \qquad (2.4)$$

while (2.2) rewrites equivalently

$$\mathcal{D}_{-m}\mathcal{D}_m = \mathcal{D}_{m+1}\mathcal{D}_{-m-1} + (2m+1). \tag{2.5}$$

Using this identity and (2.4), one can check that there hold the following intertwining relations

$$\mathcal{D}_m H_m = H_{m-1} \mathcal{D}_m, \qquad \mathcal{D}_{-m} H_{m-1} = H_m \mathcal{D}_{-m}.$$
(2.6)

These relations can be understood from a point of view of Crum-Darboux theorem [13]. According to it, if a differential operator A_n of order n annihilates n eigenstates ψ_i of a Hamiltonian H, one can construct another Hamiltonian $\tilde{H} = H - 2(\ln W(\psi_1, \ldots, \psi_n))''$, and these three operators are related by the identities

$$\tilde{H}A_n = A_n H, \qquad A_n^{\dagger} \tilde{H} = H A_n^{\dagger}.$$
 (2.7)

Here, W is the Wronskian of not obligatorily to be physical states ψ_i , $i = 1, \ldots, n$, such that $W \neq 0$. As a consequence of identities (2.7), if ψ is an eigenstate of H, $H\psi = E\psi$, $A_n\psi \neq 0$, the state $A_n\psi$ will be the eigenstate of \tilde{H} with the same eigenvalue E. And vice versa, if $\tilde{\psi}$ is an eigenstate of \tilde{H} , $\tilde{H}\tilde{\psi} = \tilde{E}\tilde{\psi}$, such that $A_n^{\dagger}\tilde{\psi} \neq 0$, the state $A_n^{\dagger}\tilde{\psi}$ will be an eigenstate of H with the the same eigenvalue \tilde{E} . This means that the Hamiltonians H and \tilde{H} will be almost isospectral. In the case of (2.6), the first order operator \mathcal{D}_m annihilates the nodeless ground state $\cosh^{-m} x$ of the Hamiltonian H_m with eigenvalue $-m^2$, and H_{m-1} has the same spectrum as H_m except the missing in it eigenvalue $-m^2$. In addition, relations (2.6) reflect the shape-invariance property of the Pöschl-Teller system. The first order Darboux transformation generated by the operator \mathcal{D}_m produces for the Hamiltonian $H = H_m$ the partner Hamiltonian \tilde{H} , which has potential of the same form but with the parameter m shifted in one, i.e. $\tilde{H} = H_{m-1}$.

By repeated application of (2.6), we can get the intertwining relations of higher order Crum-Darboux transformations [13],

$$\left(\mathcal{D}_{m-l}\dots\mathcal{D}_{m-1}\mathcal{D}_m\right)H_m = H_{m-l-1}\left(\mathcal{D}_{m-l}\dots\mathcal{D}_{m-1}\mathcal{D}_m\right),\tag{2.8}$$

$$\left(\mathcal{D}_{-m}\mathcal{D}_{-(m-1)}\dots\mathcal{D}_{-(m-l)}\right)H_{m-l-1} = H_m\left(\mathcal{D}_{-m}\mathcal{D}_{-(m-1)}\dots\mathcal{D}_{-(m-l)}\right).$$
(2.9)

In a particular case l = m - 1, relation (2.9) takes a form

$$\left(\mathcal{D}_{-m}\mathcal{D}_{-(m-1)}\dots\mathcal{D}_{-1}\right)H_0 = H_m\left(\mathcal{D}_{-m}\mathcal{D}_{-(m-1)}\dots\mathcal{D}_{-1}\right),\tag{2.10}$$

which means that the system (2.1) is almost isospectral to the free particle system. Making use of it, the scattering states of the RPT system (2.1) can be obtained from the plane wave eigenstates of the free particle,

$$\psi_m^{(\pm k)}(x) = \mathcal{D}_{-m}\mathcal{D}_{-m+1}\dots\mathcal{D}_{-1} \cdot \exp(\pm ikx).$$
(2.11)

According to (2.10) and (2.11), positive energy values of the system (2.1), $E_{m,k} = k^2$ with k > 0, are doubly degenerate, while zero energy with k = 0 corresponds to a singlet state

$$\psi_m^{(\pm 0)}(x) \equiv \psi_{m;m}(x) = \left(\mathcal{D}_{-m} \mathcal{D}_{-m+1} \dots \mathcal{D}_{-1} \right) \cdot 1.$$
 (2.12)

Function $\cosh^{-m} x$ is the zero mode of the first order operator \mathcal{D}_m . In correspondence with (2.4), it describes a bound state of the RPT system with energy $E_{m;0} = -m^2$. This observation together with the relation (2.9) taken for $l = 0, \ldots, m-2$ allow us to find the whole set of the of m bound singlet states of the RPT system (2.1),

$$\psi_{m;0}(x) = \cosh^{-m} x, \quad \psi_{m;n}(x) = \mathcal{D}_{-m} \mathcal{D}_{-m+1} \dots \mathcal{D}_{-m+n-1} \cosh^{n-m} x, \quad n = 1, \dots, m-1.$$
(2.13)

Extension of (2.13) for n = m (l = m - 1) reproduces singlet state (2.12) of the continuous part of the spectrum. The energies of all the m + 1 singlet states (2.13) and (2.12) are given then by

$$E_{m;n} = -(m-n)^2, \qquad n = 0, \dots, m.$$
 (2.14)

The double degeneration in the continuous part of the spectrum and the presence of m+1 > 1 singlet states indicate on a hidden, nonlinear supersymmetry in the system (2.1). Its corresponding supercharges can be identified easily. Applying the order 2m + 1 Crum-Darboux transformation (2.9) corresponding to l = 2m and taking into account relation (2.2), we find that the RPT system (2.1) is characterized by a local integral

$$\mathcal{A}_{2m+1} = \mathcal{D}_{-m}\mathcal{D}_{-m+1}\dots\mathcal{D}_{0}\dots\mathcal{D}_{m-1}\mathcal{D}_{m}, \qquad [\mathcal{A}_{2m+1}, H_{m}] = 0, \qquad (2.15)$$

that is a differential operator of order 2m + 1. This is a nontrivial integral of the Lax pair $(\mathcal{A}_{2m+1}, H_m)$ of the *m*-th order KdV equation [14]. It is a parity-odd operator, while the Hamiltonian (2.1) is parity-even. Identifying reflection (parity) operator R, $R\psi(x) = \psi(-x)$, as a grading operator, and integrals $Z_1 = i^{2m+1}\mathcal{A}_{2m+1}$ and $Z_2 = iRZ_1$ as Hermitian supercharges, we find that the RPT system (2.1) is characterized by a nonlinear N = 2supersymmetry

$$[Z_a, H_m] = 0, \qquad \{Z_a, Z_b\} = 2\delta_{ab}P_{2m+1}(H_m), \qquad (2.16)$$

where $P_{2m+1}(H_m)$ is a polynomial of order 2m + 1. Its explicit form can be found with the help of relation (2.5),

$$P_{2m+1}(H_m) = (H_m - E_{m;m}) \prod_{n=0}^{m-1} (H_m - E_{m;n})^2, \qquad (2.17)$$

where $E_{m;n}$ are the energies (2.14) of the singlet states. Singlet states (2.12) and (2.13) are zero modes of the supercharges Z_a , other m states annihilated by Z_a have a more intricate nature, see [19, 23].

Note that the nontrivial integral (2.15) has a sense of the integral \mathcal{D}_0 of the free particle system H_0 transferred to the RPT system H_m by means of the Crum-Darboux transformations. Indeed, multiply the relation $H_0\mathcal{D}_0 = \mathcal{D}_0H_0$ from the left by the operator $\mathcal{D}_{-m}\mathcal{D}_{-m+1}\ldots\mathcal{D}_{-1}$, and from the right by $\mathcal{D}_1\ldots\mathcal{D}_{m-1}\mathcal{D}_m$. Using on the left hand side intertwining relation (2.10), and on the right hand side its Hermitian conjugate form, we get $[\mathcal{A}_{2m+1}, H_m] = 0.$

The system composed from the two RPT systems H_m and H_l , $0 \leq l < m$, can be described by the 2×2 matrix Hamiltonian

$$\mathcal{H}_{m,l} = \begin{pmatrix} H_l & 0\\ 0 & H_m \end{pmatrix}. \tag{2.18}$$

The two RPT subsystems are almost isospectral. The subsystem H_l does not have m - l energy levels corresponding to the m - l lowest bound states of the subsystem H_m . Hence, these states form singlets of the extended system. The intertwining relations (2.9) allow us to identify the local Hermitian integrals of motion of the extended system (2.18),

$$\mathcal{X}_{m,l} = \begin{pmatrix} 0 & X_{m,l}^{-} \\ X_{m,l}^{+} & 0 \end{pmatrix}, \quad \mathcal{Y}_{m,l} = \begin{pmatrix} 0 & Y_{m,l}^{-} \\ Y_{m,l}^{+} & 0 \end{pmatrix}, \quad \mathcal{Z}_{m,l} = \begin{pmatrix} Z_{m,l}^{+} & 0 \\ 0 & Z_{m,l}^{-} \end{pmatrix}, \quad (2.19)$$

 $[\mathcal{H}_{m,l}, \mathcal{X}_{m,l}] = [\mathcal{H}_{m,l}, \mathcal{Y}_{m,l}] = [\mathcal{H}_{m,l}, \mathcal{Z}_{m,l}] = 0$, where

$$X_{m,l}^{-} = (X_{m,l}^{+})^{\dagger} = -i^{m-l} \mathcal{D}_{l+1} \mathcal{D}_{l+2} \dots \mathcal{D}_{m-1} \mathcal{D}_{m}, \qquad (2.20)$$

$$Y_{m,l}^{-} = (Y_{m,l}^{+})^{\dagger} = i^{2l+1} \mathcal{A}_{2l+1} X_{m,l}^{-},$$
(2.21)

$$Z_{m,l}^{-} = (Z_{m,l}^{-})^{\dagger} = X_{m,l}^{+} Y_{m,l}^{-} = Y_{m,l}^{+} X_{m,l}^{-} = i^{2m+1} \mathcal{A}_{2m+1}, \qquad (2.22)$$

$$Z_{m,l}^{+} = (Z_{m,l}^{+})^{\dagger} = X_{m,l}^{-} Y_{m,l}^{+} = Y_{m,l}^{-} X_{m,l}^{+} = i^{2l+1} X_{m,l}^{-} X_{m,l}^{+} \mathcal{A}_{2l+1}.$$
(2.23)

These integrals of motion mutually commute,

$$[\mathcal{X}_{m,l}, \mathcal{Y}_{m,l}] = [\mathcal{Y}_{m,l}, \mathcal{Z}_{m,l}] = [\mathcal{X}_{m,l}, \mathcal{Z}_{m,l}] = 0, \qquad (2.24)$$

and satisfy relations

$$\mathcal{Z}_{m,l} = \mathcal{X}_{m,l} \mathcal{Y}_{m,l}, \qquad (2.25)$$

$$(\mathcal{X}_{m,l})^2 = P_{\mathcal{X}}(\mathcal{H}_{m,l}), \qquad (\mathcal{Y}_{m,l})^2 = P_{\mathcal{Y}}(\mathcal{H}_{m,l}), \qquad (\mathcal{Z}_{m,l})^2 = P_{\mathcal{Z}}(\mathcal{H}_{m,l}).$$
(2.26)

In correspondence with (2.25), $P_{\mathcal{Z}}(\mathcal{H}_{m,l}) = P_{\mathcal{X}}(\mathcal{H}_{m,l})P_{\mathcal{Y}}(\mathcal{H}_{m,l})$ is a polynomial of order 2m+1 of the form (2.17) with H_m changed for $\mathcal{H}_{m,l}$, and

$$P_{\mathcal{X}}(\mathcal{H}_{m,l}) = \prod_{n=0}^{m-l-1} (\mathcal{H}_{m,l} - E_{m;n}), \quad P_{\mathcal{Y}}(\mathcal{H}_{m,l}) = P_{\mathcal{X}}(\mathcal{H}_{m,l}) \cdot (\mathcal{H}_{m,l} - E_{m;m}) \prod_{n=m-l}^{m-1} (\mathcal{H}_{m,l} - E_{m;n})^2.$$
(2.27)

Here, the roots $E_{m;n}$ of the polynomials correspond to the singlet states of the subsystem H_m , and are given by (2.14).

In addition to integrals (2.19), the extended system (2.18) has also three mutually commuting *trivial* integrals of motion,

$$\Gamma_i \in \{R, \sigma_3, R\sigma_3\}, \qquad \Gamma_i^2 = 1, \quad i = 1, 2, 3.$$
 (2.28)

Any of them can be chosen as a grading operator Γ , that classifies one of the integrals (2.19) as an even (bosonic), and the other two as odd (fermionic) generators. As a result, the set of integrals (2.19), (2.28) extended by the products of (2.28) with (2.19) generates together with Hamiltonian (2.18) a certain deformation of the su(2|2) superalgebra. Its concrete form depends on the choice of the grading operator, see [23, 24]. For the particular choice $\Gamma = \sigma_3$, the subset of integrals $\mathcal{X}_{m,l}$, $\mathcal{Y}_{m,l}$, $i\sigma_3\mathcal{X}_{m,l}$, and $i\sigma_3\mathcal{Y}_{m,l}$ is identified as the set of the odd supercharges, while $\mathcal{Z}_{m,l}$ and $\mathcal{H}_{m,l}$ are even operators. These local integrals generate a nonlinear N = 4 superalgebra, in which, in correspondence with (2.24), $\mathcal{Z}_{m,l}$ plays a role of a central charge. For m - l = 1, the integral $\mathcal{X}_{m,m-1}$ is a usual first order supercharge for the superextended system (2.18), whose conservation follows from the Darboux intertwining relations (2.6). In a generic case, the relation of commutativity of $\mathcal{X}_{m,l}$ with matrix Hamiltonian $\mathcal{H}_{m,l}$ represents just mutually conjugate intertwining relations (2.8) and (2.9) of order m-l. The integral $\mathcal{Y}_{m,l}$ corresponds to the Crum-Darboux transformations of order m+l+1that intertwine the systems H_m and $H_{-(l+1)} = H_l$.

3 Aharonov-Bohm effect on the AdS_2

In this section, we will consider a two-dimensional setting with the geometry of the AdS_2 space. The particle system will be studied in the presence of a singular magnetic flux in both classical and quantum frameworks. This will provide a valuable background for deeper understanding of the algebraic properties described in the previous section.

Consider a one-sheeted hyperboloid

$$x^{\mu}x_{\mu} = -x_1^2 - x_2^2 + x_3^2 = -\mathcal{R}^2 \tag{3.1}$$

embedded in a three-dimensional Minkowski space with metric $ds^2 = dx^{\mu}dx^{\nu}\eta_{\mu\nu}$, $\eta_{\mu\nu} = diag(-1, -1, +1)$, $\mu, \nu = 1, 2, 3$. It can be parameterized by

$$x^{1} = \mathcal{R} \cosh \chi \cos \varphi, \quad x^{2} = \mathcal{R} \cosh \chi \sin \varphi, \quad x^{3} = \mathcal{R} \sinh \chi,$$
 (3.2)

and is identified as the AdS₂ space of radius $\mathcal{R} > 0$ with the induced metric $ds^2 = \mathcal{R}^2(d\chi^2 - \cosh^2 \chi d\varphi^2)$, $-\infty < \chi < \infty$, $0 \le \varphi < 2\pi^{-1}$. Let in an ambient Minkowski space the Aharonov-Bohm vector potential is given,

$$A_1 = -\frac{\Phi}{2\pi} \frac{x_2}{x_1^2 + x_2^2}, \quad A_2 = \frac{\Phi}{2\pi} \frac{x_1}{x_1^2 + x_2^2}, \quad A_3 = 0,$$
(3.3)

that describes a "magnetic" field of a flux line along the x_3 -axis, $B_{\mu} = \epsilon_{\mu\nu\lambda}\partial^{\nu}A^{\lambda} = (0, 0, \Phi\delta^2(x_1, x_2))$. Here $\epsilon_{\mu\nu\lambda}$ is an antisymmetric tensor, $\epsilon_{123} = 1$.

3.1 Classical system

Consider a *non-relativistic* charged particle minimally coupled to the external U(1) gauge field (3.3), and confined to move on the two-dimensional surface (3.1). Taking into account

¹The AdS₂ has a topology $\mathbb{R} \times S^1$ with S^1 usually unwrapped [28]. However, for here, the S^1 -topology plays a key role.

Eqs. (3.2), (3.3), the corresponding Lagrangian of a particle of unit mass, $L = \frac{1}{2}\dot{x}_{\mu}\dot{x}^{\mu} + \frac{e}{c}A_{\mu}\dot{x}^{\mu}$, is reduced to

$$L = \frac{\mathcal{R}^2}{2} (\dot{\chi}^2 - \cosh^2 \chi \, \dot{\varphi}^2) - \alpha \dot{\varphi}, \qquad (3.4)$$

where $\alpha \equiv \frac{e\Phi}{2\pi c}$, $\dot{x}^{\mu} = dx^{\mu}/dt$, and t is an evolution parameter. The last, coupling term in (3.4) is a total time derivative. It does not effect on the classical dynamics of the particle that performs a geodesic motion on the AdS₂.

The form of the trajectories can be identified by noting that the SO(2,1)-isometry of the AdS_2 is a transitive symmetry group of the hyperboloid (3.1), i.e. any two points on the surface can be related by an appropriate SO(2,1)-transformation. The generators of this symmetry are the integrals of motion

$$J_1 = -p_\chi \sin \varphi - J_3 \cos \varphi \tanh \chi, \quad J_2 = p_\chi \cos \varphi - J_3 \sin \varphi \tanh \chi, \tag{3.5}$$

$$J_3 = p_\varphi + \alpha, \tag{3.6}$$

where $p_{\chi} = \frac{\partial L}{\partial \dot{\chi}}$, $p_{\varphi} = \frac{\partial L}{\partial \dot{\varphi}}$ are the canonical momenta, $\{\varphi, p_{\varphi}\} = 1$, $\{\chi, p_{\chi}\} = 1$. With respect to the Poisson brackets, the integrals (3.5) and (3.6) generate the (2+1)D Lorentz algebra

$$\{J_{\mu}, J_{\nu}\} = -\epsilon_{\mu\nu\lambda}J^{\lambda}.$$
(3.7)

Their conservation follows from the form of the canonical Hamiltonian, $H = p_{\chi} \dot{\chi} + p_{\varphi} \dot{\varphi} - L$,

$$H = \frac{1}{2\mathcal{R}^2} \left(p_{\chi}^2 - \frac{(p_{\varphi} + \alpha)^2}{\cosh^2 \chi} \right), \qquad (3.8)$$

that is reduced to the so(2,1) Casimir element up to a multiplicative constant

$$H = -\frac{1}{2\mathcal{R}^2}\mathcal{C}, \quad \mathcal{C} = J_{\mu}J^{\mu}.$$
(3.9)

Direct checking shows that the integrals (3.5), (3.6) satisfy identically the relation $x^{\mu}J_{\mu} = 0$ with x^{μ} given by (3.2). This also follows from the observation that the system (3.8) can alternatively be obtained in two steps. First, we reduce the ISO(2, 1)-invariant free particle system $L = \frac{1}{2}m\dot{x}_{\mu}\dot{x}^{\mu}$ to the SO(2, 1)-invariant surface given by the second class constraints $x_{\mu}x^{\mu} + \mathcal{R}^2 = 0$ and $p_{\mu}x^{\mu} = 0$ in the phase space with canonical coordinates (x^{μ}, p_{μ}) . The reduced phase space is described by the two pairs of canonical variables $(\varphi, p_{\varphi}), (\chi, p_{\chi})$, in terms of which the SO(2, 1) generators $J_{\mu} = -\epsilon_{\mu\nu\lambda}x^{\nu}p^{\lambda}$ take the form of the integrals (3.5), (3.6) with $\alpha = 0$. The case $\alpha \neq 0$ is obtained then via a subsequent canonical transformation $p_{\varphi} \rightarrow p_{\varphi} + \alpha$. Both steps do not touch the initial free case identity $x^{\mu}J_{\mu} \equiv 0$. As a result, the trajectory is determined by the intersection of the Minkowski hyperplane $x^{\mu}J_{\mu} = 0$, $J_{\mu} = const$, with the surface of the hyperboloid (3.1). Its form depends on the value of the Casimir \mathcal{C} . For $\mathcal{C} > 0$, = 0, or < 0, the trajectory is respectively an ellipse, a straight line, or a hyperbola.

Due to the Lorentzian (indefinite) metric of the AdS_2 surface, the values of the Hamiltonian (3.8) are not restricted from below. We will be interested in the quantum system reduced to certain levels of the integral J_3 , in which the spectrum is bounded from below. As it follows from (3.6) and (3.8), such a reduced system describes a one-dimensional Pöschl-Teller system. Three types of the classical trajectories of the system (3.8) correspond to a bounded periodic (C > 0), or unbounded ($C \le 0$) particle motions in a 1D classical attractive potential. In particular, straight line trajectories on the hyperboloid correspond to a zero energy motion of the Pöschl-Teller system.

Due to its algebraic background, the system (3.9) has also two discrete symmetries. They have no significant role in the classical theory, but will be of a key importance at the quantum level in the context of the supersymmetry we discuss. These symmetries are the involutive automorphisms of the so(2, 1) algebra (3.7),

$$R: (J_1, J_2, J_3) \to (-J_1, -J_2, J_3), \qquad S: (J_1, J_2, J_3) \to (-J_1, J_2, -J_3). \tag{3.10}$$

In the ambient Minkowski space they correspond to a change of a sign of a space-like coordinate x^3 , and of one of the time-like coordinates which, in correspondence with the chosen definition of S, we identify with x^2 ,

$$R: (x^1, x^2, x^3) \to (x^1, x^2, -x^3), \qquad S: (x^1, x^2, x^3) \to (x^1, -x^2, x^3). \tag{3.11}$$

In the curvilinear coordinates this corresponds to

$$R: (\chi, \varphi) \to (-\chi, \varphi), \qquad S: (\chi, \varphi) \to (\chi, -\varphi). \tag{3.12}$$

Lagrangian (3.4) is invariant under the discrete symmetry R. However, it is quasi-invariant under symmetry S, which provokes a change for a total time derivative term, $\Delta L = 2\alpha\dot{\varphi}$, that does not effect on the classical motion. In correspondence with such a quasi-invariance of Lagrangian, in order to reproduce involutive automorphism S defined by (3.10) in the presence of nonzero Aharonov-Bohm flux α , a transformation $\varphi \to -\varphi$ has to be accompanied by an additional canonical transformation $p_{\varphi} \to p_{\varphi} - 2\alpha$. At the quantum level corresponding unitary transformation is generated by the operator $U_{\alpha}(\varphi) = \exp(-2i\alpha\varphi)$. This is a well defined, 2π -periodic operator only in the case of integer and half-integer values of the flux. For $2\alpha \notin \mathbb{Z}$, as we shall see, discrete symmetry S is spontaneously broken.

3.2 Quantization and spectral properties

A canonical quantization of the system with a prescription of a symmetric ordering of noncommuting factors in quantum analog of (3.5) results in the operators

$$\hat{J}_1 = i \sin \varphi \left(\partial_{\chi} - \frac{1}{2} \tanh \chi \right) - \cos \varphi \tanh \chi \hat{J}_3,$$
 (3.13)

$$\hat{J}_2 = -i\cos\varphi\left(\partial_{\chi} - \frac{1}{2}\tanh\chi\right) - \sin\varphi\tanh\chi\hat{J}_3, \qquad (3.14)$$

$$\hat{J}_3 = -i\partial_{\varphi} + \alpha. \tag{3.15}$$

They generate the so(2,1) algebra,

$$[\hat{J}_{\mu}, \hat{J}_{\nu}] = -i\epsilon_{\mu\nu\lambda}\hat{J}^{\lambda}.$$
(3.16)

We assume that these operators act on the space of the 2π -periodic in φ wave functions $\psi(\chi, \varphi)$, where they are Hermitian with respect to a scalar product

$$(\psi_1,\psi_2) = \frac{1}{2\pi} \int_{-\infty}^{\infty} \int_0^{2\pi} \psi_1^*(\chi,\varphi)\psi_2(\chi,\varphi)d\chi d\varphi.$$
(3.17)

The quantum Hamiltonian

$$\hat{H} = -\partial_{\chi}^2 - \frac{\hat{J}_3^2 - \frac{1}{4}}{\cosh^2 \chi}$$
(3.18)

is obtained then from (3.9), where we put $2\mathcal{R}^2 = 1$, and subtract a quantum constant term $\frac{\hbar^2}{4}$, i.e. we take

$$\hat{H} = -\hat{J}_{\mu}\hat{J}^{\mu} - \frac{1}{4}.$$
(3.19)

Note that the same realization of the so(2,1) generators and of the Hamiltonian are obtained if to proceed from the definition $\hat{J}_{\mu} = -\epsilon_{\mu\nu\lambda}x^{\nu}\left(-i\partial^{\lambda} - \frac{e}{c}A^{\lambda}(x)\right)$ written in the pseudospherical coordinates $(\mathcal{R}, \chi, \varphi), \mathcal{R} > 0$. In this way we get the so(2, 1) generators in the form

$$\tilde{J}_1 = i \sin \varphi \partial_\chi - \cos \varphi \tanh \chi \hat{J}_3, \qquad \tilde{J}_2 = -i \cos \varphi \partial_\chi - \sin \varphi \tanh \chi \hat{J}_3, \qquad \tilde{J}_3 = \hat{J}_3. \quad (3.20)$$

Instead of (3.18), with the same quantum constant shift, we obtain

$$\tilde{H} = -\partial_{\chi}^2 - \tanh \chi \partial_{\chi} - (\tilde{J}_3^2 - \frac{1}{4}) \cosh^{-2} \chi.$$
(3.21)

Operators \tilde{J}_{μ} and \tilde{H} , $[\tilde{J}_{\mu}, \tilde{H}] = 0$, are Hermitian with respect to a scalar product

$$(\tilde{\psi}_1, \tilde{\psi}_2) = (2\pi)^{-1} \int_{-\infty}^{\infty} \int_0^{2\pi} \tilde{\psi}_1^*(\chi, \varphi) \tilde{\psi}_2(\chi, \varphi) \cosh \chi d\chi d\varphi.$$
(3.22)

A subsequent similarity transformation $\tilde{\psi} \to f\tilde{\psi} = \psi$, $\tilde{J}_{\mu} \to f\tilde{J}_{\mu}f^{-1} = \hat{J}_{\mu}$, $\tilde{H} \to f\tilde{H}f^{-1} = \hat{H}$ with $f = \sqrt{\cosh \chi}$ reduces the scalar product (3.22), the so(2, 1) generators (3.20), and the Hamiltonian (3.21) to (3.17), (3.13)–(3.15), and (3.18), respectively.

Since Hamiltonian (3.18) is the (shifted) so(2, 1) Casimir operator, one can choose a representation in which \hat{H} and the compact so(2, 1) generator \hat{J}_3 are diagonal. The stationary Schrödinger equation associated with the Hamiltonian (3.18) is separable in the variables χ and φ . The common wave functions of \hat{H} and \hat{J}_3 can be factorized as

$$\Psi^{\alpha}_{E,m}(\chi,\varphi) = e^{im\varphi}\psi^{\alpha}_{E,m}(\chi), \qquad m = 0, \pm 1, \pm 2, \dots,$$
(3.23)

where the superscript marks the value of the Aharonov-Bohm flux. Then we have

$$\hat{J}_3 \Psi^{\alpha}_{E,m}(\chi,\varphi) = j_3 \Psi^{\alpha}_{E,m}(\chi,\varphi), \qquad j_3 = m + \alpha, \qquad (3.24)$$

and the Schrödinger equation

$$\hat{H}\Psi^{\alpha}_{E,m}(\chi,\varphi) = E\Psi^{\alpha}_{E,m}(\chi,\varphi) \tag{3.25}$$

is reduced to

$$H_{m_{\alpha}}\psi^{\alpha}_{E,m}(\chi) = E\psi^{\alpha}_{E,m}(\chi), \qquad H_{m_{\alpha}} = -\frac{d^2}{d\chi^2} - \frac{m_{\alpha}(m_{\alpha}+1)}{\cosh^2\chi}, \qquad m_{\alpha} \equiv m + \alpha - \frac{1}{2}.$$
(3.26)

The reduced Hamiltonian $H_{m_{\alpha}}$ is just the Pöschl-Teller Hamiltonian (2.1) with parameter mand variable x changed for m_{α} and χ . Since the cases corresponding to the Aharonov-Bohm fluxes α_1 and $\alpha_2 = \alpha_1 + n$, $n \in \mathbb{Z}$, are related by a unitary transformation generated by the operator $U_{\alpha_2,\alpha_1}(\varphi) = e^{in\varphi}$, in what follows we assume without loss of generality that $0 \leq \alpha < 1$.

Define the linear combinations of the generators $\hat{J}_+ = \hat{J}_1 + i\hat{J}_2$ and $\hat{J}_- = \hat{J}_+^{\dagger}$,

$$\hat{J}_{+} = e^{i\varphi} \left(\frac{\partial}{\partial \chi} - \left(\hat{J}_{3} + \frac{1}{2} \right) \tanh \chi \right), \quad \hat{J}_{-} = e^{-i\varphi} \left(-\frac{\partial}{\partial \chi} - \left(\hat{J}_{3} - \frac{1}{2} \right) \tanh \chi \right). \quad (3.27)$$

These are the ladder operators of the so(2, 1) symmetry algebra, $[\hat{J}_3, \hat{J}_{\pm}] = \pm \hat{J}_{\pm}, \ [\hat{J}_+, \hat{J}_-] = -2\hat{J}_3$. Then for eigenstates (3.23) we have a relation

$$\hat{J}_3(\hat{J}_{\pm}\Psi^{\alpha}_{E,m}) = (j_3 \pm 1)(\hat{J}_{\pm}\Psi^{\alpha}_{E,m}).$$
(3.28)

On a subspace with $j_3 = m + \alpha$, the χ -dependent parts of the ladder operators \hat{J}_- and \hat{J}_+ correspond to the intertwining operators (2.3). The values of the parameter m_{α} are non-integer in the case $\alpha \neq \frac{1}{2}$.

Let us discuss the spectral properties of the system in dependence on the strength of the magnetic flux. First, consider the case $\alpha \neq \frac{1}{2}$. From the form of the reduced Hamiltonian (3.26) it follows that the quantum system (3.18) contains the Pöschl-Teller subsystem with repulsive potential $U(\chi) = +\gamma^2 \cosh^{-2} \chi$, $\gamma^2 > 0$. For $\alpha = 0$ this happens in the subspace with m = 0, where $\gamma^2 = 1/4$. For $0 < \alpha < 1/2$ and $1/2 < \alpha < 1$, repulsive potential appears in the subspaces with m = 0 and m = -1, where $\gamma^2 = 1/4 - \alpha^2$ and $\gamma^2 = 1/4 - (1 - \alpha)^2$, respectively. Repulsive Pöschl-Teller system has no physical states with E = 0. In accordance with Eq. (3.28), the continuous part of the spectrum of the quantum system (3.18) with $\alpha \neq \frac{1}{2}$ is described by the scattering states with E > 0. The corresponding eigenstates with $E = k^2$, k > 0, can be expressed in terms of the hypergeometric function. Any Pöschl-Teller subsystem (3.26) is characterized by a nonzero reflection coefficient [38]

$$|r|^2 = \frac{1}{1+\rho^2}, \quad \rho = \frac{\sinh \pi k}{\cos \pi \alpha}.$$
 (3.29)

In accordance with (3.19), on the scattering states with E > 0, infinite-dimensional unitary irreducible representations of the principal continuous series of the algebra $sl(2, R) \sim so(2, 1)$ [39] with $-\hat{J}_{\mu}\hat{J}^{\mu} = E + 1/4 > 1/4$ and $j_3 = \alpha + m, m = 0, \pm 1, \ldots$, are realized.

System (3.18) with $\alpha \neq 1/2$ has also bound states of certain discrete negative energies. With the help of the relation

$$\hat{H} = \hat{J}_{+}\hat{J}_{-} - (\hat{J}_{3} - 1/2)^{2} = \hat{J}_{-}\hat{J}_{+} - (\hat{J}_{3} + 1/2)^{2}, \qquad (3.30)$$

cf. (2.4) and (2.5), a part of corresponding normalizable eigenstates is identified as the states annihilated by the ladder operators \hat{J}_{-} and \hat{J}_{+} . These are

$$\Psi_{E,m}^{\alpha,-}(\chi,\varphi) = e^{im\varphi} \cosh^{-(m+\alpha-1/2)}\chi, \quad \hat{J}_{-}\Psi_{E,m}^{\alpha,-} = 0, \qquad E = -(m+\alpha-1/2)^2, \quad (3.31)$$

where m = 1, 2... for $0 \le \alpha < 1/2$, and m = 0, 1, ... for $1/2 < \alpha < 1$, and

$$\Psi_{E,m}^{\alpha,+}(\chi,\varphi) = e^{im\varphi} \cosh^{m+\alpha+1/2}\chi, \qquad \hat{J}_+\Psi_{E,m}^{\alpha,+} = 0, \quad E = -(m+\alpha+1/2)^2, \qquad (3.32)$$

with m = -1, -2, ... for $0 \le \alpha < 1/2$, and m = -2, -3, ... for $1/2 < \alpha < 1$. Infinite number of bound states with the same energy eigenvalues are obtained then by the action of the ladder operators $(\hat{J}_+)^n$ and $(\hat{J}_-)^n$, n = 1, 2, ..., on the states (3.31) and (3.32). At $\alpha = 0$ the discrete part of the spectrum reveals a symmetry with respect to the change $j_3 \to -j_3$, but it has no such a symmetry for $\alpha \neq 0$. On the states

$$(\hat{J}_{+})^{n}\Psi_{E,m}^{\alpha,-}$$
 and $(\hat{J}_{-})^{n}\Psi_{E,m}^{\alpha,+}, \quad n = 0, 1, \dots,$ (3.33)

the half-bounded infinite-dimensional unitary representations of the discrete series of the sl(2, R) [39] are realized. These representations are characterized by the value of the Casimir operator and eigenvalues of the compact generator \hat{J}_3 , which are respectively $-\hat{J}_{\mu}\hat{J}^{\mu} = -(m+\alpha)(m+\alpha-1), j_3 = m+\alpha+n, \text{ and } -\hat{J}_{\mu}\hat{J}^{\mu} = -(m+\alpha)(m+\alpha+1), j_3 = m+\alpha-n.$ The quantum spectrum of the system with $\alpha = 0$ is illustrated on Fig. 1. Note that the subspaces with j_3 and $-j_3$, $j_3 \neq 0$, are present symmetrically in the spectrum, while the subspace with $j_3 = 0$ is unpaired.



Figure 1: Spectrum of the AdS₂ system with integer Aharonov-Bohm flux ($\alpha = 0$).

Now, we will focus on the case $\alpha = 1/2$. On the subspaces with m = 0 and m = -1, the dynamics reduces to that of the one-dimensional free quantum particle, while the set of the Hamiltonians (3.26) with all the possible integer values $m_{1/2} = m$ corresponds to the family of the RPT Hamiltonians (2.1) satisfying the identity (2.2),

$$\hat{H}|_{j_3=\frac{1}{2}} = \hat{H}|_{j_3=-\frac{1}{2}} = H_0 = -\partial_{\chi}^2, \qquad \hat{H}|_{j_3=m+\frac{1}{2}} = H_m.$$
(3.34)

The eigenstates $\Psi_{E,m}^{1/2}$ with m = 1, 2... and $E = k^2 > 0$ of the Hamiltonian (3.19) are obtained from the free particle plane wave states $e^{\pm ik\chi}$ (m = 0) by the action on them of the operator $(J_+)^m = e^{im\varphi} \mathcal{D}_{-m} \mathcal{D}_{-m+1} \dots \mathcal{D}_{-1}$. This corresponds exactly to relation (2.11). The scattering states with negative values of m are produced by the action of the Hermitian conjugate operator $(J_-)^m$, m = 1, 2..., on $e^{\pm ik\chi}$. In comparison with the case $\alpha \neq 1/2$, the system has additional states with E = 0, on which the half-bounded infinite-dimensional unitary representations of the sl(2, R) are realized. These states have a form of the eigenvectors (3.33) constructed over the eigenstates of the form (3.31) and (3.32) with $\alpha = 1/2$, in which m = 0 and m = -1, respectively. The normalizable negative energy states are constructed in the same way over the eigenstates (3.31) and (3.32) with m = 1, 2... and



Figure 2: Spectrum of the AdS₂ system with half-integer Aharonov-Bohm flux ($\alpha = 1/2$).

 $m = -2, -3, \ldots$ In this case all the energy levels in the spectrum shown on Fig. 2 have a double degeneration with respect to the reflection $j_3 \rightarrow -j_3$, cf. Fig. 1.

In terms of the ladder operators, second involutive automorphism from (3.10) takes the form

$$S: (\hat{J}_{\pm}, \hat{J}_3) \to (-\hat{J}_{\mp}, -\hat{J}_3).$$
 (3.35)

Under this transformation quantum symmetry algebra (3.16) and Hamiltonian (3.19) are invariant. The described properties of the states of the quantum system mean that this discrete symmetry is spontaneously broken except the cases when the Aharonov-Bohm flux takes integer and half-integer values. In the case $\alpha = 0$ or $\alpha = 1/2$, this discrete symmetry transforms mutually the half-bounded infinite-dimensional representations of the so(2, 1)realized on the states of the discrete spectrum (including the states E = 0 when $\alpha = 1/2$).

4 Reduction and tri-supersymmetry

Let us identify now the exact analogs of the Crum-Darboux intertwining relations, and trace out how the tri-supersymmetric structure appears under the appropriate reduction of the AdS₂ system in the presence of the Aharonov-Bohm effect with the half-integer flux $(\alpha = 1/2)$. In conclusion of this section we shall comment on the case of the integer flux.

Having in mind factorization (3.23), we introduce notations $|m\rangle = e^{im\varphi}$ and $\langle \psi_1 | \psi_2 \rangle = \frac{1}{2\pi} \int_0^{2\pi} \psi_1^* \psi_2 d\varphi$, so that $\langle m | m' \rangle = \delta_{mm'}$. Taking into account explicit form of the ladder operators (3.27), we get the only nonzero matrix elements of \hat{J}_{\pm} and \hat{H} ,

$$\langle m-1|\hat{J}_{-}|m\rangle = -\mathcal{D}_{m}, \quad \langle m|\hat{J}_{+}|m-1\rangle = \mathcal{D}_{-m},$$

$$(4.1)$$

$$\langle m|\hat{H}|m\rangle = -\mathcal{D}_{-m}\mathcal{D}_m - m^2, \qquad (4.2)$$

that correspond to (2.3) and (2.4). Then, with taking into account (4.1) and (4.2), the Crum-Darboux intertwining relation (2.8) is just the matrix element of the relation $[(\hat{J}_{-})^{l+1}, \hat{H}] = 0$, $l = 0, 1, \ldots$,

$$\langle m - l - 1 | \hat{J}_{-}^{l+1} \hat{H} | m \rangle = \langle m - l - 1 | \hat{H} \hat{J}_{-}^{l+1} | m \rangle.$$
 (4.3)

The intertwining relation (2.9) is given by Hermitian conjugation, or, alternatively, is produced by the same operator relation taken between the states $\langle -m-1 |$ and $|-m+l \rangle$ with subsequent use of the identity (2.2). Relation (4.3) taken for l = 2m (or, its Hermitian conjugate) together with the identity (2.2) produce Eq. (2.15) corresponding to the nontrivial odd order integral of the RPT system H_m .

Hence, we have identified the origin of the intertwining relations associated with the RPT Hamiltonian (2.1). Since the commutativity of the operators $\mathcal{X}_{m,l}$ and $\mathcal{Y}_{m,l}$ with Hamiltonian $\mathcal{H}_{m,l}$ is reduced to the intertwining relations, one can find what matrix elements of the operators \hat{J}_{\pm}^r , with appropriately chosen integer r > 0, correspond to the operators $X_{m,l}^{\pm}$, $Y_{m,l}^{\pm}$ and $Z_{m,l}^{\pm}$, from which the integrals $\mathcal{X}_{m,l}$, $\mathcal{Y}_{m,l}$ and $\mathcal{Z}_{m,l}$, are composed. In correspondence with the discrete symmetry (3.35) we have

$$-(-i)^{m-l}\langle l|\hat{J}_{-}^{m-l}|m\rangle = -i^{m-l}\langle -l-1|\hat{J}_{+}^{m-l}|-m-1\rangle = X_{m,l}^{-}, \qquad (4.4)$$

$$-(-i)^{m+l+1}\langle -l-1|\hat{J}_{-}^{m+l+1}|m\rangle = -i^{m+l+1}\langle l|\hat{J}_{+}^{m+l+1}|-m-1\rangle = Y_{m,l}^{-}, \qquad (4.5)$$

$$(-i)^{2m+1}\langle -m-1|\hat{J}_{-}^{2m+1}|m\rangle = i^{2m+1}\langle m|\hat{J}_{+}^{2m+1}|-m-1\rangle = Z_{m,l}^{-}, \qquad (4.6)$$

$$(-i)^{2l+1}\langle -l-1|\hat{J}_{+}^{m-l}\hat{J}_{-}^{m+l+1}|l\rangle = i^{2l+1}\langle l|\hat{J}_{+}^{m+l+1}\hat{J}_{-}^{m-l}|-l-1\rangle = Z_{m,l}^{+}.$$
 (4.7)

Operators $X_{m,l}^+$ and $Y_{m,l}^+$ are obtained by Hermitian conjugation of (4.4) and (4.5). These relations are illustrated by Figures 3–7 for m = 3, l = 1.



Figure 3: Action of the operators $X_{3,1}^{\pm}$ in correspondence with (4.4).



Figure 4: Action of the operators $Y_{3,1}^{\pm}$ in correspondence with (4.5).

In order to reproduce the relations corresponding to (2.26) with the help of (4.4)–(4.7), we can use the identities

$$\hat{J}_{+}^{n}\hat{J}_{-}^{n} = \prod_{k=0}^{n-1} \left(\hat{H} + \left(\hat{J}_{3} - k - \frac{1}{2} \right)^{2} \right) \equiv P_{n}(\hat{H}, \hat{J}_{3}), \qquad \hat{J}_{-}^{n}\hat{J}_{+}^{n} = P_{n}(\hat{H}, -\hat{J}_{3}), \tag{4.8}$$



Figure 5: Action of the operator $Z_{3,1}^{-}$ in correspondence with (4.6).



Figure 6: Action of the operator $Z_{3,1}^+$ in correspondence with (4.7).

that follow from the so(2, 1) algebra and Eq. (3.30). For instance, putting in the first identity n = 2m + 1, and computing a diagonal matrix element between the states $\langle m |$ and $|m\rangle$, we reproduce the lower component case of the third relation from (2.26), i.e. $(Z_{m,l}^{-})^{2} = P_{2m+1}(H_{m})$, where the polynomial is given by Eq. (2.17).

Therefore, the reduction of the AdS_2 system with half-integer Aharonov-Bohm flux ($\alpha = 1/2$) to an eigensubspace $j_3 = m + \frac{1}{2}$ reproduces a bosonized nonlinear supersymmetry of the RPT system. On the Hilbert space composed from the two eigenspaces $j = m + \frac{1}{2}$ and $j = l + \frac{1}{2}$ with $m \neq l$ we reveal the tri-supersymmetric structure of the extended RPT system (2.18). The key role for the nontrivial nonlinear supersymmetric structure is played here by the involutive automorphism (3.35) of the so(2, 1) algebra, which is realized as a symmetry on the Hilbert space of the two-dimensional quantum system.

In the previous section we have seen that this discrete symmetry is not spontaneously broken also for integer values of the magnetic flux. In this case, however, the system is not reflectionless, and cannot be related to the one-dimensional quantum free particle system. Nevertheless, the intertwining relations corresponding to the Crum-Darboux transformations can be considered in this case as well, and they also can be related with the ladder operators. Using the intertwining relations for the case of half-integer values of the index $m = n + \frac{1}{2}$, $n \in \mathbb{Z}$, and the symmetry (2.2), we find that the operator $\mathcal{D}_{-m}\mathcal{D}_{-m+1}\ldots\mathcal{D}_{-1/2}\mathcal{D}_{1/2}\ldots\mathcal{D}_{m-1}\mathcal{D}_m$ of even order 2m + 1 = 2(n + 1) commutes with the Hamiltonian H_m . A simple calculation shows, however, that this operator reduces to a certain polynomial of order n + 1 in $H_{n+\frac{1}{2}}$. For instance, in correspondence with (2.4), $\mathcal{D}_{-1/2}\mathcal{D}_{1/2} = -H_{1/2} - 1/4$. Hence, there is no bosonized and tri- supersymmetric structure in the Pöschl-Teller system with a half-integer value of the parameter m.

5 On (super)conformal dynamical symmetry

As we observed in section 2, the nonlinear supersymmetry of the RPT system is generated by the Crum-Darboux transformations from the hidden bosonized N = 2 linear supersymmetry of the free non-relativistic particle. The latter system possesses also a *non-relativistic* conformal symmetry in the form of a dynamical so(2, 1) symmetry [29]. The same dynamical symmetry is present in some other quantum mechanical non-relativistic systems, including, in particular, the conformal mechanics model [30], and the model of a particle in the field of a magnetic vortex [32, 40]. The latter corresponds to the planar Aharonov-Bohm effect. The conformal mechanics model, like the RPT system, is related to the free particle by a Crum-Darboux transformation [41]. The conformal so(2, 1) dynamical symmetry of all these systems can be explained within the framework of the AdS/CFT correspondence, see [35] and [37].

In this section, we construct analogs of the free particle generators of the so(2, 1) symmetry for the RPT system, and investigate their properties. The hidden bosonized N = 2 linear supersymmetry of the free particle can be unified with its dynamical so(2, 1) symmetry to produce a linear bosonized superconformal osp(2|2) symmetry, see [41]. We also discuss a realization of the osp(2|2) supersymmetry in the RPT system.

Before we pass over to the discussion of the specified problem, let us make two notes. The non-relativistic AdS/CFT correspondence appears between a relativistic theory in the AdS space-time of d+3 dimensions, and non-relativistic theories in (d+1)-dimensional space-time [37]. We exploited here the relation between the non-relativistic particle system on the AdS₂, whose evolution is described by an external time variable, and the non-relativistic (1 + 1)dimensional RPT system. Next, in the AdS₂ system, the so(2, 1) isometry corrected by the magnetic flux is represented by the true, time-independent integrals of motion. Hamiltonian (3.30) is the Casimir operator of this so(2, 1) symmetry, whose generators are differential operators of the first order. In contrary, the generators of the conformal dynamical symmetry of the free particle we consider below are represented by differential operators of the second order.

The free particle is characterized by the Schrödinger symmetry [29] Lie algebra with the following nonzero commutation relations,

$$[D_0, H_0] = 2iH_0, \quad [H_0, K_0] = -4iD_0, \quad [D_0, K_0] = -2iK_0.$$
(5.1)

$$[B_0, P_0] = i \cdot 1, \quad [D_0, P_0] = iP_0, \quad [D_0, B_0] = -iB_0, \tag{5.2}$$

$$[B_0, H_0] = 2iP_0, \quad [P_0, K_0] = -2iB_0, \tag{5.3}$$

see also [42]. Here

$$P_0 = -i\frac{d}{dx}, \quad B_0 = x - 2P_0t, \quad H_0 = P_0^2, \quad D_0 = \frac{1}{2}\{P_0, B_0\}, \quad K_0 = B_0^2.$$
(5.4)

In the first relation from (5.2) the unit operator corresponds to the mass put here equal to onehalf, which is a central element of the algebra. The boost generator B_0 , and the generators of dilatations, D_0 , and special conformal transformations (expansions), K_0 , depend explicitly on time. This means that the Schrödinger symmetry is *dynamical*. Its subalgebra (5.1) is the conformal so(2, 1) dynamical symmetry.

Reflection operator R, being integral of motion, anticommutes not only with the P_0 , but also with the time-dependent integral B_0 . Therefore, the free particle can be characterized by the dynamical osp(2|2) superconformal symmetry as well. Its superalgebra is given by the so(2, 1) commutation relations (5.1) supplied with the nontrivial (anti)-commutation relations

$$\{Q_a^{(0)}, Q_b^{(0)}\} = 2\delta_{ab}H_0, \quad \{S_a^{(0)}, S_b^{(0)}\} = 2\delta_{ab}K_0, \quad \{Q_a^{(0)}, S_b^{(0)}\} = 2\delta_{ab}D_0 - \epsilon_{ab}\Sigma, \tag{5.5}$$

$$[H_0, S_a^{(0)}] = -2iQ_a^{(0)}, \quad [K_0, Q_a^{(0)}] = 2iS_a^{(0)}, \tag{5.6}$$

$$[D_0, Q_a^{(0)}] = iQ_a^{(0)}, \quad [D_0, S_a^{(0)}] = -iS_a^{(0)}, \tag{5.7}$$

$$[\Sigma, Q_a^{(0)}] = 2i\epsilon_{ab}Q_b^{(0)}, \quad [\Sigma, S_a^{(0)}] = 2i\epsilon_{ab}S_b^{(0)}.$$
(5.8)

Here

$$Q_1^{(0)} = P_0, \quad Q_2^{(0)} = iRQ_1^{(0)}, \quad S_1^{(0)} = B_0, \quad S_2^{(0)} = iRS_1^{(0)}, \quad \Sigma = -R.$$
 (5.9)

Suppose now that we have two quantum almost isospectral systems related by the Crum-Darboux intertwining relations (2.7). Let a system H possesses an integral of motion \mathcal{A} which can be time dependent, $\frac{d}{dt}\mathcal{A} = \frac{\partial}{\partial t}\mathcal{A} - i[\mathcal{A}, H] = 0$. Time-independence of the Crum-Darboux generator A_n , and the intertwining relations (2.7) allow us to construct an analog of the integral \mathcal{A} for the system \tilde{H} ,

$$\tilde{\mathcal{A}} = A_n \mathcal{A} A_n^{\dagger}, \qquad \frac{d}{dt} \tilde{\mathcal{A}} = \frac{\partial}{\partial t} \tilde{\mathcal{A}} - i[\tilde{\mathcal{A}}, \tilde{H}] = 0.$$
 (5.10)

It depends on time if and only if \mathcal{A} is time dependent as well. Identifying H and \tilde{H} with H_0 and H_m , and A_n with the operator $\mathcal{D}_{-m} \dots \mathcal{D}_{-1} \equiv X_m^{\dagger} = -i^m X_{m,0}^{\dagger}$, see Eqs. (2.3) and (2.20), we find the analogs of the even, D_0 , K_0 , and odd, $Q_a^{(0)}$, $S_a^{(0)}$, free particle integrals for the RPT system described by the Hamiltonian (2.1). We denote these analogs just by changing the index 0 for m. The integral Σ , defined in (5.9), is also the integral for the system H_m .

Let us look now what happens with the dynamical so(2, 1) symmetry in the RPT system. A direct computation gives the commutation relations

$$[D_m, H_m] = 2iH_m P_m(H_m), \quad [K_m, H_m] = 4iD_m$$
(5.11)

where in correspondence with (2.26), (2.27), $P_m(H_m) = X_m^{\dagger} X_m = \prod_{n=0}^{m-1} (H_m - E_{m;n})$. To find $[D_m, K_m]$, we use the identity $D_0^2 = H_0 K_0 + 2i D_0 + \frac{3}{4}$, and get

$$[D_m, K_m] = a_m(H_m)K_m + b_m(H_m)D_m + c_m(H_m)$$
(5.12)

where a_m , b_m and c_m are some polynomials in H_m . At this level, like in the case of the nonlinear supersymmetry analyzed in the previous sections, the Lie algebra of the conformal symmetry of the free particle (5.1) is just deformed by the polynomials in the Hamiltonian

of the corresponding RPT system. The complete algebraic structure, however, is more complicated here. Unlike the supercharges $Q_a^{(m)} = Z_a$, the dynamical integrals D_m and K_m do not commute with the Hamiltonian H_m . As a consequence, the repeated commutators produce polynomials in all the three generators H_m , D_m , K_m . To illustrate this phenomenon, consider the simplest case of m = 1. The reflectionless Pöschl-Teller Hamiltonian $H_1 = -d^2/dx^2 - 2/\cosh^2 x$ is intertwined with the free particle Hamiltonian by the first order differential operator $X_1 = \mathcal{D}_1 = d/dx + \tanh x$. The operators D_1 , H_1 and K_1 satisfy the commutation relations

$$[K_1, H_1] = 4iD_1, (5.13)$$

$$[D_1, H_1] = 2iH_1(H_1 + 1), \qquad [D_1, K_1] = -2iK_1 - 2i\{K_1, H_1\} - 3i(H_1 + 1). \tag{5.14}$$

Denoting $\mathcal{K}_1 \equiv [D_1, K_1]$, we find

$$[\mathcal{K}_1, K_1] \equiv \mathcal{M}_1 = -8\{K_1, D_1\} - 12D_1, \tag{5.15}$$

and

 $[\mathcal{M}_1, K_1] = 32i\{K_1^2, H_1\} + 32iK_1^2 + 176i\{K_1, H_1\} + 200iK_1 + 228i(H_1 + 1).$ (5.16)

From the displayed structure it is clear that despite the nonlinear character of the algebra generated by the H_1 , D_1 and K_1 , all the (repeated) commutators are polynomials in them. No new independent integrals do appear.

This ceases to be true once we incorporate the fermionic analogs of the osp(2|2) generators. Besides the deformation effect we observed above, the commutation relations between bosonic and fermionic operators produce new fermionic dynamical integrals, not reducible to the products of the fermionic generators $Q_a^{(m)}$, $S_a^{(m)}$, and of the bosonic generators. Nevertheless, the number of new fermionic integrals is finite. It depends on the value of the parameter m, and therefore, on the order of the Crum-Darboux generating operator X_m . To illustrate this phenomenon, again, restrict ourselves by the simplest case of the RPT system with m = 1. In addition to the already specified (repeated) commutation relations between bosonic operators H_1 , D_1 and K_1 , we get the following (anti)-commutation relations,

$$\{Q_a^{(1)}, Q_b^{(1)}\} = 2\delta_{ab}H_1(H_1+1)^2, \quad \{S_a^{(1)}, S_b^{(1)}\} = \delta_{ab}\left(\{K_1, H_1\} + 2(K_1+H_1+1)\right), \quad (5.17)$$

$$\{S_a^{(1)}, Q_b^{(1)}\} = \delta_{ab} \left(\{D_1, H_1\} + 2D_1\right) - \epsilon_{ab} \Sigma (3H_1 + 1)(H_1 + 1),$$
(5.18)

$$\{J_a^{(1)}, J_b^{(1)}\} = \frac{1}{8} \delta_{ab} \left(K_1 H_1 K_1 - \{K_1, H_1\} - 2(H_1 + 1) \right), \tag{5.19}$$

$$\{S_a^{(1)}, J_b^{(1)}\} = \frac{1}{4}\delta_{ab}\left(\{K_1, D_1\} + 2D_1\right) + \frac{1}{8}\epsilon_{ab}\Sigma(\{K_1, H_1\} - 2K_1 + 2(H_1 + 1)),$$
(5.20)

$$\{J_a^{(1)}, Q_b^{(1)}\} = \frac{1}{4}\delta_{ab}\left(2H_1K_1H_1 + \{K_1, H_1\} - 5H_1^2 - 4H_1 + 1\right) - \frac{1}{4}\epsilon_{ab}\Sigma\left(3\{D_1, H_1\} + 2D_1\right),$$
(5.21)

 $[K_1, Q_a^{(1)}] = 3i\{S_a^{(1)}, H_1\} + 2iS_a^{(1)}, \quad [K_1, S_a^{(1)}] = 8iJ_a^{(1)}, \quad [K_1, J_a^{(1)}] = \frac{i}{4}\{K_1, S_a^{(1)}\}, \quad (5.22)$

$$[H_1, Q_a^{(1)}] = 0, \quad [H_1, S_a^{(1)}] = -2iQ_a^{(1)}, \quad [H_1, J_a^{(1)}] = -\frac{i}{2}\{S_a^{(1)}, H_1\}, \tag{5.23}$$

$$[D_1, Q_a^{(1)}] = iQ_a^{(1)} (1 + 3H_1), \quad [D_1, S_a^{(1)}] = -\frac{i}{2} \left(2S_a^{(1)} + \{S_a^{(1)}, H_1\} \right)$$
(5.24)

$$[D_1, J_a^{(1)}] = -\frac{3i}{2} \{ J_a^{(1)}, H_1 \} - i J_a^{(1)} - i Q_a^{(1)},$$
(5.25)

$$[\Sigma, Q_a^{(1)}] = 2i\epsilon_{ab}Q_b^{(1)}, \quad [\Sigma, S_a^{(1)}] = 2i\epsilon_{ab}S_b^{(1)}, \quad [\Sigma, J_a^{(1)}] = 2i\epsilon_{ab}J_b^{(1)}.$$
(5.26)

Here

$$J_1^{(1)} = X_1^{\dagger} J_1^{(0)} X_1, \quad J_2^{(1)} = i R J_1^{(1)}, \tag{5.27}$$

are the two new fermionic integrals generated by the commutator of K_1 with $S_a^{(1)}$, see (5.22). They correspond to the dynamical integral $J_1^{(0)} = \frac{1}{8} \{K_0, Q_1^{(0)}\}$ of the free particle. From the point of view of the closed Lie superalgebraic structure of the free particle model, the operator $J_1^{(0)}$ belongs to the universal enveloping algebra of the osp(2|2).

The displayed set of the anti-commutation relations shows, nevertheless, that no other new independent integrals will be generated by the repeated anticommutation relations. Note that a similar picture corresponding to the deformation and extension of the superconformal osp(2|2) symmetry appears in the fermion-monopole system [43]. This happens also in the superconformal mechanics model when a classical boson-fermion coupling constant α is changed for $n\alpha$, where n > 1 is integer, see [44].

We do not touch here the question of relation of the deformed (and extended) conformal so(2, 1) (and superconformal osp(2|2)) dynamical symmetry of the RPT system to the so(2, 1) symmetry of the particle on the AdS₂. Let us note only that the generators of the conformal and superconformal dynamical symmetries of the RPT system could be related to the oscillator-like operators $P_+ = e^{i\varphi}\hat{J}_+$ and $P_- = P_+^{\dagger} = \hat{J}_-e^{-i\varphi}$ of the AdS₂ system. These two operators commute with the operator \hat{J}_3 , and represent the observable operators under reduction to the one-dimensional RPT system. The operators P_{\pm} are not integrals of motion for the original two-dimensional system due to a nontrivial dynamics of the angular variable φ . Nevertheless, the true and also dynamical integrals of motion of the RPT system can be obtained from them after reduction. Investigation of this aspect lies, however, beyond the scope of the present paper.

6 Discussion and outlook

We showed that the hidden bosonized nonlinear supersymmetry of the reflectionless Pöschl-Teller system and tri-supersymmetric structure of the pair of the RPT systems have an origin in the Aharonov-Bohm effect for non-relativistic particle on the AdS₂. Both supersymmetric structures are based on the two involutive automorphisms of the so(2, 1) algebra of the AdS₂ isometry with generators corrected by the magnetic flux α . One of these automorphisms, R, corresponds to a reflection of a space-like coordinate of the ambient Minkowski space. It is the symmetry for any value of the flux. Another automorphism, S, corresponds to a reflection of one of the time-like coordinates of the ambient space. This discrete symmetry is unbroken only in the case of integer and half-integer values of the flux. Classically, the system is subject to the Pöschl-Teller potential $V(\chi; J_3) = -J_3^2 \cosh^{-2} \chi$ in the noncompact AdS₂ coordinate χ . It acquires a correction term in the quantum framework, $V = -(\hat{J}_3^2 - \frac{1}{4})\cosh^{-2}\chi$. The classical free-particle dynamics in χ takes place for zero value of the axial angular momentum shifted by the magnetic flux, $J_3 = 0$. It can be recovered on the quantum level for half-integer values of the magnetic flux only. In this case, the dynamics of any partial wave of the particle on the AdS₂ is governed by the reflectionless Pöschl-Teller potential, which turns into zero in two subspaces $j_3 = \pm \frac{1}{2}$. The complete two-dimensional quantum dynamics on the AdS₂ is hence reflectionless. This is not the case for the other values of the flux. For $\alpha \neq \frac{1}{2} + n$, the dynamics of some partial waves is subject to the repulsive potential. Then this information is transmitted to all other partial wave sectors by the so(2, 1) ladder operators, which play the role of the generators of the Crum-Darboux transformations. So, the two-dimensional dynamics ceases to be reflectionless in this case.

For half-integer values of the flux, all the spectrum of the two-dimensional quantum system acquires an additional double degeneration related to the Hamiltonian symmetry $\hat{J}_3 \rightarrow -\hat{J}_3$ generated by the *S*. Such a degeneration is absent for the unpaired partial wave sector $j_3 = 0$ in the case of integer flux. It is the additional double degeneration that is behind the existence of the nontrivial nonlinear supersymmetric structure in the RPT system appearing after reduction by the compact so(2, 1) generator \hat{J}_3 . Generator of the other discrete symmetry, *R*, is transformed then into the \mathbb{Z}_2 -grading operator of the hidden bosonized supersymmetry of the RPT system. The supersymmetry generators of the order operators of the original two-dimensional system with half-integer flux.

Note that the special "magic" of half-integer values of the flux and related degeneracy were discussed in a different context in [45].

The described correspondence between the non-relativistic AdS_2 Aharonov-Bohm effect and the one-dimensional Pöschl-Teller system for a generic magnetic flux value case is somewhat of the AdS/CFT holography [27] nature. This observation deserves a deeper study that can be useful in the context of the integrable nonlinear equations.

For the peculiar case of half-integer values of the flux, we considered this aspect in the context of non-relativistic AdS/CFT [36, 37]. Namely, we showed, that the analogs of the free particle generators of the dynamical conformal, so(2, 1), and bosonized superconformal, osp(2|2), symmetries can be transferred to the RPT system by the Crum-Darboux transformation. In the case of the conformal so(2, 1) dynamical symmetry, there appears a certain nonlinear deformation of the so(2, 1) algebra. In the case of the superconformal osp(2|2) symmetry, in addition to a deformation, finite number of new fermionic integrals is generated. These dynamical symmetries of the RPT system need a further investigation.

The conformal so(2, 1) dynamical symmetry of the planar Aharanov-Bohm effect finds a natural explanation in the AdS/CFT context [35, 37]. The bound state Aharonov-Bohm effect, that corresponds to a particle confined to a circle pierced by the Aharonov-Bohm flux, is characterized by the hidden bosonized supersymmetry as well [18]. It would be interesting to look for the hidden bosonized N = 2 supersymmetry and bosonized superconformal osp(2|2) dynamical symmetry in the planar Aharonov-Bohm effect [46].

Nonlinear bosonized supersymmetry of the RPT system and tri-supersymmetry of the extended RPT system are related to the nontrivial Lax operator which is the higher order differential operator. As we saw, this operator is the first order integral $\frac{d}{dx}$ of the free particle transferred to the RPT system by the Crum-Darboux transformation. Its square gives a spectral polynomial of the RPT system, that has double roots and describes a degenerate hyperelliptic curve [14]. This degeneration originates from the infinite-period limit applied to the finite-gap periodic Lamé, or associated Lamé system. It is this limit that produces the RPT system with its unique, pure imaginary period from the double periodic model. The spectral polynomial of the associated Lamé system is non-degenerate. It was showed in [23, 24] that such a finite-gap periodic system is also characterized by the hidden bosonized nonlinear supersymmetry, while the pair of such systems is described by the tri-supersymmetric structure. In the case of the finite-gap periodic systems, Crum-Darboux transformations relate the systems with the same number of gaps. Hence, an *n*-gap ($n \in \mathbb{N}$) periodic system can not be related by Crum-Darboux transformation to the quantum free particle model, which corresponds to the simplest case of a zero-gap periodic system. Then a question appears: is there any analog of the Aharonov-Bohm effect explanation for a hidden bosonized supersymmetry and related tri-supersymmetric structure in the finite-gap periodic systems? We are going to investigate this question elsewhere.

The authors thank Peter Horvathy for stimulating communications. The work has been partially supported by CONICYT, DICYT (USACH) and by FONDECYT under grants 1050001 and 3085013.

References

- G. Pöschl and E. Teller, "Bemerkungen zur Quantenmechanik des anharmonischen Oszillators", Z. Physik 83 (1933) 143.
- [2] H. B. Thacker, C. Quigg and J. L. Rosner, "Inverse scattering problem for quarkonium systems. 1. One-dimensional formalism and methodology," Phys. Rev. D 18 (1978) 274.
- [3] Y. Alhassid, F. Gursey and F. Iachello, "Potential scattering, transfer matrix, and group theory," Phys. Rev. Lett. 50 (1983) 873.
- [4] W. Kwong and J. L. Rosner, "Supersymmetric Quantum mechanics and inverse scattering," Prog. Theor. Phys. Suppl. 86 (1986) 366.
- [5] A. O. Barut, A. Inomata and R. Wilson, "Algebraic treatment of second Pöschl-Teller, Morse-Rosen and Eckart equations," J. Phys. A 20 (1987) 4083.
- [6] A. K. Grant and J. L. Rosner, "Supersymmetric quantum mechanics and the Kortewegde Vries hierarchy," J. Math. Phys. 35 (1994) 2142 [arXiv:hep-th/9304139].
- [7] J. Goldstone and R. Jackiw, "Quantization of nonlinear waves," Phys. Rev. D 11 (1975) 1486;
 R. Jackiw, "Quantum meaning of classical field theory," Rev. Mod. Phys. 49 (1977) 681.
- [8] R. F. Dashen, B. Hasslacher and A. Neveu, "Semiclassical bound states in an asymptotically free theory," Phys. Rev. D 12 (1975) 2443.

- J. Feinberg, "Kinks and bound states in the Gross-Neveu model," Phys. Rev. D 51 (1995) 4503 [arXiv:hep-th/9408120]; "All about the static fermion bags in the Gross-Neveu model," Annals Phys. (NY) 309 (2004) 166 [arXiv:hep-th/0305240].
- [10] G. Basar and G. V. Dunne, "A Twisted Kink Crystal in the Chiral Gross-Neveu model," Phys. Rev. D 78, 065022 (2008) [arXiv:0806.2659 [hep-th]].
- [11] A. Sen, "Stable non-BPS bound states of BPS D-branes," JHEP 9808, 010 (1998)
 [arXiv:hep-th/9805019]; "Tachyon condensation on the brane antibrane system," JHEP 9808, 012 (1998) [arXiv:hep-th/9805170].
- B. Zwiebach, "A solvable toy model for tachyon condensation in string field theory," JHEP 0009, 028 (2000) [arXiv:hep-th/0008227];
 J. A. Minahan and B. Zwiebach, "Field theory models for tachyon and gauge field string dynamics," JHEP 0009, 029 (2000) [arXiv:hep-th/0008231].
- [13] V. B. Matveev and M. A. Salle, Darboux Transformations and Solitons (Springer, Berlin, 1991).
- S. P. Novikov *et al.*, Theory of Solitons (Plenum, New York, 1984);
 F. Gesztesy and H. Holden, Soliton Equations and Their Algebro-Geometric Solutions (Cambridge Univ. Press, Cambridge, 2003).
- [15] F. Cooper, A. Khare and U. Sukhatme, "Supersymmetry and quantum mechanics," Phys. Rept. 251 (1995) 267 [arXiv:hep-th/9405029].
- [16] G. Levai, "A search for shape invariant solvable potentials," J. Phys. A 22 (1989) 689.
- [17] J. O. Organista, M. Nowakowski and H. C. Rosu, "Shape invariance through Crum transformation," J. Math. Phys. 47 (2006) 122104 [arXiv:math-ph/0603056].
- [18] F. Correa and M. S. Plyushchay, "Hidden supersymmetry in quantum bosonic systems," Annals Phys. **322** (2007) 2493 [arXiv:hep-th/0605104].
- [19] F. Correa and M. S. Plyushchay, "Peculiarities of the hidden nonlinear supersymmetry of Poschl-Teller system in the light of Lamé equation," J. Phys. A 40 (2007) 14403 [arXiv:0706.1114 [hep-th]].
- [20] A. A. Andrianov, M. V. Ioffe and V. P. Spiridonov, "Higher derivative supersymmetry and the Witten index," Phys. Lett. A 174 (1993) 273 [arXiv:hep-th/9303005].
- [21] A. A. Andrianov and A. V. Sokolov, "Nonlinear supersymmetry in quantum mechanics: algebraic properties and differential representation," Nucl. Phys. B 660 (2003) 25 [arXiv:hep-th/0301062].
- [22] M. S. Plyushchay, "Deformed Heisenberg algebra, fractional spin fields and supersymmetry without fermions," Annals Phys. 245 (1996) 339 [arXiv:hep-th/9601116]; "Hidden nonlinear supersymmetries in pure parabosonic systems," Int. J. Mod. Phys. A 15 (2000) 3679 [arXiv:hep-th/9903130].

- [23] F. Correa, V. Jakubský, L. M. Nieto and M. S. Plyushchay, "Self-isospectrality, special supersymmetry, and their effect on the band structure," Phys. Rev. Lett. 101, 030403 (2008) [arXiv:0801.1671 [hep-th]].
- [24] F. Correa, V. Jakubský and M. S. Plyushchay, "Finite-gap systems, tri-supersymmetry and self-isospectrality," J. Phys. A 41 (2008) 485303 [arXiv:0806.1614 [hep-th]].
- [25] F. Correa, L. M. Nieto and M. S. Plyushchay, "Hidden nonlinear su(2|2) superunitary symmetry of N=2 superextended 1D Dirac delta potential problem," Phys. Lett. B 659 (2008) 746 [arXiv:0707.1393 [hep-th]].
- [26] Y. Aharonov and D. Bohm, "Significance of electromagnetic potentials in the quantum theory," Phys. Rev. 115 (1959) 485.
- [27] J. M. Maldacena, "The large N limit of superconformal field theories and supergravity," Adv. Theor. Math. Phys. 2 (1998) 231 [Int. J. Theor. Phys. 38 (1999) 1113] [arXiv:hep-th/9711200];
 S. S. Gubser, I. R. Klebanov and A. M. Polyakov, "Gauge theory correlators from non-critical string theory," Phys. Lett. B 428 (1998) 105 [arXiv:hep-th/9802109];
 E. Witten, "Anti-de Sitter space and holography," Adv. Theor. Math. Phys. 2 (1998) 253 [arXiv:hep-th/9802150].
- [28] O. Aharony, S. S. Gubser, J. M. Maldacena, H. Ooguri and Y. Oz, "Large N field theories, string theory and gravity," Phys. Rept. 323 (2000) 183 [arXiv:hep-th/9905111].
- [29] R. Jackiw, "Introducing scale symmetry," Phys. Today 25 (1972) 23;
 U. Niederer, "The maximal kinematical invariance group of the free Schrodinger equation," Helv. Phys. Acta 45 (1972) 802;
 C. R. Hagen, "Scale and conformal transformations in galilean-covariant field theory," Phys. Rev. D 5 (1972) 377.
- [30] V. de Alfaro, S. Fubini and G. Furlan, "Conformal invariance in quantum mechanics," Nuovo Cim. A 34 (1976) 569.
- [31] R. Jackiw, "Dynamical symmetry of the magnetic monopole," Annals Phys. 129 (1980) 183.
- [32] R. Jackiw, "Dynamical symmetry of the magnetic vortex," Annals Phys. 201 (1990) 83.
- [33] R. Jackiw and S. Y. Pi, "Classical and quantal nonrelativistic Chern-Simons theory," Phys. Rev. D 42 (1990) 3500 [Erratum-ibid. D 48 (1993) 3929].
- [34] C. Duval, G. W. Gibbons and P. Horvathy, "Celestial mechanics, conformal structures, and gravitational waves," Phys. Rev. D 43 (1991) 3907 [arXiv:hep-th/0512188].
- [35] C. Leiva and M. S. Plyushchay, "Conformal symmetry of relativistic and nonrelativistic systems and AdS/CFT correspondence," Annals Phys. 307 (2003) 372 [arXiv:hep-th/0301244].

[36] D. T. Son, "Toward an AdS/cold atoms correspondence: a geometric realization of the Schrödinger symmetry," Phys. Rev. D 78 (2008) 046003 [arXiv:0804.3972 [hep-th]];
K. Balasubramanian and J. McGreevy, "Gravity duals for non-relativistic CFTs," Phys. Rev. Lett. 101 (2008) 061601 [arXiv:0804.4053 [hep-th]];

J. L. B. Barbon and C. A. Fuertes, "On the spectrum of nonrelativistic AdS/CFT," JHEP **0809** (2008) 030 [arXiv:0806.3244 [hep-th]];

W. D. Goldberger, "AdS/CFT duality for non-relativistic field theory," arXiv:0806.2867 [hep-th];

J. Maldacena, D. Martelli and Y. Tachikawa, "Comments on string theory backgrounds with non-relativistic conformal symmetry," JHEP **0810** (2008) 072 [arXiv:0807.1100 [hep-th]];

C. P. Herzog, M. Rangamani and S. F. Ross, "Heating up Galilean holography," arXiv:0807.1099 [hep-th];

A. V. Galajinsky, "Remark on quantum mechanics with conformal Galilean symmetry," Phys. Rev. D **78** (2008) 087701 [arXiv:0808.1553 [hep-th]];

M. Schvellinger, "Kerr-AdS black holes and non-relativistic conformal QM theories in diverse dimensions," arXiv:0810.3011 [hep-th];

M. Rangamani, S. F. Ross, D. T. Son and E. G. Thompson, "Conformal non-relativistic hydrodynamics from gravity," arXiv:0811.2049 [hep-th];

A. Akhavan, M. Alishahiha, A. Davody and A. Vahedi, "Non-relativistic CFT and semi-classical strings," arXiv:0811.3067 [hep-th].

- [37] C. Duval, M. Hassaine and P. A. Horvathy, "The geometry of Schrödinger symmetry in gravity background/non-relativistic CFT," arXiv:0809.3128 [hep-th].
- [38] S. Fluegge, Practical Quantum Mechanics, (Springer, 1994), p. 94.
- [39] V. Bargmann, "Irreducible unitary representations of the Lorentz group," Ann. Math. 48 (1947) 568.
- [40] C. Duval and P. A. Horvathy, "Supersymmetry of the magnetic vortex," arXiv:0807.0569 [hep-th].
- [41] F. Correa, M. A. del Olmo and M. S. Plyushchay, "On hidden broken nonlinear superconformal symmetry of conformal mechanics and nature of double nonlinear superconformal symmetry," Phys. Lett. B 628 (2005) 157 [arXiv:hep-th/0508223].
- [42] C. Duval and P. A. Horvathy, "On Schrödinger superalgebras," J. Math. Phys. 35 (1994) 2516 [arXiv:hep-th/0508079].
- [43] F. De Jonghe, A. J. Macfarlane, K. Peeters and J. W. van Holten, "New supersymmetry of the monopole," Phys. Lett. B 359 (1995) 114 [arXiv:hep-th/9507046];
 M. S. Plyushchay, "On the nature of fermion-monopole supersymmetry," Phys. Lett. B 485 (2000) 187 [arXiv:hep-th/0005122];
 C. Leiva and M. S. Plyushchay, "Nonlinear superconformal symmetry of a fermion in the field of a Dirac monopole," Phys. Lett. B 582 (2004) 135 [arXiv:hep-th/0311150].

- [44] C. Leiva and M. S. Plyushchay, "Superconformal mechanics and nonlinear supersymmetry," JHEP 0310 (2003) 069 [arXiv:hep-th/0304257];
 A. Anabalon and M. S. Plyushchay, "Interaction via reduction and nonlinear superconformal symmetry," Phys. Lett. B 572 (2003) 202 [arXiv:hep-th/0306210].
- [45] Y. Aharonov et al., "A-B and Berry phases for a quantum cloud of charge," Phys. Rev. Lett. 73 (1994) 918 [arXiv:hep-th/9312131].
- [46] F. Correa, H. Falomir, V. Jakubský and M. S. Plyushchay, in preparation.