## POISSON LIMIT FOR ASSOCIATED RANDOM FIELDS

YURI BAKHTIN

ABSTRACT. We prove that under an easily verifiable set of conditions a sequence of associated random fields converges under rescaling to the Poisson Point Process and give a couple of examples.

## 1. INTRODUCTION AND MAIN RESULT

In this note we prove a Poisson scaling limit for a sequence of associated random fields. Let us recall that a finite family (vector)  $(X_1, \ldots, X_m)$  of random variables (r.v.'s) is called associated if for every pair of bounded and coordinatewise nondecreasing functions  $f, q : \mathbb{R}^m \to \mathbb{R}$ ,

(1) 
$$\operatorname{cov}(f(X_1,\ldots,X_m),g(X_1,\ldots,X_m)) \ge 0$$

An infinite family of r.v.'s is called associated if its every finite subfamily is associated.

The notion of association was introduced and studied in [4]. Inequalities (1) with their equivalents have been often referred to as FKG inequalities by the initials of authors of [5] who studied this type of positive correlation independently.

Associated r.v.'s arise frequently in various problems of statistical mechanics and many other areas, see numerous examples, a historic overview, theory and applications in a recent monograph [1].

Basic properties of associated random vectors: jointly independent r.v's form an associated family; monotone transformations of associated random vectors are associated, too.

A number of limit theorems for sums of associated r.v.'s have been proved, see [1] and references therein. To the best of our knowledge, no theorem on convergence to a Poisson Point Process has appeared in the literature, although some results on Poisson approximations for systems satisfying FKG inequalities can be found in [7] and references therein.

We proceed to describe the setting. We fix a dimension  $d \in \mathbb{N}$ , and for each  $n \in \mathbb{N}$ , let  $(X_j^{(n)})_{j \in \mathbb{Z}^d}$  be a weakly stationary (i.e. in the sense of first moment and covariance) associated random field. We assume that for all  $n, j, \text{ r.v. } X_j^{(n)}$  takes two values, 0 and 1, and there is a number  $\lambda > 0$  such that

$$p_n = \frac{\lambda + o(1)}{n^d},$$

where  $p_n = \mathsf{P}\{X_0^{(n)} = 1\}.$ 

We also assume that

(2) 
$$\lim_{n \to \infty} n^d \sigma(n) = 0,$$

where

$$\sigma(n) = \sum_{j \neq 0} \operatorname{cov}(X_0^{(n)}, X_j^{(n)}).$$

For any *n* we define a random measure  $\mu_n$  on  $\mathbb{R}^d$  via

$$\mu_n(A) = \sum_{j \in \mathbb{Z}^d \cap nA} X_j^{(n)},$$

where  $nA = \{nx : x \in A\}$ .

The vague topology on locally bounded Borel measures is defined by its base, the class of finite intersections of sets of the form  $\{\nu : s < \int_{\mathbb{R}^d} f d\nu < t\}$  with arbitrary nonnegative continuous function f with bounded support and  $s, t \in \mathbb{R}$ , see [8, Appendix 7].

**Theorem 1.** Under the conditions stated above, the sequence of measures  $\mu_n$  converges in distribution in the vague topology to the Poisson measure  $\mu$  with parameter  $\lambda$ .

*Proof.* By [8, Theorem 4.2], it is sufficient to check that for every continuous nonnegative function f with compact support,

$$\int_{\mathbb{R}^d} f d\mu_n \xrightarrow{Law} \int_{\mathbb{R}^d} f d\mu, \quad \text{as } n \to \infty.$$

Take a continuous function f with compact support and a number  $t \in \mathbb{R}$ , and find

$$\mathsf{E}e^{it\int fd\mu_n} = \mathsf{E}e^{it\sum_{j\in\mathbb{Z}^d} f(\frac{j}{n})X_j^{(n)}}$$

$$= \prod_{j\in\mathbb{Z}^d} \mathsf{E}e^{itf(\frac{j}{n})X_j^{(n)}} + \left| \mathsf{E}e^{it\sum_{j\in\mathbb{Z}^d} f(\frac{j}{n})X_j^{(n)}} - \prod_{j\in\mathbb{Z}^d} \mathsf{E}e^{itf(\frac{j}{n})X_j^{(n)}} \right|$$

$$= I_1(n) + I_2(n).$$

Notice that, in fact, the product in  $I_1(n)$  involves finitely many factors, and

$$I_1(n) = \prod_{j \in \mathbb{Z}^d} \left( 1 + p_n(e^{itf(\frac{j}{n})} - 1) \right).$$

Choosing the main branch of the natural logarithm ln, we can write

$$I_1(n) = \exp\left\{\sum_{j \in \mathbb{Z}^d} \ln(1 + p_n(e^{itf(\frac{j}{n})} - 1))\right\}.$$

 $\mathbf{2}$ 

Using the boundedness of f and the Taylor expansion for the logarithm we derive that

$$I_1(n) = \exp\left\{\frac{\lambda + o(1)}{n^d} \sum_{j \in \mathbb{Z}^d} (e^{itf(\frac{j}{n})} - 1)\right\} (1 + o(1)).$$

Obviously, the r.h.s converges to

$$\phi(t) = \exp\left\{\lambda \int_{\mathbb{R}^d} (e^{itf(x)} - 1)dx\right\},\,$$

the characteristic function of  $\int_{\mathbb{R}^d} f d\mu$ , and the proof will be finished as soon as we show that

(3) 
$$\lim_{n \to \infty} I_2(n) = 0.$$

To estimate  $I_2(n)$  we need Newman's inequality:

**Theorem 2** ([9]). If  $(Y_1, \ldots, Y_m)$  is a family of associated r.v.'s with finite second moment then

$$\left|\mathsf{E} e^{i\sum_{j=1}^m r_j Y_j} - \prod_{j=1}^m \mathsf{E} e^{ir_j Y_j}\right| \le \frac{1}{2} \sum_{j_1 \ne j_2} |r_{j_1} r_{j_2}| \operatorname{cov}(Y_{j_1}, Y_{j_2}),$$

for any real numbers  $r_1, \ldots, r_m$ .

Applying this inequality to  $I_2(n)$  we see that

$$\begin{split} I_{2}(n) &\leq \frac{t^{2} \|f\|_{L^{\infty}}^{2}}{2} \sum_{\substack{j_{1}, j_{2} \in \mathbb{Z}^{d} \cap n \operatorname{supp}(f) \\ j_{1} \neq j_{2}}} \operatorname{cov}(X_{j_{1}}^{(n)}, X_{j_{2}}^{(n)}), \\ &\leq \frac{t^{2} \|f\|_{L^{\infty}}^{2}}{2} |\mathbb{Z}^{d} \cap n \operatorname{supp}(f)| \, \sigma(n), \end{split}$$

where  $\operatorname{supp}(f)$  denotes the support of f, and  $|\cdot|$  denotes the number of elements. Since  $|\mathbb{Z}^d \cap n \operatorname{supp}(f)| \leq Kn^d$  for some constant K > 0 and all n > 0, (3) follows from (2).

**Remark 1.** The crucial step in the proof above is the application of Newman's inequality for associated random variables. Covariance inequalities of this type can be obtained for a wide class of dependent r.v.'s. In particular the theorem is also applicable if one replaces association by quasi-association, see [2] and proof of Theorem 2 in [9].

## 2. Examples

Let G be a finite subset of  $\mathbb{Z}^d$  for some  $d \in \mathbb{N}$ . Denote m = |G| and for each n consider an i.i.d. family  $(Y_k^{(n)})_{k \in \mathbb{Z}^d}$  of Bernoulli random variables with

$$\mathsf{P}\{Y_0^{(n)} = x\} = \begin{cases} q_n, & x = 1, \\ 1 - q_n, & x = 0, \end{cases}$$

where

4

$$q_n = \frac{1}{n^{d/m}}$$

For any finite subset H of  $\mathbb{Z}^d$  and every n, we denote

$$\chi_{H}^{(n)} = \prod_{j \in H} Y_{j}^{(n)} = \mathbf{1}_{\{Y_{j}^{(n)} = 1, \ j \in H\}},$$

and define a random field  $(X_k^{(n)})_{k\in \mathbb{Z}^d}$  via

$$X_k^{(n)} = \chi_{k+G}^{(n)},$$

where  $k + G = \{k + j : j \in G\}$ . Poisson approximations for a similar model with rectangular G was considered in [6].

Let us verify that  $X^{(n)}$  satisfies the conditions of Theorem 1. Random field  $Y^{(n)}$  is associated since it is composed of independent components. Therefore,  $X^{(n)}$  is associated being a monotone transform of the associated field  $Y^{(n)}$ . It is also stationary due to stationarity of  $Y^{(n)}$ .

For each  $n, X_0^{(n)}$  is a Bernoulli r.v. with

$$\mathsf{P}\{X_0^{(n)} = 1\} = \mathsf{P}\{Y_j^{(n)} = 1, \ j \in G\} = \left(\frac{1}{n^{d/m}}\right)^m = \frac{1}{n^d}.$$

Let us now estimate  $\sigma(n)$ . Notice that  $\operatorname{cov}(X_0^{(n)}, X_j^{(n)}) = 0$  for sufficiently large values of |j|, so that there is a number M such that for all n,

(4) 
$$\sigma(n) \le M \max_{j \ne 0} \operatorname{cov}(X_0^{(n)}, X_j^{(n)}).$$

Notice that

$$\operatorname{cov}(X_0^{(n)}, X_j^{(n)}) = \mathsf{E}\chi_{G\cup(j+G)}^{(n)} - \mathsf{E}\chi_G^{(n)}\mathsf{E}\chi_{j+G}^{(n)}.$$

Since a finite set cannot be invariant under a translation,  $|G \cup (j+G)| \ge m+1$  for any j. Therefore,

$$\operatorname{cov}(X_0^{(n)}, X_j^{(n)}) \le \frac{1}{n^{d(m+1)/m}} = o(1/n^d),$$

which, together with (4), implies (2), so that all the conditions of Theorem 1 are satisfied.

For an associated random field  $X^{(n)}$ , condition (2) means that  $X_0^{(n)}$  is asymptotically independent of the rest of the random field. There is a variety of situations that can happen if this condition is replaced with weaker restrictions on dependence. The next example illustrates the convergence to a compound Poisson point process (with nonrandom mass 2 assigned to each atom), see [3] for the definition and properties of compound Poisson point processes.

Consider d = 1, and for every n and all  $k \in \mathbb{Z}$ ,

$$X_k^{(n)} = Y_k^{(n)} \vee Y_{k+1}^{(n)} = Y_k^{(n)} + Y_{k+1}^{(n)} - Y_k^{(n)} Y_{k+1}^{(n)},$$

where  $Y^{(n)}$  is a sequence of i.i.d. Bernoulli r.v.'s with  $\mathsf{P}\{Y_0^{(n)} = 1\} = 1/n$ . Then, as an easy computation shows,  $\sigma(n) \sim 1/n$  so that (2) is violated. One can also show that the sequence of random measures  $\mu_n$  converges in distribution to  $2\mu$ , where  $\mu$  is the Poisson process with unit intensity, so that the conclusion of Theorem 1 is violated as well. Indeed, take a continuous function f with compact support, and write

$$\mathsf{E}e^{it\int_{\mathbb{R}}fd\mu_{n}} = \mathsf{E}e^{it\sum_{j\in\mathbb{Z}}f(\frac{j}{n})(Y_{j}^{(n)}+Y_{j+1}^{(n)})-it\sum_{j\in\mathbb{Z}}f(\frac{j}{n})Y_{j}^{(n)}Y_{j+1}^{(n)}}.$$

Notice that

$$\sum_{j\in\mathbb{Z}} f\left(\frac{j}{n}\right) Y_j^{(n)} Y_{j+1}^{(n)} \xrightarrow{\mathsf{P}} 0, \quad n \to \infty,$$

due to the Markov inequality, since the expectation of l.h.s. is O(1/n). Therefore, we see that

$$\begin{split} \lim_{n \to \infty} \mathsf{E} e^{it \int_{\mathbb{R}} f d\mu_n} &= \lim_{n \to \infty} \mathsf{E} e^{it \sum_{j \in \mathbb{Z}} (f(\frac{j}{n}) + f(\frac{j-1}{n})) Y_j^{(n)}} \\ &= \exp\left\{\int_{\mathbb{R}} (e^{it2f(x)} - 1) dx\right\}, \end{split}$$

by the same argument we used to analyze  $I_1(n)$ .

## References

- Alexander Bulinski and Alexey Shashkin. Limit theorems for associated random fields and related systems. Advanced Series on Statistical Science & Applied Probability, 10. World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2007.
- [2] Alexander Bulinski and Charles Suquet. Normal approximation for quasi-associated random fields. *Statist. Probab. Lett.*, 54(2):215–226, 2001.
- [3] D. J. Daley and D. Vere-Jones. An introduction to the theory of point processes. Springer Series in Statistics. Springer-Verlag, New York, 1988.
- [4] J. D. Esary, F. Proschan, and D. W. Walkup. Association of random variables, with applications. Ann. Math. Statist., 38:1466–1474, 1967.
- [5] C. M. Fortuin, P. W. Kasteleyn, and J. Ginibre. Correlation inequalities on some partially ordered sets. *Comm. Math. Phys.*, 22:89–103, 1971.
- [6] James C. Fu and Markos V. Koutras. Poisson approximations for 2-dimensional patterns. Ann. Inst. Statist. Math., 46(1):179–192, 1994.
- [7] A. Ganesh, B. M. Hambly, Neil O'Connell, Dudley Stark, and P. J. Upton. Poissonian behavior of Ising spin systems in an external field. J. Statist. Phys., 99(1-2):613–626, 2000.
- [8] Olav Kallenberg. Random measures. Akademie-Verlag, Berlin, 1976.
- [9] C. M. Newman. Normal fluctuations and the FKG inequalities. Comm. Math. Phys., 74(2):119–128, 1980.